

Effect of Ferti-Fortification of Iron and Zinc Fertilization on Quality Parameters of Mango Cv. Kesar

Abstract

The present research was accomplished on mango cv. Kesar to investigate the effect of foliar spray of iron and zinc fertilization on quality parameter of mango. The experiment was conducted in completely randomized design and replicated 3 times with 9 treatments. The significant increase in TSS and ascorbic acid content were recorded with the foliar application of 0.5 % FeSO₄ and 0.5 % ZnSO₄ which was at par with treatment T₇ (0.25 % FeSO₄ + 0.50 % ZnSO₄).

Keywords: mango, Kesar, iron, zinc, foliar spray, TSS, ascorbic acid content

INTRODUCTION

The mango is delicious fruit of tropical countries and holds a premier position amongst the commercial fruit grown in India. It is national fruit of India belong to family *annacardiaceae*. In Western India, several mango varieties viz., Alphonso, Kesar, Rajapuri, Pairi, Mankurad, Fernandin, Jamadar, Dadmiyo, etc. are commercially grown and accepted by the consumers. Out of which Kesar has been found with good yield potential, almost regular bearer, mid-season variety, having good consumers' acceptance, attractive shape, size with saffron coloured pulp and very good keeping quality. Mango play important role in balancing diet of human being by providing about 64-66 calories per 100 grams of ripe fruits. The king of fruit is nutritionally very rich, unique in flavor. From last few years, some physiological stresses and quality related issues has been raised in mango orchard. It was observed that unbalanced fertilization, micronutrient deficiencies, poor tree management and inadequate cultural practices are mainly responsible for orchard related quality issue (Ahmad and Rashid, 2003). Therefore food supplements and mineral supplements are necessary for healthy crops. According to horticulturists, only application of primary nutrients could not prove successful to produce high quality fruit in mango trees, the application of micronutrients is compulsory as well.

Foliar spray of micronutrients is the common practice to overcome the micronutrients deficiencies in order to improve the fruit quality. Nutrients are generally quickly available to the plants by the foliar application than the soil application (Bahadur *et al.*, 1998).

MATERIALS AND METHODS

The experiment material for the present investigation was carried out on 20 year old mango orchard planted at 10m×5m distance and site is located at Regional Horticultural Research Station, ASPEE College of Horticulture and Forestry, Navsari Agricultural University, Navsari (Gujarat). The experiment was conducted in completely randomized design (C.R.D) with 9 treatments and replicated 3 times. The details of experiment treatment plan employed in present investigation are as follow: Control (T₁), 0.25 % FeSO₄ (T₂), 0.50 % FeSO₄ (T₃), 0.25 % ZnSO₄ (T₄), 0.50 % ZnSO₄ (T₅), 0.25% FeSO₄ + 0.25 % ZnSO₄ (T₆), 0.25 % FeSO₄ + 0.50 % ZnSO₄ (T₇), 0.50% FeSO₄ + 0.25% ZnSO₄ (T₈), 0.50 % FeSO₄ + 0.50 % ZnSO₄ (T₉). The foliar spray of these nutrients was done at the time of flowering (21/12/2016), pea (17/02/2017) and egg stages (20/03/2017). For the preparation of Zn and Fe foliar spray solution, commercial grade of ZnSO₄.7H₂O and FeSO₄.7H₂O fertilizers were used, respectively. The required quantity of ferrous sulphate and zinc sulphate were weighed and dissolved in water and then pH of solution was adjusted to 6.0 by using saturated CaCO₃ solution.

The total soluble solids of the pulp were determined by using refractometer (0-32 °Brix). For every treatment, pulp of fruit for each replication was blended in to homogeneous mixture and then this mixture used to record five reading and average of those was considered as results.

Ascorbic acid was estimated with titrimeter method described by Ranganna (2004). Ten gram of the pulp was transferred to a 100 ml volumetric flask and volume made up with 4 % oxalic acid solution. After 30 minutes, the suspension was filtered through Whatman No. 1 filter paper. Before actual titration the 2, 6 – Dichloropheno indophenol dye solution was standardized by titrating against standard ascorbic acid solution and the dye factor was calculated. Five milliliters of the aliquot was taken from the filtrate against standardized dye solution through a burette. Titration was continued till the light pink colour persisted for 15 seconds. Ascorbic acid content was calculated adopting the following formula

$$\text{Ascorbic acid (mg/100g)} = \frac{\text{Titre} \times \text{Dye factor} \times \text{Volume made up} \times 100}{\text{Aliquot of extract taken for estimation} \times \text{Weight or Volume of sample taken for estimation}}$$

The shelf life of fruits was recorded by keeping the fruits at room temperature. The days taken from harvesting to the appearance of any spoilage symptoms or discolouration are considered as shelf life of fruits.

The mango fruits were evaluated organoleptically by five judges for colour, flavor, taste, texture. Organoleptic rating was done by following Hedonic Scale (1-9) points. Data pertaining to fruit quality assessment was subjected to statistical analysis constructing as per the method prescribed by Panse and Sukhatme (1967).

RESULTS AND DISCUSSION

The fruit quality parameters *viz.*, TSS and ascorbic acid of mango cv. Kesar were reached the level of significance due to different foliar application of micronutrients. Whereas, non-significant response was observed among the micronutrient treatments for shelf and organoleptic score.

The higher TSS (19.35 °Brix) and ascorbic acid (36.70 mg/100 g) were recorded with foliar application of 0.50 % FeSO₄ + 0.50 % ZnSO₄ (T₉). The increase in TSS and different fraction of sugar under the influence of micronutrients might be due to hydrolysis of complex polysaccharides into simple sugars synthesis of metabolites and rapid translocation of photosynthetic product and minerals from other parts of plant to developing fruits. Increased in concentration of ZnSO₄ caused the increase in TSS (Anees *et al.*, 2011) of mango. Several workers observed similar result as, and in mango, Kumar *et al.* (2017) in mandarin orange, Waskela *et al.* (2013) in guava.




Fig. 1. Effect of foliar spray of Zn and Fe on TSS and Ascorbic acid of mango cv. Kesar

Ascorbic acid content improved might be due to catalytic activity of enzymes and co-enzymes which are involve in ascorbic acid synthesis. The results of present study were confirmed by finding of Nehete *et al.* (2011) and Tulsi *et al.* (2015) in mango; Shekhar *et al.* (2010) in papaya and Meena *et al.* (2006) in ber.

CONCLUSION

It is concluded from the present study that the foliar application of 0.50 % FeSO₄ and 0.50 % ZnSO₄ can be done at flowering, pea and egg stage of fruit for getting higher TSS and ascorbic acid.

REFERENCES

- Ahmad, N and Rashid, M.(2003). Fertilizer and their use in Pakistan. *Extension Bulletin (3rd ed.)* NFDC.,Islamabad. PP.3-4.
- Anees, M.; Tahir, F.M.; Shahzad, J. and Mahmood, N. (2011). Effect of foliar application of micronutrients on the quality of mango (*Mangifera indica* L.) cv. Dashehari fruit. *Mycopath.*, **9**(1): 25-28
- Bahadur, L.; Malhi, C.S. and Singh Z. (1998). Effect of foliar and soil applications of zinc sulphate on zinc uptake, tree size, yield and fruit quality of mango. *J. Pl. Nutr.* **21**: 589-600
- Gurjar, T.; Patel, N. L.; Panchal, B. and Chaudhari, D. (2015). Effect of foliar spray of micronutrients on flowering and fruiting of Alphonso mango(*Mangifera indica* L.). *International Quarterly J. Life Sci.*, **10**(3):1053- 1056.
- Kumar, C. J.; Rajangam , J.; Balakrishnan, K. and Sampath P, M. (2017). Influence of foliar application of micronutrient on yield and quality of mandarin orange (*Citrus reticulata* Blanco.) under lower pulney hills. *International J. Agric Sci.*, **9**(17): 4151-4153.
- Meena, V. S.; Yadav, P. K. and Bhati, B. S. (2006). Effect of ferrous sulphate and borax on fruit quality of ber. *Prog.Hort.*, **38**(2): 283 – 285.
- Nehete, D. S.; Padhiar, B. V.; Shah, N. I.; Bhalerao, P. P.; Kolambe, B. N. and Bhalerao, R. R. (2011). Influence of micronutrient spray on flowering, yield, quality and nutrient content in leaf of mango cv. Kesar. *Asian J. Hort.*, **6** (1): 63 – 67.
- Panse, V. G. and Sukhatme, P. V.(1967) Statistical Methods for Agricultural Workers ICAR, New Delhi.

Ranganna, S. (2004). *“Manual of Analysis of Fruit and Vegetable Products.”* Tata McGran Hill Publ. Com. Ltd. New Delhi.

Shekhar, C.; Yadav, A. L.; Singh, H. K. and Singh, M. K. (2010). Influence of micronutrients on plant growth, yield and quality of papaya fruit (*Carica papaya* L.) cv. Washington. *Asian J. Hort.*, **5**(2):326-329

Waskela, R.S.; Kanpure, R.N., Kumawat, B.R. and Kachouli, B.K. (2013).Effect of foliar spray of micronutrients on growth, yield and quality of guava (*Psidium guajava* L.) cv. Dharidar. *International J. Agric. Sci.*, **9**(2):551- 556.

UNDER PEER REVIEW