

Assessment of Physico-chemical Properties of Soil as influenced by Different Moisture Regimes and Nitrogen Sources in Wheat Crop

Abstract

An experiment was conducted at Student's "Instructional farm of Acharya Narendra Deva University of Agriculture and Technology Kumarganj, Ayodhya (U.P.) during *rabi* season 2019-2020. The experiment was laid out in split plot design (SPD) with three replications using the wheat variety HD- 2967. The treatments comprised of four levels of irrigation *viz.* 0.6 IW/CPE ratio (I₁), 0.8 IW/CPE ratio (I₂), 1.0 IW/CPE ratio (I₃), 1.2 IW/CPE ratio (I₄), and three nitrogen sources *viz.* 100% (RDF) through urea (N₁), 50% RDF + 50% FYM (N₂), 50% RDF + 50% poultry manure (N₃). The build-up in organic carbon was found with the application of moisture regime at 1.2 IW/CPE and the minimum was received with the moisture regime at 0.6 IW/CPE during the investigation. The increment level of available nitrogen, phosphorus, potassium and zinc were found with the increasing level of moisture regime. The maximum was found with the application of moisture regime at 1.2 IW/CPE and the minimum was received with the moisture regime at 0.6 IW/CPE during the investigation. Moisture regimes 1.2 IW/CPE ratio was found best for *viz.*, p^H, EC and organic carbon, Available Nitrogen, Phosphorus, Potassium & Zinc and among the Nitrogen level, application of 50% N from urea + 50% N from Poultry manure (N₃) was found suitable for soil fertility status followed by 50% N from urea + 50% N from FYM (N₂)

Keywords: Wheat, Moisture Regime, Nitrogen Sources, Soil Fertility

Introduction

The most significant cereal crop in the world is wheat (*Triticum aestivum* L.). When it comes to production and area among the major grains, wheat tops the list and provides the primary source of nutrition for around 40 % of the world's population (Giraldo *et al.*, 2019).

In India wheat is the second-most significant grain crop and is crucial to the nation's food and nutritional security. It currently ranks second to rice as the primary food source for humans in terms of total food production (Akibode *et al.*, 2012).

Irrigation should aim to restore soil water in the root zone to a level at which crop can fully meet its evapotranspiration requirement. The amount of water to be applied at each irrigation and how often a soil should be irrigated depends, however on several factors such as the degree of soil water deficit before irrigation, soil type, crops and climatic conditions (Debaeke *et al.*, 2004). There is a positive correlation between grain yield and irrigation frequencies (Kumar *et al.*, 2012).

Nitrogen is one of the major essential elements which influence the growth, yield and quality of wheat more than any other single nutrient element (Tamim Fazily 2020). Organic manures, such as FYM, vermicompost and poultry manure act as a good reservoir of nutrients and water in the soil, which helps to improve soil structure, increase soil infiltration and promote the growth and population of beneficial soil microorganisms (Singh *et al.*, 2020).

The increase in eco-friendly production of wheat can be made possible by widespread adoption of improved Technologies of which fertilizer management particularly that of nitrogen and organic manure can play a key role.

Compost alone and in combination with chemical fertilizer in the same level reduced the soil pH, increased electrical conductivity, available phosphorus, water soluble potassium and organic matter status of soil (Kumar *et al.*, 2023). While addition of organic material to the soil such as farm yard manure (FYM) helps in maintaining soil fertility and productivity. It increases soil microbiological activities, plays key role in transformation, recycling and availability of nutrients to the crop (Singh *et al.*, 2013). It also improves the physical properties like soil structure, porosity, reduces compaction and crusting and increases water holding capacity of soil (Verma *et al.*, 2018).

Materials and methods

Site description

The field experiment was conducted during Rabi season 2019-2020 at Student's "Instructional farm of Acharya Narendra Deva University of Agriculture and Technology Kumarganj, Ayodhya Uttar Pradesh, India, on the left side of Ayodhya-Raibareilly road at a distance of 43 km away from Ayodhya district headquarter.

The local of experimental site lies between a 26.470 N latitude, 82.120 E longitude and an altitude of 113 meters from mean sea level in the gangetic alluvium of eastern Uttar Pradesh. This region falls under the sub-tropical climate of Indo-Gangetic plains of eastern Uttar Pradesh (India) having alluvial sodic soil. The climate of the area is subtropical and semi-arid with an average annual rainfall of about 1070 mm.

Wheat HD-2967 variety was taken for experiment, it is most suitable for north eastern plain zones of India, which includes a part of Uttar Pradesh also. It is an early maturity variety (135-150 days.) Its average yield is 40- 50 quintals per hectare. The experiment was laid out in split plot design (SPD) with three replications.

Scheduling of irrigation

According to the irrigation schedules for the relevant treatment, a set amount of 60 mm of water was applied to the relevant experimental plots. For better crop establishment,

uniform irrigation was applied to all treatments right away after sowing, regardless of the treatments. The USWB class 'A' open pan evaporimeter situated at the meteorological observatory, which was close to the experimental plot, was used to measure the daily pan evaporation, from which the cumulative pan evaporation data were computed. Each treatment received a set depth of irrigation water of 60 mm based on an IW: CPE ratio.

The cumulative pan evaporation (CPE) varied with IW/CPE ratio and hence irrigation based on IW/CPE applied.

$$IW/CPE = \frac{\text{Irrigation water depth (mm)}}{\text{Cumulative pan evaporation (mm)}}$$

CPE is the sum of daily pan evaporation recorded from an open pan evaporimeter. Whenever, cumulative pan evaporation value reached at respective desired level, the irrigation was given as per treatment. Effective rainfall occurred during the crop growth period was also taken into consideration by subtracting it from CPE of that treatment levels.

The time of irrigation in each plot was given on the basis of formula

$$t = \frac{ad}{q}$$

With

t is the = time of application of water

a is the = area of plot to be irrigated m²

d is the = depth of water (cm) and

q is the = discharge rate, (liter /sec)

Table 1. Treatment details

Treatment	Irrigation Water depth (IW) mm	Cumulative Pan Evaporation CPE (mm)	IW/CPE ratio
I1	60	100	0.6
I2	60	75	0.8
I3	60	60	1.0
I4	60	50	1.2

Soil sampling and proprieties

The fertility status of soil texture, bulk density, pH, Electrical conductivity and organic carbon, Soil moisture percentage before irrigation and after irrigation have been determined. Soil sampling was done before and after irrigation with the help of soil auger. Briefly soil sample were put on the moisture box for getting accurate moisture percentage.

Then soil sample were weighted in a container and dried for 24 hours at the temperature of 105⁰C in oven. In the end, the soil samples were reweigh the weight of the container was subtracted and the moisture percentage was calculated following the formula of

$$\text{Soil Moisture Content (\%)} = \frac{WS_1 - WS_2}{WS_2} \times 100$$

With,

WS₁ is the = weight of fresh soil sample and

WS₂ is the = weight of oven dried soil samples

Statistical analysis

Statistical analysis of treatment on the pattern of split plot design was carried out following the procedure as given by Fisher and Yates (1940). The degree for various factor is given in the table 2.

Table 2. Analysis of variance for different characters

<i>Sources of variation</i>	<i>D.f.</i>	<i>SS</i>	<i>MSS</i>	<i>F value calculated</i>	<i>F value tabulated</i>
<i>Replication</i>					
<i>Moisture regimes (M)</i>					
<i>Error (a)</i>					
<i>Nitrogen sources (N)</i>					
<i>M×N</i>					
<i>Error (b)</i>					
<i>Total</i>					

Note: - d. f. =Degree of freedom

SS= Sum of Square

MSS= Mean sum of Square

The standard error of mean and critical difference was calculated by the following formula:

(a) Moisture regimes

$$CD \text{ at } 5\% = \frac{\sqrt{2EMS(a)}}{r \times t} \times t_{0.05 \text{ error (a) d.f.}}$$

(b) Nitrogen sources

$$CD \text{ at } 5\% = \frac{\sqrt{2EMS(b)}}{r \times n} \times t_{0.05 \text{ error (b) d.f.}}$$


Results and Discussion

Physico-chemical Properties of Soil as influenced by Different Moisture Regimes and Nitrogen Sources

Table 3. Physico-chemical Properties of Soil as influenced by Different Moisture

Regimes and Nitrogen Sources

Treatment	Soil Texture	Bulk Density (Mg m ⁻³)	Moisture (%) Before irrigation	Moisture (%) after irrigation
Moisture regimes				
0.6 IW/CPE ratio	Silt loam	1.42	11.42	21.5
0.8 IW/CPE ratio	Silt loam	1.42	12.1	21.8
1.0 IW/CPE ratio	Silt loam	1.43	12.3	22.2
1.2 IW/CPE ratio	Silt loam	1.43	12.5	22.5
SEm±			0.36	0.56
CD at 5%			1.25	1.75
Nitrogen sources				
100% N from urea (N ₁)	Silt loam	1.42	11.20	21.23
50% N from urea +50% N from FYM (N ₂)	Silt loam	1.40	13.20	23.20
50% N from urea + 50% N from Poultry manure (N ₃)	Silt loam	1.41	13.40	23.60
SEm±			0.37	0.57
CD at 5%			1.14	1.82



Soil pH and Electrical conductivity

Data pertaining to soil pH as influenced by different moisture regimes and nitrogen sources. The reduction in pH was observed with the application of the four treatments irrigation. However no clear trend has been found with moisture regime and sources of nitrogen.

Among the nitrogen sources the data showed that the soil pH did not more effected by (100% RDF), N₁, 50% N from urea +50% N from FYM (N₂), 50% N from urea + 50% N

from Poultry manure (N₃). Table 4.

The reduction in soil pH with the application irrigation on the basis of IW/CPE ratio might be due to dilution of salts and leaching of ions beyond the root zone. The use of organic manures viz., poultry manure and FYM has also been known to help in reducing the soil pH to some extent by producing organic acids while their decomposition that may also be the reason of greater availability and mobility of nutrients mainly of micronutrients. These findings are corroborated by (Jat *et al.*, 2011).

Organic Carbon

The observation regarding the organic carbon in g kg^{-1} presented in Table 2. Revealed that a slight improvement in organic carbon was obtained with increasing level of moisture regime up to 1.2 IW/CPE ration. It could be due to the decomposition of organic residues and release of solubilized plant nutrients with maintaining the proper soil moisture levels (Cauley *et al.*, 2009).

Among the application of recommended doses of fertilizer and apply the organic manures along with inorganic fertilizer enhanced in build-up of organic carbon. The maximum organic carbon (4.89 g kg^{-1}) was obtained with the application N₃- 50% RDF+ 50% N through poultry manure followed by N₂- 50%RDF+ 50%N through FYM. The minimum was recorded with the application recommended doses of fertilizer.

Table: 4. The variation of pH, electrical conductivity (EC) and organic carbon (OC) with respect of different treatments.

Treatment	pH	EC (dSm^{-1})	OC (g kg^{-1})
Moisture regimes			
0.6 IW/CPE ratio	8.20	0.32	4.22
0.8 IW/CPE ratio	8.23	0.31	4.43
1.0 IW/CPE ratio	8.20	0.31	4.45
1.2 IW/CPE ratio	8.20	0.31	4.68
SEm±	0.17	0.004	0.37
CD at 5%	0.58	0.016	2.23
Nitrogen sources			
100% N from urea (N1)	8.05	0.31	4.44
50% N from urea +50% N from FYM (N2)	8.05	0.31	4.68
50% N from urea + 50% N from Poultry manure (N3)	8.00	0.32	4.89
SEm±	0.17	0.005	0.46
CD at 5%	0.53	0.014	1.34

Available Nitrogen, Phosphorus Potassium and Zinc of Soil as influenced by Different Moisture Regimes and Nitrogen Sources at harvest of the crop

Slight increment was received with the increasing soil moisture regime up to 1.2 IW/CPE ratio. The lower value was obtained with 0.6 IW/CPE. Table 2.


In respect of nitrogen sources which were applied as 100% RDF, 50% RDF+ 50% through FYM and 50% through RDF+ 50% Poultry Manure.

The maximum nutrients viz N, P, K and Zn were analyzed at after harvest of the crop with the application Nitrogen through 50% RDF+ 50% Poultry manure. The minimum was received with the application of 100% RDF through inorganic fertilizer. The build-up of available nutrients with the application irrigation on the basis of IW/CPE ratio might be due to dilution of salts and leaching of ions beyond the root zone. Use of organic manures viz., poultry manure and FYM has released the availability and mobility of nutrients after the decomposition mineralization organics sources of nutrients mainly of micronutrients. These findings are corroborated by (Saha *et al.* 2019).

Table: 5. Available N, P, K& Zn of Soil as influenced by Different Moisture Regimes and Nitrogen Sources

Treatment	Available Nutrients (kg/ha ⁻¹)			
	Nitrogen (kg/ha ¹)	Phosphorus (kg/ha ¹)	Potassium (kg/ha ¹)	Zinc (mg/ha ¹)
Moisture regimes				
0.6 IW/CPE ratio	210.47	16.50	250.67	0.54
0.8 IW/CPE ratio	211.64	17.07	250.80	0.55
1.0 IW/CPE ratio	211.67	17.23	254.67	0.55
1.2 IW/CPE ratio	212.83	17.57	257.00	0.56
SEm±	2.24	0.41	1.90	0.02
CD at 5%	NS	NS	6.41	0.08
Nitrogen sources				
100% N from urea (N1)	212.93	16.28	250.87	0.58
50% N from urea +50% N from FYM (N2)	214.88	16.50	250.50	0.56
50% N from urea + 50% N from Poultry manure (N3)	216.93	16.65	254.77	0.64

SEm±	2.29	0.28	2.01	0.029
CD at 5%	NS	NS	NS	NS



UNDER PEER REVIEW

CONCLUSION

Moisture regimes 1.2 IW/CPE ratio was found best for viz., p^H, EC and organic carbon, Available Nitrogen, Phosphorus, Potassium & Zinc the minimum was received with the moisture regime at 0.6 IW/CPE during the investigation and among the Nitrogen Sources, application of 50% N from urea + 50% N from Poultry manure (N₃) was found suitable for soil fertility status followed by 50% N from urea +50% N from FYM (N₂).

Conference disclaimer:

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

ACKNOWLEDGEMENT

We are thankful to the faculty of the Department of Soil Science and Agricultural Chemistry, A.N.D.U.A&T, Kumarganj, Ayodhya for facilitating us and providing all the necessary resources.

References

- Akibode, C. S., & Maredia, M. K. (2012). *Global and regional trends in production, trade and consumption of food legume crops* (No. 1099-2016-89132).
- Debaeke, P., & Aboudrare, A. (2004). Adaptation of crop management to water-limited environments. *European Journal of Agronomy*, **21**(4), 433-446.
- Fazily, T., & Hunshal, C. S. (2019). Effect of organic manures on yield and economics of late sown wheat (*Triticum aestivum*). *International journal of research & review*, **6**(1), 168-171.
- Giraldo, P., Benavente, E., Manzano-Agugliaro, F., & Gimenez, E. (2019). Worldwide research trends on wheat and barley: A bibliometric comparative analysis. *Agronomy*, **9** (7), 352.
- Jat, R. A., Wani, S. P., Sahrawat, K. L., Singh, P., & Dhaka, P. L. (2011). Fertigation in vegetable crops for higher productivity and resource use efficiency. *Indian Journal of fertilizers*, **7**(3), 22-37.

- Kumar, P. Pannu, R.K., and Khokhar, S.K 2012. Effect of organic sources of nutrition and irrigation levels on growth and yield of wheat (*Triticum aestivum* L.) *International Journal of Life Sciences Biotechnology and Pharma Research* **1** (4): 178-186
- Kumar, S., Agrawal, S., Jilani, N., Kole, P., Kaur, G., Mishra, A., ... & Tiwari, H. (2023). Effect of integrated nutrient management practices on growth and productivity of rice: A review. *The Pharma Innovation Journal*. **12**(5): 2648-2662
- McCauley, A., Jones, C., & Jacobsen, J. (2009). Soil pH and organic matter. Nutrient management module, **8**(2), 1-12.
- Saha, S., Saha, B., Seth, T., Dasgupta, S., Ray, M., Pal, B..., & Hazra, G. (2019). Micronutrients availability in soil–plant system in response to long-term integrated nutrient management under rice–wheat cropping system. *Journal of Soil Science and Plant Nutrition*, **19**,712-724.
- Singh, T. B., Ali, A., Prasad, M., Yadav, A., Shrivastav, P., Goyal, D., & Dantu, P. K. (2020). Role of organic fertilizers in improving soil fertility. *Contaminants in agriculture: sources, impacts and management*, 61-77.
- Singh, V., Singh, S. P., Singh, S., & Shivay, Y. S. (2013). Growth, yield and nutrient uptake by wheat (*Triticum aestivum*) as affected by biofertilizers, FYM and nitrogen. *Indian Journal of Agricultural Sciences*, **83**(3), 331-334.
- Verma, H.P., Sharma O.P., Yadav L.R., Yadav S.S., Shivran A.C., Kumar R., and Balwan 2018 Growth Indices and Yield of Wheat (*Triticum aestivum* L.) As Influenced by Irrigation Scheduling and Organic Manures. *Journal of Pharmacognosy and Phytochemistry*; **7**(1): 908-912