

# Effect of Integrated Nutrient Management on Growth and Yield of Beetroot (*Beta vulgaris* L.) cv. Ruby Queen

## ABSTRACT

A field experiment was carried out to study the effect of integrated nutrient management on the growth and yield of beetroot (*Beta vulgaris* L.) cv. Ruby Queen. The experiment was laid out in a randomized block design using ten treatments and three replications at the college farm of the Department of Horticulture, Tilak Dhari PG College, Jaunpur, (U.P.) during the *rabi* season of 2021-22. Integration of nutrient sources was comprised in the form of treatments *viz.* T<sub>1</sub> - 100% NPK of RDF (70:110:70 Kg/ha), T<sub>2</sub> -75% NPK of RDF + FYM (10 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ ha), T<sub>3</sub> - 75% NPK of RDF + VC (3 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ha), T<sub>4</sub> - 75% NPK of RDF + FYM (5 t/ha) + VC (1.5 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ha), T<sub>5</sub> - 50% NPK of RDF + FYM (10 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ha), T<sub>6</sub> - 50% NPK of RDF + VC (3 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ha), T<sub>7</sub> - 50% NPK of RDF + FYM (5 t/ha) +VC (1.5 t/ha) + *Azotobacter* (10 Kg/ha) + PSB (10 Kg/ha), T<sub>8</sub> - 75% NPK of RDF + FYM (10 t/ha), T<sub>9</sub> - 50% NPK of RDF + VC (3 t/ha) and T<sub>10</sub> - Control. Among all treatments, T<sub>4</sub> (75% NPK of RDF + FYM + VC + *Azotobacter* + PSB) performed best for growth and yield parameters recorded significantly minimum number of days required for 80 percent germination of seedlings (2.68 days), highest plant height (50.24 cm), most leaves per plant (13.53), maximum leaf area (3227.50 cm<sup>2</sup>), higher chlorophyll content index (21.76), maximum root length (16.23 cm), root diameter (7.73 cm), root yield per plant (220.14 g), root yield per plot (4.40 Kg), root yield per hectare (22.014 t), and highest harvest index (0.91) was observed.

**Keywords:** Beetroot, Growth, Yield, INM, *Azotobacter*, PSB

## INTRODUCTION

“Beetroot (*Beta vulgaris* L.) is usually known as chukandar. It’s one of the most valuable root vegetable crops and is consumed worldwide due to its high nutrient content. It has a diploid chromosome number of 2n=18. Beetroot is a biennial crop although it is commonly grown annually. It produces green tops and swollen roots during the first growing

season. It is grown for vegetables, salad, Juice, and other food uses (pickles). Beetroot was first described in 1557 when it was referred to as Roman beet in Germany. The crop was introduced in the USA in 1800 and is known as garden beet. In India Beetroot is mainly cultivated in Haryana, Uttar Pradesh, Himachal Pradesh, West Bengal, Maharashtra, and Tamil Nadu. It is highly productive and usually free from pests and diseases” (Ado, 1999).

“Beetroot is a rich source of protein, carbohydrates, calcium, phosphorus, and vitamin C and has 87.7% of water per 100 g of fresh weight” (Aykroyd, 1963). “Hence it is an ideal vegetable for health-conscious people” (Deuter and Grundy, 2004). The colour of beetroot is due to the presence of red violet pigments of  $\beta$  cyanine and a yellow pigment  $\beta$  xanthine. It has many nutritive properties such as antiradical properties which improve the quality of human blood and revive stress-based disorders (Pedreno and Escribano, 2000; Kugler *et al.*, 2007) drawing the attention of consumers to use it in cuisine (Kavitha *et al.*, 2014). “It is also beneficial in many clinical conditions including coronary heart disease and cancer” (Jurgen *et al.*, 2015). “It is one of the natural foods which boosts the energy in athletes as it has one of the highest nitrates and sugar contents” (Yadav *et al.*, 2016). “It received increasing attention due to possible health benefits in humans, especially its antioxidant and anti-inflammatory activities” (Georgie *et al.*, 2010 and Zielinska *et al.*, 2009).

“In modern times, chemical fertilizers have become the main source for providing the required quantities of nutrients to plants, and application aids in achieving maximum production but it must be interpreted given its harmful effects on the environment as well as increasing production costs due to its expenses” (Dadarwal *et al.*, 2009; Ranjan *et al.*, 2013). “The continuous use of chemical fertilizers often in excess, over a long time in arable land has led to the contamination of food material, environmental pollution, and depletion of soil fertility” (Singh, 1999). To reduce the cost of chemical fertilizers and their harmful effect on soil, environment, and quality, it is necessary to seek an alternative that will improve productivity, quality, and also be eco-friendly to the environment.

“Integrated nutrient management (INM) is a possible way forward to achieve sustainable crop production rather than relying on only inorganic fertilizers” (Paroha *et al.*, 2009). The use of combined organic manure, bio-fertilizers, and reduced-dose chemical fertilizers not only helps to improve crop growth, yield, and quality but also helps to maintain soil health, reduce pollution problems and increase production (Ahmad *et al.*, 2016) and ensure better food security. Keeping all the above points in view the current research work initiates to find out

the ideal combination of integrated nutrient management on growth and yield of beetroot (*Beta vulgaris* L.) cv. Ruby Queen.

## MATERIALS AND METHODS

The present experiment was laid out at the college farm of the Department of Horticulture, Tilak Dhari Post Graduate College, Jaunpur, Uttar Pradesh on 1<sup>st</sup> week of November during the *rabi* season of the year 2021-22. The experimental site is situated at an altitude of 79.75 m above mean sea level at 25°73' North latitude and 82°68' East longitude. The climate condition of Jaunpur is sub-tropical with three seasons *i.e.*, rainy, winter, and summer. During the summer season temperature reaches as high as 45°C, while in winter it falls to 5°C or even low. Most of the rainfall was received between the third week of June and the first week of October. The annual rainfall of Jaunpur is about 1097.9 mm with 46 normal rainy days. The experiment was carried out in a randomized block design using ten treatments having three replications. The treatments comprises of *i.e.*, T<sub>1</sub> - 100% NPK of RDF (70:110:70 Kg ha<sup>-1</sup>), T<sub>2</sub> - 75% NPK of RDF + FYM (10 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>3</sub> - 75% NPK of RDF + VC (3 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>4</sub> - 75% NPK of RDF + FYM (5 t ha<sup>-1</sup>) + VC (1.5 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>5</sub> - 50% NPK of RDF + FYM (10 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>6</sub> - 50% NPK of RDF + VC (3 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>7</sub> - 50% NPK of RDF + FYM (5 t ha<sup>-1</sup>) + VC (1.5 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>), T<sub>8</sub> - 75% NPK of RDF + FYM (10 t ha<sup>-1</sup>), T<sub>9</sub> - 50% NPK of RDF + VC (3 t ha<sup>-1</sup>) and T<sub>10</sub> - Control.

The variety under investigation was 'Ruby Queen' which is a high-yielding variety. Seeds of this variety were collected from a certified private seller. The soil of the experimental plot was deep, loamy sand. The site was divided into a 2 m x 1 m plot size of 30 plots. Seeds were sown in a ridge and furrow system at a depth of 1.5 cm and spacing of 45 cm × 15 cm to ensure proper spacing. To cultivate a good crop, all other recommended cultural practices were followed properly. N, P, and K were given through Urea, Single Super Phosphate, and Muriate of Potash respectively. The full dose of phosphorus and potassium and a half dose of nitrogen was applied as basal dose, as per treatment before sowing, and the remaining half dose of nitrogen was given after 30 days of sowing. Manures *viz.*, FYM and vermicompost were integrated as per treatment to respective plots before sowing. Biofertilizers (*Azotobacter* and PSB as *Bacillus megaterium*) were inoculated to seeds before sowing as a seed treatment.

The observations of Growth parameters and yield parameters were recorded on five randomly selected and tagged plants from each treatment in all replications.

Growth parameters were the number of days required for 80% germination of the seedlings, plant height (cm) which was measured from ground level to the tip of the longest leaf, number of leaves per plant, leaf area (cm<sup>2</sup>) which was measured using non-destructive methods by recording the observations of length and width of the leaf at the middle portion of each leaf and chlorophyll content index measured using spad meter and recorded data on 25 DAS, 45 DAS, and 65 DAS. Yield parameters were the root length (cm), root diameter (cm), root yield per plant (g plant<sup>-1</sup>), root yield per plot (Kg plot<sup>-1</sup>), root yield per hectare (t ha<sup>-1</sup>), and harvest index. The statistical analysis of the variance of the data in respect of growth and yield components of beetroot was done according to the standard procedure for factorial randomized block design given by Panse & Sukhatme (1985).

## **RESULT AND DISCUSSION**

### **GROWTH PARAMETERS**

**The number of days required for 80 % germination:** The days required for 80 % germination of beetroot seedlings were significantly influenced by integrated nutrient combinations as presented in table 1. Among all the treatments T<sub>4</sub> (75% NPK of RDF + FYM (5 t ha<sup>-1</sup>) + VC (1.5 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>)) resulted in early germination (2.68 days) and it was at par with T<sub>7</sub> (3.14 days). The results show early germination in the combined application of N, P, K, FYM, VC, *Azotobacter*, and PSB as *Bacillus megaterium*, which was due to the combined effect of all these nutrients which enhanced the availability of all macro and micronutrients in the rhizosphere also of its high content of nitrogen, phosphorus, and potassium may help in early germination of seedlings (Warman, 1986). The findings agreed with those of Gyewali *et al.* (2020). Similar results were also obtained by Jagadeesh *et al.* (2018) and Mounika *et al.* (2021) in beetroot.

**Table 1: Effect of integrated nutrient management on number of days required for 80 % germination of beetroot seedlings cv Ruby Queen**

Treatments (T)	Number of days required for 80 % germination (days)
T <sub>1</sub> -100% NPK	3.98
T <sub>2</sub> -75% NPK + FYM + <i>Azotobacter</i> + PSB	3.28
T <sub>3</sub> -75% NPK + VC + <i>Azotobacter</i> + PSB	3.21
T <sub>4</sub> -75% NPK + FYM + VC + <i>Azotobacter</i> + PSB	2.68
T <sub>5</sub> -50% NPK + FYM + <i>Azotobacter</i> + PSB	3.67
T <sub>6</sub> -50% NPK + VC + <i>Azotobacter</i> + PSB	3.51
T <sub>7</sub> -50% NPK + FYM + VC + <i>Azotobacter</i> + PSB	3.14
T <sub>8</sub> -75% NPK + FYM	4.05
T <sub>9</sub> -50% NPK + VC	4.08
T <sub>10</sub> -Control	4.81
<b>CD at 5 %</b>	<b>0.49</b>
<b>SE (m) ±</b>	<b>0.16</b>

**Plant height:** The data on plant height at 25, 45, and 65 days after sowing was significantly increased by the treatment as shown in table 2. At 25 DAS the highest plant height was recorded in T<sub>4</sub> (15.77 cm) and it was at par with T<sub>7</sub> (15.42 cm) and T<sub>3</sub> (15.19 cm) but significantly superior to all other treatments. At 45 DAS and 65 DAS T<sub>4</sub> treatments were significantly superior to all other treatments with heights of 36.74 cm and 50.24 cm respectively. The treatment T<sub>4</sub> (75% NPK of RDF + FYM + VC + *Azotobacter* + PSB) was recorded at the highest plant height in every stage. The positive effect of treatment (T<sub>4</sub>) on plant height could be due to the readily available required nitrogen, which is a component of amino acids, nucleotides, nucleic acids, a number of co-enzymes, auxins, cytokinin and alkaloids that resulted in increased cell elongation, cell enlargement, and cell division, Also, higher plant height could be due to certain growth promoting substances secreted by the *Azotobacter* and PSB, which improved better transportation of water, uptake and deposition of nutrients which positively affect the better root development. The results are in conformity with the findings of Singh *et al.* (2017) and Shanu *et al.* (2019) in carrot, Dlamini *et al.* (2020), and Mounika *et al.* (2021) in beetroot.

**Number of leaves per plant:** The data recorded on the number of leaves per plant was significantly influenced by the various integrated nutrient management treatments at different stages of plant growth. The results are presented in table 2. At 25 DAS, the maximum number of leaves per plant was observed in T<sub>4</sub> (6.40) which was at par with T<sub>7</sub> (6.20). At 45

DAS T<sub>4</sub> (10.47) recorded the highest value which was at par with T<sub>7</sub> (10.07) and 65 DAS, the Maximum value was recorded in T<sub>4</sub> (13.53) which was significantly superior to all other treatments. T<sub>7</sub> (12.53) was comparable with T<sub>2</sub> (12.07) and T<sub>3</sub> (12.20). The maximum number of leaves per plant at all growth stages was recorded in T<sub>4</sub> treatment with 75% NPK of RDF + FYM + VC + *Azotobacter* + PSB which was due to the timely supply of all nutrients in the rhizosphere, resulted in a greater number of leaves per plant. Besides, it might be due to the same reasons that registered the highest plant height over other treatments. Kumar *et al.* (2007) has also advocated Microbial based IPNS for higher growth of rice. These results are in accordance with that of Shanu *et al.* (2019) in carrot, Dlamini *et al.* (2020), and Mounika *et al.* (2021) in beetroot.

**Table 2: Effect of integrated nutrient management on plant height (cm) and Number of leaves per plant at different growth stages of beetroot cv. Ruby Queen**

Treatments (T)	Plant height (cm)			Number of leaves per plant		
	25 DAS	45 DAS	65 DAS	25 DAS	45 DAS	65 DAS
T <sub>1</sub>	13.62	30.56	39.61	5.20	8.80	11.13
T <sub>2</sub>	14.75	33.23	44.95	5.73	9.87	12.07
T <sub>3</sub>	15.19	33.97	44.71	5.80	9.80	12.20
T <sub>4</sub>	15.77	36.74	50.24	6.40	10.47	13.53
T <sub>5</sub>	14.20	31.33	43.28	5.60	9.27	11.40
T <sub>6</sub>	14.07	31.03	42.63	5.73	9.13	11.47
T <sub>7</sub>	15.42	34.63	46.79	6.20	10.07	12.53
T <sub>8</sub>	13.52	31.02	40.74	5.53	9.00	11.40
T <sub>9</sub>	12.90	29.67	39.29	5.40	8.93	11.00
T <sub>10</sub>	10.55	26.43	34.89	5.07	8.20	10.13
CD at 5 %	<b>0.73</b>	<b>1.59</b>	<b>2.27</b>	<b>0.33</b>	<b>0.58</b>	<b>0.63</b>
SE (m) ±	<b>0.24</b>	<b>0.53</b>	<b>0.76</b>	<b>0.11</b>	<b>0.19</b>	<b>0.21</b>

**Leaf area:** Leaf area was found significant influence by the integrated nutrient management treatments. The results are presented in table 3. At 25 DAS T<sub>4</sub> recorded a maximum leaf area (270.11 cm<sup>2</sup>) which was at par with T<sub>7</sub> (253.59 cm<sup>2</sup>). At 45 DAS highest leaf area was recorded in T<sub>4</sub> (1649.52 cm<sup>2</sup>) and it was comparable with T<sub>7</sub> (1609 cm<sup>2</sup>). At 65 DAS maximum leaf area was recorded in T<sub>4</sub> (3227.50 cm<sup>2</sup>) and it was at per with T<sub>7</sub> (3136.76 cm<sup>2</sup>), followed by T<sub>3</sub> treatment (3069.09 cm<sup>2</sup>), and T<sub>2</sub> (3052.43 cm<sup>2</sup>). The data revealed that maximum leaf area at 25, 45, and 65 days after sowing was recorded in T<sub>4</sub> treatment with 75% NPK of RDF + FYM + VC + *Azotobacter* + PSB may be due to the same treatment registered the highest plant height and a greater number of leaves per plant, resulted in

maximum leaf length and width of the leaf led to higher leaf area at all growth stages. Similar results were also reported by Jaisankar (2018) in radish.

**Chlorophyll content index:** The chlorophyll content index was found to be significant to the integrated nutrient management treatment combinations as shown in table 3. At 25 DAS the highest value was recorded in T<sub>4</sub> (13.88) and it was at par with T<sub>7</sub> treatment (13.35), T<sub>3</sub> (13.19), and T<sub>2</sub> (13.06). It was the minimum in the T<sub>10</sub> treatment (8.99). At 45 DAS Maximum value was recorded in the T<sub>4</sub> treatment (17.44). At 65 DAS maximum value was recorded in T<sub>4</sub> (21.76) treatment and was at par with T<sub>7</sub> (20.84) and T<sub>3</sub> (20.62). T<sub>4</sub> treatment with 75 % NPK of RDF + FYM + Vermicompost + *Azotobacter* + PSB recorded the highest value in every stage. It might be due to readily available macro and micronutrients, particularly nitrogen supplied by FYM, vermicompost, and biofertilizers, and an important constituent of chlorophyll. These results are in accordance with that of Gairola *et al.* (2009) in spinach beet.

**Table 3: Effect of integrated nutrient management on leaf area (cm<sup>2</sup>) and Chlorophyll content index at different growth stages of beetroot cv. Ruby Queen**

Treatments (T)	leaf area (cm <sup>2</sup> )			Chlorophyll content index		
	25 DAS	45 DAS	65 DAS	25 DAS	45 DAS	65 DAS
T <sub>1</sub>	200.33	1404.78	2809.85	12.01	15.10	18.41
T <sub>2</sub>	236.77	1555.37	3052.43	13.06	16.12	20.39
T <sub>3</sub>	239.09	1528.47	3069.09	13.19	16.26	20.62
T <sub>4</sub>	270.11	1649.52	3227.50	13.88	17.44	21.76
T <sub>5</sub>	222.00	1465.76	2961.73	12.40	15.58	18.86
T <sub>6</sub>	230.34	1456.78	2996.93	12.31	15.81	19.61
T <sub>7</sub>	253.59	1608.22	3136.73	13.35	16.50	20.84
T <sub>8</sub>	209.72	1425.11	2871.15	12.31	15.33	19.37
T <sub>9</sub>	201.35	1409.26	2797.64	10.21	14.21	17.37
T <sub>10</sub>	162.48	1201.81	2348.32	8.99	12.09	15.58
<b>CD at 5 %</b>	<b>24.97</b>	<b>64.46</b>	<b>121.11</b>	<b>1.01</b>	<b>1.22</b>	<b>1.26</b>
<b>SE (m) ±</b>	<b>8.34</b>	<b>21.53</b>	<b>40.45</b>	<b>0.34</b>	<b>0.41</b>	<b>0.42</b>

## YIELD PARAMETERS

**Root length and Root diameter:** Root length and root diameter were significantly influenced by the integrated nutrient management treatments. The results are presented in table 4.

The highest root length (16.23 cm) was recorded in the T<sub>4</sub> treatment and it was on par with the T<sub>7</sub> treatment (15.67 cm), T<sub>3</sub> (15.20 cm), and T<sub>2</sub> (15.067 cm), whereas the lowest value

was recorded in T<sub>10</sub> treatment (9.27 cm) with control and the highest root diameter (7.73 cm) was recorded in T<sub>4</sub> treatment which was at par T<sub>7</sub> (7.49 cm), T<sub>2</sub> (7.27 cm) and T<sub>3</sub> (7.17 cm). T<sub>1</sub> with 100% NPK of RDF recorded a root diameter of 6.13 cm which is comparable with T<sub>9</sub> (5.99 cm) while the lowest was recorded in T<sub>10</sub> (4.74) with control. It might be due to enhanced cell division and quick cell multiplication resulted in better plant growth in all aspects and more translocation of photosynthates from leaves (source) to root (sink) led to increased root length and diameter. Similar results were also reported by Mali *et al.* (2018) in radish, Ingole *et al.* (2018), and Dlamini *et al.* (2020) in beetroot.

The results pertaining to root yield per plant (g plant<sup>-1</sup>), root yield per plot (Kg plot<sup>-1</sup>), root yield per hectare (t ha<sup>-1</sup>), and harvest index were significantly influenced by the integrated nutrient management treatments combinations as results are presented in table 4.

**Root yield:** The highest value of root yield per plant was recorded in T<sub>4</sub> (220.14 g plant<sup>-1</sup>) treatment which is significantly superior to all, followed by the T<sub>7</sub> treatment (191.98 g plant<sup>-1</sup>). It was significantly minimum (96.77 g plant<sup>-1</sup>) in T<sub>10</sub> treatment with control.

Root yield per plot is significantly superior to treatments with T<sub>4</sub> recorded as the highest value (4.40 Kg plot<sup>-1</sup>) and superior among others, followed by T<sub>7</sub> treatment (3.84 Kg plot<sup>-1</sup>), whereas it was minimum in T<sub>10</sub> (1.94 Kg plot<sup>-1</sup>) treatment with control. T<sub>1</sub> (100% NPK of RDF) recorded a lower value (2.97 Kg plot<sup>-1</sup>) among treatments combination and was only better than T<sub>9</sub> (2.74 Kg plot<sup>-1</sup>).

Maximum root yield per hectare was recorded (22.014 t ha<sup>-1</sup>) in the T<sub>4</sub> treatment which was significantly superior to anyone followed by the T<sub>7</sub> treatment (19.20 t ha<sup>-1</sup>) and was statistically significant with each other, while it was minimum in T<sub>10</sub> (9.68 t ha<sup>-1</sup>) with control. T<sub>1</sub> (100% NPK of RDF) recorded 14.87 t ha<sup>-1</sup> which is a lower value than all treatments and only better than T<sub>9</sub> (13.72 t ha<sup>-1</sup>) (fig. 1).


**Harvest index:** The highest harvest index (0.91) was recorded in T<sub>4</sub> which was significantly superior to all other treatments, followed by T<sub>7</sub> (0.84). T<sub>1</sub> with 100% NPK of RDF recorded a harvest index of 0.73 and T<sub>10</sub> with control resulted lowest (0.64).

The data shows that T<sub>4</sub> treatment with 75% NPK of RDF + FYM + VC + *Azotobacter* + PSB recorded the highest value in root yield per plant (g plant<sup>-1</sup>), root yield per plot (Kg plot<sup>-1</sup>), root yield per hectare (t ha<sup>-1</sup>), and harvest index parameters which might be due to the maximum root length and root diameter recorded over other treatments. The present investigation was consistent with other reports by Pathak *et al.* (2018), and Monika *et al.*

(2018) in radish. The present findings are comparable with that of Jagadeesh *et al.* (2018) and Mounika *et al.* (2021) in beetroot.

**Table 4: Effect of integrated nutrient management on root length (cm), root diameter (cm), root yield per plant (g plant<sup>-1</sup>), root yield per plot (Kg plot<sup>-1</sup>), root yield per hectare (t ha<sup>-1</sup>), and harvest index of beetroot *cv.* Ruby Queen**

Treatments (T)	Root length (cm)	Root diameter (cm)	Root yield (g plant <sup>-1</sup> )	Root yield (Kg plot <sup>-1</sup> )	Root yield (t ha <sup>-1</sup> )	Harvest index
T <sub>1</sub>	13.633	6.127	148.663	2.973	14.866	0.727
T <sub>2</sub>	15.067	7.267	171.727	3.435	17.173	0.813
T <sub>3</sub>	15.200	7.173	184.170	3.683	18.417	0.833
T <sub>4</sub>	16.233	7.727	220.143	4.403	22.014	0.907
T <sub>5</sub>	14.933	6.747	152.293	3.046	15.229	0.787
T <sub>6</sub>	14.767	6.810	159.287	3.186	15.929	0.800
T <sub>7</sub>	15.667	7.487	191.977	3.840	19.198	0.840
T <sub>8</sub>	14.300	6.630	153.907	3.078	15.391	0.747
T <sub>9</sub>	13.133	5.993	137.150	2.743	13.715	0.720
T <sub>10</sub>	9.267	4.743	96.767	1.935	9.677	0.637
<b>CD at 5 %</b>	<b>1.225</b>	<b>0.597</b>	<b>23.727</b>	<b>0.475</b>	<b>2.373</b>	<b>0.046</b>
<b>SE (m) ±</b>	<b>0.409</b>	<b>0.199</b>	<b>7.924</b>	<b>0.158</b>	<b>0.792</b>	<b>0.015</b>



**Fig. 1: Effect of integrated nutrient management on root yield (t ha<sup>-1</sup>) of beetroot *cv.* Ruby Queen**

## CONCLUSION

Based on the findings of the current study among all the different integrated nutrient treatment combinations, T<sub>4</sub> treatment with 75% NPK of RDF + FYM (5 t ha<sup>-1</sup>) + VC (1.5 t ha<sup>-1</sup>) + *Azotobacter* (10 Kg ha<sup>-1</sup>) + PSB (10 Kg ha<sup>-1</sup>) proven to be the best treatment to give maximum results on growth parameters and yield parameters of beetroot (*Beta vulgaris* L.) cv Ruby Queen. Therefore, it may be concluded that the beetroot producers suggested to apply T<sub>4</sub> treatment in their field to get maximum production and profits.

### Conference disclaimer:

Some part of this manuscript was previously presented in the conference: 6th International Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding:

<https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

## REFERENCES

- Ado, P.O., 1999. Beetroot Cultivation. *Beetroot and Eggplant Newsletter*, 18: 21 – 24. 8(1):81-83.
- Ahmad, I., Ahmad, B., Ali, S., Fang, H.Q., Liu, T., Kamran, M., Wu, Y., Rahman M. U. and Faisal, S. (2016). Organic and Inorganic Fertilization Influenced on Yield and Quality of Sugar Beet Genotypes. *Russian Agricultural Sciences*, 42(3- 4): 218–223
- Aykroyd, U.R. (1963) Indian Council of Medical Research, Special Report. *Vegetables, National Book Trust India*, New Delhi, 42, 188-191.
- Dlamini, V. C., Nxumalo, K. A., Masarirambi, M. T., Wahome, P.K., Oseni T. O and Zwane, M. G. (2020). Effects of Cattle Manure on the Growth, Yield, Quality, and Shelf Life of Beetroot (*Beta vulgaris* L.) Cv. Detroit Dark Red. *Journal of Experimental Agriculture International*, 42(1): 93-104.
- Deuter, P. and Grundy, T. (2004). Beetroot Commercial Production and Processing. Agency for Food and Fibre Sciences. *Holland Horticultural Limited Partnership*. P. 1 – 4.
- Dadarwal, R. S., Jain N. K. and Singh, D. (2009). Integrated management in baby corn (*Zea mays*). *Indian Journal of Agricultural Sciences*, 79(12): 1023–1025.

- Gairola, S., Umar S. and Suryapani, S. (2009). Nitrate accumulation, growth, and leaf quality of spinach beet as affected by NPK fertilization with special reference to potassium. *Indian Journal of Science and Technology*, 2: 35-39.
- Georgiev, V. G., Weber, J., Kneschke, E. M., Denev, P. N., Bley, T and Pavlov, A. I. 2010. Antioxidant activity and phenolic content of betalain extracts from intact plants and hairy root cultures of the red beetroot (*Beta vulgaris*). *Journal of Plant Foods for Human Nutrition*. 65(2): 105-111.
- Gyewali, B., Maharjan, B., Rana, G., Pandey, R., Pathak, R., and Poudel, P. R. (2020). Effect of different organic manures on growth, yield, and quality of radish (*Raphanus sativus*). *SAARC Journal of Agriculture*, 18(2): 101-114.
- Ingole, V.S., Wagh, P.K., Nagre, P.K and Bharad, S.G. 2018. Effect of biofertilizer and organic manure on yield and yield contributing characters of beetroot (*Beta vulgaris* L.). *International Journal of Chemical Studies*. 6(5): 1226-1228.
- Jagadeesh, C., Madhavi, M., Prasad, S and Padmaja, V.V. 2018: Effect of organic manures on growth and yield attributes of beetroot cv. crimson globe. *International Journal of Current Microbiology and Applied Sciences*. 7(11): 3538-3553.
- Jaisankar, P. 2018. Effect of Integrated Nutrient Management on Growth and yield of radish (*Raphanus sativus* L.) cv. Pusa Chetki. *Int.J.Curr.Microbial.App Sci*. 7(11): 461-466.
- Jurgen, W., Gundula, W., Stefan, H., Pinar, U., Peter L. and Ulrike, M. (2015). Composition characteristics of commercial beetroot products and beetroot juice prepared from seven beetroot varieties grown in Upper Austria. *Journal of Food Composition and Analysis*, 42: 46-55.
- Kavitha, R., Ravichandran, P., Saw, N.M.M.T., Gabr, A.M.M., Abdel, R.R., Ahmed, R., Knorr, D. and Smetanska, I. 2014. Effects of different encapsulation agents and drying process on the stability of betalains extract. *J. Food Sci. Technol.*, 51(9): 2216–2221.
- Kugler, F., Stintzing, F.C. and Carle, R. 2007. Evaluation of the antioxidant capacity of betalainic fruits and vegetables. *J.Appl. Bot. Food Qual.*, 81(1): 69-76.
- Kumar, N., Singh R., Misra, B.N. and Chandra, K. K. (2007). Influence of Organic Manure, Biofertilizer and Nitrogen Levels on Nutrients Uptake, Soil Properties and Yield of Rice. *Indian Journal of Fertilizers*, 3 (7): 45.

- Mali, D. L., Virendra, S., Sarolia, D. K., Suresh, K. T., Akshay, C and Dhakar, R. 2018. Effect of organic manures and biofertilizers on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese white. *International Journal of Chemical Studies*. 6(2):1095-1098.
- Monika, S., Sharma, A. K and Shilpa. 2018. Integrated Nutrient Management of Pea-Capsicum-Radish Cropping Sequence in Sub-Temperate Climate of North-Western Himalayas. *Int. J. Curr. Microbiol. App.Sci*. 7(3): 538-545.
- Mounika, D. Lakshminarayana, J. Srinivas, G. Sathish and P. Gouthami (2022). Influence of Integrated Nutrient Management in Beet Root (*Beta Vulgaris* L.) Cv. Crimson Globe for Growth and Yield Under Alkaline Conditions. *Biological Forum – An International Journal*, 14(1): 464-469.
- Pathak, M., Tripathy, P., Dash, S. K., Sahu, G.S and Pattanayak, S. K. 2018. Efficacy of biofertilizer, organic and inorganic fertilizer on yield and quality of radish (*Raphanus sativus* L.). *International Journal of Chemical Studies*. 6(4): 1671-1673.
- Pedreno, M. A., & Escribano, J., “Studying the oxidation and antiradical activity of betalain from beetroot,” *Journal of Biological Education*, 35, pp.49–59, 2000.
- Panse, V. G. and Sukhatme, P. V. 1967. Statistical methods for Agricultural workers. Indian Council of Agricultural Research, New Delhi.
- Ranjan, J. K., Ahmed, N., Das, B., Ranjan, P. and Mishra, B. K. (2013). Green technology for production of baby corn (*Zea mays* L.) under north-west Himalayan conditions. *International Journal of Chemical Technology Research*, 5(2): 880–885.
- Paroha, S., Chandra, K. K., Yadav, R. (2009). Integrated effect of biofertilizers (AM, Azotobacter & PSB) fertilizers on growth and nutrient acquisition by *Tectona grandis*. *Journal of Tropical Forestry*, 25 (1/2): 54-61.
- Shanu, V., D. Lakshminarayana, P. Prasanth and Saida Naik, D. 2019. Studies on the Effect of Integrated Nutrient Management (INM) on Growth and Yield Parameters of Carrot (*Daucas carota* L.) cv. Kuroda Improved under Southern Telangana Conditions. *Int.J.Curr.Microbiol.App.Sci*. 8(04): 2786-2791.

- Singh, D. P., Kumar, S., Sutanu, M. and Vijay, P. K. (2017). Studies on integrated nutrient management on growth, yield, and quality of carrot (*Daucus carota* L.). *International Journal of Agriculture Sciences*, 51(8): 2187-2188.
- Singh, G. B. 1999. Sustainable use of micronutrients in agriculture. Proceed on the national symposium on the management of micronutrient deficiencies in alluvial plains. 4-7.
- Warman P.R., The effect of fertilizer, chicken manure, and dairy manure on timothy yield, tissue composition and soil fertility, *Agricultural Wastes*, 10.1016/0141-4607(86)90074-0, 18, 4, (289-298), (1986).
- Yadav, M., Chawla, H., Parle, M. and Sharma, K. (2016). Beetroot: A health-promoting functional food. *Inventi Rapid: Nutraceuticals*. (1), 1-5.
- Zielinska, P. M., Olejnik, A., Dobrowolska, Z and Grajek, W. 2009. In vitro effects of beetroot juice and chips on oxidative metabolism and apoptosis in neutrophils from obese individuals. *Phytotherapy Research*. 23(1):49-55.