

**Original Research Article**  
**EFFECT OF INTEGRATED NUTRIENT MANAGEMENT (INM)  
PRACTICES ON GROWTH, YIELD AND ECONOMICS OF SWEET  
PEPPER UNDER OPEN CONDITION AT MID HILL OF HIMACHAL  
PRADESH**

**ABSTRACT**

The present investigation entitled “**Effect of Integrated Nutrient Management (INM) practices on growth, yield and economics of Sweet Pepper under open condition at mid hill of Himachal Pradesh**” was carried out at the Research farm of the School of Agriculture, Abhilashi University, Mandi (H.P.) during *Kharif* season 2022. The experiment was laid out in a randomized block design with three replications. The experiment consisted of two Bulky organic manures (FYM and Vermicompost) and three chemical fertilizers (Nitrogen, Phosphorus and Potassium). The result revealed that maximum plant height (30.33 cm, 42.47 cm and 55.03 cm at 30, 45 and 60 DAT respectively), number of branches per plant (3.10, 3.43 and 4.03 at 30, 45 and 60 DAT respectively), number of leaves per plant (31.50, 32.60 and 35.03 at 30, 45 and 60 DAT respectively), plant spread (27.60 and 34.50 cm), minimum number of days taken to 50 per cent flowering (36.01), fruit length (8.03 cm), fruit diameter (6.73 cm), average fruit weight (72.07 g), maximum number of fruits per plant (13.13), maximum number of flowers per plant (14.47), number of pickings (4.33), yield per plant (1.01 kg), fruits yield per plot (19.21 kg/plot) and fruits yield per hectare (350.34 q/ha) were recorded with the conjoint use of 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost in treatment T<sub>7</sub>. Treatment T<sub>7</sub> also showed higher nutrient availability in the soil with the residual nutrient status of nitrogen (269.97 kg/ha), phosphorus (40.03 kg/ha) and potassium (292.33 kg/ha). Maximum gross return (₹ 595578/ha), net return (₹ 485173/ha) and maximum B: C ratio (1:5.39) was obtained in T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). Thus, application of optimum doses of NPK along with organic manures was found highly beneficial for plant growth and yield of sweet pepper.

**Keywords:** *Integrated Nutrient Management (INM), sweet pepper, B: C ratio and yield.*

**1. INTRODUCTION**

Sweet pepper (*Capsicum annum* L.) is an important crop belongs to the family Solanaceae. It is an annual, day neutral plant grown extensively in sub- temperate climate throughout the world. British colonists introduced sweet pepper to India in the 19th century and the first successful large scale cultivation was done in the Shimla hills area, that is why named as “Shimla mirch”. It is second most consumed vegetable crop worldwide. In India, sweet pepper occupies an area of 37 thousand hectares with a production of 586 thousand metric tonnes [1]. In Himachal Pradesh, sweet pepper is being cultivated as cash crop in the sub- temperate areas and covers an area of 2.85 thousand hectare with production of 48.86 metric tonnes [1]. Himachal Pradesh is the top producer of fresh sweet pepper fruits to the plains during the summer and rainy seasons which provides large returns to small and marginal hill farmers.

Sweet pepper is cultivated as summer crop in the hilly areas where as autumn- winter crop in the plains of India. It can be grown in well- drained, loamy soils rich in organic matter. It is susceptible to frost and temperatures between 26 - 28 °C during the day and 16 - 18 °C at night is ideal for its growth and development. Extreme temperatures, such as those below 15 °C and above 30 °C, greatly hinder growth and fruit set. High relative humidity is good for growth, development, fruit set, and yield, but it also encourages the spread of foliar and fruit diseases Hazra et al. [2].

In India especially in Himachal Pradesh, in spite of its great potential the yield per unit area is quite poor when compared to the developed countries. This is mostly because there are no adequate production technologies, particularly for the nutrient management. Only one or two nutrients are supplied to the crop when chemical fertilisers are used exclusively without the use of organic manure. On the other hand, organic manures are known to have beneficial effect on soil health but its low

**Comment [Dr1]:** Effect Of Integrated Nutrient Management (INM) Practices On Growth, Yield And Economics Of Sweet Pepper Under Open Condition At Mid Hill Of Himachal Pradesh

**Comment [Dr2]:** vermicompost

**Comment [Dr3]:** nutrients i.e nitrogen, phosphorus and potassium.

**Comment [Dr4]:** delete

**Comment [Dr5]:** delete

**Comment [Dr6]:** Delete

**Comment [Dr7]:** This treatment

**Comment [Dr8]:** Delete

**Comment [Dr9]:** Delete

**Comment [Dr10]:** delete

**Comment [Dr11]:** delete

**Comment [Dr12]:** 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost

**Comment [Dr13]:** Shimla Mirch

**Comment [Dr14]:** Delete

**Comment [Dr15]:** in

**Comment [Dr16]:** Delete

nutritional content and abundant supply makes it difficult to use on a larger scale. So combined application of organic and inorganic fertilizers is considered ideal for obtaining the sustainable crop production. Keeping this in view, the present experiment is proposed to be undertaken to find out the effect of Integrated Nutrient Management (INM) practices on the growth, yield and economics of sweet pepper under open condition at mid hill of Himachal Pradesh.

Comment [Dr17]: Delete and add limited supply

Comment [Dr18]: Delete and add was

Comment [Dr19]: Delete

Comment [Dr20]: Integrated Nutrient Management (INM) Practices on The Growth, Yield and Economics of Sweet Pepper under Open Condition At Mid Hill of Himachal Pradesh.

## 2. METHODOLOGY

### 2.1 Experimental site

The present investigation was conducted during the *Khari* season 2022 at Research farm of the School of Agriculture, Abhilashi University, Mandi (H.P). The experimental farm situated at 31°33'32"N latitude and 77°00'47"E longitudes with the elevation of 1404 m above mean sea level. Soil at the experimental site was sandy clay loam, neutral in reaction, low in available nitrogen, medium in available potassium and high in available phosphorus.

Comment [Dr21]: Delete at and add of

Comment [Dr22]: Give the values

### 2.2 Treatment and crop culture

The experiment was laid out in Randomized Block Design (RBD) Factorial with three replications comprising of seven treatment combination of manure and liquid manure, the details of treatment combination is given in Table 1. Seeds of sweet pepper cv. 'Royal Wonder' were sown in the nursery in raised beds under polyhouse condition in 1.5 x 1 x 0.15 m seed beds at Research Farm of Abhilashi University on 1<sup>st</sup> April, 2022 and transplanting was done on 12<sup>th</sup> May, 2022. The plot size was 3 x 1.8 m<sup>2</sup> and a spacing of 60 cm x 45 cm was followed. After 20 days of transplanting, five plants were tagged randomly of each plot for taking observation. FYM and vermicompost were applied before sweet pepper was sown. N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied through urea, single super phosphate (SSP) and muriate of potash (MOP) respectively according to the treatment plan. The half dose of N and full doses of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O in the form of chemical fertilizer were applied at the sowing time in each plot in accordance with the different treatments.

Table 1. Details of treatments

Symbols	Treatment combinations
T <sub>1</sub>	Control (without fertilizers)
T <sub>2</sub>	100% RDF through inorganic fertilizers
T <sub>3</sub>	100% Recommended dose of FYM
T <sub>4</sub>	100% Recommended dose of vermicompost
T <sub>5</sub>	75% RDF through inorganic fertilizer + 25% through FYM
T <sub>6</sub>	75% RDF through inorganic fertilizer + 25% through vermicompost//
T <sub>7</sub>	50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost

\*RDF: Recommended Dose of Fertilizer (100 N: 75 P: 55 K kg/ha)

\*\* FYM: Farm Yard Manure

Comment [Dr23]: 100:75:55Kg N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O /ha

### 2.3 Soil Analysis

The soil samples from 0–15 cm depth were randomly collected at five places in each plot before and after experimentation. The soil samples were thoroughly mixed and composite samples were obtained. The soil samples were air-dried, ground and passed through 2-mm sieve. Soil samples were analyzed for pH with 1:2.5 soil water suspension glass electrode pH method Jackson [3], available nitrogen content was analyzed with alkaline permanganate method Subbiah and Asija [4], available phosphorus content was determined with Olsen method of extraction with 0.5 NaHCO<sub>3</sub> (pH 8.5) Olsen et al. [5], and available potassium content was estimated with Ammonium acetate extraction method of Jackson [3].

Comment [Dr24]: Delete at and add from

### 2.4 Statistical Analysis

The statistical analysis was carried out by using the statistical package OPSTAT as per suggested by Gomez and Gomez [6]. Economical analysis was carried out by calculating the amount of material needed for one hectare, calculating annual cost of production by considering present market rate. The results have been interpreted on the basis of 'F' test value and critical difference (CD) was calculated at 5 % level of significance. The analysis of variance was calculated as follows:

Comment [Dr25]:

Comment [Dr26]: Of data

Table 2. Analysis of variance

Source of variation	Degree of freedom	Sum of squares	Mean sum of square	Variance ratio ("F" Value)
Replication (R)	(r-1)=2	Sr	RSS/(r-1) = Mr	Mr/Me
Treatments (t)	(t-1)=6	St	TrSS/(t-1) = Mt	Mt/Me
Error	(r-1)(t-1)=12	Se	Err.SS/(r-1)(t-1) = Me	

Where,

r = Number of replications

t = Number of treatments

Me = Mean sum of square due to error

df = Degree of freedom

The standard error of mean (SEm) and critical difference (CD) for comparing the mean of any two treatments were computed as follows:

$$SEm = (Me/r)^{1/2}$$

$$SE(d) = (2 Me/r)^{1/2}$$

$$CD = SE(d) \text{ "t" value at error degree of freedom.}$$

Comment [Dr27]: Write in correct form

### 3. RESULTS AND DISCUSSION

#### 3.1 Chemical properties of soil

Integrated use of inorganic and organic sources of nutrients had significant effect on soil pH, available N, P and K content (kg/ha) in soil as mentioned in Table-3.

##### 3.1.1 Soil pH

Soil reaction (pH) is the primary element which affects the availability of nutrients in soil. The pH range of 5.5-6.5 is optimum for availability of most of the nutrient elements for pepper crop. Maximum pH value (6.13) was observed in T<sub>5</sub> (75% RDF through inorganic fertilizer + 25% through FYM) while minimum in T<sub>2</sub> (100% RDF through inorganic fertilizers) i.e. 5.68. The pH in the plots treated with a combination of organic and inorganic nutrients moved slightly towards neutrality, indicating that the application of organic amendments had a positive impact on soil health Nandi et al. [7]. Similar findings were obtained by Jamir et al. [8].

##### 3.1.2 Available Nitrogen (kg/ha)

Maximum available nitrogen (269.97 kg/ha) was recorded in plots supplied with (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) treatment T<sub>7</sub> which might be due to the release of additional nitrogenous compounds into the soil due to the presence of vermicompost, which may have enhanced the soil's capacity for retaining nutrients Vijaya and Seethalakshmi [9]. Also these results are in close conformity with Atal [10] and Jamir et al. [8].

##### 3.1.3 Available Phosphorus (kg/ha)

As far as available phosphorus is concerned maximum available phosphorus (40.03 kg/ha) was recorded in T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). The availability of phosphate was improved by combining inorganic fertiliser with organic manure, such as vermicompost and farm yard manure, which was also observed by Vimera et al. [11]. Raturi et al. [12] and Sharma et al. [13] expressed the similar views.

##### 3.1.4 Available Potassium (kg/ha)

The data pertaining to available potassium content (292.33 kg/ha) was maximum in treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). The increase in potassium content might be due to application of organic manures such as vermicompost in bulk quantity and their subsequently slow mineralization process Chetri et al. [14]. Another possible reason might be due to increase in K availability caused by a change in the equilibrium in the soil from relatively exchangeable K to soluble K Prativa and Bhattarai [15]. The above findings are in accordance with Atal [10].

Comment [Dr28]: Delete K and add potassium

Comment [Dr29]: Delete K and add potassium

Comment [Dr30]: Delete K and add potassium

**Table 3. Effect of integrated Nutrient Management (INM) practices on chemical properties of soil**

Treatments	Soil pH	Available N (kg/ha)	Available P (kg/ha)	Available K (kg/ha)
T <sub>1</sub> : Control (without fertilizers)	6.06	217.18	16.07	123.88
T <sub>2</sub> : 100% RDF through inorganic fertilizers	5.68	261.13	38.20	245.93
T <sub>3</sub> : 100% Recommended dose of FYM	5.71	234.13	27.13	145.07
T <sub>4</sub> : 100% Recommended dose of vermicompost	6.10	252.13	31.07	257.53
T <sub>5</sub> : 75% RDF through inorganic fertilizer + 25% through FYM	6.13	249.07	24.03	190.87
T <sub>6</sub> : 75% RDF through inorganic fertilizer + 25% through vermicompost	5.86	259.27	34.07	179.93
T <sub>7</sub> : 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost.	5.94	269.97	40.03	292.33
SE(m) ±	8.68	3.10	3.09	2.60
CD at 5%	N/A	9.65	9.62	8.09

### 3.2 Plant growth attributes

Improvement in growth characters is considered to be pre-requisite to increased yield as showed in Table-3.

#### 3.2.1 Plant height (cm)

Height of the plant is one of the key variables impacting yield and harvest duration particularly for plants with an indeterminate growth habit. Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded a maximum plant height of 42.47 cm and 55.03 at 45 and 60 DAT respectively and 31.53 cm at 30 DAT of treatment T<sub>2</sub> (100% RDF through inorganic fertilizers). The increase in plant height may be attained due to the release of the fixed nitrogen, hence increasing the concentration and availability of nitrogen in the root zone. Nitrogen accelerates the synthesis of protein, plant growth and its development was obtained Fawzy et al. [16]. Similarly it was observed by Malik et al. [17], Lal and Kanaujia [18], Jamir et al. [8] and Shilpa et al. [19].

#### 3.2.2 Number of branches per plant

Branching in sweet pepper is a natural phenomenon but if allowed that usually induce more fruit production at the expense of size *i.e.* more fruits with more number of branches. The maximum number of branches per plant (3.10, 3.43 and 4.03) was produced at 30, 45 and 60 DAT respectively from T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). While minimum number of branches per plant (1.67, 1.80 and 2.27) was produced at 30, 45 and 60 DAT respectively from T<sub>1</sub> (Control) treatment. The reason for the enhancement of branches production might be due to the direct effect of higher amount of inorganic nitrogen, which is a component of protein and chlorophyll molecules which might have increased the foliage of the plants and thus improved photosynthesis Kumar and Dhar [20]. Alike results was observed by Malik et al. [17], Shilpa et al. [19].

#### 3.2.3 Number of leaves per plant

Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded a maximum number of leaves *i.e.* 31.50, 32.60 and 35.03 at 30, 45 and 60 DAT respectively and was found significantly superior to all other treatments. Maximum number of leaves might be due to fact that application of NPK and vermicompost provided adequate Nitrogen which is associated with high photo synthetic activity and vigorous vegetative growth Pariari

Comment [Dr31]: Delete

Comment [Dr32]: Plant

Comment [Dr33]: also

Comment [Dr34]: Delete

and Khan [21]. The results of present investigation in terms of vegetative growth are also in concordance with the findings reported earlier by Lal and Kanaujia [18] and Jamir et al. [8].

UNDER PEER REVIEW

**Table4. Effect of Integrated Nutrient Management (INM) practices in growth parameters of sweet peppers**

Treatments	Plant height (cm)			Number of branches			Number of leaves			Plant spread (cm)		Days to 50% flowering
	30 DAT	45 DAT	60 DAT	30 DAT	45 DAT	60 DAT	30 DAT	45 DAT	60 DAT	45 DAT	60 DAT	
T <sub>1</sub> : Control (without fertilizers)	25.27	37.43	48.17	1.67	1.80	2.27	22.40	25.77	26.70	22.47	29.70	45.01
T <sub>2</sub> : 100% RDF through inorganic fertilizers	31.53	41.53	54.53	2.97	3.27	3.87	29.83	31.40	33.17	27.57	33.97	38.67
T <sub>3</sub> : 100% Recommended Dose of FYM	27.10	37.98	52.30	1.77	2.27	3.23	24.03	26.90	27.50	23.87	32.23	43.97
T <sub>4</sub> : 100% Recommended Dose of vermicompost	28.50	40.53	49.20	2.53	2.97	2.80	23.73	27.70	28.60	26.43	30.90	41.03
T <sub>5</sub> : 75% RDF through inorganic fertilizer + 25% through FYM	27.53	39.50	54.01	2.40	2.83	3.30	28.93	30.30	31.40	27.20	33.07	40.33
T <sub>6</sub> : 75% RDF through inorganic fertilizer + 25% through vermicompost	28.77	41.07	54.07	2.70	3.00	3.37	27.50	30.80	32.01	27.50	33.67	38.96
T <sub>7</sub> : 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost	30.33	42.47	55.03	3.10	3.43	4.03	31.50	32.60	35.03	27.60	34.50	36.01
SE(m) ±	0.45	0.54	0.69	0.04	0.04	0.04	2.00	1.40	1.59	1.14	0.92	1.53
CD at 5%	1.42	1.69	2.15	0.14	0.13	0.13	6.24	4.36	4.95	3.54	2.85	4.76

### 3.2.4 Plant spread (cm)

Plant spread is an important growth parameter which consequently affect yield per plant. Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded maximum plant spread (27.60 and 34.50 cm) at 45 and 60 DAT respectively. The increase in plant spread was attributed due to increase in fertilizer and organic manure application, which may increase amount of nutrients like nitrogen, phosphorus and potassium in plants, increasing the formation of plant metabolites that supported the development of plant tissues Malik et al. [17]. Similar result was also obtained by Raturi et al. [12].

### 3.2.5 Days to 50 per cent flowering

Early flowering results in early fruit development, which benefits in timing the crop. The number of days for inducing flowering varied from 36 to 45 days. The minimum day (36.01) for appearance of 50 per cent flowering was observed in treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). It seems that with increase in the levels of nutrient application the number of days taken for 50 per cent flowering showed decreasing trend. Another aspect that could contribute to nutritional integration that results in earlier flowering is quicker photosynthesis and better translocation of photosynthates towards the flower bud initiation Naidu et al. [22] and Prativa and Bhattarai [15].

## 3.3 Yield attributes

Integrated application of chemical fertilizers and organic manures increased yield and yield attributing characters of sweet pepper (Table-4).

### 3.3.1 Fruit length (cm)

Fruit length is also an important yield characteristic that influences fruit size as well as yield per hectare. The treatment module T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded maximum fruit length (8.03 cm) whereas the treatment T<sub>1</sub> (Control) recorded minimum fruit length (5.67 cm). Sufficient quantity of the fertilizers *i.e.* nitrogen, phosphorus and specially potassium, fulfilled the need of plants to attain more vigour, flowering and fruit development which produced fruits of high quality and significant size (length) Dubey et al. [23]. Similar results were obtained by Malik et al. [17], Chetri et al. [14] and Lal and Kanaujia [18].

Comment [Dr35]: Delete

Comment [Dr36]: Delete

### 3.3.2 Fruit diameter (cm)

Fruit diameter is an important yield character which affects fruit size and consequently yield. Analysis of the data shows that the interaction effect of integrated nutrient management practices on sweet pepper recorded maximum fruit diameter (6.73 cm) in T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost). It may be due to improved soil physical, chemical and biological properties and addition of nutrients through vermicompost and chemical fertilizers which in turn enhanced growth and led to the production of blocky fruits in the sweet pepper Lal and Kanaujia [18]. Similar are the findings of Malik et al. [17], Dubey et al. [23], Ngupok [24] and Raturi et al. [12].

### 3.3.3 Number of flowers per plant

Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded a maximum number of flowers *i.e.* 14.47 and was found significantly superior to all other treatments. This may be due to translocation of nutrients to the aerial parts of the plant. Vermicompost contains (3% N), (1% P<sub>2</sub>O<sub>5</sub>) and (1.5% K<sub>2</sub>O). The solubilization of plant nutrients caused by the addition of vermicompost leads to increase NPK uptake and resulted in maximum number of flowers per plant in sweet pepper Shiva et al. [25]. Alike results was obtained by Bhattarai et al. [26], Chetri et al. [14] and Jamir et al. [8].

### 3.3.4 Number of fruits per plant

This is one of the most important characters contributing directly towards yield. The data on different levels of nutrients revealed that T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) produced maximum number of fruits per plant (13.13) and minimum number of fruits per plant (8.93) was recorded in T<sub>1</sub> (Control). The increase in number of fruits per plant as results of integrated application of organic and inorganic fertilizers may be due to higher organic sources, higher organic matter build up, balanced C:N ratio and availability of nutrients Malik et al. [17]. Superiority of vermicompost over other organic sources could be attributed to its nutritional richness which results in better growth, more number of fruits when used in combination with organic

or inorganic sources Lal and Kanaujia [18]. Similar observations were also made by other workers like Chetri et al. [14], Ngupok [24].

### 3.3.5 Average fruit weight (g)

Fruit weight is an important yield parameter for obtaining maximum yield per hectare. The treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) which produced highest fruit length and fruit diameter, also observed maximum fruit weight 72.07 g followed by T<sub>6</sub> (75% RDF through inorganic fertilizer + 25% through vermicompost) which is 69.30 g. It appears from the findings of Suthar [27], that supply of nutrients from conjoint application of organic and inorganic sources *i.e.* vermicompost and chemical fertilizer improved the partitioning of photo-assimilates from source to sink (leaf to fruit) thereby increased fruit weight. Similar results have also been reported by Chetri et al. [14], Lal and Kanaujia [18] and Raturi et al. [12].

### 3.3.6 Fruit yield per plant (kg)

The treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) which produced highest fruit yield per plant *i.e.* 1.01 kg whereas the lowest fruit yield per plant was observed in control treatment T<sub>1</sub> (Control) *i.e.* 0.46 kg. This might be due to continuous availability of more nutrients in higher amount and better utilization by plants Chetri et al. [14]. These results are supported by Bhattarai et al. [26] and Raturi et al. [12].

### 3.3.7 Fruit yield per plot (kg)

Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded highest fruit yield per plot of 19.21 kg/plot which was found significantly superior to other treatment combinations. The increased fruit yield with conjoint application of organic manures and inorganic fertilizers may be due to large uptake and effective utilization of nutrients by plants Dass et al. [28]. These results are in line with the findings of Bhattarai et al. [26], Pariari and Khan [21], Dubey et al. [23], Shilpa et al. [19] and Sharma et al. [13].

### 3.3.8 Fruit yield per hectare (q)

The main objective of cultivation is to have maximum yield for better returns. Among the treatments, treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded highest fruit yield per hectare of 350.34 q/ha which was found significantly superior to other treatment combinations. While the control treatment T<sub>1</sub> having the lowest yield attributes had attained the lowest fruit yield of 165.21 q/ha. It is seen that the treatment having the maximum number of fruits per plant, fruit length, fruit diameter and average fruit weight had attained the highest fruit yield. Similar are the findings of Bhattarai et al. [26], Dubey et al. [23] and Sharma et al. [13].

### 3.3.9 Number of pickings

The maximum number of picking was observed in treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) *i.e.* 4.33 which was statistically at par with treatment T<sub>5</sub> (75% RDF through inorganic fertilizer + 25% through FYM) which is 4.01 and T<sub>2</sub> (100% RDF through inorganic fertilizers) which is 4.00. While the minimum number of picking was done in treatment T<sub>1</sub> (Control) which is 2.33.

**Comment [Dr37]:** Delete Superior and add higher

**Comment [Dr38]:** Plant growth coupled with yield contributing parameters like ....

**Table 5: Effect of integrated nutrient management practices in yield parameters of sweet pepper**

Treatments	Fruit length (cm)	Fruit diameter (cm)	No. of flowers per plant	No. of fruits per plant	Average fruit weight (g)	Number of pickings	Fruit yield per plant (kg)	Fruit yield per plot (kg)	Fruit yield per hectare (q)
T <sub>1</sub> : Control (without fertilizers)	5.67	5.03	9.53	8.93	49.87	2.33	0.46	9.12	165.21
T <sub>2</sub> : 100% RDF through inorganic fertilizers	6.90	5.60	11.37	10.63	62.38	4.00	0.68	13.36	246.23
T <sub>3</sub> : 100% Recommended Dose of FYM	6.57	5.07	10.87	9.97	51.22	2.67	0.52	10.34	189.23
T <sub>4</sub> :100% Recommended Dose of vermicompost	6.67	5.40	11.07	10.23	55.98	3.00	0.56	11.62	213.61
T <sub>5</sub> : 75% RDF through inorganic fertilizer + 25% through FYM	7.40	6.13	13.53	11.40	64.62	4.01	0.76	14.89	274.69
T <sub>6</sub> : 75% RDF through inorganic fertilizer + 25% through vermicompost	7.43	6.53	12.73	11.97	69.30	3.67	0.85	16.71	308.12
T <sub>7</sub> : 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost.	8.03	6.73	14.47	13.13	72.07	4.33	1.01	19.21	350.34
SE (±)	0.36	0.36	0.59	0.56	1.40	0.43	0.03	0.65	12.15
CD <sub>0.05</sub>	1.13	1.13	1.83	1.74	4.37	1.34	0.10	2.04	37.85

**Comment [Dr39]:** Fruit yield per plot can be deleted as already data has been converted on per hectare basis

### 3.4 Economics

#### 3.4.1 Cost of cultivation (₹/ha)

A perusal of data revealed that highest cost of production ₹ 143310 was incurred in treatment T<sub>4</sub> (100% Recommended Dose of FYM) because of the higher cost of vermicompost, whereas lowest (₹ 93310) was observed in treatment T<sub>1</sub> (Control).

#### 3.4.2 Gross return (₹/ha)

The maximum gross return per hectare amounting to ₹ 595578 was obtained in treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) followed by T<sub>6</sub> (75% RDF through inorganic fertilizer + 25% through vermicompost) which is ₹ 523804 and T<sub>5</sub> (75% RDF through inorganic fertilizer + 25% through FYM) which is ₹466973. On the other hand, the minimum gross return (₹ 280857) was recorded for treatment T<sub>1</sub> (Control).

#### 3.4.3 Net return (₹/ha)

The highest net return (₹ 485173) was recorded in T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) followed by T<sub>6</sub> (75% RDF through inorganic fertilizer + 25% through vermicompost) i.e. ₹ 414851 and T<sub>5</sub> (75% RDF through inorganic fertilizer + 25% through FYM) i.e. ₹ 368020. Whereas, the lowest net return (₹ 187546) was observed in T<sub>1</sub> (Control).

#### 3.4.4 B: C ratio

In overall, maximum B: C ratio (1:5.39) was recorded in T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) followed by B: C ratio (1:4.80) in T<sub>6</sub> (75% RDF through inorganic fertilizer + 25% through vermicompost) and T<sub>5</sub> (75% RDF through inorganic fertilizer + 25% through FYM) which is 1:4.71. Whereas minimum B: C ratio (1:3.00) was calculated in T<sub>1</sub> (Control), which may be due to the lower cost of production and comparable gross income in T<sub>1</sub> (Control). The reason for increased profit and highest Benefit: Cost ratio in treatment T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) is due to maximum fruit weight, healthy and better fruit size and higher net returns as compared to other treatments.

**Table 6: Effect of integrated Nutrient Management (INM) practices on economics of sweet pepper production**

Treatments	Cost of cultivation (₹ ha <sup>-1</sup> )	Gross Return (₹ ha <sup>-1</sup> )	Net Return (₹ ha <sup>-1</sup> )	Benefit: Cost ratio
T <sub>1</sub> : Control (without fertilizers)	93310	280857	187546	1:3.00
T <sub>2</sub> : 100% RDF through inorganic fertilizers	97499	418591	321091	1:4.29
T <sub>3</sub> : 100% Recommended Dose of FYM	103310	321691	218380	1:3.11
T <sub>4</sub> : 100% Recommended Dose of vermicompost	143310	363137	219826	1:2.53
T <sub>5</sub> : 75% RDF through inorganic fertilizer + 25% through FYM	98952	466973	368020	1:4.71
T <sub>6</sub> : 75% RDF through inorganic fertilizer + 25% through	108952	523804	414851	1:4.80

vermicompost				
T <sub>7</sub> : 50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost.	110404	595578	485173	1:5.39

Comment [Dr40]: Add SEM± and CD

#### 4. CONCLUSION

It can be concluded that, among different treatment, T<sub>7</sub> (50% RDF through inorganic fertilizer + 25% through FYM + 25% through vermicompost) recorded significantly higher values of growth characters and yield attributing characters followed by treatment T<sub>6</sub> (75% RDF through inorganic fertilizer + 25% through vermicompost). However, in terms of benefit: cost ratio, gross return, net return as well as the chemical properties of soil, treatment T<sub>7</sub> was found utmost. The highest cost of production ₹ 143310 was incurred in treatment T<sub>4</sub> (100% Recommended Dose of vermicompost) because of the higher cost of vermicompost, so it is suggested that cost of production in T<sub>4</sub> can be reduced if the farmers produce vermicompost by utilizing plenty of organic matter available at their own farm(s).

Comment [Dr41]: Delete

Comment [Dr42]: As well as

Comment [Dr43]: And also

Comment [Dr44]: Improved

Comment [Dr45]: Delete

#### References

1. Anonymous. Department of Agriculture and Farmers Welfare, horticulture Statistics; 2022. Available: <https://agricoop.nic.in/en/StatHortEst>
2. Hazra P, Chattopadhyah A, Karmakar K and Dutta S. Modern technology in vegetable production. New India publishing Agency, Pitam Pura, New Delhi 2011;10:413.
3. Jackson M.L. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi. 1967;183-192.
4. Subbiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soils. Current Science. 1956;25:259-260
5. Olsen SR, Cole CV, Watanable FS, Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. Circular of U.S. Department of Agriculture science. 1954;939:19.
6. Gomez KA and Gomez AA. Statistical Procedures for Agricultural Research. 2nd Edition, John Wiley and Sons, New York, 1984;680 p.
7. Nandi A, Patnaik SK, Prasad G, Dash SK, Mishra HN. Response of brinjal to integrated nutrient management in acidic sandy loam soil. All India coordinated vegetables improvement project. Department of horticulture, Orissa. University of Agriculture & Technology, Bhubaneswar; 2012.
8. Jamir T, Rajwade VB, Prasad VM, Lyngdoh C. Effect of organic manure and chemical fertilizers on growth and yield of sweet pepper (*Capsicum annum* L.) Hybrid Indam Bharath in shade net condition. *International Journal of Current Microbiology and Applied Sciences* 2017;6:1010-1019.
9. Vijaya KS, Seethalakshmi S. Response of eggplant (*Solanum melongena* L.) to integrated nutrient management amended soil. *International Journal of Scientific & Engineering Research* 2011;2:1-8.
10. Atal. Studies on effect of organic nutrient sources on bell pepper (*Capsicum annum* L.) production. M.Sc. Thesis. YSP University of horticulture and forestry, Palampur, H.P, India; 2017.
11. Vimera K, Kanaujia SP, Singh VB and Singh PK. Effect of integrated nutrient management on growth and yield of king chilli under foothill condition of Nagaland. *Journal of the Indian Society of Soil Science* 2012;60:45-49.
12. Raturi HC, Uppal GS, Singh SK and Kachwata DS. Effect of organic and inorganic nutrient sources on growth, yield and quality of bell pepper (*Capsicum annum* L.) grown under polyhouse condition. *Journal of Pharmacognosy and Phytochemistry* 2019;8:1788-1792.

13. Sharma N, Shukla YR, Singh K and Mehta DK. Soil fertility, Nutrient uptake and yield of bell pepper as influenced by conjoint application of organic and inorganic fertilizers. *Communications in Soil Science and Plant Analysis* 2020;51:1626-1640.
14. Chetri DA, Singh AK and Singh VB. Effect of integrated nutrient management on yield, quality and nutrient uptake by capsicum (*Capsicum annuum*) cv. California wonder. *Journal of Soils and Corporation* 2012;22:44-48.
15. Prativa KC and Bhattarai BP. Effect of integrated nutrient management on the growth yield and soil nutrient status in tomato. *Nepal Journal of Science and Technology* 2011;12:23-28.
16. Fawzy ZF, Behairy AG and Shehata SA. Effect of Potassium Fertilizer on Growth and Yield of Sweet pepper plants (*Capsicum annuum* L.). *Journal of Agriculture Reserach* 2010;2:599-610.
17. Malik AA, Chattoo MA, Sheemar G and Rashid R. Growth, yield and fruit quality of sweet pepper hybrid SH-SP- (*Capsicum annuum* L.) as affected by integration of inorganic fertilizers and organic manures (FYM). *Journal of Agricultural Technology* 2011;7:1037-1048.
18. Lal S. and Kanaujia SP. Integrated nutrient management in capsicum under low cost polyhouse condition. *Annals of Horticulture* 2013;6:170-177.
19. Shilpa AK, Sharma, Thakur S and Sharma K. Integrated Nutrient Management of Peppers (*Capsicum annuum* L.) in 21st Century - A Review. *International Journal Current Microbiology Applied Science* 2017;6:227-237.
20. Kumar A and Dhar S. Evaluation of organic and inorganic sources of nutrients in maize (*Zea mays*) and their residual effect on wheat (*Triticum aestivum*) under different fertility levels. *Indian Journal of Agriculture Science* 2010;80:364-71.
21. Pariari A and Khan S. Integrated nutrient management of chilli (*Capsicum annuum* L.) in Gangetic alluvial plains. *Journal of Crop and Weed* 2013;9:128-130.
22. Naidu AK, Kushwah SS, Mehta AK and Jain PK. Study of organic, inorganic and biofertilizers in relation to growth and yield of tomato. *Jawaharlal Nehru Krishi Vishavidalay Research Journal* 2002;35:36-37.
23. Dubey AK, Devi S, Pranjali SR, Yogesh K, Ajay KV and Sandip KC. Effect of NPK on plant growth, yield and quality of capsicum (*Capsicum annuum* L.) under shade net condition. *International Journal of Current Microbiology and Applied Sciences* 2017;6:1085-1091.
24. Ngupok O, Warade SD, Das S, and Raghavan M. Effect of NPK on Growth, Yield and Quality of Hybrid Capsicum (*Capsicum annuum* L. var. *grossum*) under Protected Condition. *International Journal of Chemical Studies* 2018;6:451-453.
25. Shiva KN, Srinivasan V, Zachariah TJ and Leela NK. Integrated nutrient management on growth, yield and quality of paprika alike chillies (*Capsicum annuum* L.) *Journal of Spices and Aromatic Crops* 2015;24:92-97.
26. Bhattarai DR, Poudyal KP, Pokhrel S. Effect of *Azotobacter* and nitrogen levels on fruit yield and quality of bell pepper. *Nepal Journal of Science and Technology* 2011;12:29-34.
27. Suthar S. Impact of vermicompost and composted farmyard manure on growth and yield of garlic (*Allium sativum* L.) field crop. *International Journal of Plant Production* 2009;3:27-38.
28. Dass A, Lenka NK, Patnaik US and Sudhishri S. Integrated nutrient management for production, economics and soil improvement in winter vegetables. *International Journal of Vegetable Science* 2008;14:104-20.

UNDER PEER REVIEW