

## **Bactericidal and bacteriostatic effects of four different parts of 2 plants *Moringa* extracts against the bacteria *Vibrio cholerae* and *Salmonella typhi*, and some phytochemical properties**

### **ABSTRACT**

**Background and Objectives:** Typhoid fever and cholera are two major health challenges in many developing countries. The use of extracts of the plants *Moringa* for their treatment is often done without perfect knowledge of the antimicrobial properties of each part of plant. This study aims to evaluate the bactericidal and bacteriostatic effects of extracts of *M. oleifera* and *M. stenopetala* against the bacteria *S. typhi* and *V. cholera* which cause typhoid fever and cholera respectively.

**Materials and Methods:** Four different parts (leaves, seeds, stem, and roots) of each plant species were dried and then ground. The minimum inhibitory concentrations (MICs) and minimum bactericidal concentrations (MBCs) of the extracts of each part of the plants against the considered bacteria were determined by dilution method. Their phytochemical profiles were determined by chromatographic and mass spectral analyses.

**Results:** It has been noted that MICs and MBCs varied from one part to another of the same *Moringa* species, and from one bacterium to another for the same extract. With *M. oleifera*, the lowest MBC (0.078 mg/mL) against *S. typhi* and *V. cholerae* came from seeds and roots extracts, respectively. With *M. stenopetala*, the lowest MBCs against *S. typhi* (0.156 mg/mL) and *V. cholerae* (0.312 mg/mL) were from seeds extracts. Extracts of different parts of *M. oleifera* have bactericidal effects against *S. typhi*, and bacteriostatic effects against *V. cholerae*. Extracts from all considered parts of *M. stenopetala* have bacteriostatic effects against both bacteria. Although some phytochemicals were found in both *Moringa*, others were specific to one species or to a given part.

**Conclusion:** In case of co-infection by both bacteria, *M. oleifera* roots extracts could be the most recommended in the treatment because of their lowest MIC and MBC. With *M. stenopetala*, roots and seeds extracts could be the most recommended because of their relatively lower MIC and MBC, respectively.

**Key words:** plants *M. oleifera* and *M. stenopetala*, plant part's extract, antimicrobial activities, *S. typhi* and *V. cholerae*, MICs and MBCs, phytochemicals

## 1. INTRODUCTION

Many people on earth suffer from water-borne diseases due to the lack of appropriate drinking water. Among these diseases, typhoid and cholera are two major health challenges in many African and Asian countries [1, 2]. Typhoid is caused by the bacteria named *Salmonella typhi* and *S. paratyphi*, while cholera is caused by *Vibrio cholera* [2, 3]. The problem is that there is an increasing resistance to antibiotics by the 3 bacteria species above. The 3 bacteria *S. typhi*, *S. paratyphi* and *V. cholerae* essentially live in water and humans often become contaminated by drinking polluted water. In modern medicine, both diseases are treated by the consumption of the appropriate antibiotics. Due to the low socio-economic level and the often high cost of antibiotics, people in many developing countries use several local plants to treat these diseases.

These plants are supposed to contain certain substances or chemical elements which would act against the microorganisms at the origin of these diseases. The synthesis of these substances or chemical elements would depend not only on the genetic properties of the plant, but also on the mineralogical and petrographic properties of the crop soils, as well as meteorological parameters such as temperature, precipitation, insolation, light intensity and degree of humidity [4]. The nature, properties and concentrations of these chemical elements would vary depending on the part of the plant [4, 5].

Medicinal plants include among others *Moringa oleifera* Lam., and *M. stenopetala* (Baker f.) Cufod. Both species belong to the family of Moringaceae, order of Capparidales, class of Dicotyledonae, subphylum of Angiospermae, phylum of Spermatophyta, kingdom of Plantae, domain of Eukaryota [6-8]. The parts of each plants above used to treat each disease varies from one region of the world to another, as well as from one traditional healer (doctor) to another. Their leaves, flowers, seeds, stem barks or roots are often used to treat typhoid fever and cholera, and the set-up include decoction, infusion, maceration [7, 9]. A variety of liquid is often used by traditional healers to prepare plant extract. Water and alcohol are regularly used [10].

Effective treatment of an infectious disease depends on the mode of action of the antimicrobial agent against the infectious agent. The antimicrobial agent can use bacteriostatic or bactericidal mechanism to fight against the pathogen. Bacteriostatic antimicrobial agent limit just the growth of bacteria. Its actions are reversible. In contrast, bactericidal antimicrobials kill their target bacteria [11, 12].

In modern medicine, the decision to use a bacteriostatic or bactericidal drug depends on the type of infection and the immune status of the patient. In traditional herbal medicine, there is in

most cases insufficient knowledge of the mode of action of the extract from each part of the plant concerned. The perfect knowledge of the mechanism of action of the extract of a precise part of a medicinal plant would not only quickly eradicate the microorganism responsible of the disease, but would also allow the patient to save money while shortening the duration of treatment. It would also make it possible to avoid sensitization of the microorganism which often leads to resistance of the germ concerned to antimicrobial agents.

The use of *Moringa* extracts to fight against *Salmonella* and *Vibrio* bacteria is often done without perfect knowledge of the antimicrobial properties of each plant part. Little is known about the differences in extract activities from one part of the plant to another to fight against the same bacterium, as well as from one bacterium to another for the extract of the same plant part. The present study aims to evaluate the bactericidal or bacteriostatic effect of extracts of several parts of *M. oleifera* and *M. stenopetala* to fight against the bacteria *S. typhi* and *V. cholera* which respectively cause typhoid fever and cholera in humans.

## **2. MATERIALS AND METHODS**

### **2.1 Harvesting the parts of the plant and the microorganisms used**

The plant species *Moringa oleifera* Lam. and *M. stenopetala* (Baker f.) Cufod. are cultivated in Yaoundé (Cameroon, Central Africa). The plant names have been checked with <http://www.theplantlist.org/tpl1.1/search?q=Moringa> web site (accessed on 28<sup>th</sup> May 2023). They were identified at the National Herbarium at Yaounde (Cameroon) and registered at No. 8241/SRF/Cam for *M. oleifera* and 67475/HNC for *M. stenopetala*. The parts of plant considered were the seeds, leaves, roots, and stem. Each of them was dried and then ground.

The microorganisms *V. cholerae* and *S. typhi* were isolated from the wells of Yaoundé (Cameroon, Central Africa) by the membrane filter method (cellulose ester membranes, porosity 0.45 µm). The agar culture media used were TCBS (Bio-Rad) and Wilson-Blair media (Oxoid) for isolation of *V. cholerae* and *S. typhi* respectively. Incubations were done at 37 °C for 24hrs. Each bacterial specie was subsequently identified using conventional biochemical tests [13, 14].

### **2.2 Preparation of *M. oleifera* and *M. stenopetala* powders**

The leaves, the bark of the roots and stems were cut up and dried in the shade, at laboratory temperature ( $25 \pm 2$  °C) for 30 days. The dried seeds were pulped, and the almond also dried at laboratory temperature ( $25 \pm 2$  °C) for 30 days. The different parts of the 2 chosen plants have been

dried, crushed and then ground. Their crushing was done in an autoclave-sterilized porcelain mortar around the flame of the Bunsen beak. The sterility of the powder was then checked by the negative search for the 2 bacteria considered.

To check the absence of *S. typhi*, 5 g of powder from each part of the plant were dissolved in 5 ml of sterile physiological water. After homogenization, 1 ml of this suspension was taken and diluted in 10 ml of sterile physiological water, and the whole was then filtered on a Millipore membrane (porosity 0.45  $\mu\text{m}$ ). This membrane was then deposited on the Wilson-Blaird agar culture medium contained in a Petri dish. Then the whole was incubated at 37° C for 24 hours. The absence of colony formed unit reflected the absence of contamination of the powder by this bacterial species. For the control of the absence of *V. cholerae*, the same protocol was used. The culture medium used was TCBS agar. The absence of colony formed unit reflected the absence of contamination of the powder by this bacterium.

### **2.3 Preparation of the methanoic plant extract**

The same protocol was used for each plant material. Thirty grams (30 g) of plant material, was extracted with methanol at room temperature for 24 hours (1 g of plant in 10 mL of methanol), repeated three times. After filtration, the solvent was removed under vacuum to furnish a crude extract [15].

### **2.4 Preparation of plant extracts for chromatographic analyses and mass spectral determination**

The method used was that described by other authors [16, 17]. From the crude methanoic extract prepared above of the different parts of the plant *M. oleifera* on one hand and *M. stenopetala* on the other hand, 5 mg of material were diluted in 5 mL of Methanol. Then the solution was filtered first through a 0.45  $\mu\text{m}$  porosity membrane, and secondly through a 0.22  $\mu\text{m}$  porosity membrane. Part of the filtrate was transferred to an HPLC vial for analysis.

Data was processed using Xcalibur software (Thermo Fisher Scientific Inc., MA, USA). The identification of all compounds described was carried out using the negative ionization mode.

Mass spectrum data were registered in a standard Excel format. These data included the exact mass of each compound. Each compound detected was characterized. Its potential raw formula was then calculated using Xcalibur software, which takes into account its relative

abundance, its molar masses and its retention time. The names of chemical compounds were determined by comparison with analytical standard or according to literature data [18-21].

## **2.5 Preparation of plant extracts for antimicrobial activities**

The protocol used was that described by Ashraf et al [22]. A stock solution of extract of 2.5 mg/mL concentration was prepared in a volume of 5 mL of 8.5 g/L NaCl sterile solution. The mixture obtained was homogenized in a vortex, and then filtered through a filter membrane (porosity 0.2  $\mu\text{m}$ ).

For antimicrobial activity, the standard antibiotic chosen was Gentamicin. The microbial inoculum was prepared from a pure 18-24 hours' culture on standard agar, and then adjusted at standard solution using a densitometer (density 0.5 on the Mc Farland scale) containing approximately  $1.5 \times 10^8$  CFU/mL, and diluted to 1/100 for the saline dilution test [23]. The minimum inhibitory concentrations (MICs) and minimum bactericidal concentrations (MBCs) were then determined.

## **2.6 Determination of the minimum inhibitory concentrations (MICs)**

The method of 96 wells plate was used and this has been described by others [12, 24]. A sterile Muller Hinton nutrient broth was used. From the bacterial suspension indicated above, bacterial inoculum containing approximately  $10^6$  CFU/mL was prepared by dilution. From this suspension, 100  $\mu\text{L}$  was inoculated into each well. The serial dilutions of extract ranging from 1.25 to 0.0045 mg/mL were then introduced in 96-well microplates. A sterility control well and a growth control well were also done. Plates were then incubated at 37 ° C for 18 to 24 hours. After incubation, 40  $\mu\text{L}$  of the colorless iodinitrotetrazolium chloride (INTC) solution were introduced into each of the wells of the plate. INTC reagent is an indicator of microbial growth. About 30 minutes after its addition, the appearance of a red color indicates the cells growth. The unstained well and which contained the lowest concentration of the extract corresponds to the MIC. This was done in triplicate. The antibiotic Gentamicin was used for positive control, and the negative control was at the absence of plants extract.

## **2.7 Determination of the Minimum Bactericidal Concentrations (MBCs)**

The protocol used was that indicated by other authors [24-26]. 50  $\mu\text{L}$  of extract taken from the wells corresponding to the MICs obtained were added. Also, 50 $\mu\text{L}$  of extract was taken from the wells that preceded those with MIC. The sterility checks consisted of 200  $\mu\text{L}$  of Mueller Hinton

broth. The positive control consisted of Gentamicin, treated like the extracts. The negative control was the medium without extract. Incubation was at 37°C for 18 - 24hrs. The CMB was the lowest concentration of extract with no visible microbial growth (around 99,9% of cells have been killed). The tests were done in triplicate.


## **2.8 Determination of the MBC/MIC ratios**

The ratio obtained aimed to conclude whether extracts tested were bacteriostatic or bactericidal. When this ratio was  $> 4$ , extract was considered bacteriostatic. When it was  $\leq 4$ , then extract was considered bactericidal [24, 26].

## **3. RESULTS**

### **3.1 Minimum inhibitory concentrations (MICs) and minimum bactericidal concentrations (MBCs) of each part of plants extracts against *S. typhi* and *V. cholerae***

For each of the 2 bacterial species; *S. typhi* and *V. cholerae*, the mean values of MICs and MBCs obtained with the extracts of each part of *M. oleifera* and *M. stenopetala* were calculated. They are presented in Fig. 1. For both bacterial species Gentamicin has been chosen as the reference antibiotic. It has a MIC of 1.562 mg/mL to fight against *S. typhi* and 3.130 mg/mL to fight against *V. cholerae*. With plants, it was noted that the MICs values of extracts parts of *M. oleifera* against *S. typhi* varied from 0.625 mg/mL to 0.019 mg/mL. The highest value was registered with leaves and stem extracts, while the lowest was recorded with seeds extract. The MIC value with roots extract was 0.078 mg/mL. Against *V. cholerae*, the MICs with extract from this plant species varied from 0.156 mg/mL to 0.005 mg/mL. The highest value was registered with leaves and stem extracts while the lowest value was recorded with roots extract. With the seeds extract the MIC against *V. cholerae* was 0.009 mg/mL (Fig. 1).



**Fig. 1.** Variation in MICs and MBCs of leaves, seeds, stem, and roots extracts of *M. oleifera* (**M. oleif**) and *M. stenopetala* (**M. steno**) as well as Gentamicin (**Genta**) against *S. typhi* and *V. cholerae*

When using *M. stenopetala* extracts, the MICs against *S. typhi* varied from 0.039 mg/mL to 0.004 mg/mL. The highest value was registered with leaves extract while the lowest value was recorded with roots extract. The MICs of stem and seeds extracts was 0.019 mg/mL. Against *V. cholerae*, MICs with *M. stenopetala* extracts varied from 0.078 mg/mL to 0.019 mg/mL. The highest MIC value was recorded with leaves extracts while the lowest value was obtained with stems and roots extracts. With seeds extracts, the MIC value was 0.039 mg/mL (Fig. 1).

For both bacterial species, Gentamicin chosen as the reference antibiotic has an MBC of 12.5 mg/mL against *S. typhi* and 6.25 mg/mL against *V. cholerae*. The MBCs when using *M. oleifera* extracts against *S. typhi* varied from 1.250 mg/mL to 0.078 mg/mL. The highest value was obtained with leaves and stem extracts and the lowest value with seeds extracts. With roots extracts, it was 0.312 mg/mL. In the presence of *V. cholerae*, the MBCs with *M. oleifera* extracts also varied from from 1.250 mg/mL to 0.078 mg/mL. The highest was registered with leaves and stem extracts while the lowest was obtained with roots extracts. With seeds extracts the MBC was 0.625 mg/mL (Fig. 1).

The MBCs values when using *M. stenopetala* extracts, to fight against *S. typhi* varied from 0.625 mg/mL to 1.156 mg/mL. The highest was obtained with leaves and stems extracts while the lowest was recorded with seeds extracts. With roots extracts, the MBC against *S. typhi* was 0.312 mg/mL (Fig. 1). When using *M. stenopetala* extracts, the MBCs in the presence of *V. cholerae* varied from 1.250 mg/mL to 0.312 mg/mL. The highest value was recorded with stems and roots extracts while the lowest value was registered with seeds extracts. With leaves extracts, the MBC obtained was 0.625 mg/mL (Fig. 1).

### **3.2 Effect of the extract of each part of *M. oleifera* and *M. stenopetala* against *S. typhi* and *V. cholerae***

From the values of the MBC/MIC ratios, the antibacterial properties of each extract from each *Moringa* species and on each microorganism considered were determined. These different properties are presented in Table 1. It was noted that extracts of each part from the two *Moringa* species are bacteriostatic against *V. cholerae*. Whereas against *S. typhi*, extract of each considered part of *M. oleifera* is bactericidal, and extracts from each part of *M. stenopetala* are bacteriostatic (Table 1). It should be noted that Gentamicin as our control is bactericidal against *V. cholerae* but bacteriostatic against *S. typhi* (Table 1).

**Table 1.** Effect of the extract of each part (leaves, seeds, stem and roots) of *M. oleifera* and *M. stenopetala* against *S. typhi* and *V. cholerae*

Part of the plant species and standard antibiotic used		Bacteria species and extract's effect	
Plant part	Plant species	<i>S. typhi</i>	<i>V. cholerae</i>
Leaves	<i>M. oleifera</i>	Bactericidal	Bacteriostatic
	<i>M. stenopetala</i>	Bacteriostatic	Bacteriostatic
Stem	<i>M. oleifera</i>	Bactericidal	Bacteriostatic
	<i>M. stenopetala</i>	Bacteriostatic	Bacteriostatic
Roots	<i>M. oleifera</i>	Bactericidal	Bacteriostatic
	<i>M. stenopetala</i>	Bacteriostatic	Bacteriostatic
Seeds	<i>M. oleifera</i>	Bactericidal	Bacteriostatic
	<i>M. stenopetala</i>	Bacteriostatic	Bacteriostatic
	Gentamicin	Bacteriostatic	Bactericidal

### 3.3 Comparison amongst the MICs and MBCs values for each bacteria specie considered for the 4 parts of *M. oleifera* and *M. stenopetala*

A comparison of the MICs and MBCs among the extract from the 4 parts of each the *Moringa* species against each bacterial species was made. The result is presented in Table 2. It can be noted that MICs of extract from different parts of *M. oleifera* significantly differed ( $P < 0.05$ ) against *V. cholerae* and *S. typhi*. However, MICs of extract from different parts of *M. stenopetala* do not significantly differ from one part of the plant to another ( $P > 0.05$ ) against both bacteria species. MBCs of extract from different parts the *Moringa* species do not also significantly differ against the 2 bacteria species (Table 2).

**Table 2.** “P” value of the comparison for each considered bacteria species, of the MICs and MBCs values of the 4 parts of the plant

Plant species and cells species		Extracts' concentrations	
Plant species	Bacterial species	MIC	MBC
<i>M. oleifera</i>	<i>V. cholerae</i>	<b>P= 0.025*</b>	P=0.392
	<i>S. typhi</i>	<b>P= 0.013*</b>	P= 0.392
<i>M. stenopetala</i>	<i>V. cholerae</i>	P= 0.435	P= 0.392
	<i>S. typhi</i>	P= 0.835	P=0.392

\* : Significant difference ( $P < 0.05$ ) ; ddl=3

### 3.4 Phytochemical profiles of extracts

The chemical profiles of each extract have been established by mass spectra in negative ionization mode. The chemical compounds identified from extracts of leaves, seeds, roots and stems of *M. oleifera* and *M. stenopetala* are presented in Table 3.

Concerning *M. oleifera*, some compounds have been identified in several parts of the plant. Saccharose (glycoside) and Glucomoringinin (Glucosinolate) for example, were identified in leaves, seeds, stems and roots extracts. The O-acetylshanzhiside methyl ester was found in leaves, stems and roots extracts. The Glucotropaeolin was identified in stems and roots extracts. On the other hand, some compounds have only been found in certain specific parts of the plant. This is for example Neochlorogenic acid, Isoquercetin, Kaempferol-3-O-glucoside and Kaempferol acetyl glucoside which were found only in leaves extracts. It is the same for Quinic acid and Catechin which were found only in stems extracts (Table 3).

Compounds identified are representative of 7 different classes (Table 3): heteroside, glucosinolate, phenolic acid, iridoid, flavonoid, flavonoid heteroside and carboxylic acid. The heteroside and glucosinolate identified were found in the 4 parts of the plants considered. The phenolic acid identified was found only in leaves. The iridoid identified was found in leaves, stems and roots extracts. The flavonoids identified were found in leaves and stems. The flavonoids

heterosides identified were found only in leaves. The carboxylic acid identified was found only in stems (Table 3).

As in the case of *M. oleifera*, the extracts of *M. stenopetala* also show the presence of same compounds in different part of the plant. This is the case for Sucrose and Glucomoringin. Others are present only in certain parts of the plant, like Quercetin-3-O-rutinoside and Neochlorogenic acid which are only present in the leaves. Glucoconringiin-2-hydroxy-2-méthylpropylglucosinolate is only found in the seeds. Acetyl-4-(L-rhamnopyranosyloxy)benzyl glucosinolate and 1,2-Di-O-galloyl-6-O-cinnamoyl- $\beta$ -D-glucose are only in the stem. Glucose is present only in the roots. Glucotropaeolin is present only in the stem and roots. Citric acid and O-acetylshanzhiside methyl ester are present only in the leaves, stem and roots. All compounds identified are of 6 different classes: heteroside, glucosinolate, phenolic acid, iridoid, flavonoid and flavonoid-heteroside (Table 3).

Some main differences in the chemical profiles of the extracts of the 2 *Moringa* species have been highlighted. In leaves it is the presence of Isoquercetin (Flavonoid), Kaempferol-3-O-glucoside and Kaempferol-acetyl glucoside (Flavonoid heteroside) only in *M. oleifera*, and the presence of Quercetin-3-O-rutinoside (Flavonoid heteroside), only in *M. stenopetala*. In stems extracts, it is the presence of Quinic acid (Carboxylic acid) and Catechim (Flavonoid) only in *M. oleifera* and that of Acetyl-4-( $\alpha$ -L-rhamnopyranosyloxy)benzyl glucosinolate (Glucosinolate), 1,2-Di-O-galloyl-6-O-cinnamoyl- $\beta$ -D-glucose and Citric acid (organic acid) only in *M. stenopetala*. In roots, it is the presence of Glucose (heteroside) and Citric acid (organic acid) only in *M. stenopetala*. In seeds Glucoconringiin-2-hydroxy-2-méthylpropylglucosinolate has been found in *M. stenopetala* and not in *M. oleifera* (Table 3).

**Table 3.** Phytochemical profile of MeOH extracts of each part considered of *M. oleifera* and *M. stenopetala* (comparison with analytical standard and according literature data [18-21])

Part of the plant considered	Plant species and chemical compound identified									
	<i>M. oleifera</i>					<i>M. stenopetala</i>				
	Rt (min)	M-H <sub>exp</sub> (m/z)	Compound formula	Compound name	Class of molecule	Rt (min)	M-H <sub>exp</sub> (m/z)	Compound formula	Compound name	Class of molecule
Leaves	3.99	341.1086	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside	3.96	341.1086	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside
	8.99	570.0956	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate	6.84	191.0189	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	Citric acid	Organic acid
	11.34	353.0875	C <sub>16</sub> H <sub>18</sub> O <sub>9</sub>	Neochlorogenic acid	Phenolic acid	9.01	570.0956	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate
	14.13	447.1508	C <sub>19</sub> H <sub>28</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid	11.32	353.0875	C <sub>16</sub> H <sub>18</sub> O <sub>9</sub>	Neochlorogenic acid	Phenolic acid
	18.55	463.0883	C <sub>21</sub> H <sub>20</sub> O <sub>12</sub>	Isoquercetin	Flavonoid	14.11	447.1508	C <sub>19</sub> H <sub>28</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid
	22.12	447.0931	C <sub>21</sub> H <sub>20</sub> O <sub>11</sub>	Kaempferol-3-O-glucoside	Flavonoid heteroside	16.98	609.1465	C <sub>27</sub> H <sub>30</sub> O <sub>16</sub>	Quercetin-3-O-rutinoside	Flavonoid heteroside
	26.65	489.1037	C <sub>23</sub> H <sub>22</sub> O <sub>12</sub>	Kaempferol acetyl glucoside	Flavonoid heteroside	/	/	/	/	/
Seeds	4.04	341.1093	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside	3.98	341.1086	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside
	8.78	570.0966	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate	7.29	390.0533	C <sub>11</sub> H <sub>21</sub> NO <sub>10</sub> S <sub>2</sub>	Glucoconringiin-2-hydroxy-2-méthylpropyl-glucosinolate	Glucosinolate
	/	/	/	/	/	8.91	570.0957	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate
	3.85	191.0547	C <sub>7</sub> H <sub>12</sub> O <sub>6</sub>	Quinic acid	Carboxylic acid	3.98	341.1087	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside

<b>Stems</b>	4.00	341.1046	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside	6.86	191.0185	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	Citric acid	Organic acid
	9.03	570.0955	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate	8.98	570.0959	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate
	12.89	408.0426	C <sub>14</sub> H <sub>28</sub> NO <sub>9</sub> S <sub>2</sub>	Glucotropaeolin	Glucosinolate	11.66	612.1067	C <sub>22</sub> H <sub>31</sub> NO <sub>15</sub> S <sub>2</sub>	Acetyl-4-( $\alpha$ -L-rhamnopyranosylox) benzyl glucosinolate	Glucosinolate
	13.09	289.0718	C <sub>15</sub> H <sub>14</sub> O <sub>6</sub>	Catechin	Flavonoid	12.26	612.1066	C <sub>29</sub> H <sub>25</sub> O <sub>15</sub>	1,2-Di-O-galloyl-6-O-cinnamoyl- $\beta$ -D-glucose	Acyl glucoside
	14.11	447.1506	C <sub>19</sub> H <sub>28</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid	12.80	408.0428	C <sub>14</sub> H <sub>28</sub> NO <sub>9</sub> S <sub>2</sub>	Glucotropaeolin	Glucosinolate
	/	/	/	/	/	14.12	447.1509	C <sub>19</sub> H <sub>28</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid
<b>Roots</b>	3.99	341.1087	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside	3.23	179.0554	C <sub>6</sub> H <sub>12</sub> O <sub>6</sub>	Glucose	Heteroside
	9.04	570.0956	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate	3.98	341.1087	C <sub>12</sub> H <sub>22</sub> O <sub>11</sub>	Saccharose	Heteroside
	12.85	408.0426	C <sub>14</sub> H <sub>28</sub> NO <sub>9</sub> S <sub>2</sub>	Glucotropaeolin	Glucosinolate	6.85	191.0186	C <sub>6</sub> H <sub>8</sub> O <sub>7</sub>	Citric acid	Organic acid
	14.12	447.1507	C <sub>19</sub> H <sub>19</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid	9.02	570.0959	C <sub>20</sub> H <sub>29</sub> NO <sub>14</sub> S <sub>2</sub>	Glucomoringinin	Glucosinolate
	/	/	/	/	/	12.77	408.0428	C <sub>14</sub> H <sub>28</sub> NO <sub>9</sub> S <sub>2</sub>	Glucotropaeolin	Glucosinolate
	/	/	/	/	/	14.11	447.1509	C <sub>19</sub> H <sub>28</sub> O <sub>12</sub>	O-acetylshanzhiside methyl ester	Iridoid

#### 4. DISCUSSION

This study shows that some compounds in the extracts of *Moringa* species, are found in all parts of the plant, while others are only localized in some specific parts (Tables 3). These qualitative and probably quantitative variations (because of the variations in retention times), would be at the origin of the apparent variations of the MICs and MBCs from one part of the plant species to another.

Four compounds identified in *M. stenopetala* extracts were absent in *M. oleifera*. They are Quercetin-3-O-rutinoside, Citric acid, Acetyl-4-( $\alpha$ -L-rhamnopyranosyloxy)benzyl glucosinolate, and glucose. Five compounds identified in *M. oleifera* extracts, were absent in the *M. stenopetala* extracts. They are Isoquercetin, Kaempferol-3-O glucoside, Kaempferol acetyl glucoside, Quinic acid and Catechim. Each of the chemical compound has its own properties towards microorganisms.

Quinic acid which is a carboxylic acid was found in *M. oleifera* extract, unlike extracts from *M. stenopetala* (Tables 3). Carboxylic acids are known to facilitate the destruction of microbial membranes and cell walls. They act through the inhibition of peptidoglycan synthesis [27]. They can also inhibit bacterial growth through the inhibition of DNA/RNA replication, for example the inhibitory activity against DNA gyrase (an essential enzyme that controls the topological state of DNA replication) [28, 29]. Carboxylic acids also target horizontal gene transfer inhibition, quorum sensing inhibition, inhibition of efflux pumps and inhibitors of metabolic routes [29].

Isoquercetin was isolated only in *M. oleifera* and not in *M. stenopetala* extracts (Table 3). This flavonoid is a naturally occurring polyphenol that has antioxidant, anti-proliferative, anti-carcinogenic, neuroprotection, anti-inflammatory, antibacterial and antiviral properties [30].

The flavonoids Kaempferol-3-O glucoside, Kaempferol acetyl glucoside and Catechim have been found only in *M. oleifera* and the phenolic acid Neochlorogenic acid has been found in leaves extracts of both *Moringa* species (Table 3). Phenolic products can also inhibit the synthesis of nucleic acids of both gram negative and gram-positive bacteria although this antibacterial effect varies with chain length of the molecules [31]. In other circumstances, they can form a complex with bacterial cell wall thus inhibiting the microbial growth [32]. The anti-microbial mechanisms of flavonoids and phenolic compounds include the inhibition of cell envelop synthesis, nucleic acid synthesis, bacterial motility, electron transport chain and ATP synthesis, bacterial toxins, bacterial quorum sensing, biofilm formation, bacterial enzyme-dependent virulence, bacterial efflux pumps, cytoplasmic membrane function causing marked increase in potassium loss from cells, the membrane disruption, among others [33-35].

Quercetin-3-O-rutinoside and Citric acid were found only in *M. stenopetala* extracts. They respectively destroy the normal functioning of the endoplasmic reticulum and increase the intracellular acidity leading to the destruction of the bacterial cell [36, 37].

Traditionally, non-ionized citric acid is considered more antimicrobial, presumably due to the combined effects of the molecule and the acidic environment in which it occurs [37]. It is believed that at low pH, citric acid in its un-charged and un-dissociated state can freely cross the microbial membrane. Once inside the cytoplasm, it dissociates into citric acid anions and protons leading to the acidification of the intracellular media, while causing functional and structural damage to the cell [38-40].

Some compounds as Glucomoringin, Neochlorogenic acid and O-acetylshanzhiside methyl ester have been found in the extracts of both *Moringa* species. Glucomoringin has been found in most parts of the plant studied. The antibiotic activity of this glucosinolate on *S. aureus* has been reported [21, 41]. Its bioactivation with myrosinate enzyme leads in the increasing power of the compound [41]. O-acetylshanzhiside methyl ester which is an iridoid has been detected in leaves, stems and roots extracts. Cytotoxic activities of iridoids have been indicated by many authors [42].

The differences in the resistance mechanisms to antimicrobial agents between *S. typhi* and *V. cholerae* could also be at the origin of the relative differences observed in the MICs and MBCs against the two bacteria species (Fig. 1), as well as the effect of the extract of each part of each *M. oleifera* and *M. stenopetala* against each bacteria species (Table 1).

It is indicated that *S. typhi* resists to antimicrobials by the following ways: inactivation of the antimicrobial agent and efflux or transport of the antimicrobial, modification of the antimicrobial target site and reduction of the permeability of the antimicrobial agent [43]. These resistance mechanisms can be plasmid mediated or chromosome mediated [44, 45].

Antimicrobial resistance in *V. cholerae* is mostly due to acquisition of resistance genes from closely or distantly related microbial species through horizontal gene transfer [46]. *V. cholerae* can also acquire resistance to antimicrobial compounds by any of the following seven mechanisms : i) modification of the antibiotic target site, ii) antibiotic target replacement, iii) antibiotic target protection, iv) inactivation of the antibiotic by hydrolysis or chemical modification, v) preventing access to the target site by changing membrane permeability, vi) actively exporting antibiotics from bacterial cell, and vii) resistance by absence of antibiotic target [47, 48]. Both, spontaneous mutation and horizontal gene transfer could contribute to all the resistance mechanisms [48].

In the present study, the difference in behavior between the 2 pathogens would also vary according to the concentration of each antimicrobial agent identified, although this concentration

was not determined in the extract of each part of the plant. The concentration of the antimicrobial agent is often said to be very closely related to the kinetics of its activity on the target bacterium [49, 50]. This could also partly explain the variations in the recorded MICs and MBCs, as well as the variation in the extract activity from one microorganism to another.

## 5. CONCLUSION

The MICs and MBCs of the *M. oleifera* and *M. stenopetala* extracts against *S. typhi* and *V. 16holera* relatively vary on one hand from one part of the plant to another for the same *Moringa* species, then from one bacterium species to the other for the same extract on the other hand. The leaves, seeds, stem, and roots extracts of *M. oleifera* have bactericidal effect against *S. typhi*, and bacteriostatic effect against *V. 16holera*. Extracts from all parts of *M. stenopetala* have bacteriostatic effects against each of the 2 bacterial species. These differences would be linked on one hand to the relative phytochemical differences amongst the parts of the 2 plant species, and on the other hand to the relative differences between the antimicrobial resistance mechanisms of the 2 bacteria. Although some phytochemicals were found in both *Moringa* species, others were specific to one species or to a given plants part. In case of co-infection by the 2 bacterial species, *M. oleifera* extracts could be the most recommended in the treatment, especially root extracts, because of their lowest MIC and MBC. With *M. stenopetala*, root and seed extracts could be the most recommended because of their relatively lower MIC and MBC, respectively. It seems important to explore in each region of the world, the antimicrobial properties of the extracts of different parts of each medicinal plant with respect to each microorganism or group of microorganisms which cause the diseases whose said extracts are used to treat the symptoms in Humans. This would increase the effectiveness of herbal medicine against the germs concerned and would also contribute to the fight against their resistance to antimicrobial agents.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

No animal or human samples were used in the study.

## 6. REFERENCES

1. Olusola LF, Baba J, Muhammad IL. Medicinal plants for the treatment of malaria and typhoid diseases. J. Medicinal Plants Research. 2023 ; 17 (1): 16-27. <https://doi.org/10.5897/JMPR2022.7249>
2. WHO. Typhoid fever. <https://www.who.int/fr/news-room/fact-sheets/detail/typhoid>. 2023, (Accessed 28th May 2023).
3. WHO. Cholera. <https://www.who.int/news-room/fact-sheets/detail/cholera>. 2022, (Accessed 13th June 2022).
4. Li Y, Kong D, Fu Y, Sussma, MR, Wu H. The effect of developmental and environmental factors on secondary metabolites in medicinal plants. Plant Physiology Biochem. 2020;148: 80-89. <https://doi.org/10.1016/j.plaphy.2020.01.006>
5. Islam Z, Rashadul Islam SM, Hossen F, Mahtab-ul-Islam K, Hasan MR, Karim R. *Moringa oleifera* is a prominent source of nutrients with potential health benefits. Int. J. Food Sci. 2021, Article ID 6627265, 11 p. <https://doi.org/10.1155/2021/6627265>
6. Flores B, Ramirez E, Moncada A, Salinas N, Fischer R, Hernandez C, Mora-Sanchez B, Mora-Sanchez B, Sheleby-Elias J, Jiron W, Balcazar JL. Antimicrobial effect of *Moringa oleifera* seed powder against *Vibrio cholerae* isolated from the rearing water of shrimp (*Penaeus vannamei*) postlarvae. Lett. Appl. Microbiol. 2022; 74 (2): 238-246. <https://doi.org/10.1111/lam.13604>
7. Pareek A, Pant M, Mohan Gupta M, Kashania P, Ratan Y, Jain V, Pareek A, Chaturgoon AA. *Moringa oleifera*: An updated comprehensive review of its pharmacological activities, ethnomedicinal, phytopharmaceutical formulation, clinical, phytochemical, and toxicological aspects. Int. J. Mol. Sci. 2023; 24, 2098. <https://doi.org/10.3390/ijms24032098>
8. Velez-Gavilan J. *Moringa oleifera* (horse radish tree). CABI Compendium. 2023, <https://doi.org/10.1079/cabicompendium.34868>
9. Akwa TE, Nguimbous SP. Common plants used in the treatment of typhoid fever, their active components and toxicity related issues: A review. Nat. Res. Human Health. 2021; 1 (1): 36-42. <https://doi.org/10.53365/nrfhh/141241>
10. Hadis M, Gebreyohannes Y, Gameda N. Potential therapeutic uses of *Moringa stenopetala*: a scoping review. J. Global Health Science. 2020; 2 (2), e26. <https://doi.org/10.35500/jghs.2020.2.e26>

11. Baquero F, Levin BR. Proximate and ultimate causes of the bactericidal action of antibiotics - Perspectives. *Nature Reviews Microbiology*. 2021; 19 (2): 123-132. <https://doi.org/10.1038/s41579-020-00443-1>
12. FMS-EUCAST. French Microbiology Society-European Committee on Antimicrobial Susceptibility Testing/Comité de l'Antibiogramme de la Société Française de Microbiologie. Recommandations 2022. <https://www.sfm-microbiologie.org/boutique/comite-de-lantibiogramme-de-la-sfm-casfm/>
13. Holt JG. Bergey's manual of determinative bacteriology. 1994, Williams and Wilkins eds, Baltimore.
14. Rigby J, Elmerhebi E, Diness Y, Mkwanda C, Tonthola K, Galloway H, Miles R, Henrion MYR, Edwards T, Gauld J, Msefula C, Johnston R, Nair S, Feasey N, Elvis NC. Optimized methods for detecting *Salmonella typhi* in the environment using validated field sampling, culture and confirmatory molecular approaches. *J. Appl. Microbiol.* 2021 ; 132 : 1503-1517. <https://doi.org/10.1111/jam.15237>
15. Mellon FA, Bennett RN, Holst B, Williamson G. Intact glucosinolate analysis in plant extracts by programmed cone voltage Electrospray LC/MS: Performance and comparison with LC/MS/MS methods. *Analytical Biochem.* 2002 ; 306 : 83-91. <https://doi.org/10.1006/abio.2002.5677>
16. Aubert P, Guinobert I, Blondeau C, Bardot V, Ripoche I, Chalard P, Neunlist M. Basal and spasmolytic effects of a hydroethanolic leaf extract of *Melissa officinalis* L. on intestinal motility: An ex vivo study. *J. Medicinal Food.* 2019; 22(7): 653-662. <https://doi.org/10.1089/jmf.2018.0154>
17. Gainche M, Ogeron C, Ripoche I, Senejoux F, Cholet J, Decombat C, Delort L, Berthon JY, Saunier E, Caldefie Chezet F, Chalard P. Xanthine oxidase inhibitors from *Filipendula ulmaria* (L.) Maxim and their efficient detections by HPTLC and HPLC analyses. *Molecules.* 2021; 26: 1939. <https://doi.org/10.3390/molecules26071939>
18. Lalas S, Tsaknis J, Sflomos K. Characterisation of *Moringa stenopetala* seed oil variety "Marigat" from island Kokwa. *Eur. J. Lipid Sci. Technol.* 2003; 105: 23-31. <https://doi.org/10.1002/EJLT.200390002>
19. Lin H, Zhu H, Tan J, Wang H, Wang Z, Li P, Zhao C, Liu J. Comparative analysis of chemical constituents of *Moringa oleifera* leaves from China and India by ultra-performance liquid

- chromatography coupled with quadrupole-time-of-flight mass spectrometry. *Molecules*. 2019 ; 24 (5). <https://doi.org/10.3390/molecules24050942>
20. Xu YB, Chen GL, Guo MQ. Antioxidant and anti-inflammatory activities of the crude extracts of *Moringa oleifera* from Kenya and their correlations with flavonoids. *Antioxidants*. 2019; 8: 296. <https://doi.org/10.3390/antiox8080296>
  21. Xiong Y, Shahid Riaz Rajoka M, Zhang M, Zhendan H. Isolation and identification of two new compounds from the seeds of *Moringa oleifera* and their antiviral and anti-inflammatory activities. *Nat. Prod. Res.* 2022; 36 (4): 974-983. <https://doi.org/10.1080/14786419.2020.1851218>
  22. Ashraf AM, Abdulaziz AAA, Khalid SA, Turki MD, Essam NS, Marwah MB. Antimicrobial activity of some plant extracts against bacterial strains causing food poisoning diseases. *Saudi J. Biol. Sci.* 2018; 25: 361-366. <https://doi.org/10.1016/j.sjbs.2017.02.004>
  23. Andrews JM. Determination of minimum inhibitory concentrations. *J. Antimicrob. Chemotherapy.* 2001; 48 (Suppl. S1) : 5-16. [https://doi.org/10.1093/jac/48.suppl\\_1.5](https://doi.org/10.1093/jac/48.suppl_1.5)
  24. Mogana R, Adhikari, Tzar MN, Ramliza R, Wiart C. Antibacterial activities of the extracts, fractions and isolated compounds from *Canarium patentinervium* Miq. against bacterial clinical isolates. *BMC Complementary Medicine Therapies.* 2020 ; 20 : 55. <https://doi.org/10.1186/s12906-020-2837-5>
  25. Naïtali M, Dubois-Brissonnet F. Développement des microorganismes pathogènes dans les aliments. In: *Risques microbiologiques alimentaires*. 2017, Lavoisier, Paris, France.
  26. Okou OC, Yapo SES, Kporou KE, Baibo GL, Monthaut, S, Djaman AJ. Évaluation de l'activité antibactérienne des extraits de feuilles de *Solanum torvum* Swartz (Solanaceae) sur la croissance in vitro de 3 souches d'entérobactéries. *J. Appl. Biosci.* 2018 ; 122 : 12282-12290. <https://doi.org/10.4314/jab.v122i1.8>
  27. Yue Y, Chen C, Zhong K, Wu Y, Gao H. Purification, fermentation optimization, and antibacterial activity of Pyrrole-2-carboxylic acid produced by an endophytic bacterium, *Bacillus cereus* ZBE, isolated from *Zanthoxylum bungeanum*. *Indust. Engin. Chem. Res.* 2022; 61: 1267-1276. <https://doi.org/10.1021/acs.iecr.1c04164>
  28. Tomasic T, Katsamakas S, Hodnik Z, Ilas J, Brvar M, Solmajer T, Montalvao S, Tammela P, Banjanac M, Ergovic G, Anderluh M, Masic LP, Kikelj D. Discovery of 4,5,6,7-tetrahydrobenzo[1,2-d] thiazoles as novel DNA gyrase inhibitors targeting the ATP-binding site. *J. Med. Chem.* 2015; 58 (14): 5501-21. <https://doi.org/10.1021/acs.jmedchem.5b00489>

29. Casillas-Vargas G, Ocasio-Malave C, Medina S, Morales-Guzman C, Garcia Del Valle R, Carballeira NM, Sanabria-Rios DJ. Antibacterial fatty acids: An update of possible mechanisms of action and implications in the development of the next-generation of antibacterial agents. *Progr. Lipid Res.* 2021 ; 82 : 101093. <https://doi.org/10.1016/j.plipres.2021.101093>
30. Narendra Jaisinghani R. Antibacterial properties of quercetin. *Microbiol. Res.* 2017; 8: 6877. <https://doi.org/10.4081/mr.2017.6877>
31. Cabezas-Pizarro J, Redondo-Solano M, Umana-Gamboa C, Arias-Echandi ML. Antimicrobial activity of different sodium and potassium salts of carboxylic acid against some common foodborne pathogens and spoilage-associated bacteria. *Revista Argentina de Microbiologia.* 2018 ; 50 (1) : 56-61. <https://doi.org/10.1016/j.ram.2016.11.011>
32. Kozyra M, Biernasiuk A, Malm A. Analysis of phenolic acids and antibacterial activity of extracts obtained from the flowering herbs of *Carduus acanthoides* L. *Acta Polonicae Pharmaceutica-Drug Res.* 2017 ; 74 (1) : 161-172. PMID : 29474772
33. Ngoufack Tagousop C, Tamokou JD, Endeguele Ekom S, Ngnokam D, Voutquenne Nazabadioko L. Antimicrobial activities of flavonoid glycosides from *Graptophyllum grandulosum* and their mechanism of antibacterial action. *BMC Complementary and Alternative Medicine.* 2018 ; 18 : 252. <https://doi.org/10.1186/s12906-018-2321-7>
34. Bouarab-Chibane L, Forquet V, Lantéri P, Clément Y, Léonard-Akkari L, Oulahal N, Degraeve P, Bordes C. Antibacterial properties of polyphenols: characterization and QSAR (quantitative structure-activity relationship) models. *Frontiers Microbiol.* 2019 ; 10 : 829. <https://doi.org/10.3389/fmicb.2019.00829>
35. Biharee A, Sharma A, Kumar A, Jaitak V. Antimicrobial flavonoids as a potential substitute for overcoming antimicrobial resistance. *Fitoterapia.* 2020 ; 146 : 104720. <https://doi.org/10.1016/j.fitote.2020.104720>
36. Cuyette J, Cherubin P, Serrano A, Taylor M, Abedin F, O'Donnell M, Burrell H, Tatulian SA, Teter K. Quercetin-3-Rutinoside blocks the disassembly of cholera toxin by protein disulfide isomerase. *Toxins (Basel).* 2019; 11 (8): 458. <https://doi.org/10.3390/toxins11080458>
37. Burel C, Kala A, Purevdorj-Gage L. Impact of pH on citric acid antimicrobial activity against Gram-negative bacteria. *Lett. Appl. Microbiol.* 2020; 72: 332-340. <https://doi.org/10.1111/lam.13420>

38. Mani-Lopez E, Garcia S, Lopez-Malo A. Organic acids as antimicrobials to control Salmonella in meat and poultry products. *Food Res. Int.* 2012; 45: 713-721. <https://doi.org/10.1016/J.FOODRES.2011.04.043>
39. Zhou C, Wang F, Chen H, Meng L, Qiao F, Liu Z, Hou Y, Wu C, Fan Y, Liu L, Wang S, Wang Y . Selective antimicrobial activities and action mechanism of micelles self-assembled by cationic oligomeric surfactants. *ACS Appl. Mater Inter.* 2016; 8: 4242-4249. <https://doi.org/10.1021/acsami.5b12688>
40. Kundukad B, Udayakumar G, Grela E, Kaur D, Rice SS, Kjelleberg S, Doyle PS. Weak acids as an alternative anti-microbial therapy. *Biofilm.* 2020; 2. <https://doi.org/10.1016/j.bioflm.2020.100019>
41. Galuppo M, De Nicola GR, Iori R, Utri PD, Bramanti P, Mazzou E. Antibacterial activity of glucomoringin bioactivated with myrosinase against two important pathogens affecting the health of long-term patients in hospitals. *Molecules.* 2013; 18: 14340-14348. <https://doi.org/10.3390/molecules181114340>
42. Liu YH, Wu PQ, Hu QL, Pei YJ, Qi FM, Zhang ZX, Fei DQ. Cytotoxic and antibacterial activities of iridoids and sesquiterpenoids from *Valeriana jatamansi*. *Fitoterapia.* 2017; 123: 73-78. <https://doi.org/10.1016/j.fitote.2017.09.011>
43. Ugboko H, De N. Mechanisms of antibiotic resistance in *Salmonella typhi*. *Int. J. Curr. Microbiol. Appl. Sci.* 2014; 3 (12): 461-476. Corpus ID: 57264492
44. Willey J, Sherwood L, Wolverton C. Prescott s Microbiology. 2013, McGraw-Hill, 9th ed., New York.
45. Ghannoum MA, Rice LB. Antifungal agents : mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *ASM J. Clin. Microbiol. Rev.* 2020 ; 12(4). <https://doi.org/10.1128/CMR.12.4.501>
46. Verma J, Bag S, Saha B, Kumar P, Ghosh TS, Dayal M, Senapati T, Mehra S, Dey P, Desigamani A, Kumar D, Rana P, Kumar B, Maiti TK, Sharma NC, Bhadra RK, Mutreja A, Nair GB, Ramamurthy T, Das B. Genomic plasticity associated with antimicrobial resistance in *Vibrio cholerae*. *Proc. Natl. Acad. Sci. USA.* 2019; 116: 6226-6231. <https://doi.org/10.1073/pnas.1900141116>
47. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic resistance. *Nat. Rev. Microbiol.* 2015; 13: 42-51. <https://doi.org/10.1038/nrmicro3380>

48. Das B, Verma J, Kumar P, Ghosh A, Ramamurthy T. Antibiotic resistance in *Vibrio 22holera*: Understanding the ecology of resistance genes and mechanisms. *Vaccine*. 2020; 38: A83-A92. <https://doi.org/10.1016/j.vaccine.2019.06.031>
49. Levison ME, Levison JH. Pharmacokinetics and pharmacodynamics of antibacterial agents. *Infect Dis. Clin. North Am.* 2009; 23 (4): 791-815. <https://doi.org/10.1016/j.idc.2009.06.008>
50. Werth BJ. Overview of antibacterial drugs. <https://www.msmanuals.com/professional/infectious-diseases/bacteria-and-antibacterial-drugs/overview-of-antibacterial-drugs>. 2022; (Accessed 25<sup>th</sup> June 2023).

UNDER PEER REVIEW