

## Original Research Article

# Isolation and characterization of *Lactobacillus* strains with probiotic potential from dairy products

### ABSTRACT

**Aims:** This research aims to isolate and characterize from dairy products, in particular yoghurts, cultivated *Lactobacillus* strains with potential for use as probiotics in poultry farming.

**Study design:** This research consists of four parts, an introduction with literature review, a description of the materials and methods used, finally the results obtained and their discussion.

**Place and Duration of Study:** Central Veterinary Laboratory of Kinshasa, specifically in the Bacteriology and Animal Services departments, between April and September 2022.

**Methodology:** A yogurt was cultured in order to observe the presence of Lactobacilli in the M.R.S (Mans, Rogosa and Sharp) Agar culture medium; this culture revealed the growth of Lactobacilli. It was even proceeded to the serial dilution of these strains, the results had also revealed the growth of Lactobacilli in the 6 test tubes used.

**Results:** The results obtained showed a significant increase in weight (positive effect on the immune system) in the poultry of the experimental group and an absence of pathogenicity after an incubation period (inhibitory effect of strains of Lactobacilli on those of *Escherichia coli* and *Streptococcus sp*) whereas in the control group, the poultry showed a lower weight gain compared to the experimental group and developed colibacillosis associated with other effects such as weakness, fever and angina after an incubation period.

**Conclusion:** The positive results of Lactobacilli strains observed in poultry in the experimental group show the potential of these strains to be used as probiotics in poultry farming to improve poultry health through the inhibitory effect of these strains on pathogenic *Escherichia coli* and *Streptococcus sp*. This opens up safe alternatives to the use of these strains to combat the misuse of antibiotics by reducing their use in human and animal care.

*Keywords: Probiotics, prebiotics, antibiotics, poultry, yoghurt.*

- **INTRODUCTION**

Lactic acid bacteria by their acidifying, flavoring and texturizing properties are widely used in milk-derived products and their probiotic properties are very useful for health, in fact, they improve digestive functions and have a very positive effect on the intestinal microflora [1]. Products derived from traditional lactic fermentation have experienced considerable development in recent years thanks to the interest that consumers find on the organoleptic, nutritional, therapeutic and even hygienic level because of their acidity. The biodiversity of these lactic acid bacteria involved in this process is a fundamental factor for the preservation of the typicality and original characteristics of products [2] and without disadvantages for lactose intolerant consumers.

Dairy products, thanks to their rich and diversified composition, have very interesting therapeutic effects and regular consumption of these products can have an anticarcinogenic [3], antimicrobial [4], immunological effect very important [5] thanks to their very low final pH between 4.3 and 4.4 [6,7] without forgetting the beneficial effect on the intestinal tract [8].

In order to improve the quality and avoid the disappearance in the more or less long term [9] of traditional dairy products, the latter deserve a transfer of production on a semi-industrial or industrial scale to be known and marketed on a larger scale. This will make it possible to standardize the various stages of their production process, to better control their quality and to reinforce the added value of their properties.

Hence the idea of isolating lactic acid bacteria from traditional products because the interest of these strains lies in their ability to produce specific metabolites, especially those responsible for flavor, which may be different from those produced by industrial strains [2]. Then use them in a semi-industrial process to obtain other fermented products with better hygienic quality and very advanced therapeutic, especially than in recent years; research has intensified to find probiotic-based therapies for the treatment of various chronic diseases [10].

- **Objective**

The overall purpose of this research is to isolate and characterize from dairy products, in particular yoghurts, cultivated *Lactobacillus* strains with potential for use as probiotics in poultry farming.

- **MATERIAL, METHOD AND EXPERIMENTAL DETAILS**

The field work took place at the Central Veterinary Laboratory in Kinshasa, specifically in the Bacteriology and Animal Services departments. Two valves were fitted out to carry out the experiments on poultry, one had been used for the experimental group with strains of Lactobacilli (GELB) and for the control group (GT). Poultry species were marked as follows: P1GELB, P2GELB, P3GELB, P4GELB, P5GELB, P6GELB & P7GELB (for the experimental group) and P1GT, P2GT, P3GT, P4GT, P5GT, P6GT & P7GT (for the control group). P=Chicken; 1, 2, 3, 4, 5, 6, 7=number of hens in their respective group; G=Group; E=Experimental and T=Control. During the experiment, these hens, which were of the same breed (Sasouat) and the same age (3 months), were fed poultry feed P1 and P2 with a consumption of 100g per hen per day, i.e. 1400g per day for the set of poultry.

- **Methodology**

The methodology consisted in isolating and characterizing Lactobacilli with a dairy product containing potential strains of these microorganisms in order to prepare inoculums on the one hand and prepare strains of pathogenic germs (*Escherichia coli* and *Staphylococcus sp*) to carry out inoculations in poultry in order to observe the related effects.

The samples of the dairy product (yoghurt) were cultured on MRS Agar, poured into petri dishes heated to the oven at 37.5 ° C for 72 hours. Each sample was read after the incubation period.

- **Experimental details**

In order to fortify the Lactobacilli that grew, the seeded boxes were placed in an anaerobic jar, incubated again at room temperature of 37°C for 72 hours in a CO<sub>2</sub>-enriched atmosphere. Other cultured strains of Lactobacilli were subjected to serial dilution to test the effectiveness of this medium (MRS Agar) in order to observe the growth of Lactobacilli.

The different species of *Lactobacillus* strains have been characterized as Lactobacilli casei by their origin (dairy product used) through electron microscopic observation. The chickens of the two groups were weighed before the *lactobacillus* and pathogenic germs of *Escherichia coli* and *Staphylococcus sp* inoculation sessions using a manual scale. In the experimental species, three sessions of inoculation of fortified lactobacilli in peptone water took place within a week, every 72 hours with 2 ml of inoculum per head at each session and administered by the route oral so as to travel through the digestive tract until it reaches the target, that is to say the colon of the poultry where they exert the beneficial effect on health. Inoculation and sampling were done using a sterile single-use syringe.

After having administered the strains of Lactobacilli to the experimental species, a series of three sessions of inoculation of the strains of *Escherichia coli* and *Staphylococcus sp* were carried out in the two groups, in the space of a week, every 72 hours with 2 ml of inoculum per species at each session. To enhance the probiotic effect, honey was also used as a prebiotic during these inoculation sessions at a rate of 2 ml per species in the experimental group. After these different inoculations, the poultry was weighed again, four weeks after the inoculations, taking into account the incubation period for colibacillosis (from 1 to 3 weeks). The presence of lactobacilli was sought in the stools to confirm or not, the effectiveness of the latter during their passage through the digestive tract of the animal. Alongside this search for Lactobacilli in the stool, we should also look for the presence of *Escherichia coli* and *Staphylococcus sp* which was noted.

Stool samples were cultured in Petri dishes containing MRS Agar culture medium to isolate potential lactobacilli, while for *Escherichia coli* and *Staphylococcus sp*, their presence was to be observed in the stools of hens in the control group n' who have not benefited from the inoculation of lactobacillus strains. The results obtained after these various manipulations and experiments will be presented and discussed below.

• **RESULTS AND DISCUSSION**

**3.1. Results**

**3.1.1. Initial poultry weights**

These weights obtained by species in the two groups are shown in the following Table 1:

Table 1. Initial weights of poultry before inoculations

Experimental Lactobacillus Group (GELB)		Control Group (WG)	
Species	Weight (g)	Species	Weight (g)
P1GELB	1160	P1GT	1405
P2GELB	1190	P2GT	945
P3GELB	875	P3GT	825
P4GELB	1370	P4GT	1365
P5GELB	1270	P5GT	1365
P6GELB	1460	P6GT	1495
P7GELB	1330	P7GT	975

- Average weight (Experimental group):  $\mu_1 = 1236.4 \pm 189.8g$ ;  $S_2^1 = 36056.9g$
- Average weight (control group):  $\mu_2 = 1196.4 \pm 270.7g$ ;  $S_2^2 = 73280.9g$

The fourteen hens were fed under the same conditions for 2 months. Before starting the inoculations, all the hens were weighed. There was no significant difference in weight between the two groups because according to the Student's test on the difference of two means applied, the  $t_{cal}$  was lower than the  $t_{tab}$ , i.e.:  $t_{cal} < t_{tab}$  :  $0.320 < 2.179$ .

**3.1.2. Culture of dairy products (yogurt)**

The results related to the growth of dairy products are reported in table 2 below:

**Table 2.** Inoculation of dairy products

Petri Dishes	Growth
1	+++
2	+++
3	+++
4	++
5	+
6	+
7	+

Legend:

- High growth: +++
- Average growth: ++
- Low growth: +

This dairy product (yogurt) was used for its potential to have strains of lactobacilli capable of growing easily in the culture medium (MRS Agar).


### 3.1.3. Serial dilution of Lactobacilli strains

The results obtained after serial dilution of the sources of Lactobacilli are given in table 3 below:

**Table 3.** Serial dilution of Lactobacilli strains

	Petri Dishes					
	1	2	3	4	5	6
Peptone water	9 ml	9 ml	9 ml	9 ml	9 ml	9 ml
Diluent	1 ml	1 ml	1 ml	1 ml	1 ml	1 ml
Dilution rate	$10^{-1}$	$10^{-2}$	$10^{-3}$	$10^{-4}$	$10^{-5}$	$10^{-6}$

This table shows the serial dilution carried out in 6 test tubes with the addition of 1 ml of diluent (direct sample of Lactobacillus strains after inoculation on MRS Agar) in 9 ml of peptone water contained in each of these tubes.



**Fig. 1.** Growth of lactobacilli in petri dishes.

This figure shows the growth of Lactobacilli in inoculated petri dishes after serial dilution from the direct sample (yogurt). When reading the cultures in these dishes, growth was noted after 48 hours under CO<sub>2</sub> (one decimal dilution):

- Box 1: >300 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-1</sup>;
- Box 2: >300 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-2</sup>;
- Box 3: >250 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-3</sup>;
- Box 4: 120 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-4</sup>;
- Box 5: 37 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-5</sup>;
- Box 6: >3 colonies/box/mm<sup>3</sup> with a dilution rate of 10<sup>-6</sup>.

**3.1.4. Weight after poultry inoculations**

These weights obtained by species in the two groups are given in the following table 4:

Table 4. Poultry weight after inoculations.

Experimental Lactobacillus Group (GELB)		Control Group (GT)	
Species	Weight (g)	Species	Weight (g)
P1GELB	2065	P1GT	1780
P2GELB	1905	P2GT	1410
P3GELB	1945	P3GT	1725
P4GELB	1935	P4GT	1510
P5GELB	1760	P5GT	1415
P6GELB	1700	P6GT	1200
P7GELB	1780	P7GT	1320

- Average weight (Experimental group):  $\mu_1 = 1870 \pm 127.9g$ ;  $S_2^1 = 16366.6g$
- Average weight (control group):  $\mu_2 = 1480 \pm 209.7g$ ;  $S_2^2 = 44008.3g$

The results of the two groups were subjected to the hypothesis test on the difference of two means to see if this difference was significant or not. For this, the Student's t test (Comparison of two means observed on two independent samples) showed  $t_{cal} > t_{tab}$  (i.e.:  $t_{cal} > t_{tab} : 4.199 > 2.179$ ) thus indicating a significant difference between the two groups of hens whereas at the start the hens had almost the same weight (Student's t test). The increase in weight was due to the administration of Lactobacilli. After the 3 sessions of inoculation of Lactobacillus strains to the experimental species (GELB), a marked improvement in the

immunity was observed in the poultry of this group, characterized by the absence of disease (**Table 5**) compared to that of the control group (GT).

### 3.1.5. Seeding of stool collected by swabs from poultry in both groups

From the search for the presence or absence of lactobacilli in poultry feces (indication of the inhibitory effect of lactobacilli against strains of *Escherichia coli* and *Staphylococcus sp*) of both groups and after seeding these samples in petri dishes based on M.R.S Agar, it appears from reading these different dishes that only in the dishes containing the stool of the experimental group (GELB) than lactobacilli grew while in the stool boxes of the control group (GT), there was no growth (**Table 5**). This proves that these are the administered strains that have resisted crossing the entire digestive tract of the experienced hens.

**Table 5.** Results of inoculation of stools taken by swabs in poultry in the two groups.

Experimental Lactobacillus Group (GELB)		Control Group (GT)	
Species	Presence of LB	Species	Presence of LB
P1GELB	+++	P1GT	- - -
P2GELB	+++	P2GT	- - -
P3GELB	+++	P3GT	- - -
P4GELB	+++	P4GT	- - -
P5GELB	+++	P5GT	- - -
P6GELB	+++	P6GT	- - -
P7GELB	+++	P7GT	- - -

Legend: high growth (+++); lack of growth (- - -); LB (Lactobacilli)

**Figure 2** below also illustrates the growth of these germs in petri dishes. This figure shows optimal growth of Lactobacilli with more than 300 colonies per dish. Inoculation of pathogenic strains of *Escherichia coli* and *Staphylococcus sp* to experimental and control species.



**Fig. 2.** Growth of *Lactobacilli* colonies in petri dishes inoculated from stools collected by swabs from poultry in the experimental group (GELB)

**Table 6.** Signs observed in poultry of the control group (GT) after inoculation with strains of *Escherichia coli*.

WG species	Signs observed
P1GT	Weakness, fever, angina
P2GT	Weakness, whitish diarrhoea, angina
P3GT	Weakness, angina, drowsiness
P4GT	Weakness, white diarrhea, fever
P5GT	Weakness, drowsiness, white diarrhea
P6GT	Weakness, white diarrhea, angina
P7GT	Affaiblissement, diarrhée blanche, somnolence, décès



After the 3 sessions of inoculation of the strains of *Escherichia coli* and *Staphylococcus sp* to the experimental species (GELB) and to the control species (GT), the hens were observed for three weeks to see the reaction to *E. coli* and *S. sp*.

It appears from this period that no hen in the experimental group (A) developed or showed signs of the disease, whereas in the control species (group B), the following signs were observed: whitish diarrhoea, weakness and drowsiness (Table 6) and the diagnosis of colibacillosis had been established by the veterinary service. One of the seven hens in the control group (GT) had died as a result of this colibacillosis and the six others were treated by the laboratory for the appropriate care.

### 3.2. Discussion

Lactic acid bacteria make up more than 93% of the intestinal flora and play a key role in our metabolism: they are the sentinels that keep pathogens in small numbers and safe, which allows them to live in symbiosis in our body without harm. For 20 or 30 years, efforts to improve human and animal health have focused on the modulation pathways of the intestinal flora by the addition of live microbes now called probiotics. The main probiotics used are lactic acid bacteria, including bifidobacteria, lactobacilli and lactococci but can also be yeasts such as *Saccharomyces cerevisiae*. The "health benefit claims" related to probiotics are the maintenance of a normal and healthy intestinal flora, protection against infections, alleviation of lactose intolerance, stimulation of the immune system, reduction of blood lipids thus preventing the onset of cardiovascular diseases in human health and improving zootechnical parameters in animal husbandry

In addition, bifidobacteria and lactobacilli can be stimulated by indigestible food ingredients such as oligosaccharides collectively called prebiotics. Pre and probiotics can be combined in a food called a symbiotic. The contribution of probiotics in poultry feed is a favorable alternative to the use of antibiotics as growth factors which is currently contested following the application of multiple resistances. In addition, probiotics contribute effectively to improving livestock performance by maintaining a stable and healthy digestive flora [11]. It is for this reason that the present work wanted to verify the effects of lactobacilli on the health of poultry by inoculating them first alone and then associated with the strains of *Escherichia coli* and *Staphylococcus sp* in the experimental group (GELB).

The control group (GT) received only the inoculation of strains of *Escherichia coli* and *Staphylococcus sp* (pathogenic strains). By strengthening the microbial ecosystem of poultry, Patterson and Burkholder [12] demonstrated that probiotics contribute to the immune defense and protect chickens against the consequences of stress such as vaccination and temperature changes. Improvements in terms of weight gain and feed efficiency have thus been observed following the consumption of probiotics [13,14]. Such was the case with this work where the hens tested experienced significant weight gain. The role of probiotics is to colonize the intestine and thus prevent its colonization by enteric pathogens causing diarrhea [15].

This effect is probably attributable to the action of bacterial probiotics on the intestinal mucosa and which can be justified in the case of this work, since the addition of lactobacilli in poultry feed had an effect of preventing colibacillosis from settle. It is well known that, especially in monogastrics, bacterial probiotics can modify the permeability of the intestinal mucosa, activate immune cells and prevent adhesion of pathogens to the intestinal mucosa [16,17]. For antibiotic-associated diarrhea, probiotics have been shown to be useful as a preventative treatment, and potentially, they can be used to alleviate signs and symptoms once antibiotic-induced diarrhea has occurred [18,19,20].

The use of probiotics seems to offer an alternative solution in reducing the use of antibiotics thanks to their beneficial effects on the health of poultry. It emerges from this experiment that the majority of hens in

the experimental group (GELB) show a greater weight gain than those in the control group; after subjecting this difference in mean weight to Student's t-test, we found that the difference was significant. This difference in weight gain in the experimental group (GELB) reflects the beneficial effect of lactobacilli (probiotics) on the immunity of poultry which is generally manifested by a considerable weight gain compared to the control group (GT) which does not benefit from this immunomodulatory effect of lactobacilli. The authors have also affirmed this in previous works, in particular [13,14,21].

The fermented milk products (yoghurts) used had shown a potential to provide strains of lactobacilli likely to be used, after culture, as probiotics. The inoculation of the pathogenic strains of *Escherichia coli* and *Staphylococcus sp* in the two groups (GELB & GT) gave different effects: in the experimental group, resistance was noted in the hens translated by the absence of symptoms and the disease until the end of the experiment, despite the prolonged incubation period. On the other hand, in the control group, after the incubation period, symptoms of disease were observed and subsequently, the death of a hen of this group (P1GT) was recorded. In the experimental group, the inhibitory effect of lactobacilli on pathogenic strains of *Escherichia coli* and *Staphylococcus sp* played in favor of poultry.

The efficacy of lactobacilli strains through the digestive tract of the host (the animal) must also be demonstrated by the presence of these strains in the faeces of the animal as discussed in the probiotic selection criteria according to Klaenhammer and Kullen [22,23,24,25]. The results obtained with the stool showed the presence of lactobacilli in the species of the experimental group while nothing was observed in the species of the control group. Thus the strains of lactobacilli administered in adequate quantities demonstrated their effectiveness in poultry of the experimental group because on the one hand, they resisted acidity along the digestive tract and on the other hand, they exerted an inhibitory effect on the pathogenic strains of *Escherichia coli* administered to these species. That's how they ended up in the stool.

The fact that we subsequently had to resort to antibiotics to treat the hens in the control group who had fallen ill, whereas the experimental group had just benefited from Lactobacilli to resist the disease clearly shows the ecological role played by Lactobacilli (probiotics) to inhibit the effects of strains of *Escherichia coli* and *Staphylococcus sp* to prevent disease. This aspect is very important economically in poultry farming because prevention costs less than curative care; even if we can treat with antibiotics, we must fear the phenomena of resistance and the by-products of antibiotics in the environment with all the possible consequences.

The results obtained in this work are only preliminary in relation to the actual use of probiotics in care. The most used probiotics are represented by two genera: *Lactobacillus* and *bifidobacterium* [26,27,28,29,30] but in the As part of this work, we were only able to exploit the *Lactobacillus*. However, other studies may continue in the future with bifidobacterium strains. This work has made it possible to highlight *Lactobacillus* and to assess their effects on avian health in order to consider their use in care in the long term with a view to reducing the use of antibiotics.

## • CONCLUSION

This work has made it possible to isolate and characterize from dairy products (yoghurts) strains of Lactobacilli with probiotic potential. The results obtained showed that these strains have the ability to give positive effects on poultry health by modulating the immune system through their passage through the animal's digestive tract, characterized by resistance to pathogenic strains (*Escherichia coli* and *Staphylococcus sp*) on the one hand and weight gain on the other. The addition of fructooligosaccharides, as a prebiotic substance, to the culture medium of these bacteria improves the growth and acidifying power of all lactic strains. It would also be interesting to continue this study on larger numbers of hens, and over a longer period of time in order to be able to better specify the effects of the prebiotic. The interest in the application of probiotic bacteria in the food field has increased considerably in recent years and will certainly continue in the future. Obtaining indigenous probiotic strains remains an important objective in terms of research as well as in terms of applications in human and animal health.

## REFERENCE

- Gorbach S.L. The discovery of L. GG. Nutrition Today. 1996.
- Ayad A.H.E et al. Flavour forming abilities and amino acid requirements of Lactococcus lactis strains isolated from artisanal and non-dairy origin, International Dairy Journal, 1999 ; 9 : 725-735.
- Nagira T. et al. Suppression of UVC-induced cell damage and enhancement of DNA repair by the fermented milk, kefir, Cytotechnology, 2002 ; 40 : 125-137.
- Santos A., San Mauro M., Sanchez M., Torres J.M and Marguina D. The antimicrobial properties of different strains of Lactobacillus spp. Isolated from kefir, System and Applied Microbiology, 2003 ; 26 : 434-437.
- Vinderola C.G et al. Immunomodulating capacity of kefir, Journal of Dairy Research, 2005 ; 72 : 195-202.
- Fontan G., Martinez S., Franco I., Carballo J. Microbiological and chemical changes during the manufacture of kefir made from cows' milk, using a commercial starter culture, International Dairy Journal, 2006 ; 16 : 762-767.
- Kwak H.S, Park S.K, Kim D.S. Biostabilization of kefir with a nonlactose-Fermenting Yeast. Journal of Dairy Science, 1996 ; Vol.79 N°6.
- Urdaneta J, Barrenetxe J.P, Aranguren A, Irigoyen F, Marzo F, Ibàñez C. Intestinal beneficial effects of kefir-supplemented diet in rats, Nutrition Research 2007 ; 27 : 653-658.
- Benkerroum N et Tamime A.Y. Technology transfer of some Moroccan traditional dairy products (Iben, jben and smen) to small industrial scale, Food Microbiology, 2004 ; 21 : 399-413.
- Katja Z et Andreja G. Modelling of batch kefir and bio-products processing, 2010 ; 88 : 55-60.
- Amara Sabrina. Effets probiotiques des bactéries lactiques sur le poulet de chair. Mémoire de Master en Biotechnologie. Option : écosystèmes microbiens complexes. Faculté des Sciences. Université d'Oran. Algérie. 2010 ; 187 pages.
- Patterson, J. A. and Burkholder K. M. "Application of prebiotics and probiotics in poultry production." Poultry Science 2003 ; 82(4): 627-631.
- Jin, L. Z. et Ho Y. W. "Effects of adherent Lactobacillus cultures on growth, weight of organs and intestinal microflora and volatile fatty acids in broilers." Animal Feed Science and Technology 1998 ; 70(3): 197-209.
- Simon, O. et Jadamus A. "Probiotic feed additives - effectiveness and expected modes of action." Journal of Animal and Feed Sciences 2001 ; 10: 51-67.

- Krehbiel C.R, Rust S.R, Zhang G., Gilliland S.E. Bacterial direct-fed microbials in ruminant diets : Performance response and mode of action. *Journal of Animal Science*, volume 81, Issue 14\_suppl\_2 February 2003, Pages E120-E132, [http://doi.org/10.2527/2003.8114\\_suppl\\_2E120x](http://doi.org/10.2527/2003.8114_suppl_2E120x).
- Kaur, I. P., Kuhad A. "Probiotics: Delineation of Prophylactic and Therapeutic Benefits." *Journal of Medicinal Food* 2009 ; 12(2): 219-235.
- Kaur I.P and Kuhad A. Probiotics : Delineation of prophylactic and therapeutic benefits. *Journal of medical food*. 2003 : 12(2) : 45-47.
- Arvola T, Laiho K, Torkkeli S, Mykkänen H, Salminen S, Maunula L, Isolauri E. Prophylactic *Lactobacillus GG* reduces antibiotic-associated diarrhea in children with respiratory infections: a randomized study. *Pediatrics*, 1999 ; 104(5): 1
- Vanderhoof JA, Whitney DB, Antonson DL, Hanner TL, Lupo JV, Young RJ. *Lactobacillus GG* in the prevention of antibiotic-associated diarrhea in children. *J Pediatr*, 1999 ; 135: 564-568.
- Armuzzi A, Cremonini F, Bartolozzi F, Canducci F, Candelli M, Ojetti V, Cammarota G, Anti M, De Lorenzo A, Pola P, Gasbarrini G, Gasbarrini A : The effect of oral administration of *Lactobacillus GG* on antibiotic-associated gastrointestinal side-effects during *Helicobacter pylori* eradication therapy. *Aliment Pharmacol Ther*, 2001 ; 15(2): 163-169.
- Tannock, G. W. "Effect of dietary and environmental stress on the gastrointestinal microbiota." In: Hentges, D. (Ed.), *Human intestinal microflora health and disease*. Academic Press. New York, NY. 1983 ; pp. 517-539.
- Klaenhammer, T.R. and M.J. Kullen. "Selection and design of probiotics." *International Journal of Food Microbiology*. 1999 ; 50(1-2): 45-57.
- Saarela, M. et Mogensen G. "Probiotic bacteria : safety, functional and technological properties." *Journal of Biotechnology*. 2000 ; 84(3): 197-215.
- Ouwehand, A.C., Salminen S. "Probiotics: an overview of beneficial effects." *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*. 2002 ; 82(1-4): 279-289.
- Gueimonde, M. and S. Salminen. "New methods for selecting and evaluating probiotics." *Digestive and Liver Disease*. 2006 ; (38): S242-S247.
- Marteau P., Shanahan F. Basic aspects and pharmacology of probiotics: an overview of pharmacokinetics, mechanisms of action and side effects. *Best Practices and Research Clinical Gastroenterology*, 2003 ; 17, 725-740.
- Mercenier A., Pavan S., Pot B. Probiotics as biotherapeutic agents: Present knowledge and future prospects. *Current Pharmaceutical Design*, 2002 ; 8, 99-110.
- Coconnier MH, Lievin V, Bernet-Camard MF, Hudault S, Servin AL. Antibacterial effect of the adhering human *Lactobacillus acidophilus* strain LB. *Antimicrob Agents Chemother*, 1997 ; 41: 1046-52.
- Matsuzaki T, Chin J. Modulating immune responses with probiotic bacteria. *Immunol Cell Biol*, 2000 ; 78: 67-73.
- Andersson H, Asp N-G, Bruce A, Roos S, Wadstrom T, Wold AE : Health effects of probiotics and prebiotics: A literature review on human studies. *Scand J Nutr*, 2001 ; 45: 58-75.
- Bodganov S., Jurendic T. et al. Honey for nutrition and health : A review. *J am Coll Nutr* 2008 ;27 : 677-89.