

Original Research Article

Monitoring and Control of Particle Matter concentration's Traffic-Related Air Pollution Using Low-Cost Mobile Sensors at Six Intersections in Yaoundé, Cameroon

ABSTRACT

Particle pollution has a major influence on public health in Central African cities. Measuring the levels of pollution to which populations are exposed is difficult because only a few African countries have an air quality monitoring network in place. Yet, given the specific anthropogenic sources prevalent in African countries, as well as the predicted increase in their emissions in the next years if no laws are implemented, solutions must be developed. The methodology utilised in this study allows diverse research teams to estimate the population's exposure levels to urban particle pollution at a cheaper cost. In this investigation, we collected data concentrations using Air Master. The spatial mapping of pollutants shows that the Mvog-Mbi crossroad and the Education crossroad are the most polluted points among the six measurement points, with PM_{2.5} concentrations around 145-170 $\mu\text{g.m}^{-3}$. Regarding PM₁₀ and PM₁, the Mvog-Mbi crossroads is the point where concentration are highest, fluctuating respectively between 150-180 $\mu\text{g.m}^{-3}$, 41-48 $\mu\text{g.m}^{-3}$. At some locations, the particle ratio surpasses the WHO recommended range of 0.5-0.8. (Mvog-Mbi crossroads, Education crossroads, Poste Centrale, Hilton). These are the high-risk zones for those with respiratory difficulties. At some locations, the particle ratio surpasses the WHO recommended range of 0.5-0.8 (Mvog-Mbi crossroads, Education crossroads, Poste Centrale, Hilton). Regardless of particle size, these pollutants in air readings are greater and beyond the World Health Organization's recommended values. Furthermore, this investigation provides critical information on Douala's pollution levels, which can be a major source of disease in the city.

Keywords: Urban air quality; Low-Cost Mobile Sensors; Particulate Matter; air pollution; Yaoundé.

1. INTRODUCTION

People's mobility in urban areas today is one of the primary causes of poor air quality in metropolitan regions. The rapid urbanization of cities increases the need for transportation, which lowers air quality. Traffic-Related Air Pollution (TRAP) is the leading cause of air pollution in cities and towns around the world [1]. Intersections in urban settings are places where traffic is dense for much of the day and are

positioned as one of the hotspots of pollutants [2]. Gaseous pollutants and particle pollutants are two types of pollutants that harm the environment in metropolitan areas [3, 4]. Numerous studies illustrate the impact of high-concentration exposure. Furthermore, exposure to contaminated air causes the development and worsening of respiratory and cardiovascular disorders, lung cancer, and a variety of other ailments [5]. Air quality management entails public awareness efforts about the impacts of exposure, measurement campaigns to determine the levels of gaseous or particle pollutants, and compliance with gaseous emissions requirements (Euro standards) on automobiles (two wheels, three wheels, etc.). Several countries are vulnerable to all of these air quality management requirements. These are mainly Sub-Saharan African (SSA) countries that lack the necessary resources to adequately study and manage air quality deterioration. Conventional air pollution monitors are expensive, and underdeveloped countries such as Cameroon lack the resources to set up the necessary number of monitoring stations to measure air pollution. The advancement of miniaturized sensor technologies provides an intriguing low-cost alternative approach for monitoring air pollution [6-8]. These sensors have various advantages over traditional monitoring stations, including the fact that they are modular and may be adapted to meet a variety of needs, including mobile air pollution monitoring [7]. Some scientific studies in Africa and elsewhere have already made use of low-cost mobile measurement instruments [6, 7, 9]. Intersections in metropolitan areas have higher levels of road pollution than other areas. To that purpose, Hounbégnon et al. [2], found that crossroads are the most polluting spots in cities. He contrasts the amounts of particle pollution in a crossroads (many lanes) and a single-lane road in his study. In 2014, Antonel and Chaudhry [10] used a fixed measurement station to study the amount of particulate pollution in the city of Yaoundé and discovered that the concentration levels of particulate matter in the air surpassed WHO standards. A few years later, Nducol et al. [4], observed that concentrations in the air in the city of Yaoundé surpassed WHO standards. The evaluation of air pollutants in cities along highways offers the right amounts of emissions generated by automobiles as well as the exposure levels to commuters, walkers, and those who live or work near these routes. To the best of our knowledge, no traffic-related mobile air pollution monitoring study has been published in Yaoundé city.

Due to the high cost of obtaining and maintaining air quality measuring equipment, the paucity of air pollution monitoring networks, even in the most crowded cities, can be compensated for by the installation of low- and medium-priced sensors offering easy and inexpensive alternatives. In contrast to industrialized countries, most developing-country cities lack the infrastructure to monitor urban air concentrations in order to effectively combat air pollution. Due to a lack of permanent monitoring stations for long-term measurements, financial resources for measurement campaigns, and information on air quality monitoring, air quality monitoring in Africa remains a challenge [11]. Routine monitoring of PM does not exist in most African cities and only a few studies have reported annual average levels of PM_{10} and $PM_{2.5}$. In most of the studies we found, monitoring campaigns were conducted for less than a year and often for various durations of less than 24 hours per day [12]. The deficiency of air pollution monitoring networks, even in the most urbanized cities, due to the high cost of acquisition and

maintenance of air quality measurement equipment, can be compensated by the installation of low- and medium-cost sensors offering simple and low-cost alternatives. Given the diversity of anthropogenic sources and their potency, which is expected to increase if no regulations are taken [13], there is a need to expand studies on pollution in African cities. Several works already exist on air quality in Africa [2, 3, 6]. Most of this work has been made possible by the use of less expensive measuring devices [6, 14].

Although accessible, these sensors cannot be deployed always and everywhere. To overcome this problem, an existing experimental methodology based on the use of low-cost sensors in intensive measurement campaigns and a geostatistical processing tool is used to assess particulate air pollution in Yaoundé. In this study, the data were collected in six crossroads of the city of Yaoundé during the month of August 2022. The work of Choudhary. A et al [15], shows that the crossroads are considered as hotspots in urban areas. The different measurement sites were chosen in the interest of having an idea of the level of particulate matter concentrations in the city of Yaoundé. Specifically, this study quantifies the level of PM₁₀, PM_{2.5} and PM₁ concentrations and their spatial and temporal distribution and assesses the traffic-related air pollution on road-intersections in the city of Yaoundé, Cameroon.

2. MATERIAL AND METHODS

2.1 Study area

The capital of Cameroon is the city of Yaoundé (3.52° N, 11.31° E). It covers an area of 180 km² for a population that has grown from 1.8 million inhabitants in 2006 to 4.3 million inhabitants in 2020 [16]. According to Matchebou et al. [17], population growth has happened concurrently with vehicle fleet growth, which is predicted to have increased from 152,000 vehicles in 2006 to 216,000 in 2015. Passenger cars, buses, coaches, trucks and vans, trailers, semi-trailers, and motorcycles comprise the vehicle fleet. Motorcycles are typically purchased new, whereas four-wheelers are typically purchased used. According to the Cameroonian Ministry of Transport, 92% of the Cameroonian car fleet in 2015 was made up of used vehicles that were more than 15 years old [3]. Yaoundé's traffic is congested, especially during rush hour, resulting in traffic bottlenecks. The latter increase journey time, resulting in greater energy consumption and considerable exhaust emissions. It should be mentioned that automobiles in Yaoundé run on fossil fuels such as gasoline and diesel. The studies will be conducted within the framework of this study in the city of Yaoundé's major crossroads, which are areas with high traffic and population density. This is the poste centrale (3°51'38"N, 11°31'13"E), du carrefour Mvog Mbi (3°51'04"N, 11°31'18"E), palais de justice(3°51'32"N, 11°31'05"E), Carrefour Hylton (3°51'53"N, 11°30'58"E), carrefour éducation (3°53'25"N, 11°30'17"E) and Camair (3°51'54"N, 11°31'18"E).



Fig.1. The location of the data collection site

2.2. Data collection and other material needed

In this study, a simple, low-cost mobile air quality testing instrument named "Air master" was utilized to acquire all of the particle concentration data required. At each measurement station, particulate matter was monitored for three days. A GPS was utilized to obtain the geographical coordinates of the various measurement stations (Fig.2b); the Extech (thermo-anemometer) was used to detect the wind direction (Fig.2c). The Air master additionally measures temperature and relative humidity (Fig.2a). The wind speed was measured using an anemometer (Fig.2d). The measuring campaign was carried out in the city of Yaoundé during the month of August 2022, where we took measurements on three separate city intersections from 7 a.m. to 6 p.m. (Fig.1).



Fig. 2. The different devices needed for this study (a) Air analyser and charger, (b) Garmin type GPS, (c) Exttech, (d) Anemometer.

The acquired raw data were then organised in Excel software. SPSS was used to evaluate the categorized data, which was then reported as tables and graphs using descriptive statistics (central and dispersion indexes).

2.3. Meteorological data and surrounding pollution

Temperature, relative humidity, and wind velocity are evaluated at local scales to assess the effects of meteorological conditions on PM concentration levels observed at each site throughout the sample period (Table 1). The wind rose allows you to choose your favourite wind direction at each of the six collection stations. The winds blow northwest at Camair and Poste Centrale. The wind is blowing from the east at the Mvog-Mbi interchange, while it is blowing from the south-east at the Education interchange. There are two preferred directions for the two remaining sites (Hilton junction and Palais de Justice), South-East and North-East.

After analyzing the field data, it was discovered that the winds at the Education junction have a low percentage of calm wind, i.e. 44.44%, compared to the Mvog-Mbi junction, which has 72%. This is due to the fact that the carrefour Education is placed at a higher height than the other sites.

Table 1. Average variation of some meteorological parameters at the different measurement locations (August 2022)

Sites of measurement	Relative humidity (%)	Temperature (°C)	Wind Speed (Km/h)
Carrefour Education	75.33 ± 31	24.92 ± 16	2.61 ± 0.32
Hilton	76.17 ± 19	24.69 ± 31	2.31 ± 0.22
Palais de Justice	76.27 ± 32	24.62 ± 10	1.68 ± 0.72
Camair	76.65 ± 18	25.47 ± 16	1.73 ± 0.2
Poste centrale	77.65 ± 39	25.51 ± 05	1.81 ± 0.39
Carrefour Mvog Mbi	71.63 ± 23	26.78 ± 21	1.51 ± 0.52

UNDER PEER REVIEW




Fig.3. Wind rose : (a) carrefour Hilton, (b) carrefour Mvog-Mbi, (c) Palais de Justice, (d) Camair, (e) Poste Centrale, (f) carrefour Education

2.4. Spatial mapping

Geospatial Information Systems (GIS) use interpolation to provide a robust method of geographical analysis. In previous publications, the most generally utilized methods, kriging (geostatistical method) and reverse weighting of IDW (deterministic method) have been compared [8]. The megacity of Delhi revealed that kriging and IDW approaches are linked with errors of approximately 22% and 24%, respectively [18].

Furthermore, as part of the DACCIWA Program, Bahino et al. [19] investigated the geographical distribution of gaseous pollutants (NO₂, SO₂, NH₃, HNO₃, and O₃) in Abidjan (Ivory Coast) and Cotonou (Benin) during the dry season. The later study compared kriging with IDW and discovered that the IDW method is associated with reduced uncertainty than the kriging method.

Furthermore, our measurement network (number of points and geographic locations) is based on the network of Bahino et al. [19]; thus, in this study, the IDW method is chosen and applied to the acquired data. ArcGIS version 10.3 software (ESRI Inc.) is utilised since it integrates geostatistical data processing and map editing. In brief, IDW interpolation predicts concentrations around a measured point using the technique shown in Eq (1).


$$Z_{p_j} = \sum_{i=1}^N \lambda_i \times Z_{0_i} \quad (1)$$

Where Z_{0_i} is the observed value at position i form geometrical mean of measured values at each site during the different measurement campaigns. Z_{p_j} is the expected value for position j ; N is the number of measured sample points that will be considered in the prediction around the expected location; λ_i are weighting factors used for each measured value at location. They decrease as the separation distance increases.

3. RESULTS AND DISCUSSION

3.1. Hourly variations of measured particulate matter concentrations

Because of their growing impact on human health, particularly respiratory disease and cardiovascular difficulties, PM_{2.5} pollutants have recently become the focus of intensive research worldwide. The current study considers the importance of monitoring ambient air quality because acute particle pollution can cause poor visibility and a significant increase in respiratory illnesses. The experimental details revealed a similar pattern in PM_{2.5} concentrations as in PM₁. By 11 a.m., the concentrations reached their peak in Camair (230.50 µg/m³) (Fig.4a and Fig.4b). Figure 4 depicts the various variances based on the hours for the other sites.



UNDER F...





Fig. 4. Hourly average variation of PMs at different measurement sites (a) PM_{10} ;(b) $PM_{2.5}$ and (c) PM_1

Because the particles are similar in nature due to their larger size when compared to the particle size already considered, the results for PM_{10} and TSP have been considered in the same section. While the bulk of these pollutants is smaller particles, the scenario for PM_{10} and TSP is similar. Most research on PM_{10} and TSP has found that $PM_{2.5}$ particles account for about 38-77% of the proportion of these two pollutants [20]. These contaminants are greater in size than other particles and, as a result, can be deposited due to their increased dispersion rate and deposition capacity. The movement of automobiles causes a steady dispersion of dust from the road surface into the surrounding air. In August, the relative humidity of the air is higher, causing these contaminants to stagnate in the air. Furthermore, fog in the winter provides these toxins with a bigger surface area to persist in the air. Camair ($220 \mu\text{g}/\text{m}^3$) had the highest maximum average for PM_{10} , followed by Carrefour Mvog Mbi ($175 \mu\text{g}/\text{m}^3$), Hilton ($1104.40 \mu\text{g}/\text{m}^3$), Poste Centrale ($140 \mu\text{g}/\text{m}^3$), Carrefour Education ($135 \mu\text{g}/\text{m}^3$), and Palais de Justice ($85 \mu\text{g}/\text{m}^3$).

Data on particulate matter (PM_1) in the research area's ambient air quality, the data was statistically examined. According to the statistics, the highest average PM_1 concentration was found at Carrefour Mvog Mbi ($39.35 \mu\text{g}/\text{m}^3$) (Table 2). PM_1 particles come from a variety of natural and manmade sources, and they are the consequence of chemical transformations in the gas phase, heterogeneous, and multiphase reactions [21]. Because of their small size, these particles can travel a greater distance and stay in the lower atmosphere for a longer amount of time [22]. Figure 5 illustrates a box plot of the variance of each particle size at the various measurement locations; it is clear that the roundabout Mvog Mbi has higher particle concentrations than all other areas where monitoring was conducted.




Fig. 5. Box plot of concentrations for different measurement site: PM₁₀ (a); PM_{2.5} (b) and PM₁ (c)

Table2: Average variation of PMs at the respective measurement sites

Sites of measurement	PM10 [Min-Max] ($\mu\text{g}/\text{m}^3$)	PM2.5 [Min-Max] ($\mu\text{g}/\text{m}^3$)	PM1 [Min-Max] ($\mu\text{g}/\text{m}^3$)
Carrefour Education	77.20 \pm 22.29 [52-135]	71.42 \pm 30.36 [48-162]	28.45 \pm 6.89 [17-44]
Hilton	72.77 \pm 42.46 [35-171]	60.38 \pm 36.48 [33-144]	31.17 \pm 07.39 [20-48]
Palais de Justice	59.91 \pm 12.26 [36-78]	54.91 \pm 14.31 [38-80]	33.96 \pm 13.59 [21-69]
Camair	78.07 \pm 55.93 [31-212]	79.00 \pm 67.89 [28-231]	29.02 \pm 16.19 [17-79]
Poste centrale	78.96 \pm 30.32 [45-140]	71.38 \pm 31.61 [36-125]	32.70 \pm 09.24 [23-50]
Carrefour Mvog Mbi	126.69 \pm 41.02 [63-175]	121.51 \pm 31.02 [65-164]	39.35 13.71 [27-80]

3.2. $\text{PM}_{2.5}/\text{PM}_{10}$, $\text{PM}_1/\text{PM}_{10}$ and $\text{PM}_1/\text{PM}_{2.5}$ ratio distribution

Table 3 shows the ratio of particulate matter sizes at the various measurement sites. The ratio's high values suggest that the majority of PM pollution is caused by combustion sources, while low values show that air pollution is caused by natural and dust sources [4]. As a result, the larger the ratio, the greater the dangers of PM pollution. The obtained ratio values were interpreted for various site appreciation scenarios. Figure 6 informs us that the highest ratio values are those of $\text{PM}_{2.5}/\text{PM}_{10}$ which is about 0.89 ± 0.09 . This number is less than one (1). Because this value is less than one (1), we can conclude that road dust is the source of the majority of particle pollution. This resuspension can contribute as much as to 50% of the particle emissions from road traffic [23]. Despite this, the high range of $\text{PM}_{2.5}$ and PM_{10} (fine and coarse particles) ratios examined in this study was $0.79 \pm 0.09 - 0.89 \pm 0.09$, which was higher than the WHO guideline standard of 0.5 - 0.80 [24]. This ratio is nevertheless a health risk for pregnant women and their foetus, as well as infants under the age of five, whose respiratory airways have high surface surfaces and absorption capacity for fine particulate matter.

Table 3. $PM_{2.5}/PM_{10}$, PM_1/PM_{10} and $PM_1/PM_{2.5}$ ratio values

Sites of Measurement	$PM_{2.5}/PM_{10}$ [Min-Max]	PM_1/PM_{10} [Min-Max]	$PM_1/PM_{2.5}$ [Min-Max]
Carrefour Education	0.86 ± 0.06 [0.70-0.95]	0.40 ± 0.16 [0.17-0.85]	0.44 ± 0.18 [0.17-0.92]
Hilton	0.81 ± 0.66 [0.71-0.96]	0.50 ± 0.24 [0.16-0.94]	0.62 ± 0.25 [0.16-0.93]
Palais de Justice	0.79 ± 0.09 [0.58-0.91]	0.53 ± 0.18 [0.30-0.99]	0.60 ± 0.19 [0.30-0.96]
Camair	0.79 ± 0.11 [0.61-0.95]	0.45 ± 0.16 [0.12-0.68]	0.51 ± 0.22 [0.11-0.79]
Poste centrale	0.83 ± 0.66 [0.67-0.92]	0.43 ± 0.11 [0.32-0.68]	0.50 ± 0.16 [0.26-0.82]
Carrefour Mvog Mbi	0.89 ± 0.09 [0.64-0.97]	0.33 ± 0.13 [0.17-0.59]	0.34 ± 0.14 [0.17-0.68]





Fig. 6. Ratio of particles in the different measurement locations

3.3. Pollution spatial mapping

Figure 7 represents the contours of the particulate matter concentration values obtained from extrapolation of the Air Master readings. The scale bar depicts nine distinct color concentrations going from less dense (blue) to denser (pink). The geostatistical distribution of PM_{10} is depicted in Figure 7a. According to the colors on the map in Figure 7a, the most polluting location is the carrefour Mvog-Mbi, which has concentrations up to $180 \mu\text{g.m}^{-3}$, followed by the carrefour education, which has a concentration of around $135 \mu\text{g.m}^{-3}$. Concerning Figure 7a, the least dense areas in comparison to the other two are Poste centrale, Hilton, Palais de Justice, and Camair, with concentrations ranging from 60 to $100 \mu\text{g.m}^{-3}$. Figure 7b shows that the Mvog-Mbi and Carrefour Education crossroads had the highest $PM_{2.5}$ concentration thresholds as compared to other sites, with values ranging from 140 to $170 \mu\text{g.m}^{-3}$. Nevertheless, in terms of PM_1 , the Mvog-Mbi crossroads remains the location with the highest pollutant threshold, around $48 \mu\text{g.m}^{-3}$, followed by Carrefour Hilton with a concentration of $37 \mu\text{g.m}^{-3}$ (Figure 7c).







Fig. 7. Spatial variation of the relative difference between (a) PM_{10} , (b) $PM_{2.5}$ and (c) PM_1 in Yaoundé (Cameroon), the black triangle representing each measurement site

In general, the Mvog-Mbi intersection has the highest concentrations of PM_{10} , $PM_{2.5}$, and PM_1 . This is understandable given that the Mvog-Mbi crossroads is in the heart of a market. We know that markets, such as carpentry, painting houses, road traffic, wood burning for catering, and so on, are sources of particulate matter.

4. CONCLUSION

Particulate matter in the environment is a severe environmental concern to human health, particularly for vulnerable groups such as children. We used a real-time PM detector to quantify ambient particulate matter concentrations such as PM_1 , $PM_{2.5}$, and PM_{10} in this investigation (Air Master). Because of the high traffic density at these spots, the results suggest that traffic intersections are the most polluted places in urban regions. Because of the high traffic density, the intersections can be considered hotspots. The varied particle ratios indicate that the majority of the fine particles originate from dust sources. Turbulence caused by vehicle movement resuscitates this dust. According to Rehn et al. [23], road traffic contributes up to 50% of particle emissions. According to this study, the Carrefour Mvog Mbi intersection is one of the most polluted in the city of Yaoundé; this high concentration might be justified by the closeness of a market near this crossroads. The closeness of a market near this intersection can explain the high concentration. At the same time, the markets are brimming with alternative particle sources (crushing machines, carpentry, painting, etc.). During the research period, however, mean PM_{10} and $PM_{2.5}$ concentrations surpassed the WHO air quality guideline and national standard.

REFERENCES

1. Matz C.J., E. M., Hocking R., Seenundun S., Charman N. and Edmonds N (2019). Human health effects of traffic-related air pollution (TRAP): a scoping review protocol. *Systematic Reviews*, 8(1), 223. doi: DOI: 10.1186/s13643-019-1106-5
2. Houngbégnon, P., Ayivi-Vinz, G., Lawin, H., Houessionon, K., Tanimomon, F.,

- Kêdoté, M., Agueh, V. (2019). Exposure to PM 2.5 Related to Road Traffic: Comparison between Crossroads and Outside of Crossroads at Cotonou, Benin. *Open Journal of Air Pollution*, 8, 108-117.
3. Adiang, C. M., Monkam, D., Lenouo, A., Njeugna, E., & Gokhale, S. (2017). Evaluating impacts of two-wheeler emissions on roadside air quality in the vicinity of a busy traffic intersection in Douala, Cameroon. *Air Quality, Atmosphere & Health*, 10(4), 521-532.
 4. Nasser Nducol, Y. F. T. S., Susan Younui Yakum-Ntaw, Saidou, Joseph Dika Manga, Juliette Cathérine Vardamides, Yerima Abba Hamadou & Augustin Simo. (2020). Ambient air particle mass concentrations in the urban area of the capital city of Yaoundé (Cameroon, Central Africa): monthly and seasonal behaviour. *International Journal of Environmental Analytical Chemistry*. doi: 10.1080/03067319.2020.1715378
 5. Pope III, C Arden Schwartz, Joel Ransom, & R. M. (1992). Daily mortality and PM10 pollution in Utah Valley. *Archives of Environmental Health: An International Journal*, 47(3), 211-217.
 6. Ngo, N. S., Asseko, S. V. J., Ebanega, M. O., Allo'o Allo'o, S. M., & Hystad, P. (2019). The relationship among PM2.5, traffic emissions, and socioeconomic status: Evidence from Gabon using low-cost, portable air quality monitors. *Transportation Research Part D: Transport and Environment*, 68, 2-9. doi: <https://doi.org/10.1016/j.trd.2018.01.029>
 7. Senarathna M., P. S., Jayaratne R., Weerasooriya R., Morawska L. and Bowatte G. (2022). Measuring Traffic-Related Air Pollution Using Smart Sensors In Sri Lanka: Before And During A New Traffic Plan. *Geography, Environment, Sustainability*, 3(15), 27-36.
 8. Sylvain Gnamien, Véronique Yoboué, Cathy Liousse, Money Osohou, Sékou Keita, Julien Bahino, . . . Diaby, L. (2020). Particulate Pollution in Korhogo and Abidjan (Cote d'Ivoire) during the Dry Season. *Aerosol and Air Quality Research*, 21. doi: <https://doi.org/10.4209/aaqr.2020.05.0201>
 9. Ngangmo YC, Adiang CM , Choudhary A & M. D. (2023) Road traffic-induced particle matter dispersion in a calm wind environment at the main

roundabout in Douala, central Africa. *Journal of Air Pollution and Health*, 8(1), 59-76.

10. Antonel, J., & Chowdhury, Z. (2014). Measuring ambient particulate matter in three cities in Cameroon, Africa. *Atmospheric Environment*, 95, 344-354.
11. Amegah, A. K., & Agyei-Mensah, S. . (2017). Urban air pollution in Sub-Saharan Africa: Time for action. *Environmental Pollution*, 220, 738-743.
12. Elisaveta, P., Darby, J., Nicole, V. C., & Patrick, K. (2013, September). Particulate Matter Monitoring in Urban Africa: A review. *In ISEE Conference Abstracts*.
13. Liousse, C., & Galy-Lacaux, C. (2010). Pollution urbaine en Afrique de l'Ouest. *La Météorologie*.
14. Sow, B., Tchanche, B., Fall, I., Souaré, S., & Mbow-Diokhané, A. . (2021). Monitoring of Atmospheric Pollutant Concentrations in the City of Dakar, Senegal. *Open Journal of Air Pollution*, 10(01), 18.
15. Choudhary, A., and Gokhale, S. (2019) On-road measurements and modelling of vehicular emissions during traffic interruption and congestion events in an urban traffic corridor. *Atmospheric Pollution Research*, 10(2), 480-492.
16. <https://www.populationdata.net/pays/cameroun/>
17. Matchebou, A., Yamba, J., & Tatiéte, T. T. . (2008). Impact du parc automobile sur la congestion du trafic et la pollution de l'air dans la ville de Yaoundé. *Ghardaïa, Algeria*, 245.
18. Shukla, K., Kumar, P., Mann, G.S., Khare, M. . (2020). Mapping spatial distribution of particulate matter using Kriging and Inverse Distance Weighting at supersites of megacity Delhi. *Sustainable Cities Soc* 54,, 101997. doi: <https://doi.org/10.1016/j.scs.2019.101997>
19. Bahino, J., Yoboué, V., Galy-Lacaux, C., Adon, M., Akpo, A., Keita, S., Liousse, C., Gardrat, E., Chiron, C., Osohou, M., Gnamien, S., Djossou, J. . (2018). A pilot study of gaseous pollutants measurement (NO₂, SO₂, NH₃, HNO₃ and O₃) in Abidjan, Côte d'Ivoire: contribution to an overview of gaseous pollution in African cities. *Atmos. Chem. Phys*, 18, 5173–5198. doi: <https://doi.org/10.5194/acp-18-5173-2018>
20. Kulshrestha, A., Satsangi, P., Masih, J., & Taneja, A. . (2009). Metal concentration

of PM_{2.5} and PM₁₀ particles and seasonal variations in urban and rural environment of Agra, India. *Science of the total environment*, 407(24), 6196–6204. doi: <https://doi.org/10.1016/j.scitotenv.2009.08.050>

21. Hamed, A., Birmili, W., Joutsensaari, J., Mikkonen, S., Asmi, A., Wehner, B. (2010). Changes in the production rate of secondary aerosol particles in Central Europe in view of decreasing SO₂ emissions between 1996 and 2006. *Atmospheric Chemistry and Physics*, 10(3) 1071–1091. doi: <https://doi.org/10.5194/acp-10-1071-2010>
22. Kappos, A. D., Bruckmann, P., Eikmann, T., Englert, N., Heinrich, U., Höppe, P & Wichmann, H. E. (2004). (2004). Health effects of particles in ambient air. *International journal of hygiene and environmental health*, 207(4), 399-407. doi: <https://doi.org/10.1078/1438-4639-00306>
23. Rehn, L. P., A. Waked, A. Charron, C. Piot, J.-L. Besombes, N. Marchand & G. Guillaud Estimation de la contribution des émissions.
24. Kanee, R. B., Adeyemi, A., Edokpa, D.O., and Ede, P.N. (2020). Particulate Matter-Based Air Quality Index Estimate for Abuja, Nigeria: Implications for Health. *Journal of Geoscience and Environment Protection*, 8, 313-321. doi: <https://doi.org/10.4236/gep.2020/85019>