

“Effect of Poultry manure, Vermicompost and Boron on growth and yield of Maize”

ABSTRACT:

A field experiment titled “Effect of Poultry Manure, Vermicompost and Boron on Growth and Yield of Maize” (*Zea mays* L.) was conducted during *Rabi* 2022 at Crop Research Farm, Department of Agronomy, Naini Agriculture Institute SHUATS, Prayagraj Uttar Pradesh. The experiment was laid out in Randomized Block Design with ten treatments which are replicated thrice. Results obtained that combined application of Poultry Manure 50% (1t/ha) along with Vermicompost 50% (2 t/ha) and Boron – 5% (Treatment) significantly increased higher plant height (234.45 cm), dry weight (135.84 g/plant) and also yield attributes like effective cobs per plant (2.00), Seeds per cob (302.67), Seed index (24.69 g), Grain yield (6.73t/ha), stover yield (15.20 t/ha).

Keywords: *Maize, vermicompost, Poultry manure, boron, growth parameters, yield attributes.*

INTRODUCTION:

Maize (*Zea mays* L.) is one of the most versatile crops grown throughout the tropical as well as temperate regions of the world. Every month of the year, a crop of maize is planted and harvested somewhere in the world. It is sometimes referred to as the "QUEEN OF CEREALS" since no other cereal on the planet has such enormous potential. After rice and wheat, it is the third most significant cereal crop in the world. India ranks eighth in the world for maize production and fifth for acreage. From loamy sand to clay loam, maize can be cultivated successfully in a variety of soil types. Currently, 170 nations produce maize, with an average global productivity of 5.75 t/ha. This results in a total production of around 1147.7 million metric tonnes of maize from an area of 193.7 million hectares. (FAOSTAT, 2020).

India accounts for around 4% of the world's total maize acreage and 2% of its production, placing it fourth in both area and production among the nations that cultivate maize. It occupies an area

of 9.38 million ha, contributes almost 9% of the country's food supply, produces 28.8 million tonnes annually, and has an average yield of 3.07 tonnes per hectare (**Anonymous 2018**). According to anonymous (2016), Himachal Pradesh has 294 thousand ha, 784.29 thousand tonnes, and 2.68 tonnes /ha of land. Maize has greater nutritional value as it contains about 72% starch, 10% proteins, 4.8% oil, 8.5%, fiber, 3% sugar and 1.7% ash (**Chaudhary, 1993**). A substantial amount of maize is utilized as feed for poultry and cattle. 61% of maize produced worldwide is used as animal feed, 17% as food, and 22% as industrial raw material. Because of its significant involvement (83%) in the feed, starch, and biofuel industries, it has become a recognized "industrial crop" on a global scale.

The availability of boron (B), an essential nutrient for optimal growth of higher plants, in the soil and irrigation water is a key factor in determining agricultural production. **Saleem et.al., (2011)**. B shortage has various consequences on a wide range of vascular plant functions, including root extension, indole acetic acid oxidase activity, sugar translocation, carbohydrate metabolism, nucleic acid synthesis, and pollen tube formation (**Goldbach and Wimmer, 2007; Saleem et.al., 2011**). When plants were exposed to salinity, their B concentration reduced (**Holloway and Alston, 1992; Wimmer et.al., 2001**).

Poultry manure is an excellent organic fertilizer, as it contains high nitrogen, phosphorus, potassium and other essential nutrients. When compared to chemical fertilizer, it enhances soil structures, nutrient retention, aeration, soil moisture holding capacity, and water infiltration by adding organic matter to the soil (**Deksissa et.al., 2008**). Additionally, it was found that compared to other organic manure sources, poultry dung more readily provides P to plants (**Garg and Bahla, 2008**). Poultry manure is a valuable fertilizer and can serve as a suitable alternate to chemical fertilizer. Poultry manure application registered over 53% increases of N level in the soil, from 0.09% to 0.14 % and exchangeable cations increase with manure application (**Boateng et.al., 2006**). In agriculture, the main reasons for applying PM include the organic amendment of the soil and the provision of nutrients to crops (**Warren et.al., 2006**).

Vermicompost is an excellent alternative to synthetic fertilizers since it contains more N, P, and K than regular heap manure (**Srivastava and Beohar, 2004**). Vermicomposting aids in enhancing and preserving the soil's fertility. Vermicompost gives the soil a black hue and helps to regulate the soil's temperature by doing so. The of vermicompost helps to improves and conserves the fertility of soil. Vermicompost imparts a dark colour of the soil and thereby help to maintain the temperature of soil. Vermicompost is one of the manure used by the farmer in growing crops because of early availability and presence of almost all the nutrients required by plants.

MATERIAL AND METHODS:

The experiment conducted to know the “Effects of Poultry manure, Vermicompost and Boron on growth and yield of Maize” was carried out at Crop Research Farm of Sam Higginbottom University, Prayagraj, Uttar Pradesh in 2022. The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 7.4), low level of organic carbon (0.51%), available N (108.69 Kg/ha), P (80.5 kg/ha), K (83.3 kg/ha). The treatments combination are T1: Poultry Manure - 100% (2 t/ha) + Boron - 1%, T2: Poultry Manure - 100% (2 t/ha) + Boron - 3%, T3: Poultry Manure - 100% (2 t/ha) + Boron -5%, T4: Vermicompost - 100% (4 t/ha) + Boron - 1%, T5: Vermicompost - 100% (4 t/ha) + Boron - 3%, T6: Vermicompost - 100% (4 t/ha) + Boron - 5%, T7: Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 1%, T8: Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3%, T9: Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%, T10: Control – 120:60:40 (N: P: K) Kg/ha are used. To facilitate sowing, the experimental field was thoroughly ploughed and harrowed and brought to fine tilth. Stubbles and weeds were picked up from the field and the land was levelled with the help of a rake and the plots were demarcated according to layout. Manures were applied at 4-5cm deep furrows were made along the seed rows with a hand hoe. Manures used for application were vermicompost, and neem cake Quantity of each manure required for each plot was calculated as per the treatment combinations. Manures like Vermicompost and neem cake were applied uniformly in seed furrows as basal dose. Maize, Variety (NK 7720) was selected for sowing which can be matured around 85- 90 days. Line sowing was done manually. Seeds were covered with soil immediately after sowing the seeds. The sowing adopted was (60 x 20 cm) and the seeds were drilled at 3-4 cm depth.

RESULTS AND DISCUSSION:

Plant height: At 100 DAS there was a significant difference among the treatments. However, maximum plant height (234.45 cm) was observed in the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% and Poultry Manure - 100% (2 t/ha) + Boron - 3% (233.16 cm), Poultry Manure - 100% (2 t/ha) + Boron - 5% (229.94 cm) were found statistically at par with Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%. The maximum value of plant height was achieved at higher uptake of B due to which translocation of photosynthesis was improved from leaf to grain Akhtar *et.al.*, (2003). The increase in plant height with PM was mainly due to the reason of more availability of nutrients by PM throughout the growing season. Warren *et.al.*, (2006). Vermicompost increases in plant height and LAI with the use of organic sources consequently enhanced the dry matter/plant Jayaprakash *et.al.*, (2004)

Plant Dry weight: maximum plant dry weight (135.84 g) was observed in the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%. However, in Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% (133.50 g) was statistically at par with Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron –5%. **Ahmed *et.al.*, (2011)** also reported that dry matter yield increased significantly with B up to 2.0 kg/ha. Exogenous application of B reportedly responded to increasing plant dry matter (**Oyinlola 2007**). Increased biomass production by the application of boron is thought due to the role of boron in cell elongation, photosynthesis and transpiration.

Number of cobs per plant: Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% (2.00) was recorded maximum significant value of the number of cobs per plant and the minimum was recorded in control (1.28). However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% (2.00) was statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%. The application of boron activates several enzymes that are involved in carbohydrate metabolism, protein synthesis and pollen formation **George *et.al.*, (2002)**.

Seed index (g): Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% (24.69 g) was recorded maximum significant value of test weight and the minimum was recorded in control (21.48 g). However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% (24.23 g), was statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%.

Zhang *et.al.*, (1998) who observed increase in grain weight with increased level of Poultry manure which could be due to balanced supply of food nutrients throughout development of the maize plant and concluded that maximum grain weight (g) can be obtained by the high rate of poultry manure application.

Number of Seeds per Cob: Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% (302.67) was recorded maximum significant value of the number of Seeds per cob and the minimum was recorded in control (221.33). However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% was statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%.

The increase in number of grains per row may be attributed to the availability of more nitrogen and other nutrients from **PM** required for plant development up to cob formation (**Farhad 2009**).

Grain yield (t/ha)

Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% (6.73 t/ha) was recorded higher significant value of grain yield. However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% (6.40 t/ha) were found statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%.

The boron application has been reported to increase grain yield up to 33% **Gunnes *et.al.*, (2003)**. **Rahim *et.al.*, 2004** reported increased dry weight plant, a number of grains cob and grain weight cob in maize plants which support the findings of the present study that boron application significantly increased various yield attributes of maize hybrids. Significantly increased with application of poultry manure which resulted in an overall increase in grain yield. (**Zhang *et.al.*, 1998**) who stated that maximum grain yield can be obtained by the high rate of poultry manure application.

Stover yield (t/ha)

Higher number of stover yield is in Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron–5% (15.20 t/ha) was recorded. However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron –1% (14.98 t/ha), Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 3% (15.11 t/ha) were found statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%.

Farhad *et.al.*, (2009) and Zhang *et.al.*, (1998) that the maximum biological yield can be obtained by the high rate of poultry manure application that due to the proper and balanced supply of nutrients to the plants throughout the growth period.

Harvest index (%)

Significantly higher harvest index is recorded in Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% of (30.68 %). However, Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 1% (29.43%), Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron –3% (29.75%) were found statistically at par with the Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5%.

Many researchers (**Renukadevi *et.al.*, 2003; Wrobel *et.al.*, 2006; Malhi *et.al.*, 2007**) suggested the application of boron for enhancing the yield of crop plants. Boron helps in better uptake and utilization of nitrogen which helps in enhancing the photosynthetic activity, vegetative growth, accumulation of metabolites in reproductive organs and the overall yield (**Rizk and Abdo 2001**).

Vermicompost is a good substitute to commercial fertilizers and has more N, P and K content than the normal heap manure (**Srivastava and Beohar, 2004**). The increase in grain and stover yield, weight of cob/ plant, number of grains /cob and test weight of maize due to application of vermicompost might be attributed mainly to higher content of available nutrients in vermicompost, presence of beneficial micro flora such as nitrogen fixers, phosphate solubalizers, VAM fungi and higher activity of dehydrogenase enzyme in soil. The finding of this investigation close conformity with finding of **Ramesh *et.al.*, (2008)** and **Meena *et.al.*, (2013)**.

Poultry manure is rich source of nutrients like Nitrogen and Phosphorous and thus improving soil properties, enhancing availability of nutrients and thereby important for development. optimum plant growth and development **Dauda *et.al.*, (2008)**

CONCLUSION

In the light of above study, it is concluded that maize yield is increased with application of treatment 9 Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron – 5% and this treatment has proved to be better option for getting higher productivity. Since the findings are based on research done in one season in Prayagraj it may be repeated for confirmation

References:

1. Anonymous. 2018. IndiaAgriStat.com; <https://eands.dacnet.nic.in> (8t h March, 2021).
2. Ahmed N, Abid M, Ahmad F, Amanullah M, Javaid Q, AliMA (2011) Impact of boron fertilization on dry matter production and mineral constitution of irrigated cotton. Pak J Bot 43 (6): 2903—2910.
3. Boateng, A.S., J. Zickermann and M. Kornahrens. 2006. Poultry manure effect on growth and yield of maize. West Africa J. App. Eco., 9: 1-11.
4. Chaudhary, A.R.1993. Maize in Pakistan. Punjab Agriculture Coordination Board, Univ. Agric., Faisalabad.
5. Dauda, S. N., F. A. Ajayi and E. N. Dor. 2008. Growth and yield of watermelon (*Citrulluslunatus*) as affected by poultry manure application. J. Agric. & Social Sci., 4 (3): 121 124.

6. Deksissa T., I. Short and J. Allen (2008). Effect of soil amendment with compost on growth and water use efficiency of Amaranth. In: Proceedings of the UCOWR/NIWR annual conference:
7. FAOSTAT. 2020. Crops. <http://www.fao.org/faostat/en/#data/QC> (22nd December 2020). International water resources: challenges for the 21st century and water resources education, July 22 – 24, 2008, Durham, NC.
8. Farhad, W., M. F. Saleem, M. A. Cheema and H. M. Hammad. 2009. Effect of poultry manure levels on the productivity of spring maize (*Zea mays* L.) *The J. Ani. & Plant Sci.*, 19(3): 122-125.
9. Garg, S and G.S. Bahla. 2008. Phosphorus availability to maize as influenced by organic manures and fertilizer P associated phosphatase activity in soil. *Bio-resource Tech.*, 99(130): 5773-5777.
10. Güneş, A., M. Alpaslan, A. Inal, M. S. Adak, F. Eraslan and N. Çiçek. 2004. Effects of boron fertilization on the yield and some yield components of bread and durum wheat. *Turkish Journal of Agriculture and Forestry*, 27: 329-335.
11. Jayaprakash, T.C., Nagalakar, V.P., Pujari, B.T. and Shetty, R.A. 2004. Dry matter and its accumulation pattern in maize as influenced by organics and inorganics. *Karnataka Journal of Agricultural Sciences* 17(2): 327–29.
12. Malhi SS, Gan Y and Raney JP. 2007. Yield, seed quality, and sulfur uptake of Brassica oilseed crops in response to sulfur fertilization. *Agronomy Journal* 99: 570- 577
13. Meena, B.P., Kumar, A., Meena, S.R., Dhar, S., Rana, D.S. and Rana, K.S. 2013. Effect of different organic sources and levels of nutrients on growth and yield behaviour of popcorn (*Zea mays*) and potato (*Solanum tuberosum*) sequence. *Indian Journal of Agronomy*, 58(4): 474-479.
14. Oyinlola E.Y., (2007) Effect of boron fertilizer on yield and oil content of three sunflower cultivars in the Nigerian savanna; *Journal of Agronomy* 6 (3): 421 – 426.
15. Rahim, M., H. Ali and T. Mahmood. 2004. Impact of nitrogen and boron application on growth and yield of maize (*Zea mays* L.) crop. *Journal of Research in Science*, 15: 153-157.
16. Ramesh, P., Singh, M. and Subbarao, A. 2008. Organic farming its relevance to the Indian context. *Current Science*, 88: 561-569.
17. Rizk WM and Abdo FA. 2001. The response of two moong bean cultivars to zinc, manganese and boron II. Yield and chemical composition of seeds. *Bulletin Faculty of Agriculture Cairo Univ.* 52: 467-477.

18. Renukadevi A, Savithri P and Andi K. 2003. Sources, levels and methods of boron application on the dry matter production, yield attributes and yield of sunflower (*Helianthus annuus*) crop. *Crop Research Hisar - The Pherobase* 25: 436-440.
19. Saleem M., Y.M. Khanif, F. Ishak, A.W. Samsuri and B. Hafeez, 2011. Importance of Boron for Agriculture Productivity: A Review *International Research Journal of Agricultural Sciences and Soil Sciences*. **1**(8): 293-300
20. Warren, J. G., S. B. Phillips, G. L. Mullins, D. Keahey and C. J. Penn. 2006. Environmental and production consequences of using alumamended poultry litter as a nutrient source for corn. *Journal of Environment. Quality.*, 35: 172-182.
21. Wimmer, M.A., K.H. Muehling, A. Lauchli, P.H. Brown and H.E. Goldbach, 2001. Interaction of Salinity and Boron Toxicity in Wheat (*Triticum aestivum* L.). In: *Plant Nutrition: Food Security and Sustainability of Agro Ecosystems Through Basic and Applied Research*, Horst, W.J., M.K. Schenk, A. Burkert, N. Claassen, H. Flessa, W.B. Frommer, H. Goldbach, H.W. Olf and V. Romheld (Eds.). Springer, Netherlands, ISBN: 978-0-7923-7105-2, pp: 426-427.
22. Zhang, H., D. Smeal and J. Tomko. 1998. Nitrogen fertilizer value of feed lot manure for irrigated corn production. *Journal of Plant Nutrition.*, 21: 287-296.

Table 1: Effect of Poultry manure, vermicompost and boron on growth and yield attributes of maize.

		100 DAS							
S. No	Treatments	Plant height (cm)	Plant dry weight	Cobs/ Plant (No)	Seeds/ Cob (No)	Seed index (g)	Grain yield (t/ha)	Stover Yield (t/ha)	Harvest index (%)
1.	Poultry Manure-100% (2 t/ha) + Boron 1%	226.80	130.83	1.60	232.67	21.80	4.82	13.84	25.83
2.	Poultry Manure-100% (2 t/ha) + Boron 3%	228.80	131.76	1.80	245.67	21.91	5.08	13.95	26.69
3.	Poultry Manure-100% (2 t/ha) + Boron 5%	227.30	132.28	1.80	253.33	22.01	5.27	14.29	26.94
4.	Vermicompost-100% (4 t/ha) + Boron 1%	224.60	129.66	1.21	260.67	22.61	5.52	14.46	27.62
5.	Vermicompost-100% (4 t/ha) + Boron 3%	224.60	129.95	1.24	271.67	23.02	5.83	14.57	28.57
6.	Vermicompost-100% (4 t/ha) + Boron 5%	226.90	131.25	1.21	277.00	23.64	6.05	14.82	28.98
7.	Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron 1%	230.94	132.17	1.90	284.67	23.92	6.25	14.98	29.43
8.	Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron 3%	234.90	133.50	2.00	293.60	24.23	6.40	15.11	29.75
9.	Poultry Manure 50% (1t/ha) + Vermicompost 50% (2 t/ha) + Boron 5%	236.45	135.84	2.00	302.60	24.69	6.73	15.20	30.68
10.	Control - 120: 60: 40 (N:P: K) kg/ha	223.43	127.95	1.60	232.67	21.80	4.69	13.72	25.47
F- test		S	S	S	S	S	S	S	S
SEm(±)		2.23	0.81	0.02	3.13	0.20	0.13	0.07	0.43
CD(p=0.05)		6.63	2.42	0.07	9.30	0.58	0.38	0.22	1.54

