

Original Research Article

**Effect of enriched TSP polymer seed coating on germination physiology of greengram  
Var. CO8**

**Abstract**

During 2022, the research was carried out under laboratory condition at the Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore, to find out the effect of TSP polymer seed coating on germination physiology of [greengramgreen gram](#). The seeds were coated with TSP polymer, TSP polymer with preservatives, TSP polymer with PGRs and TSP polymer with preservatives and PGRs @ 6 g/ kg of seeds. Among the various constituents of polymer, TSP polymer with preservative (0.1% sodium sorbate) and PGR (1.5 ppm BRs)performed betterwith good speed of germination and seedling vigour.

**Keywords:** TSP polymer, Sodium sorbate, GA<sub>3</sub>, BRs

**Introduction**

Green gram (*Vignaradiata*) is third important pulse crop in the country after redgram and [blackgramblack gram](#). [GreengramGreen gram](#) is an excellent source of high-quality protein (25%),Riboflavin. Thiamine and vitamin C with high digestibility. Being a leguminous crop has capacity to fix the atmospheric nitrogen (30-40 kg N/ha). India is the major producer of green gram in the world, and it is grown in almost all the states. [GreengramGreen gram](#) is mainly cultivated in rainfed condition in marginal lands mostly as an intercrop with less care and prone to number of disease and pest. All together causes hardship in maintaining population per unit area.Poor population lead to reduction in yield and might be the major cause for less productivity. There are number of seed management technique which augment the germination under wide range of environmental condition. Through which it is possible to maintain plant population per unit area there by chances for improving the yield.

Seed enhancement technique is thepost-harvest treatment which helps to improve the germination and growth of seedlings (Taylor *et al.*,1998). Many shotgun approaches are being used for seed enhancement for the last three decades, which include seed priming, seed pelleting, hardening, pregermination, magnetic stimulation, film coating, encrusting etc.,Seed coating refers to the application of nutrients, protectant and other ingredients to improve the

performance of seed in all the environmental condition (Ni, 1997). Polymer coating improves germination due to increase in the rate of imbibition where the fine particles in the coating act as a 'wick' or moisture attracting material which help the seed to establish a good contact with soil. Due to their exceptional qualities, such as good coating and barrier performance, biodegradability, and low weight, natural or bio polymers or plant-based seed coating formulations are recommended as suitable for continuous use. Biopolymer is a polymerized substance produced from natural resources, biological materials, or by living organisms. Superabsorbent hydrogels (SHs) made of polysaccharides have shown promise as alternatives since they are biopolymers that are abundant in nature, nontoxic, typically inexpensive, and inherently biodegradable and biocompatible (Kamath and Par, 1993).

Tamarind seed polysaccharide (TSP) is a polysaccharide explored widely as a bioadhesive. It is a mucilaginous polysaccharide derived from the seed kernel of *Tamarindus indica* Linn. belongs to the family Fabaceae. Tamarind seed polysaccharide (TSP) based polymer is considered ideal for formulating immediate and sustained release of plant stimulants and protectant due to their high thermal and hemo stability, safety, non-toxic, hydrophilic and gel forming nature (Baveja, 1988; Joseph *et al.*, 2012). Black gram seeds coated with TSP polymer (6g) + 0.3g humic acid + 0.2 g ascorbic acid + 0.5 ml of Zimmu leaf extract /kg performed better and recommended as presowing seed treatment (Sivasakthi *et al.*, 2022)

Formatted: Font: Italic

Plant Growth Regulators (PGRs) or phytohormones are defined as small, simple chemicals produced naturally by plants to regulate their growth and development. Among the phytohormones GA<sub>3</sub> was studied abundantly for their biological role in plants and referred as cell elongation hormone (Jones *et al.*, 1983). GA<sub>3</sub> regulates germination by counteracting ABA in seed.

Formatted: Font: Italic

Brassinosteroids (BRs) are a new type of polyhydroxy steroidal phytohormones with significant growth-promoting influence. It is essentially important for plant development and growth. BR signalingsignalling promote cell expansion and cell division, and plays a role in etiolation and reproduction. BRs also exhibit synergistic effect with other phytohormones. It cross ~~talk~~ talk with auxin, gibberellins, abscisic acid, ethylene, salicylic acid and jasmonic acid to promote plant growth and metabolism (Choudhary *et al.*, 2012)

One of the highly consumed food additives is food preservatives. Preservatives are used to increase the shelf life of foods by preventing microorganism growth and to control food spoilage (Kefi *et al.*, 2022, Zhang *et al.*, 2021). Sorbate and benzoates are the most common

preservatives that are used in a wide range of foods including mayonnaise and salad dressings (Amirpouret *et al.*, 2015, Chaleshtoriet *et al.*, 2018). Sodium sorbate (E 201), which is the sodium salt or sorbic acid, is widely used as food preservatives on cheese, meat, ketchup, mayonnaise and marmalade said by Potter 1984; Warth 1985. Revenlifornana and Nascimento (2019) reported that SMB (0.5% m/v) and combination of SMB (0.25% m/v) were the most effective in the controlling the browning in processed tomato.

## Materials and Methods

The laboratory experiment was conducted at the Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore. Green gram seeds var. CO 8 was procured from the Department of Pulses, Tamil Nadu Agricultural University, Coimbatore, TamilNadu. Tamarind seeds were procured from tamarind mandi<sup>y</sup> at Coimbatore, Tamil Nadu.

### Preparation of seed coating TSP polymer

TSP polymer was prepared according to Sivasakthiet *et al.*, (2022) which served as basic source for other treatments. To improve the shelf life and to enrich the polymer it was added with preservative ~~ana~~ PGR. For enriching, five sets of 6 g of polymer were weighed. First set was added with 0.12 ml of 0.1 % Sodium sorbate which served as T<sub>2</sub>, second set was added with 0.06 ml of 30 ppm GA<sub>3</sub>, which served as T<sub>3</sub>. Third set was added with 0.18 ml of 1.5 ppm BRs which served as T<sub>4</sub>, Fourth set was added with 0.12 ml of 0.1 % Sodium sorbate + 0.06 ml of 30 ppm GA<sub>3</sub> which served as T<sub>5</sub>, and final set was added with 0.12 ml of 0.1 % Sodium sorbate + 0.18 ml of 1.5 ppm BRs which served as T<sub>6</sub>.

After adding of preservative and PGRs, for proper mixing, the polymer was stirred in a mechanical stirrer for 20 min then preserved for further use.

### Seed treatment

From each treatments required quantity of TSP polymer (6 g/kg of seeds) was weighed and dissolved in 15 ml of distilled water (Sivasakthiet *et al.*, 2022). After dissolving the polymer, the seeds were coated with respective treatments. While coating, care was taken for uniform coating. After coating the coated seeds were dried for 5 hours.

### Treatment details

T<sub>0</sub>- Absolute control

T<sub>1</sub> - TSP polymer 6 g/kg of seeds

T<sub>2</sub> - TSP polymer 6 g + 0.12 ml of 0.1 % Sodium sorbate

T<sub>3</sub> - TSP polymer 6 g + 0.06 ml of 30 ppm GA<sub>3</sub>

T<sub>4</sub> - TSP polymer 6 g + 0.18 ml of 1.5 ppm BRs

T<sub>5</sub> - TSP polymer 6 g + 0.12 ml of 0.1 % Sodium sorbate + 0.06 ml of 30 ppm GA<sub>3</sub>

T<sub>6</sub> - TSP polymer 6 g + 0.12 ml of 0.1 % Sodium sorbate + 0.18 ml of 1.5 ppm BRs

The coated seeds along with control were subjected to germination study and the following observation viz., seed germination (%) (ISTA, 2013), speed of germination (Maguire, 1962), root and shoot length (Sivasakthiet *al.*, 2021), dry matter production (g seedlings<sup>-10</sup>) (ISTA, 2013) and vigour index II was computed using the following formula as per Abdul-Baki and Anderson, 1973.

Vigour index II = Germination (%) x Drymatter production (g seedling<sup>-10</sup>).

The data obtained were subjected to statistical analysis using Completely Randomized Block Design (CRD).

## Result

The results revealed that the seeds coated with TSP polymer + Preservatives and PGRs showed significant difference for seed germination and seedling vigour. Among all the treatments maximum speed of germination was observed in T<sub>4</sub>(25.05), it was on par with T<sub>3</sub> (24.98), T<sub>6</sub> (24.85), and T<sub>5</sub> (24.74). Though statistical difference was noticed for germination percentage, the numerical difference between treatments were minimum. It ranged from 90 (T<sub>0</sub>) to 96 (T<sub>4</sub> and T<sub>6</sub>) with mean of 94%. Treatments T<sub>3</sub>, T<sub>5</sub> (95%) and T<sub>4</sub> and T<sub>6</sub> (96%) were on par with each other; and control recorded significantly low germination which was on par with T<sub>2</sub> and T<sub>3</sub>(93%).

Maximum root length was observed in T<sub>6</sub> (20.09 cm) it was on par with all treatments except T<sub>0</sub>(16.79 cm). Maximum shoot length was observed in T<sub>6</sub> (21.40 cm) it was on par with T<sub>5</sub>(21.31), T<sub>4</sub>(21.04), T<sub>3</sub>(20.85), and T<sub>2</sub>(19.64) and T<sub>1</sub> (19.28).

Maximum dry matter production was observed in T<sub>6</sub> (0.193) followed by T<sub>5</sub>(0.192), T<sub>4</sub> (0.191) and T<sub>3</sub> (0.189), all the treatments were on par with each other. Maximum vigour index

was observed in T<sub>6</sub>,T<sub>5</sub>,T<sub>4</sub>,T<sub>3</sub>(18) and T<sub>2</sub> (17), It was on par with each other except T<sub>0</sub> (15) (Table 1).

**Table 1 : Effect of seed coating with TSP Polymer + Preservatives and Plant growth regulators on seed germination physiology of greengramgreen gram**

Treatment	Speed of germination	Germination (%)	Root length (cm)	Shoot length (cm)	DMP ( g seedlings <sup>-10</sup> )	Vigour index II
T0	21.84	90	16.79	18.08	0.165	15
T1	22.72	93	17.93	19.28	0.177	16
T2	22.80	93	18.36	19.64	0.180	17
T3	24.98	95	19.56	20.85	0.189	18
T4	25.05	96	19.68	21.04	0.191	18
T5	24.74	95	19.95	21.31	0.192	18
T6	24.85	96	20.09	21.40	0.193	18
<b>MEAN</b>	<b>23.85</b>	<b>94</b>	<b>18.17</b>	<b>19.46</b>	<b>0.177</b>	<b>17</b>
SED	0.3387	1.0437	2.2631	2.4368	0.0222	2.0310
CD (0.05)	0.7044	2.1705	4.7064	5.0675	0.0461	4.2237

Formatted: Font: Bold

T<sub>0</sub>- Absolute control, T<sub>1</sub>-TSP polymer (6 g/kg of seeds), T<sub>2</sub>-TSP polymer (6 g) + 0.12 ml of 0.1 % Sodium sorbate, T<sub>3</sub> - 0.06 ml of 30 ppm GA<sub>3</sub>, T<sub>4</sub> -0.18 ml of 1.5 ppm BRs T<sub>5</sub> - 0.12 ml of 0.1 % Sodium sorbate + 0.06 ml of 30 ppm GA<sub>3</sub>and T<sub>6</sub> - 0.12 ml of 0.1 % Sodium sorbate + 0.18 ml of 1.5 ppm BRs

### Discussion

High quality seeds plays major role in maximizing the productivity through assured rapid establishment under wide range of environmental condition. Most of the species are sensitive

to drought and water logging. The most affected stage is seed germination and seedling establishment. Seed germination can be ~~augumented~~ augmented through various seed enhancement techniques. Among the number of seed enhancement techniques, seed coating is considered as one of the best. Because it augments the germination without modifying the physiological status on the seeds, where other treatment like priming there was a threat of storing the seeds with good viability even for months

In this present study ~~greengram~~ green gram seeds coated with TSP polymer constituting preservative and plant growth regulator performed a little better than the control and TSP polymer alone coated seeds. Since in the study freshly harvested seeds with high germination percentage (90%) was used a jump in seedling character was not observed. The highest speed of germination is the indicator of high vigour. The speed of germination along with high germination percentage enable the seeds to establish faster and make the seedling to utilize the resource for further development and yield. The initial established vigour contributed a lot for yield and resultant seed quality. Dry matter production is an indicator of how tightly cells are packed in a tissue and number of cells, which contribute for biomass. In the present study seeds coated with T<sub>6</sub> and T<sub>4</sub> (96%) both showed high values for seed germination but considering all other factors viz., seedling length, dry matter production and vigour index T<sub>6</sub> excelled than T<sub>4</sub>.

The reason for improvement in germination and vigour attributes may be due to availability of carbon source, from the TSP polymer and phytohormone Brassinosteroids (BRs). Tamarind seeds are rich in polysaccharide, contains about 65% of ~~non-fibre~~ non-fibre carbohydrates; many essential amino acids, like isoleucine, leucine, lysine, methionine, phenylalanine and valine. In addition to this, seeds are also good source of essential fatty acids, minerals particularly calcium, magnesium, phosphorous and potassium (Joseph *et al.*, 2012). BRs regulate root meristem size and lateral root development in a concentration-dependent manner. A low concentration of BRs promotes root growth, whereas a high concentration of BRs inhibit root growth (Gupta *et al.*, 2015; Lee *et al.*, 2015). In the present study, besides improving root length, a ~~net-work~~ network of roots are visualized in T<sub>6</sub> (figure 1). The secondary roots may help the plants to absorb soil nutrients and contribute for better establishment, growth and yield.



**T<sub>0</sub> . Absolute control**

**T<sub>6</sub> – TSP polymer (6g) +  
0.18 ml of 0.15 ppm  
Brassinosteroid + 0.12 ml  
of 0.15 % sodium sorbate**

Fig 1 :A network of roots are visualized in T<sub>6</sub>

## Conclusion

Coating of seeds with TSP polymer (6g) + 0.18 ml ~~of 0.15~~of 0.15 ppm Brassinosteroid+ 0.12 ml ~~of 0.15 of 0.15% sodium~~sodium sorbate had a positive effect on seed germination and seedling vigour of ~~green gram~~green gram. It improves the seedling efficiency, qualitatively and quantitatively and also substitute the synthetic ~~polymers, that~~polymers, which in turns helps in reduction of environmental pollution.

## Reference

Abdul-Baki AA, Anderson JD. (1973). Vigour deterioration of soybean seeds by multiple criteria. *Crop Science*. ~~1973~~13:630-633.

Formatted: Font: Italic

Taylor, A., Allen, P., Bennett, M., Bradford, K., Burris, J., & Misra, M. (1998). Seed enhancements. *Seed Science Research*, 8(2), 245-256. doi:10.1017/S0960258500004141

Formatted: Font: Italic

Sivasakthi S. (2010) Developmental of natural seed coating polymer to improve the planting value of crop seeds. Ph.D. (Agri.) Thesis, Tamil Nadu Agricultural University, Coimbatore, India, 2010.

ISTA. (2013) International rules for seed testing. Switzerland: The International Seed Testing Association (ISTA). 2013.

Sivasakthi, S., Renganayaki, P. R., & Sundareswaran, S. (2022). Influence of TSP polymer coating and biostimulants on seed germination behavior and seedling vigour of black gram var. VBN 8.

Formatted: Font color: Auto

Kamath KR, Park K. (1993) Biodegradable hydrogels in drug delivery. *Advances in Drug Delivery Review*. 1993;11:59-84

Kefi, B. B., Baccouri, S., Torkhani, R., Koumba, S., Martin, P., & M'hamdi, N. (2022). Application of Response Surface Methodology to Optimize Solid-Phase Extraction of Benzoic Acid and Sorbic Acid from Food Drinks. *Foods*, 11(9), 1257.

Formatted: Font color: Auto

Zhang, S., Luo, L., Sun, X., & Ma, A. (2021). Bioactive peptides: a promising alternative to chemical preservatives for food preservation. *Journal of Agricultural and Food Chemistry*, 69(42), 12369-12384.

Brown, R. F., & Mayer, D. G. (1986). A critical analysis of Maguire's germination rate index. *Journal of seed technology*, 101-110.

Formatted: Font color: Auto

Maguire, J. D. (1962). Speed of germination-aid in selection and evaluation for seedling emergence and vigor. *Crop Sci.*, 2, 176-177.

Baveja, S. K., Rao, K. R., Singh, A., & Gombar, V. K. (1988). Release characteristics of some bronchodilators from compressed hydrophilic polymeric matrices and their correlation with molecular geometry. *International journal of pharmaceutics*, 41(1-2), 55-62.

Joseph, J., Kanchalochana, S. N., Rajalakshmi, G., Hari, V., & Durai, R. D. (2012). Tamarind seed polysaccharide: A promising natural excipient for pharmaceuticals. *International Journal of Green Pharmacy (IJGP)*, 6(4).

Amirpour, M., Arman, A., Yolmeh, A., AkbariAzam, M., &Moradi-Khatoonabadi, Z. (2015). Sodium benzoate and potassium sorbate preservatives in food stuffs in Iran. *Food Additives & Contaminants: Part B*, 8(2), 142-148.

Chaleshtori, F. S., Arian, A., &Chaleshtori, R. S. (2018). Assessment of sodium benzoate and potassium sorbate preservatives in some products in Kashan, Iran with estimation of human health risk. *Food and chemical toxicology*, 120, 634-638.

Potter NN (1984). *Food Science, Food additives*. AVI Publishing Co, New York, pp 685–697

Gudesblat, G. E., &Rusinova, E. (2011). Plants grow on brassinosteroids. *Current opinion in plant biology*, 14(5), 530-537.

Tozetto, L. M., Nascimento, R. F. D., Oliveira, M. H. D., Van Beik, J., &Canteri, M. H. G. (2019). Production and physicochemical characterization of craft beer with ginger (*Zingiberofficinale*). *Food Science and Technology*, 39, 962-970.

Warth, A. D. (1985). Resistance of yeast species to benzoic and sorbic acids and to sulfur dioxide. *Journal of Food Protection*, 48(7), 564-569.

Joseph, J., Kanchalochana, S. N., Rajalakshmi, G., Hari, V., &Durai, R. D. (2012). Tamarind seed polysaccharide: A promising natural excipient for pharmaceuticals. *International Journal of Green Pharmacy (IJGP)*, 6(4).

Gupta, G., Parihar, S. S., Ahirwar, N. K., Snehi, S. K., & Singh, V. (2015). Plant growth promoting rhizobacteria (PGPR): current and future prospects for development of sustainable agriculture. *J MicrobBiochemTechnol*, 7(2), 096-102.

Choudhary, S. P., Yu, J. Q., Yamaguchi-Shinozaki, K., Shinozaki, K., & Tran, L. S. P. (2012). Benefits of brassinosteroid crosstalk. *Trends in plant science*, 17(10), 594-605.

Müssig, C. (2005). Brassinosteroid-promoted growth. *Plant biology*, 110-117.

Jones, R. L., & Kaufman, P. B. (1983). The role of gibberellins in plant cell elongation. *Critical Reviews in Plant Sciences*, 1(1), 23-47.

Formatted: Font: Italic, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Italic, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Italic, Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font color: Auto