

Original research article.

POTENTIAL HEALTH RISK EVALUATION OF HEAVY METAL EXPOSURE FROM CONSUMING BRANDED SEASONINGS IN ENUGU, NIGERIA

Comment [DMF1]: A table with the information in the label of the branded seasonings is missing

Abstract

Heavy metals are inherently non-biodegradable elements of the earth's crust that collect and remain in the ecosystem in perpetuity due to both human and natural activity. Their contamination of seasonings remains a public health concern because of the frequency of use. Prolonged exposure to these heavy metals is a severe health risk worldwide. This study evaluated the potential health risk of heavy metal exposure from consuming branded seasonings in Enugu, Nigeria. Thirty (30) branded seasonings were selected by simple random sampling and grouped into five (6) according to their batches. Atomic absorption spectroscopy was used to analyze arsenic, cadmium, lead and nickel. The level of arsenic in group S_A(1.28±0.21mg/kg), S_B(1.73±0.31mg/kg), S_C(1.13±0.17mg/kg), S_D(2.11±0.20mg/kg) were observed to be above (1.00mg/kg), the WHO maximum permissible limit. Cadmium levels in S_A(0.36±0.01mg/kg) were observed to be above(0.30 mg/kg), the WHO maximum permissible limit. The estimated daily intake, hazard quotient and hazard index of the metals were within the normal range. The presence of heavy metals above WHO permissible limits in branded seasonings should be a public health concern. To protect the population's health, producers, retailers, and vendors of branded seasonings should be informed of the hazards of exposing their products to heavy metal contamination. Continuous surveillance of branded seasonings for heavy metal contamination is recommended.

Comment [DMF2]: A table with the information about the labels, composition, ingredients, any legal authority or sanitary registration label, numbers, etc., is missing.

1. INTRODUCTION

Seasonings are a group of diversely composed and functioning edible products. Numerous definitions of seasonings can be found in the literature. Codex Alimentarius lists salts, seasonings, spices, soups, sauces, salads, and protein supplements among the ingredients added to food to improve scent and flavor [1]. Most scientists classify seasonings as products of plant origin, albeit with a more complex composition than seasoning mixtures [2]. They are composed of dried seed, fruit, root, bark, or other plant parts mainly used as aromatic additives enhancing the sensory values of food or keeping food fresh [3]. In addition to plant-based constituents, seasonings may also contain salt, monosodium glutamate, and citric acid [2].

Comment [DMF3]: Without proper characterization and definition of the composition of each seasoning, this is not a correct term to use.

Despite all the benefits of these products, it is believed that seasonings harm the biological system, probably due to contamination. Contamination may arise from several different factors, including alterations in the amount and quality of components, the use of a raw material lacking

a bioactive substance, biogenic or chemical environmental contaminants, Pesticide residues, mycotoxins, and heavy metals, as well as unintentional contaminants including floral debris, ashes and others [4, 5, 6].

The toxicity of heavy metals to human health and the environment has received much attention. Plants are the primary conduit for heavy metals from polluted soil to people. Metals can easily contaminate plants because they can emanate from the type of soil used for cultivation, the fertilizers applied, and the source of irrigation water [7]. Heavy metals in extremely low quantities may be hazardous due to their poor excretion rates via the kidneys [8]. Heavy metals enter our food chains through soils [9, 10]. Heavy metals have piqued the attention of researchers owing to their persistent, bio accumulating, and biomagnifying properties [11, 12]. Heavy metals cause oxidative stress in aerobic cells by increasing reactive oxygen species (ROS). It has been linked to changes in the nervous system, cognitive impairment, infertility, foetal harm, preterm birth, cardiovascular disease, bronchitis, and coughing [13]. Cancers of the lung and bladder are the most common sites where it's carcinogenic effects manifest [14, 15]. Due to excessive levels of lead, the Food and Drug Administration of the United States (US FDA) reported in September 2013 that distributors of turmeric powder, including a business based in Bangladesh, were voluntarily recalling their products [16].

Commercially produced branded food seasonings are preferred currently for convenience; it is easy to use. Information regarding the raw material used in the production of seasonings is supposed to be made available in the containers or packs of such seasonings. Regrettably, this information is sometimes not present on the containers of such seasonings. It may be quite challenging to spot adulterations in this all-important and every day-use product, which calls for sophisticated analytical techniques [17].

Due to a paucity of such information and because most research focused on natural food spices, it is critical to examine the heavy metal contents, estimated daily intake, total hazard quotients and hazard index in commercially produced branded food seasonings sold in Enugu markets. This study assesses the possible health risk of heavy metal exposure from consuming branded seasonings in Enugu, Nigeria.

2. MATERIALS AND METHODS

2.1 Subject area

The study was conducted in Enugu State, situated in the Southeastern region of Nigeria. Enugu is a metropolitan city notable for its coal extraction, with few industrial sites and numerous markets. This city has a population of approximately 722,664; a total area of 556 km²; latitudes of 6.4584°N and 7.5464°E; and a warm climate with an annual average temperature of 26.3°C and an average yearly rainfall of 1,730_{mm}. Enugu State residents work in various sectors, including government, business, the arts, and agriculture. The map in Figure 1 [18] presents where the branded seasonings were purchased.

Comment [DMF4]: Information in the label, components of the seasonings, data available in the label about sanitary registry, or sanitary authority or any surveillance agency for quality/safety and if there is nothing about that information, indicate that the information is "not available"




Fig. 1. Map of Enugu Municipality

2.1.1 Sample selection and analysis

The branded seasonings were selected by simple random sampling. Every fourth seasoning as was placed on the table was selected, as all the branded seasonings were placed on a table. The selected seasonings were then bought from the vendors at the different markets in Enugu municipality. A total of thirty (30) commercially produced branded seasonings were purchased from four different markets in Enugu municipality. The selected seasonings were grouped into seven (6). Each group constituted of five (5) batches (replicates?) as followed:

Group A- This consisted of five different batches of stew seasoning (SA).

Group B- This consisted of five different batches of local soup seasoning (SB)

Group C- This consisted of five different batches of Beef seasoning (SC)

Group D- This consisted of five different batches of Chicken seasoning (SD)

Group E- This consisted of five different batches of Jollof rice seasoning (SE)

Group F- This consisted of five different batches of fried rice seasoning (SF)

Their expiry dates were used in place of the batch numbers as some of the batch numbers were blurred.

The selected seasonings were used for the analysis of heavy metals (arsenic, cadmium, lead and nickel) using flame atomic absorption spectroscopy as described [19] at the National Centre for Energy Research and Development, University of Nigeria Nsukka.

Comment [DMF5]: Any reference of sanitary agency? Any local authority of sanitary control to indicate the reference in the origin of the seasonings? Any labelling to be incorporated as additional information? (or additional table with the available information about the seasonings, and their components?

Comment [DMF6]: ? Seven? Six?

Comment [DMF7]: 5 batches, how many samples were analysed per batch? N= ?

Comment [DMF8]: Not available text for checking the procedure. Please, describe briefly the protocol. The processing of the samples, the extraction/digestion process, and the analytical equipments.

2.1.2. Determination of Hazard quotient and Hazard index of the selected seasonings

$$DIM (mgkg^{-1} Day^{-1}) = \frac{X (mgkg^{-1}) \times Y (kg)}{Z (kg)} \quad (1) [20]$$

where DIM refers to the Dietary intake of metal, X is the level of heavy metal contained in the seasoning as determined by atomic absorption in this study Y (0.01) is the ingestion rate while Z (24 for child and 60 for adult) [21].

The hazard quotient (HQ) was calculated using the ratio of the daily intake of metal (DIM) to the reference dose for heavy metals.

$$Hazard\ quotient = \frac{Daily\ intake\ of\ metal}{RfD} \quad (2) [22]$$

Where: R_fD is the oral reference for each heavy metal investigated.

The hazard index (HI) was expressed as the total sum of the hazard quotients (HQs) of the heavy metals.

$$Hazard\ index = HQ1 + HQ2 + HQ3 + HQ4 + \dots + HQ_n \quad (3) [23]$$

2.2 Statistical data analysis

The statistical analysis software, Graph Pad prism version 7 (Graph Pad Software Inc., USA) was used to analyse the data from this study. Values were expressed as Mean ± Standard deviation. Student's t-test was used to compare continuous variables such as concentration of heavy metals in the seasonings and WHO permissible limits [24]. P-value of <0.05 was considered statistically significant.

Comment [DMF9]: N=?

3.0 RESULTS AND DISCUSSION

The levels of heavy metals in seasonings consumed by Enugu metropolis residents as sold in their markets have been brought to focus. The findings of this study showed an increase level of arsenic in SA (1.28±0.21)mg/kg, SB (1.73±0.31)mg/Kg, SD (1.13±0.17)mg/kg and SE (2.11±0.20)mg/kg in comparison with the WHO maximum permissible limit (1.00) mg/kg [24] (Table 1). The beauty of this study is that it is the first to report the level of arsenic in seasonings in Nigeria. This study's results agree with the observation of Pakistan and Bangladesh studies that revealed the presence of arsenic in food spices [6, 25,]. The level of arsenic found in both studies was higher than the findings in this current study. The finding of their study could be because they worked on natural spices, technological conditions or environmental contamination. Arsenic ingestion, even though at low-level may cause non carcinogenic effects, such as sore pharynx and irritated airways, vertigo, vomiting, and discolouration of skin lesions and tumours [26]. In addition, chronic consumption may result in neurobehavioral and neuropathic changes, memory loss, a decrease in intellectual level, infertility disorder, foetal injury, premature delivery, cardiovascular diseases, chronic cough, and chronic bronchitis [13]. In most cases, its carcinogenic effects manifest as lung and bladder cancer [14, 27].

The findings of this study showed an increased cadmium level in SA (0.36 ± 0.01)mg/kg in comparison with the WHO maximum permissible limit (Table 1). These findings are lower than those of [6,24,28] and are higher than that observed in the work of Nnorom et al [29]. Variability in outcomes may be attributable to variations in raw materials used in manufacturing, technological processes, and distribution networks. Colourants may also contribute to cadmium contamination of spices. Cadmium (Cd) causes chronic human diseases. It predominantly affects the liver, kidneys, testicles, pancreas, and bones [30]. Cd^{2+} may also inhibit the liver enzymes catalase, glutathione peroxidase, and superoxide dismutase [28]. In addition to damaging the skeletal system by interfering with the metabolism of Ca, Zn, Cu, Fe and Mg [31]. Cd may also play an initial role in the development of cancer in the breasts, pancreas, lungs, prostate, kidneys, nasopharynx, and genitalia [32]. Cadmium levels in SA, SC, SD, SE, and SF were below the WHO maximum permissible limit[24] (Table 1).

The levels of other heavy metals such as Lead (Pb) and Nickel assayed in this study were all within the WHO maximum permissible limit as observed in this study. Pb primarily affects the brain, the kidneys, the liver, and the reproductive system [33]. It may induce a decline in IQ and perceptual ability. It may also increase nervous system hyperactivity [30]. Moreover, the presence of Pb in the blood causes bladder cancer [35]. Nickel can induce a variety of pathological effects in humans. The chronic accumulation of nickel and nickel compounds in the human body may be responsible for a variety of adverse health effects, including pulmonary fibrosis, renal and cardiovascular diseases, and respiratory cancer [32].

A health risk assessment was conducted to determine if there may be a link between the intake of branded seasoning and the overall well-being of the study population. The EDIs of the heavy metals as a consequence of seasoning intake are shown in Table 2. The EDI ranged from $1.03E-05$ to $9.17E-06$ in adults and $1.00E-05$ to $8.33E-07$ in juveniles. All EDIs were lower than the TDIs [35], indicating that the individual consumption of branded seasonings sold in Enugu metropolis markets may not pose any health hazards to the population.

The Hazard quotient(HQ) of all the heavy metals in this study were less than 1 ($HQ < 1$) as shown in Table 3 in this study. HQ values less than 1 ($HQ < 1$) indicate that the cumulative consumption of that substance will not induce an adverse effect but with the tendency of adverse effect when it is above 1 ($HQ > 1$); the same applies also to HI. The HI of all spices were less than 1.

The HI of the heavy metals, the source of contamination and methods to eradicate them have been described as a constant focus of research [36]. It is necessary to periodically check the branded seasonings for these heavy metals contamination in order to rule out any potential chronic heavy metal toxicity in the population.

4.0 CONCLUSION

This study revealed an arsenic level above the WHO maximum permissible limit in SA, SB, SD and SE. The cadmium levels in S_A was above the WHO maximum permissible limit.

However, the Nickel and Lead levels in the selected branded seasonings were below the WHO permissible limits. The risk assessment indicates that there is little cause for concern. The presence of these contaminants in branded seasonings should be a public health concern. To protect the population's health, producers, retailers, and vendors of branded seasonings should be informed of the hazards of exposing their products to heavy metal contamination. Continuous surveillance on branded seasonings for heavy metal contamination is recommended.

Conflict of interest- None

Table 1. Analytical result of the heavy metals(mg/kg) in branded seasonings in Enugu, Nigeria

Samples		Pb	As	Ni	Cd
WHOMPL		10.00	1.00	1.50	0.30
SA	Mean±SD	0.15±0.04	1.28±0.21	0.62±0.13	0.36±0.01
SB	Mean±SD	0.18±0.02	1.73±0.31	0.38±0.12	0.24±0.01
SC	Mean±SD	0.17±0.03	0.98±0.08	0.07±0.02	0.08±0.01
SD	Mean±SD	0.10±0.02	1.13±0.17	0.79±0.12	0.02±0.01
SE	Mean±SD	0.10±0.02	2.11±0.20	0.55±0.02	0.12±0.03
SF	Mean±SD	0.15±0.07	0.75±0.16	0.62±0.80	0.05±0.01

*WHO MPL=World Health Organization Maximum Permissible Limit

SA= Stew seasoning

SB=Local soup seasoning

SC=Beef seasoning

SD=Chicken seasoning

SE=Jollof rice seasoning

SF=Fried rice seasoning

Pb=Lead

As= Arsenic

Ni=Nickel

Cd=Cadmium

Table 2. Estimated daily intake (EDI) (mg/kg⁻¹) of the branded seasonings in Enugu,

Samples	Pb		As		Ni		Cd	
	Adults	Children	Adult	Children	adult	Children	Adult	Children
SA	2.50E-06	6.20E-05	2.13E-05	5.33E-05	1.03E-05	2.58E-05	6.00E-06	1.50E-05
SB	3.00E-06	7.50E-06	2.88E-05	7.20E-05	6.33E-06	1.58E-05	4.00E-06	1.00E-05
SC	2.80E-06	7.00E-06	1.63E-05	4.08E-05	1.17E-06	2.92E-06	1.33E-06	3.33E-06
SD	1.67E-06	4.17E-06	1.88E-05	4.71E-05	1.32E-05	3.29E+05	3.33E-07	8.33E-07
SE	1.67E-06	4.17E-06	3.52E-05	8.79E-05	9.17E-06	2.29E-05	2.00E-05	5.00E-06
SF	2.50E-06	6.25E-06	1.25E-05	3.13E-05	1.03E--05	2.58E-05	8.33E-07	2.08E-06

Nigeria

Comment [DMF10]: Change unit to µg/kg to present cleaner tables. Use, S.I. units

SA= Stew seasoning

SB=Local soup seasoning

SC=Beef seasoning
 SD=Chicken seasoning
 SE=Jollof rice seasoning
 SF=Fried rice seasoning
 Pb=Lead
 As= Arsenic
 Ni=Nickel
 Cd=Cadmium

Table 3. Hazard quotient (HQ) and (HI) of the branded seasonings in Enugu, Nigeria

Metals	RfD	SA		SB		SC		SD		SE		SF	
		Adults	Children	Adults	Children	Adults	Children	Adults	Children	Adults	Children	Adults	Children
Nickel	0.02	5.15E-04	1.29E-04	3.16E-04	7.90E-04	5.85E-05	1.46E-04	6.60E-04	1.60E-03	4.85E-04	1.14E-03	5.15E-04	1.29E-03
Cadmium	0.001	6.00E-03	1.50E-02	4.00E-03	1.00E-02	1.30E-03	3.30E-03	3.33E-04	8.33E-04	2.00E-04	5.00E-03	8.33E-04	2.08E-03
Lead	0.0035	7.14E-04	1.70E-02	8.57E-04	2.10E-03	8.09E-04	2.00E-03	4.77E-04	2.00E-03	4.77E-04	1.00E-03	7.14E-04	1.70E-02
Arsenic	0.0003	7.10E-02	2.10E-02	9.60E-02	2.00E-03	5.40E-02	3.60E-02	6.20E-03	1.57E-02	3.50E-02	2.93E-02	1.25E-02	1.04E-01
HI		7.80E-02	5.30E-02	1.00E-01	4.50E-02	5.62E-02	4.14E-02	7.67E-03	2.00E-02	7.20E-02	3.64E-02	1.40E-02	1.24E-01

Comment [DMF11]: Use the unit more properly, 5.15 x 10-4, etc., not in the arbitrary unit of the software E-04...etc.

SA= Stew seasoning
 SB=Local soup seasoning
 SC=Beef seasoning
 SD=Chicken seasoning
 SE=Jollof rice seasoning
 SF=Fried rice seasoning
 HI=Hazard Index
 RfD=Oral Reference Dose

REFERENCES

1. García-Casal, M.; Peña-Rosas, J.P.; Malavé, H.G. Sauces, Spices, and Condiments: Definitions, Potential Benefits, CONSUMPTION Patterns, and Global Markets. *Ann. N. Y. Acad. Sci.* **2016**, *1379*, 3–16. Available online: <http://www.innocua.net/web/download-5387/garc-a-casal-et-al-2016-annals-of-the-new-york-academy-of-sciences.pdf> (accessed on 19 August 2021).
2. Smiechowska, M.; Nowak, J.; Nowak, J.; Skotnicka, M. Spices and Seasoning Mixes in European Union—Innovations and Ensuring Safety. *Foods* **2021**, *10*, 2289. <https://doi.org/10.3390/foods10102289>
3. Kowalski, G. The Safety Assessment of Toxic Metals in Commonly Used Herbs, Spices, Tea, and Coffee in Poland. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5779. <https://doi.org/10.3390/ijerph18115779>
4. Kucharska-Ambrozej, K.; Karpinska, J. The application of spectroscopic techniques in combination with chemometrics for detection adulteration of some herbs and spices. *Microchem. J.* **2020**, *153*, 104278.

5. Karadaş C, Kara D (2012) Chemometric approach to evaluate trace metal concentrations in some spices and herbs. *Food Chem* 130(1): 196–202
6. Baig, J. A., Bhatti, S., Kazi, T. G., & Afridi, H. I. (2019). Evaluation of arsenic, cadmium, nickel and lead in common spices in Pakistan. *Biological trace element research*, 187, 586-595.
7. Nnari, N., Ow hoeke, E. & Naze, A. (2020). Health risk of selected inorganic contaminants in some food spices sold in markets within port harcourt metropolis, Nigeria. *Asian Journal of Research in Biology*, 3(1), 15-21.
8. Jaeyoung Shim, Taeyoung Cho, Donggil Leem, Youngmi Cho & Changhee Lee (2018): Heavy metals in spices commonly consumed in Republic of Korea, *Food Additives & Contaminants: Part B*, DOI: 10.1080/19393210.2018.1546772
9. Margenat, A., Matamoros, V., Díez, S., Cañameras, N., Comas, J., Bayona, J.M., 2019. Occurrence and human health implications of chemical contaminants in vegetables grown in peri-urban agriculture. *Environ. Int.* 124, 49–57
10. Khalid, S., Shahid, M., Niazi, N.K., Murtaza, B., Bibi, I., Dumat, C., 2017. A comparison of technologies for remediation of heavy metal contaminated soils. *J. Geochem. Explor.* 182, 247–268.
11. Zakaria, Z.; Zulkafflee, N.S.; MohdRedzuan, N.A.; Selamat, J.; Ismail, M.R.; Praveena, S.M.; Tóth, G.; AbdullRazis, A.F. Understanding Potential Heavy Metal Contamination, Absorption, Translocation and Accumulation in Rice and Human Health Risks. *Plants* 2021, 10, 1070. <https://doi.org/10.3390/plants10061070>
12. Pandiyan, J., Mahboob, S., Govindarajan, M., Al-Ghanim, K.A., Ahmed, Z., Al-Mulhm, N., Jagadheesan, R., Krishnappa, K., 2021. An assessment of level of heavy metals pollution in the water, sediment and aquatic organisms: a perspective of tackling environmental threats for food security. *Saudi J. Biol. Sci.* 28 (2), 1218– 1225.
13. Milton, A.H., Rahman, M., 2002. Respiratory effects and arsenic contaminated well water in Bangladesh. *Int. J. Environ. Health Res.* 12 (2), 175–179.
14. Cohen, G., Levy, I., Yuval, Kark, J. D., Levin, N., Witberg, G., ...& Gerber, Y. (2018). Chronic exposure to traffic-related air pollution and cancer incidence among 10,000 patients undergoing percutaneous coronary interventions: A historical prospective study. *European journal of preventive cardiology*, 25(6), 659-670.
15. Sanyal, T., Bhattacharjee, P., Paul, S., & Bhattacharjee, P. (2020). Recent advances in arsenic research: significance of differential susceptibility and sustainable strategies for mitigation. *Frontiers in public health*, 8, 464.
16. U.S. Food and Drug Administration. 2013. Recalls PRAN brand turmeric powder due to elevated levels of lead. [accessed 2013 Oct 04]. www.fda.gov/Safety/Recalls/ucm371206.htm
17. Wadood, S. A., Boli, G., Xiaowen, Z., Hussain, I., & Yimin, W. (2020). Recent development in the application of analytical techniques for the traceability and authenticity of food of plant origin. *Microchemical Journal*, 152, 104295.
18. Ubachukwu NN, Phil-Eze PO, Emeribe CN. Analysis of household hazardous wastes awareness level in Enugu metropolis. *Academic Journal of Interdisciplinary Studies*. 2014;3(1):369.

Formatted: Spanish (Spain, International Sort)

19. Oluremi OI, Solomon AO, Saheed AA, Fatty acids, metal composition and physico-chemical parameters of IgbemoEkiti rice bran oil. *Journal of Environmental Chemistry and Ecotoxicology*. 2013;5(3):39-46.
20. USEPA (US Environmental Protection Agency), Exposure Factors Handbook General Factors. EPA/600/P-95/002Fa, vol. I. Office of Research and Development, National Center for Environmental Assessment, US Environmental Protection Agency. Washington, DC; 2010.
21. Orisakwe OE, Ozoani HA, Nwaogazie IL, Ezejiolor AN. Probabilistic health risk assessment of heavy metals in honey, *Manihotesculenta*, and *Vernoniaamygdalina* consumed in Enugu State, Nigeria. *Environ Monit Assess*. 2019; 191(7):424.
22. USEPA (US Environmental Protection Agency), Exposure Factors Handbook General Factors. EPA/600/P-95/002Fa, vol. I. Office of Research and Development, National Center for Environmental Assessment, US Environmental Protection Agency. Washington, DC; 2010
23. USEPA (US Environmental Protection Agency). Risk assessment guidance for superfund. Human Health Evaluation Manual Part A, Interim Final, vol. I. Washington (DC), United States Environmental Protection Agency; EPA/540/1-89/002; 1989.
24. WHO (2006) WHO guidelines for assessing quality of herbal medicines with reference to contaminants and residues. World Health Organization, Geneva, Switzerland
25. Mahtab AliMollah*, Golam Rabbany, M Nurunnabi, ShaMd.ShahanShahriar and Sayed M A Salam. Health Risk Assessment of Heavy Metals Through Six Common Spices of MohanpurUpazila of Rajshahi District, Bangladesh. *Glob J Nutri Food Sci*. 3(5): 2022. GJNFS. MS.ID.000574. DOI: 10.33552/GJNFS.2022.03.000574.
26. Signes-Pastor, A.J., Carey, M., Meharg, A.A., 2016. Inorganic arsenic in rice-based products for infants and young children. *Food Chem*. 191, 128–134.
27. Umar, M. A., Saliyu, Z. O., & Pepper, C. (2014). Heavy metals content of some spices available within FCT-Abuja, Nigeria. *International Journal of Agricultural and food science*, 4(1), 66-74.
28. Nnorom, I. C., Osibanjo, O., & Ogugua, K. (2007). Trace heavy metal levels of some bouillon cubes, and food condiments readily consumed in Nigeria. *Pakistan journal of nutrition*, 6(2), 122-127.
29. Fu, Z., Xi, S., 2020. The effects of heavy metals on human metabolism. *Toxicol. Mech. Methods* 30 (3), 167–176.
30. Kim, H.S., Kim, Y.J., Seo, Y.R., 2015. An overview of carcinogenic heavy metal: molecular toxicity mechanism and prevention. *J. Can. Prevent*. 20 (4), 232. Kumar, V., Thakur, R.K., 2018. Health risk assessment of heavy metals via dietary intake of vegetables grown in wastewater irrigated areas of Jagjeetpur, Haridwar India. *Arch. Agricult. Environ. Sci*. 3 (1), 73–80.
31. Genchi G, Carocci A, Lauria G, Sinicropi MS, Catalano A. Nickel: Human Health and Environmental Toxicology. *Int J Environ Res Public Health*. 2020 Jan 21;17(3):679. doi: 10.3390/ijerph17030679. PMID: 31973020; PMCID: PMC7037090.
32. Brewer, G.J., Prasad, A.S. eds., 2020. Essential and Toxic Trace Elements and Vitamins in Human Health. Academic Press.

Formatted: Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International Sort)

33. Awadalla, A., Mortada, W.I., Abol-Enein, H., Shokeir, A.A., 2020. Correlation between blood levels of cadmium and lead and the expression of microRNA-21 in Egyptian bladder cancer patients. *Heliyon* 6 (12).
34. Baars AJ, Theelen RMC, Janssen PJCM, Hesse JM, van Apeldoorn MV, Meijerink MV. Re-evaluation of human-toxicological maximum permissible risk levels, Microsoft Word - 711701025.doc (openrepository.com); 2001.
35. Barnaby R, Liefeld A, Jackson BP, Hampton TH, Stanton BA. Effectiveness of table top water pitcher filters to remove arsenic from drinking water. *Environ. Res.* 2017;158:610–615.
36. Wongsasuluk P, Chotpantarat S, Siritwong W, Robson M. Heavy metal contamination and human health risk assessment in drinking water from shallow groundwater wells in an agricultural area in UbonRatchathani province, Thailand. *Environ Geochem Hlth.* 2013;36:169–182.