

## IMPORTANCE OF OKRA (*Abelmoschus esculentus* L.) AND IT'S PROPORTION IN THE WORLD AS A NUTRITIONAL VEGETABLE

### ABSTRACT

Okra belong to malvaceae family, a commercial vegetable crops. It is extensively distributed in tropical, subtropical, and warm temperate parts of the world and is native to Ethiopia. It is essential to human nutrition and a good source of total minerals, vitamins, calcium, potassium, enzymes, and other nutrients that are frequently lacking in developing country diets. Additionally, reports of its medicinal value in treating curingulcers and providing relief from haemorrhoids. Okra has found medical application as a plasma replacement or blood volume expander and also useful in genito-urinary disorders, spermatorrhoea and chronic dysentery. The fruits of okra crop bringing into commercial production have reawakened beneficial interest.

### KEY WORDS

*Abelmoschus esculentus*, Biochemical composition, diseases, insects, okra, origin, production.

### INTRODUCTION

Okra is also known as lady's fingers, bhindi in India, krajiab kheaw in Thailand, okra plant, ochro, okoro, quimgombo, quingumbo, gombo, kopi arab, kacang bendi, and bhindi in South East Asia. Okra is also known as *Abelmoschus esculentus* (L.) Moench in many English-speaking nations. However, it is referred to as bamia, banya, or bamieh in the Middle East and gumbo in the South. Okra is referred to as quiabo in Portuguese and Angola, quimbombo in Cuba, gombo commun, gombo, gumbo in France, mbamia and mbinda in Sweden, and okra in Japan [13], [30]. In Nigerian Igbo, Taiwan is referred to as qiu kui [53]. It belongs to the genus *Abelmoschus* and family Malvaceae. Okra's geographic origin is contested, with claims that it originated in South Asia, Ethiopia, and West Africa. Around the world, the plant is grown in tropical, subtropical, and warm temperate climates [38]. Although okra may be grown on a variety of soil types, well-drained fertile soils with sufficient organic matter produce a high yield [7]. In the tropics, the crop is extensively grown all year round. Okra is a nutrient-dense vegetable that is crucial in addressing the market's lacklustre demand for vegetables [6]. The world's total area under cultivation in 2009–2010 was 0.43 million hectares, and output totaled 4.54 million tonnes, with India producing 5784 thousand tonnes of okra with a yield of 11.1 tons/hectare [23]. In comparison to other developing countries' yields of 9.7 to 10 tonnes per hectare, the yield is quite low. India contributed largest producer (67.1%), abide by Nigeria (15.4%) and Sudan (9.3 %).

### OKRA'S GEOGRAPHIC ORIGIN AND DISTRIBUTION

Previously, the genus *Hibiscus*, section *Abelmoschus* of the family Malvaceae, contained the okra plant or lady's finger [32]. It was therefore suggested that the section *Abelmoschus* be elevated to the status of separate genus. In the taxonomy and modern literature, the widespread use of *Abelmoschus* was soon approved [22]. The calyx of the *Hibiscus* genus is

spathulate, has five small teeth, is connate to the corolla, and becomes caducous after flowering [27, 59]. By the 12th century B.C., the ancient Egyptians were cultivating okra, which had its origins somewhere near Ethiopia. Its cultivation became widespread in North Africa and the Middle East [3, 60]. The most comprehensively recorded research of the genus *Abelmoschus* are those that Borssum and colleagues [11] and Bates [9] conducted on its taxonomic revision. At the International Okra Workshop conducted at the National Bureau of Plant Genetic Resources (NBPGR) in 1990, a modern classification was chosen using van Borssum Waalkes' classification as a starting point. Eight species are the most often accepted, despite the fact that over 50 have been described [24]. Okra is cultivated all throughout the world, although it is more common in nations with tropical and subtropical climates [8, 50]. This crop may be cultivated as a garden crop or on a huge commercial farm [49]. Many nations, including India, Japan, Turkey, Iran, Western Africa, Yugoslavia, Bangladesh, Afghanistan, Pakistan, Myanmar, Malaysia, Thailand, India, Brazil, Ethiopia, Cyprus, and the Southern United States, cultivate okra plants for commercial purposes [43, 10, 44].

### **STRUCTURE AND PHYSIOLOGY**

In tropical and warm temperate areas of the world, *Abelmoschus esculentus* is grown for its fibrous fruits or pods that contain spherical, white seeds. The seeds are soaked the night before sowing in culture, at a depth of 1-2 cm. Between six days (wet seeds) to three weeks, germination takes place. Seedlings need a lot of water. Within a week of the fruit being pollinated, the seed pods must be collected since they quickly turn fibrous and woody. The immature fruits are collected and consumed as vegetables(41). It is one of the world's most heat- and drought-tolerant vegetable species, and it can grow on thick clay soils and intermittent moisture but frost, can harm the pods.

### **CYTOGENETIC RELATIONSHIP OF OKRA**

The number of chromosomes and ploidy levels of various species of the genus *Abelmoschus* varied significantly from one another. Within *A. esculentus*, the chromosomal numbers are  $2n = 72, 108, 120, 132,$  and  $144$ , which are in a regular succession of polyploids with  $n = 12$  [15]. *A. angulosus* has the lowest number ( $2n=56$ ), whereas *A. manihot* var. *caillei* has the largest number (almost 200) of known chromosomes ( 54, 55,56).

### **BIOCHEMICALLY RELATIONSHIP**

The okra fruit is mostly consumed raw or cooked, and it is a significant source of vitamins A, B, and C, minerals, iron, and iodine, as well as an essential vegetable source of viscous fibre [3,26,35]. However, it is also said to be low in salt, saturated fat, and cholesterol. Okra pods contain the following nutrients per 100 g of edible portion: water 88.6 g, energy 144.000 kJ (36 kcal), protein 2.10 g, carbohydrate 8.20 g, fat 0.20 g, fibre 1.70 g, Ca 84.00 mg, P 90.00 mg, Fe 1.20 mg, -carotene 185.00 g, riboflavin 0.08 mg, thiamin 0.04 mg, niacin 0.60 mg, ascorbic acid 47.00 mg. Okra is rich in protein, carbohydrates, and vitamin C [17,21,30,42, 50] and is an essential part of the human diet [ 25,50]. Young, immature okra pods may be eaten in a variety of ways and are crucial to consume as fresh fruits [40]. There have also

been reports of Fe, Zn, Mn, and Ni [36]. Okra is a valuable source of vitamins, calcium, potassium, and other minerals that are frequently missing in the diets of poor nations [24]. The freshest okra pods that are seven days old contain the highest concentration of nutrients [5]. Okra leaves have the following nutritional values per 100 g of edible portion: 81.50 g of water, 235.00 kJ (56.00 kcal) of energy, 4.40 g of protein, and 0.60 g of fat. 11.30 g of carbohydrates, 2.10 g of fibre, 532.0 mg of calcium, 70.00 mg of phosphorus, 0.70 mg of iron, 59.00 mg of ascorbic acid, 385.00 g of  $\beta$ -carotene, 0.25 mg of thiamin, 2.80 mg of riboflavin, and 0.20 mg of niacin were found in the study [21],[61]. Mucilage is the primary form of carbohydrates [33], [28]. Additionally edible are the flower and leaf buds [16]. About 20% of the proteins and 20% of the oil in okra seeds are proteins [17], [14]. Potential hypocholesterolemic effects of okra seed oil exist [46]. Okra has a very high potential for widespread planting for both cake and edible oil [47]. Cereal flour might potentially be fortified with okra seed flour [2]. Okra oil was determined to be appropriate for usage as a biofuel in a 2009 research [18]. When making gur or brown sugar from sugarcane juice, the roots and stems of okra are utilised to clarify the juice [13].

One of the main elements in the production of okra is the occurrence of insect pests. Numerous insect pests affect the crop, but the shoot and fruit borer, *Earias vittella* (Fabricius), and *Earias Insulana* are the most dangerous since they have the upper hand by directly harming vulnerable fruits. Fruit borer damage ranges from 88 to 100 percent [45]. When compared to healthy okra fruits, the usual number of seeds per fruit was reduced by 16.47 percent, while the number of stained seeds increased by 200 percent and the number of damaged seeds decreased by 18.70 percent [58]. After a rainstorm, the prevalence of fruit borers often occurs in a humid environment. Individual eggs are laid by the female adult on leaves, while flower buds are laid on soft fruits. Before the production of fruits, little brown caterpillars pierce the top stalk and feed there. *Amrasca biguttula biguttula* (Ishida), a leafhopper, and *Earias* spp., a shoot and fruit borer, are serious pests that may wreck havoc and cause significant damage. Leafhopper alone was responsible for 32.06%–40.84% [57]. Fruit output was reduced by 50% as a result of shoot and fruit borer [12]. Fruit is unsuitable for human consumption when fruit and shoot borer larvae penetrate into shoots during the vegetative development stage and subsequently in blooms and fruits. Insecticide usage has given crops instant respite and appears to have benefitted farmers among other suggested pest management methods. The usage of chemicals is rising quickly and will keep doing so unless some trustworthy alternative control techniques are created for the same reason. In Asia, 95% of the population uses pesticides.

## **DISEASES OF OKRA**

### **Yellow Vein Mosaic Virus (YVMV)**

The Yellow Vein Mosaic Virus is the cause causative agent. This is the most significant and harmful viral disease that affects okra crops at all stages of growth. The diseased plants produce fruits that are malformed, tiny, rough, and light yellow to white in colour. If the plants contract the disease within 20 days of germination, there will be a 50–100% loss in production and quality [20], [48], [51].

### **Cercospora Leaf Spot**

Agent responsible for Cercospora Leaf Spot is *Cercospora hibisci*, *C. malayensis*, and *A. abelmoschi*. Three species of *Cercospora* in India cause leaf marks in okra. *C. abelmoschi* creates sooty black, angular dots, and *C. malayensis* causes brown, irregular patches. The harmed leaves wilt, roll and then fall. The leaf spots are frequent during wet seasons and severely defoliate the leaves [14] [37].

### **Fusarium Wilt Causative agent**

Vasoinfectum *Fusarium oxysporum* f. sp. Everywhere that okra is actively produced, there is a dangerous disease called fusarium wilt. The fungus colonises the vascular system, invades the roots, and so prevents water from moving through the plant. The illness is transmitted by interculture operations and is carried in the soil.

### **Powdery Mildew**

Powdery mildew is brought on by *Erysiphe cichoracearum*, *Sphaerotheca fuliginea*, and these two organisms are also responsible for its cause. While the latter has just been recorded from Bangalore, the former's sickness is most prevalent in okra-growing regions [29].

### **MEDICAL PROPERTY OF OKRA CROP**

Haemorrhoids and ulcers have been treated using its therapeutic benefits, according to reports [1]. Numerous publications related to herbal and traditional medicine make mention to the 1898 claim that unspecified plant components have diuretic qualities [19]. Okra has found use in medicine as a blood volume expander or plasma replacement [52] [34], [31], [4], [62]. It is also an excellent source of iodine, which is helpful in the treatment of uncomplicated goiter, as well as other compounds with medicinal use [35]. It is highly helpful in treating chronic dysentery, spermatorrhoea, and genitourinary diseases [39]. According to experiments carried out in China, an alcohol extract of okra leaves may be able to reduce protein urea, enhance renal function, reduce protein free radicals, and treat renal tubular-interstitial disorders. Numerous texts pertaining to herbal and conventional medicine make mention to the 1898 claim [19] that unspecified plant components have diuretic qualities. Studies focusing on okra extract as a diabetic treatment are currently being research.

### **CONCLUSION**

Okra (*Abelmoschus esculentus* (L.) Moench) is a very important medicinal plant with several pharmacological uses. It has been used as an ingredient in many herbal formulations that are used to treat various illnesses, particularly the regulation of blood pressure, fat, diabetes, chronic dysentery genito-urinary disorders, simple goitre, and ulcer, in addition to having the aforementioned nutritional, medical, and industrial properties.

## Reference

1. Adams C. F., Nutritive value of American foods in common units, U.S. Department of Agriculture, Agric Handbook. 425, pp 29, (1975).
2. Adelokun O.E., Oyelade O.J., Ade-Omowaye B.O., Adeyemi I.A., Van M., Influence of pre-treatment on yield, chemical and antioxidant properties of Nigerian okra seed (*Abelmoschus esculentus* Moench) flour: DOI: 10.1016/j.fct.2008.12.023, (2008).
3. Adeboye O.C., Oputa C.O., Effect of galex on growth and fruit nutrient composition of Okra (*Abelmoschus esculentus* (L.) Moench) Int.J.Agric. 18(1,2):1-9, (1996).
4. Adetuyi F.O., Osagie A.U., Adekunle A.T., Effect of Postharvest Storage Techniques on the Nutritional Properties of Benin Indigenous Okra *Abelmoschus esculentus* (L) Moench. Pakistan J. Nutrit. 7: 652-657, (2008).
5. Agbo A.E., Gnakri D., Beugre G.M., Fondio L., Kouame C., Maturity degree of four okra fruit varieties and their nutrients composition. Elect. J. Food Plant Chem. 5:1-4, (2008).
6. Ahmed K.U., Pal-Phul O., Shak-Shabji., (In Bengali) 5th ed. Mrs Mumtaj Kamal Mirpur,Dhaka, Bangladesh, pp:400, (1995).
7. Akinyele B.O., Temikotan T., International Journal of Agricultural Research, 2: 165 – 169, (2007).
8. Arapitsas P., Identification and quantification of polyphenolic compounds from okra seeds and skins. Food Chem. 110:1041-1045, (2008).
9. Bates D. M., Notes on the cultivated Malvaceae 2, *Abelmoschus*. *Baileya*.16. 99-112, (1968).
10. Benjawan C., Chutichudet P., Kaewsit S., Effect of green manures on growth yield and quality of green okra (*Abelmoschus esculentus* L) har lium cultivar. Pakistan J. Biological Sci. 10: 1028-1035, (2007).
11. Borssum W., Van I., Malesian Malvaceae revised. *Blumea* 14:1- 251, (1966).
12. Brar K.S., Arora K.S., Ghai T.R., Losses in fruit yield of okra due to *Earias* spp. as influenced by dates of sowing and varieties. J Insect Sci 7(2):133–135, (1994).
13. Chauhan D.V.S., Vegetable production in India, Ram Prasad and Sons, India (1972).
14. Charrier A., Genetic resources of genus *Abelmoschus* Med. (Okra). IBPGR, Rome. Siesmonsma, J.S. 1991. International Crop Network Series. Report of an international workshop on okra genetic resources. IBPGR, Rome. 5:52-68, (1984).
15. Datta P.C., Naug A., A few strains of *Abelmoschus esculentus* (L.) Moench their karyological in relation to phylogeny and organ development. Beitr. Biol. Pflanzen. 45: 113-126, (1968)
16. Doijode S.D., Seed storage of horticultural crop. Food Product Press, New York, USA, (2001).
17. Dilruba S., Hasanuzzaman M., Karim R., Nahar K., Yield response of okra to different sowing time and application of growth hormones. J. Hortic. Sci. Ornamental Plants 1: 10-14, (2009).
18. Farooq A., Umer R., Muhammad A., Muhammad N., "Okra (*Hibiscus esculentus*) seed oil for biodiesel production". Applied Energy 87 (3): 779–785, (2010).
19. Felter., Harvey Wickes., Lloyd., John Uri., King's American Dispensatory., Retrieved 27 November 2011, (1898).

20. Givord L., Denboer L., Insect transmission of okra mosaic virus in the Ivory Coast. *Annals Appl. Biol.* 94: 235-241, (1980).
21. Gopalan C., Sastri S.B.V., Balasubramanian S., Nutritive value of Indian foods, National Institute of Nutrition (NIN), ICMR, India, (2007).
22. Hochreutimer B.P.G., Centres of origin for family Malvaceae. *Candolla* . 2:79, (1924).
23. Indian Horticulture database, Ministry of Agriculture, Government of India, (2011).
24. International Board for Plant Genetic Resources IBPGR, International Workshop on Okra Genetic Resources, Rome: International Board for Plant Genetic Resources, (1991).
25. Kahlon T.S., Chapman M.H., Smith G.E., In vitro binding of bile acids by okra beets asparagus eggplant turnips ree beans carrots and cauliflower. *Food Chem.* 103: 676-680, (2007).
26. Kendall C.W.C., Jenkins D.J.A., A dietary portfolio: maximal reduction of low-density lipoprotein cholesterol with diet. *Current Atherosclerosis Reports* 6:492-498, (2004).
27. Kundu B.C., Biswas C., Anatomical characters for distinguishing *Abelmoschus* spp. and *Hibiscus* spp. *Proc. Indian Sci. Cong.* 60: 295-298, (1973).
28. Kumar R., Patil M.B., Patil S.R., Paschapur M.S., Evaluation of *Abelmoschus esculentus* mucilage as suspending agent in paracetamol suspension. *Intern. J. PharmTech Res.* 1: 658-66, (2009).
29. Kumar S., Dagnoko S., Haougui A., Ratnadass A., Pasternak D., Kouame C., Okra (*Abelmoschus* spp.) in West and Central Africa: potential and progress on its improvement. *African J. Agric. Res.* 5: 3590-3598, (2010).
30. Lamont W., Okra a versatile vegetable crop. *Hort. Technol.* 9: 179-184, (1999).
31. Lengsfeld C., Titgemeyer F., Faller G., Hensel A., Glycosylated compounds from okra inhibit adhesion of *Helicobacter pylori* to human gastric mucosa. *J. Agric. Food Chem.* 52:1495-1503, (2004).
32. Linnaeus C., *Species Plantarum*. Vol I & II. Stockholm, (1753). 13. Medikus F.K., Ueber einige Kunstliche Geschlechter aus der Malvenfamilie, den der Klasse der, Monadelphien. 45-46, (1787).
33. Liu I.M., Liou S.S., Lan T.W., Hsu F.L., Cheng J.T., Myricetin as the active principle of *Abelmoschus moschatus* to lower plasma glucose in streptozotocin-induced diabetic rats. *Planta Medica* 71: 617-621, (2005).
34. Markose B.L., Peter K.V., Okra review of research on vegetable and tuber crops. Kerala Agricultural University Press, Kerala, India, (1990).
35. Moaward F.G., Abdelwhab B.M., Abdelnahun F.M., Shehaya F.W., *Annual of Agricultural science*, 21: 603 – 613, (1984).
36. Moyin-Jesu E.I., Use of plant residues for improving soil fertility pod nutrients root growth and pod weight of okra *Abelmoschus esculentum* L. *Bioresour. Tech.* 98: 2057-2064, (2007).
37. Moekchantuk T., Kumar P., Export okra production in Thailand. Inter-country programme for vegetable IPM in South & SE Asia phase II Food & Agriculture Organization of the United Nations, Bangkok, Thailand, (2004).

38. National Research Council, "Okra". Lost Crops of Africa: Volume II: Vegetables. Lost Crops of Africa. 2. National Academies Press. ISBN 978-0-309-10333-6. Retrieved 2008-07-15, (2006-10-27).
39. Nandkarni K.M., Indian Meteria Medica. Nadkarni and Co Bombay, (1927)
40. Ndunguru J., Rajabu A.C., Effect of okra mosaic virus disease on the above-ground morphological yield components of okra in Tanzania. *Scientia Horticulturae* 99: 225-235, (2004).
41. Okra Seed", Retrieved 10-17, (2012).
42. Owolarafe O.K., Shotonde H.O., Some physical properties of fresh okra fruit. *J. Food Engin.* 63: 299-302, (2004).
43. Purseglove J.W., Tropical crop dicotyedons. Language Book Society Longman, London, UK, (1987).
44. Qhureshi Z., Breeding investigation in bhendi (*Abelmoschus esculentus* (L.) Moench). Master Thesis, University of Agriculture Sciences, GKVK, Bangalore, (2007).
45. Radake S.G., Undirwade R.S., Seasonal abundance and insecticidal control of shoot and fruit borer, *Earais* spp. on okra, *Abelmoschus esculentus* (L.). *Indian J. Ent.*, 43: 283-287, (1981).
46. Rao P.S., Rao P.U., Serikeran B., Serun cholesterol, triglycerides and total total fatty acid of rates in response to okra (*Hibiscus esculentus*) seed oil. *JAOCA* 68:433, (1991).
47. Rao P.U., Chemical composition and biological evaluation of okra (*Hibiscus esculentus*) seeds and their kernels. *Qual. Plant Food Human Nutr.*, 35:389-396, (1985).
48. Rashid M.H., Yasmin L., Kibria M.G., Mollik A., Hossain S., Screening of okra germplasm for resistance to yellow vein mosaic virus under field conditions. *Pakistan J. Plant Pathol.* 1: 61-62, (2002).
49. Rubatzky V.E., Yamaguchi M., World vegetables: principles, production, and nutritive values. Chapman and Hall, New York, USA, (1997). International Crop Network Series, Report of and nutritive values. Chapman and Hall, New York, USA, (1997).
50. Saifullah M., Rabbani M.G., Evaluation and characterization of okra (*Abelmoschus esculentus* L. Moench.) genotypes. *SAARC J. Agric.* 7: 92-99, (2009).
51. Sastry K.S.M., Singh S.J., Effect of yellow-vein mosaic virus infection on growth and yield of okra crop. *Indian Phytopathol.* 27: 294-297, (1974).
52. Savello P.H., Martin F.W., Hill J.M., Nutritional composition of okra seed meal. *J. Agric. Food Chem.* 28: 1163-1166, (1980).
53. Siemonsma J.S., Kouame C., *Abelmoschus esculentus* (L.) Moench. Wageningen Agricultural University, Wageningen, Netherlands, (2000).
54. Siemonsma J.S., La culture du gombo (*Abelmoschus* spp) legume fruit. Thesis Univ Wageningen, the Netherlands, (1982)
55. Siemonsmo J.S., West African okra. Morphological and cytological indications for the existence of amatural amphiploid of *Abelmoschus esculentus* (L.) Moench and *A. Manihot* (L.) Medikus. *Euphytica.* 31(1): 241-52, (1982).
56. Singh H.B., Bhatnagar A., Chromosome number in an okra from Ghana. *Indian J. Genet. Plant Breed.* 36:26-27, (1975).
57. Singh G., Brar K.S., Effects of dates of sowing on the incidence of *Amrasca biguttula biguttula* (Ishida) and *Earias* species on okra. *Indian J Ecol* 21(2):140-144, (1994).
58. Sinha S.N., Sharma S.P., Chakrabarti A.K., Effect of spotted bollworm infestation on seed quality of okra. *Seed Res.*, 6:161-164, (1978).

59. Terrell E.E., Winters H.F., Change in scientific names for certain crop plants. Hort.Sci. 9: 324-325, (1974).
60. Tindall H. D., Vegetable in the tropics Macmillan press ltd London first edition, (1983).
61. Varmudy V., Marketing survey need to boost okra exports. Department of economics, Vivekananda College, Puttur, Karnataka, India, (2011).

UNDER PEER REVIEW