

Evaluation of genetic diversity through D² statistic in Chickpea (*Cicer arietinum* L.)

Abstract

The present investigation was conducted at the Agriculture Research Farm, Department of Genetics and Plant Breeding, College of Agriculture, RVSKVV, Gwalior. Madhya Pradesh, India. The objective of the investigation was to assess the genetic diversity among 71 different chickpea genotypes in relation to their yield and its attributing traits. The experimental design employed was a complete randomized block design with two replications. A comprehensive set of observations was made on twelve distinct yield accrediting traits from five randomly selected plants within each genotype. Based on D² Statistics analysis, the 71 chickpea genotypes were classified into 26 distinct clusters. Conspicuously, the cluster with the highest numbers of genotypes was designated as cluster one. A noteworthy finding emerged from the analysis of intra-cluster distances, with cluster 16 displaying the greatest distance within its constituents. The evaluation of inter-cluster distances revealed significant dissimilarity between clusters 22 and 26, suggesting presence of considerable genetic differentiation between these clusters. Conversely, the inter-cluster distance was minimal between clusters 2 and 4, indicating a closer genetic relationship between these two groups. In terms of the genetic diversity analysis, it became evident that the yield related traits exerting the most substantial influence on the overall genetic divergence among the 71 chickpea genotypes were biological yield per plant, 100-seed weight, and the numbers of pods per plant. In contrast, the numbers of effective pods per plant contributed minimally to the overall genetic divergence. Based on the findings from both inter-cluster distances and individual performance assessments (*per se* performance), two specific genotypes viz., ICCV 201207 and SAGL 22-121, were short out as promising for inclusion in a hybridization programme. These findings contribute to a deeper understanding of chickpea genetic variability and lay the groundwork for further breeding programmes aimed to enhance chickpea crop productivity.

Key words: Chickpea, D² statistics, Diversity, Genotype, Clusters, Yield

Comment [B1]: Major revision is needed. There are two types of corrections; one is **Compulsory corrections** for its acceptance as full paper and other is **Optional corrections** for the improvement of the paper.

Comment [B2]: First capital letter for each word

Comment [B3]: Compulsory correction: Uniform style of the International Journal of Environment and Climate Change need to be thoroughly followed for the whole manuscript, as mentioned in the given 'SDI-Paper-template' of the Journal ;

Comment [B4]: ABSTRACT

Comment [B5]: Details

Comment [B6]: cluster 1

Comment [B7]:relationship among the genotypes between....

Comment [B8]: You must explain in results and discussion part.

Or you must need to add Table 1 as instructed.

Introduction

Chickpea (*Cicer arietinum* L.), commonly known as garbanzo bean or Bengal gram, is an ancient and globally significant leguminous crop cultivated for its nutritious seeds [1-2]. Belonging to the Fabaceae family, chickpea holds historical and cultural importance, being cultivated and consumed across various regions for millennia [3-4]. It is a self-pollinated crop [5] and known as King of pulses. Chickpea is a cool season legume with a genome size of ~738 mb [6] and originated in south eastern Turkey and Syria. Two primary categories of chickpea germplasm are recognized: "Kabuli" and "Desi." Kabuli chickpea (*Cicer arietinum* var. *macrosperma*) is characterized by its sizeable creamy-colored seeds, accompanied by white flowers, and lacks anthocyanin pigmentation. In contrast, Desi chickpea (*Cicer arietinum* var. *microsperma*) is distinguished by its smaller seeds of diverse colors, accompanied by purplish flowers, and contains anthocyanin pigments [7]. The diminished productivity of chickpea is attributed to a combination of different biotic and abiotic factors, as substantiated by recent research [8-19].

The consumption of chickpeas enjoys popularity across numerous global regions, primarily attributed to its remarkable nutritional excellence. Within the chickpea seed matrix, a harmonious composition of essential components prevails, encompassing 50–58% carbohydrates, 15–22% protein, 7–8% moisture, 3.8–10.20% fat, and <1% micronutrients [20-21; 10]. The average protein content found in chickpeas is approximately 18%, with kabuli chickpeas containing around 18.4% ranging from 16.2% to 22.4%, and desi chickpeas contain about 18.2% in range of 15.6% to 21.4%. This protein content surpasses that of lentils and field peas [22]. As a significant member of the legume family, chickpea holds a pivotal position among pulses. Notably devoid of cholesterol, it boasts remarkable nutritional value, being rich in dietary fiber (DF), vitamins, and minerals. Although its methionine content ranges between 1.3% to 1.6% and cysteine content containing sulfur is around 2.5% to 3.0%, these sulfur-containing amino acids are typically found in relatively lower amounts. However, when consumed alongside cereals deficient in lysine amino acid, chickpea's nutritional profile becomes complementary, ensuring a balanced diet [23].

Assessing genetic diversity through the measurement of genetic distance proves to be an invaluable tool in crop breeding [24-40] including chickpea [4-5], aiding in the selection of

Comment [B9]: 1. INTRODUCTION

Arial, Bold, 11 font, caps

Please follow the given 'SDI-Paper-template' of the Journal for the whole manuscript;

Comment [B10]: Please give reference here

Comment [B11]: make space

suitable parent plants to facilitate novel genetic recombination and enhance grain yield [41]. The presence of genetic diversity stands as a pivotal factor in the formulation of effective and fruitful breeding initiatives [22]. Employing cluster analysis serves as a suitable approach to delineate familial relationships and genetic affinities among genotypes, enabling the quantification of genetic distance between them. D^2 statistic, a variant of generalized distance, pioneered by Mahalanobis [42], offers a means to evaluate genetic diversity among genotypes. The prosperity of any crop enhancement programme is intricately linked to the genetic diversity within the existing germplasm pool [43-44, 24-40]. Subsequently, a quantitative assessment of the genetic diversity present within a population assists plant breeders in the meticulous selection of desirable parental candidates for breeding undertakings. In the present investigation, the extent of genetic diversity present in a set of seventy-one chickpea genotypes for various yield attributing traits were evaluate by means of D^2 statistic for further enhancing our understanding of the species for genetic enhancement.

Materials and method

Experimental material

In this investigation, Mahalanobis (D^2) statistics was utilized to analyze genetic diversity present among 71 distinct chickpea genotypes.

Experimental site

The experiment was conducted at the Agriculture Research Farm, Department of Genetics and Plant Breeding, College of Agriculture, RVSKVV, Gwalior, Madhya Pradesh, India.

Experimental design and management

The genotypes were evaluated under a randomized block design with two replications during the *Rabi* seasons of 2021 and 2022. The observations were recorded, with a pooled data of both the seasons and further analyzed. Each genotype was planted across four rows, each spanning a length of 3 meters. The spatial arrangement maintained a row-to-row distance of 30 units and a plant-to-plant distance of 15 cm. The entire cultivation process adhered to recommended agronomic practices.

Data collection

Comment [B12]: 2. MATERIALS AND METHOD

Arial, Bold, 11 font, caps

Comment [B13]: ARIAL, BOLD, 11 FONT

First capital letter for each word

Comment [B14]: ARIAL, BOLD, 11 FONT

First capital letter for each word

Comment [B15]: ARIAL, BOLD, 11 FONT

First capital letter for each word

Comment [B16]: ARIAL, BOLD, 11 FONT

First capital letter for each word

The recorded data encompassed twelve distinct morpho-physiological attributes, including days to 50 % flowering, days to maturity, plant height, numbers of primary and secondary branches per plant, numbers of pods per plant, numbers of effective pods per plant, numbers of seeds per pod, 100-seed weight, harvest index, biological yield per plant and seed yield per plant. To ensure comprehensive observations, a random selection of five plants was made from each replication of every genotype, serving as the foundation for data collection across all measured traits.

Statistical analysis

The multivariate analysis, specifically employing the D^2 statistics proposed by Mahalanobis [42], has emerged as a robust and indispensable tool for elucidating the extent of differentiation observed among biological populations at the genotypic level [43]. Through this analytical approach, one can effectively assess the relative contributions of distinct components to the overarching divergence phenomenon, spanning both inter and intra-cluster levels. Remarkably, the D^2 statistical analysis, as pioneered by Mahalanobis, was complemented by the construction of a dendrogram, which not only offers a comprehensive visual representation but also provides insights into the intricate genetic relationships inherent within the investigated populations. For the rigorous execution of statistical computations and dendrogram construction, the specialized Statistical software WINDOSTAT version 9.2 provided by INDOSTAT Services, Hyderabad, India was harnessed, thereby ensuring a heightened level of precision, robustness, and reliability in the resulting analytical outcomes.

Results and Discussion

Clustering of genotypes

D^2 statistical analysis was employed to assess the genetic divergence among 71 genotypes. The Tocher's method [44] was used to form clusters based on the calculated D^2 values. Consequently, the 71 genotypes were categorized into four main clusters and 22 minor clusters. Among these clusters, cluster 1 was found to be polygenotypic, encompassing 32 genotypes. On the other hand, Clusters 11, 13, and 16 were oligogenotypic, comprising more than two genotypes each. The remaining clusters consisted of only one genotype, rendering them monogenotypic (Fig. 1). Cluster 1 stood out with its significant count of 32 genotypes, indicating a limited genetic

Comment [B17]: ARIAL, BOLD, 11 FONT

Comment [B18]: 3. RESULTS AND DISCUSSION

Comment [B19]: Optional not compulsory:
Please add analysis of variance or and estimation of genetic parameters, if possible.

Comment [B20]: ARIAL, BOLD, 11 FONT

First capital letter for each word

Comment [B21]: Please send this sentence to statistical analysis part like

.....The Tocher's method [44] was used to form clusters based on the calculated D^2 values. Through this analytical.....

Comment [B22]: Compulsory correction:
Please give a separate table. Title of the Table: Clustering pattern among 71 genotypes of chickpea by Tocher's Method. It will contains cluster no., no. of genotypes and genotypes (its name) in separated columns.

The title may be as –

Table 1. The distribution of 71 chickpea genotypes into different clusters based on D^2 statistic.

This is must for the acceptance of this paper.

divergence among them. This similarity could potentially be attributed to their shared ancestry in the base population from which they evolved. It is worth noting that the convergence of similar phenotypes within a cluster might arise from focused unidirectional selection on specific traits or a cluster of linked traits across different geographical regions.

These factors could encompass a range of influences such as natural and artificial selection pressures, the exchange of breeding materials, genetic drift, and environmental variability. As such, the intricate interplay of these multifaceted forces appears to play a pivotal role in shaping and sustaining the observed genetic diversity landscape. In a congruent investigation, Gupta *et al.* [45] conducted a genetic divergence study encompassing 25 chickpea genotypes, ultimately arranging them into seven clusters through the utilization of Tocher's method. Koinain *et al.* [46] undertook a study involving thirty diverse chickpea genotypes, classifying them into eight distinct clusters, Rajkumar *et al.* [47] partitioned a collection of 100 distinct chickpea genotypes into sixteen clusters.

Intra and inter cluster divergence D^2 values

When examining a range of important agronomic traits like yield potential, disease resistance, growth duration, and seed quality, the Euclidean distance method is frequently employed. This method facilitates the identification of hierarchical patterns within the analyzed accessions. By grouping these accessions based on specific traits, it enables the identification of promising pairs for crossbreeding purposes in the context of chickpea studies [4-5]. This approach serves to assess both the likeness and distinctiveness among different accessions, while also quantifying the extent of a particular trait's expression. Past research, exemplified by studies conducted by Syed *et al.* [48] and Malik *et al.* [49], has demonstrated the effectiveness of this method in characterizing and evaluating chickpea accessions.

The average D^2 values for both intra and inter-cluster distances, along with the nearest clusters based on D^2 values are presented in Table 1 and Fig. 2. Intra-cluster distances spanned from zero to 30.50. Cluster 16 exhibited the highest intra-cluster distance at 30.50, followed by cluster 13 (25.02), cluster 11 (22.82), and cluster 1 (19.7). Other clusters had an intra-cluster distance of 0.00 due to their single-genotype composition. Furthermore, inter-cluster distances ranged between 6.59 to 186.56. The most substantial inter-cluster distance was observed between cluster 22 and 26 (199.44), succeeded by cluster 24 and 26 (186.56), cluster 16 and cluster 19

Comment [B23]: ARIAL, BOLD, 11 FONT
.....

Comment [B24]: Table 2

Comment [B25]: cluster 26

Comment [B26]: cluster 26

(153.40), and cluster 16 and cluster 20 (153.37). Conversely, the smallest inter-cluster distances were observed between cluster 2 and cluster 4 (6.59), cluster 3 and cluster 6 (7.54), and cluster 2 and cluster 5 (8.71). These extensive inter-cluster distances signify that the genotypes within these clusters exhibit a wide array of genetic diversity. They hold promising potential for utilization in chickpea improvement through hybridization programmes. To enhance the likelihood of generating favorable recombinants in segregating generations, crossing between genetically diverse genotypes from clusters with substantial inter-cluster distances would be advantageous. These are the similar work done by Janghel *et al.* [50], non-hierarchical Euclidean cluster analysis categorized 60 chickpea genotypes into seven distinct clusters. Katkani *et al.* [51] identified nine clusters among forty-two desi chickpea lines through divergence analysis. Moreover, Mihoariya *et al.* [4] grouped eighty-three chickpea genotypes into twelve clusters after evaluating the nature and extent of genetic divergence.

Cluster mean values

The cluster mean values for diverse traits have been presented in Table 2 revealing significant differences among the studied traits. The data highlights specific clusters with distinct trait averages. Cluster 25 displays the highest mean for days to 50% flowering (61.17), whilst cluster 19 exhibits the lowest (49.50). Cluster 2 records the maximum mean for days to maturity (121.50), contrasting with the lowest mean in cluster 17 (112.17). In terms of plant height, cluster 21 demonstrates the highest mean (75.67), whereas cluster 19 reflects the minimum (42.99). The numbers of primary branches per plant is maximized in cluster 24 (4.87), whereas cluster 10 displays the least (2.55). For the numbers of secondary branches per plant, cluster 18 contributed the highest mean (10.88), while cluster 10 showing the lowest (6.37). Regarding pod-related traits, cluster 25 comprehends the highest mean for the numbers of pods per plant (55.50), while cluster 19 presents the lowest (32.17). Similarly, cluster 25 demonstrates the maximum mean for the numbers of effective pods per plant (52.17), whereas cluster 19 showing the minimum (29.17). Cluster 9 displays the highest mean for the numbers of seeds per pod (1.73), while clusters 24 consisted the lowest (1.18). For 100- seed weight, cluster 12 exhibits the highest mean (29.77), contrasting with the lowest mean in cluster 6 (17.41). Harvest index is maximized in cluster 18 (68.57), whereas cluster 6 presents the lowest (40.59). In terms of yield, cluster 26 had the highest mean for biological yield per plant (51.07), In contrast, cluster 23

Comment [B27]: ARIAL, BOLD, 11 FONT.....

Comment [B28]: Table 3

Comment [B29]: unit?

Comment [B30]: unit?

Comment [B31]: unit?

indicates the lowest value (29.14). On the other hand, cluster 13 exhibits the highest mean for seed yield per plant (25.59), while cluster 22 reveals the lowest (13.82).

Comment [B32]: unit?

Comment [B33]: unit?

Comment [B34]: unit?

Likewise, Gediya *et al.* [52] also conducted a study focusing on 13 distinct characters within the realm of chickpea. The observed cluster mean values exhibited a broad spectrum of averages across the clusters for various traits. Notably, cluster XV showed the highest mean values for numbers of pods (136.00) and seeds per plant (201.87), while registering the lowest mean value for the first pod height (33.87 cm). Within cluster XII the highest mean values were recorded for seed yield per plant (38.77 g) and harvest index (54.89%). Cluster X displayed the highest cluster mean value for 100-seed weight (42.34 g), whereas cluster XVI exhibited the highest mean average for plant height (105.87 cm). Conversely, cluster XIV exhibited the lowest mean worths for days to 50 percent flowering (44.00 days) and days to maturity (107.33 days), while simultaneously showcasing the highest mean value for seeds per pod (1.93). Cluster I displayed the highest mean value for primary branches per plant (3.13), whereas cluster VI exhibited the highest mean value for numbers of secondary branches per plant (12.70). In the study conducted by Reddy *et al.*[53], cluster II exhibited the highest mean values for a range of traits, including the numbers of secondary branches per plant, pod count per plant, seed count per pod, and seed count per plant. Cluster IV, conversely, displayed the highest mean potential for the numbers of primary branches per plant, biological yield per plant, and seed yield per plant. Furthermore, the mean values for cluster V were most pronounced in traits like plant height and harvest index. Cluster VI demonstrated its highest mean values in the context of 100 - seed weight, whilst cluster VII showcased an early disposition for traits such as days to 50% flowering, days to 50% pod setting, and days to maturity.

Comment [B35]: cluster 15 or cluster XV?

Please make it similar for all cases in the whole manuscript.

Comment [B36]: cluster 12 or cluster XII?

Because in previous paragraph, you mention it as cluster 12.

Please make it similar for all cases in the whole manuscript.

Comment [B37]:

Comment [B38]:

Comment [B39]:

Comment [B40]:

Comment [B41]: cluster ?

Comment [B42]: Cluster ?

Comment [B43]: cluster ?

Comment [B44]: Cluster ?

Comment [B45]: No space as 100-seed weight

Comment [B46]: cluster ?

Contribution of individual characters towards genetic divergence

Comment [B47]: ARIAL, BOLD, 11 FONT.....

The percentage contributions to genetic divergence attributed to various characters are presented in (Table 3, Fig. 3). Among these, the character biological yield per plant exhibited the most substantial contribution at 38.07%, followed by 100-seed weight (28.01%), numbers of pods per plant (14.57 %), plant height (6.4%), seed yield per plant (6.24 %), harvest index (2.37%), numbers of primary branches per plant (1.77%), days to 50% flowering (0.76 %), numbers of seeds per pod (0.64%), days to maturity (0.56 %), numbers of secondary branches per plant (0.52%) and numbers of effective pods per plant (0.08%).

Comment [B48]: in Table 4 and Fig. 3.

Consistent with earlier studies, Pahre *et al.* [54], Kuldeep *et al.* [55]), and Thakur *et al.* [56] reported similar trends wherein 100-seed weight, numbers of pods per plant, and days to 50% flowering emerged as the primary contributors to the observed genetic diversity. This alignment underscores the significance of these specific traits in driving the divergence documented within the chickpea population.

Conclusion

In this ongoing divergence analysis, assessing the proportional influence of each trait on genetic divergence offers valuable insights that can aid plant breeders in discerning and selecting superior genotypes from the existing germplasm pool. These chosen genotypes hold the potential to serve as optimal parent candidates in forthcoming crop enhancement initiatives. Concurrently, the present investigation unveils a significant level of diversity within the examined assemblage of 71 chickpea genotypes. This diversity is particularly driven by the traits of biological yield per plant, 100-seed weight, and the numbers of pods per plant. Among the clusters, cluster I stands out as the most populous, encompassing 32 genotypes. Furthermore, the analysis underscores a substantial inter-cluster disparity, prominently observed between clusters 22 and 26. The genotypes within these distinct clusters hold potential significance as parental candidates for generating transgressive segregants in future breeding endeavors. This elucidates their viability in augmenting the genetic variability of chickpea cultivars.

References

1. Asati R, Tripathi MK, Tiwari S, Yadav RK, Tripathi N. Molecular breeding and drought tolerance in chickpea. *Life (Basel)*. 2022;12(11):1846. doi: [10.3390/life12111846](https://doi.org/10.3390/life12111846). Pubmed:[36430981](https://pubmed.ncbi.nlm.nih.gov/36430981/).
2. Tiwari PN, Tiwari S, Sapre S, Babbar A, Tripathi N, Tiwari S, Tripathi MK. Screening and selection of drought-tolerant high-yielding chickpea genotypes based on physio-biochemical selection indices and yield trials. *Life (Basel)*. 2023;13(6):1405. doi: [10.3390/life13061405](https://doi.org/10.3390/life13061405), Pubmed:[37374187](https://pubmed.ncbi.nlm.nih.gov/37374187/).
3. Gupta S, Arora S, Mirza N, Pande S, Latha M, Rathore A et al. The chickpea genomic web resource: visualization and analysis of the Desi-type *Cicer arietinum* nuclear genome for comparative exploration of legume genomes. *Plant Cell Physiol*. 2019;60(1):e12-.

Comment [B49]: Optional suggestion not compulsory: Please add a separate Table and sub-heading like-

Table 5. Eigene Values, contribution of variance and factor loading for the Principal Component Analysis for different traits. **if possible**

....for explaining the Principal component analysis (PCA) that provides expression of the maximum contributor towards the variability.

It may improve the paper.

Comment [B50]: 4. CONCLUSION

Arial, Bold, 11 font, left aligned, caps

Comment [B51]: 5. REFERENCES

Arial, Bold, 11 font, left aligned, caps

Comment [B52]: Compulsory: Uniform style of the International Journal of Environment and Climate Change need to be thoroughly followed, as mentioned in the given 'SDI-Paper-template' of the Journal; especially for References section.

4. Mihoariya M, Tiwari S, Yadav RK, Asati R, Solanki RS, Tiwari PN et al. Genetic variability and diversity analysis for yield and its associated traits in chickpea (*Cicer arietinum* L.). *Curr J Appl Sci Technol.* 2023;42(16):17-33. doi: [10.9734/cjast/2023/v42i164127](https://doi.org/10.9734/cjast/2023/v42i164127).
5. Ningwal R, Tripathi MK, Tiwari S, Yadav RK, Tripathi N, Solanki RS et al. Assessment of genetic variability, correlation and path coefficient analysis for yield and its attributing traits in chickpea (*Cicer arietinum* L.). *J Pharm Innov.* 2023a;12(3):4851-9.
6. Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG et al. Draft genome sequence of chickpea (*Cicer arietinum* L.) provides a resource for trait improvement. *Nat Biotechnol.* 2013;31(3):240-6. doi: [10.1038/nbt.2491](https://doi.org/10.1038/nbt.2491), Pubmed:[23354103](https://pubmed.ncbi.nlm.nih.gov/23354103/).
7. Berger JD, Ali M, Basu PS, Chaudhary BD, Chaturvedi SK, Deshmukh PS et al. Genotype by environment studies demonstrate the critical role of phenology in adaptation of chickpea (*Cicer arietinum* L.) to high and low yielding environments of India. *Field Crops Res.* 2006;98(2-3):230-44. doi: [10.1016/j.fcr.2006.02.007](https://doi.org/10.1016/j.fcr.2006.02.007).
8. Sahu VK, Tiwari S, Tripathi MK, Gupta N, Tomar RS, Yasin M. Morphophysiological and biochemical traits analysis for Fusarium wilt disease using gene-based markers in desi and Kabuli genotypes of chickpea (*Cicer arietinum* L.). *Indian J Genet.* 2020a;80:16.
9. Sahu VK, Tiwari S, Gupta N, Tripathi MK, Yasin M. Evaluation of physiological and biochemical contents in desi and Kabuli chickpea. *Legume Res.* 2020b. doi: [10.18805/LR-4265](https://doi.org/10.18805/LR-4265).
10. Gupta N, Tiwari S, Tripathi MK, Bhagyawant SS. Antinutritional and protein-based profiling of diverse desi and wild chickpea accessions. *Curr J Appl Sci Technol.* 2021;40(6):7-18. doi: [10.9734/cjast/2021/v40i631312](https://doi.org/10.9734/cjast/2021/v40i631312).
11. Jain N, Babbar A, Kumawat S, Yadav RK, Asati R. Correlation and path coefficient analysis in the promising advance chickpea lines. *J Pharm Innov.* 2022;11(5):2124-8.
12. Solanki RS, Babbar A, Tripathi N. Genetic diversity analysis in Kabuli chickpea (*Cicer arietinum* L.) genotypes based on quantitative traits and molecular markers. *Bangladesh J Bot.* 2022;51(3):581-7. doi: [10.3329/bjb.v51i3.62005](https://doi.org/10.3329/bjb.v51i3.62005).
13. Tripathi N, Tripathi MK, Tiwari S, Payasi DK. Molecular breeding to overcome biotic stresses in soybean: update. *Plants (Basel).* 2022;11(15):1967. doi: [10.3390/plants11151967](https://doi.org/10.3390/plants11151967), Pubmed:[35956444](https://pubmed.ncbi.nlm.nih.gov/35956444/).

14. Asati R, Tripathi MK, Yadav RK, Tiwari S, Chauhan S, Tripathi N et al. Morphological description of chickpea (*Cicer arietanum* L) genotypes using DUS characterization. *Int J Environ Clim Change*. 1341;2023(9)(9):1321-41. doi: [10.9734/ijecc/2023/v13i92361](https://doi.org/10.9734/ijecc/2023/v13i92361).
15. Yadav RK, Tripathi MK, Tiwari S, Tripathi N, Asati R, Patel V et al. Breeding and genomic approaches towards development of Fusarium wilt resistance in chickpea. *Life (Basel)*. 2023a;13(4):988. doi: [10.3390/life13040988](https://doi.org/10.3390/life13040988), Pubmed:[37109518](https://pubmed.ncbi.nlm.nih.gov/37109518/).
16. Rajput S, Jain S, Tiwari S, Sikarwar RS, Tripathi N, Barela A, Tripathi MK. Evaluation of gene-based markers against Fusarium wilt disease in chickpea (*Cicer arietinum* L.). *Curr J Appl Sci Technol*. 2023a;42(6):26-32. doi: [10.9734/cjast/2023/v42i64074](https://doi.org/10.9734/cjast/2023/v42i64074).
17. Ningwal R, Tripathi MK, Tiwari S, Asati R, Yadav RK, Tripathi N, Yasin M. Identification of polymorphic SSR marker and diversity analysis in a set of desi chickpea genotypes, *Biological Forum-An [International journal]*. 2023b;15(3):45-51.
18. Rajput S, Jain S, Tiwari S, Barela A, Chauhan S, Tiwari PN et al. Biochemical characterization of chickpea (*Cicer arietinum* L.) genotypes. *Plant Cell Biotechnol Mol Biol*. 2023b;24(3-4):1-9. doi: [10.56557/pcbmb/2023/v24i3-48239](https://doi.org/10.56557/pcbmb/2023/v24i3-48239).
19. Tutlani A, Kumar R, Banshidhar KumariS, Chouhan S. Correlation and path analysis for yield and its phenological, physiological, morphological and biochemical traits under salinity stress in chickpea (*Cicer arietinum* L.). *Int J Bio-Resource Stress Manag*. 2023;14(6):878-90. doi: [10.23910/1.2023.3519](https://doi.org/10.23910/1.2023.3519).
20. Jukanti AK, Gaur PM, Gowda CLL, Chibbar RN. Nutritional quality and health benefits of chickpea (*Cicer arietinum* L.): a review. *Br J Nutr*. 2012;108;Suppl 1:S11-26. doi: [10.1017/S0007114512000797](https://doi.org/10.1017/S0007114512000797), Pubmed:[22916806](https://pubmed.ncbi.nlm.nih.gov/22916806/).
21. USDA; 2021. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/174288/nutrients>.
22. Upadhyaya HD, Bajaj D, Narnoliya L, Das S, Kumar V, Gowda CLL, et al. Genome-wide scans for delineation of candidate genes regulating seed-protein content in chickpea. *Front Plant Sci*. 2016;7:302. doi: [10.3389/fpls.2016.00302](https://doi.org/10.3389/fpls.2016.00302), Pubmed:[27047499](https://pubmed.ncbi.nlm.nih.gov/27047499/).
23. Heuzé V, Tran G, Boudon A, Bastianelli D, Lebas F. Chickpea (*Cicer arietinum*); 2015. <https://www.feedipedia.org/node/319>.
24. Barfa D, Tripathi MK, Kandalkar VS, Gupta JC, Kumar G. Heterosis and combining ability analysis for seed yield in Indian mustard [*Brassica juncea* (L.) Czern & Coss.]. *Ecol Environ Conserv*. 2017;23;Suppl:75-83.

25. Baghel R, Sharma AK, Tiwari S, Tripathi MK, Tripathi N. Genetic diversity analysis of Indian mustard (*Brassica spp.*) germplasm lines using SSR molecular markers. *Int J Curr Microbiol Appl Sci.* 2020;9(12):137-43. doi: [10.20546/ijcmas.2020.912.018](https://doi.org/10.20546/ijcmas.2020.912.018).
26. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Trivedi HK. Morphological and molecular screening of soybean genotypes against yellow mosaic virus disease. *Legume Res.* 2020;(OF). doi: [10.18805/LR-4240](https://doi.org/10.18805/LR-4240).
27. Rajpoot NS, Tripathi MK, Tiwari S, Tomar RS, Kandalkar VS. Characterization of Indian mustard germplasm on the basis of morphological traits and SSR markers. *Curr J Appl Sci Technol.* 2020;39:300-11. doi: [10.9734/cjast/2020/v39i4831234](https://doi.org/10.9734/cjast/2020/v39i4831234).
28. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Gupta N, Sharma A. Morphological and physiological performance of Indian soybean [*Glycine max (L.) Merrill*] genotypes in respect to drought. *Legume Res.* 2021;(Of). doi: [10.18805/LR-4550](https://doi.org/10.18805/LR-4550).
29. Sharma A, Tripathi MK, Tiwari S, Gupta N, Tripathi N, Mishra N. Evaluation of soybean (*Glycine max L.*) genotypes on the basis of biochemical contents and anti-oxidant enzyme activities. *Legume Res.* 2021;(Of). doi: [10.18805/LR-4678](https://doi.org/10.18805/LR-4678).
30. Shyam C, Tripathi MK, Tiwari S, Tripathi N. Genetic components and diversity analysis in Indian mustard [*Brassica juncea (Linn.) Czern & Coss*] based on different morpho-physiological traits. *Curr J Appl Sci Technol.* 2021a;40(20):34-57. doi: [10.9734/CJAST/2021/v40i2031462](https://doi.org/10.9734/CJAST/2021/v40i2031462).
31. Shyam C, Tripathi MK, Tripathi N, Tiwari S, Sikarwar RS. Genetic variations in fatty acids and oil compositions among 188 Indian Mustard *Brassica juncea (Linn.) Czern & Coss* genotypes. *Curr J Appl Sci Technol.* 2021b;40(46):9-28. doi: [10.9734/CJAST/2021/v40i4631629](https://doi.org/10.9734/CJAST/2021/v40i4631629).
32. Verma K, Tripathi MK, Tiwari S, Tripathi N. Analysis of genetic diversity among *Brassica juncea* genotypes using morpho-physiological and SSR markers. *Int J Curr Microbiol Appl Sci.* 2021;10(1):1108-17. doi: [10.20546/ijcmas.2021.1001.134](https://doi.org/10.20546/ijcmas.2021.1001.134).
33. Yadav PK, Tiwari S, Kushwah A, Tripathi MK, Gupta N, Tomar RS, Kandalkar VS. Morpho-physiological characterization of bread wheat genotypes and their molecular validation for rust resistance genes Sr2, Sr31 and Lr24. *Proc Indian Natl Sci Acad.* 2021;87(3):534-45. doi: [10.1007/s43538-021-00049-y](https://doi.org/10.1007/s43538-021-00049-y).

34. Shyam C, Tripathi MK, Tiwari S, Tripathi N, Sikarwar R. S. Morphophysiological variations and genetic components analysis in Brassica juncea (Linn.) Czern & Coss. In book: Research Developments. Interface Sci Technol. 2022;1:98-126. doi: [10.9734/bpi/rdst/v1/6009F](https://doi.org/10.9734/bpi/rdst/v1/6009F).
35. Sharma S, Tripathi MK, Tiwari S, Solanki RS, Chauhan S, Tripathi N et al. The Exploitation of genetic variability and trait association analysis for diverse quantitative traits in bread wheat (*Triticum aestivum* L.). Curr J Appl Sci Technol. 2023a;42(8):19-33. doi: [10.9734/cjast/2023/v42i84087](https://doi.org/10.9734/cjast/2023/v42i84087).
36. Sharma S, Tripathi MK, Tiwari S, Solanki RS, Chauhan S, Tripathi N et al. Discriminant function analysis for yield improvement in bread wheat (*Triticum aestivum* L.). The Pharm Innov J. 2023b;12(5):224-32.
37. Shrivastava A, Tripathi MK, Solanki RS, Tiwari S, Tripathi N, Singh J, Yadav RK. Genetic correlation and path coefficient analysis of yield attributing parameters in Indian mustard. Curr J Appl Sci Technol. 2023;42(7):42-58. doi: [10.9734/cjast/2023/v42i74079](https://doi.org/10.9734/cjast/2023/v42i74079).
38. Makwana K, Tiwari S, Tripathi MK, Patel V. Morphological diversity analysis of foxtail millet (*Sateria italica* (L.) Beauv.) using qualitative and quantitative traits. Scientists. 2023;2(2):319-26.
39. Rajpoot P, Tripathi MK, Solanki RS, Tiwari S, Tripathi N, Chauhan S et al. Genetic variability and multivariate analysis in pearl millet (*Pennisetum glaucum* (L.) R. Br.) germplasm lines. The Pharm Innov J. 2023;12(4):216-26.
40. Yadav PK, Tripathi MK, Tiwari S, Chauhan S, Tripathi N, Sikarwar RS et al. Genetic components and variability assessment for grain yield and its accrediting traits in maize (*Zea mays* L.). IJECC. 2023;13(9):772-84. doi: [10.9734/ijecc/2023/v13i92298](https://doi.org/10.9734/ijecc/2023/v13i92298).
41. Poudel A, Thapa DB, Sapkota M. Assessment of genetic diversity of bread wheat (*Triticum aestivum* L.) genotypes through cluster and principal component analysis. Int J Exp Res Rev. 2017;11:1-9.
42. Mahalanobis PC. On the generalized distance in statistics. Proc Natl Inst Sci India. 1936;2(1):49-55.
43. Sneath PH, Sokal RR. Numerical taxonomy: the principles and practice of numerical classification. W H Freeman; 1973.

44. Rao CR. Advanced statistical methods in biometrical research. New York: Wiley & Sons; 1952.
45. Gupta S, Kaushal R, Kumar R, Sharma S. Genetic divergence studies in chickpea (*Cicer arietinum* L.). *Int J Plant Sci.* 2016;11(1):84-7.
46. Koinain S, Dikshit N, Dantu PK. Genetic diversity studies in chickpea (*Cicer arietinum* L.). *Int J Curr Microbiol Appl Sci.* 2015;4(6):438-44.
47. Rajkumar B, Ranjekar PK, Gupta VS. Genetic diversity studies reveal the genetic relationships among elite chickpea genotypes. *Crop Sci.* 2015;55(3):1133-42.
48. Syed MA, Islam MR, Hossain MS, Alam MM, Amin MN. Genetic divergence in chickpea (*Cicer arietinum* L.). *Bangladesh J Agric Res.* 2012;37(1):129-36. doi: [10.3329/bjar.v37i1.11184](https://doi.org/10.3329/bjar.v37i1.11184).
49. Malik SR, Shabbir G, Zubir M, Iqbal SM, Ali A. Genetic diversity analysis of morphogenetic traits in Desi chickpea (*Cicer arietinum* L.). *Int J Agric Biol.* 2014;16:956-60.
50. Janghel DK, Kumar K, Sunil R, Chhabra AK. Genetic diversity analysis, characterization and evaluation of elite chickpea (*Cicer arietinum* L.) genotypes. *Int J Curr Microbiol Appl Sci.* 2020;9(1):199-209. doi: [10.20546/ijcmas.2020.901.023](https://doi.org/10.20546/ijcmas.2020.901.023).
51. Katkani D, Babbar A, Upadhyay S, Goyal V; 2022. Computation of genetic variability and divergence analysis in advance breeding lines of chickpea. In: *Biological Forum—An International Journal* (Vol. 14, No. 2, pp. 611-617).
52. Gediya LN, Patel DA, Parmar DJ, Patel R, Rahevar P. Assessment of genetic diversity of chickpea genotypes using D2 statistics. *Int J Chem Stud.* 2018;6(4):3177-81.
53. Reddy MA, Lal GM. Assessment of genetic diversity for different quantitative traits in chickpea (*Cicer arietinum* L.). *J Pharm Innov J.* 2021; 10(9):21391:2135.
54. Parhe SD, Harer PN, Nagawade DN 2014. Investigation of genetic divergence in chickpea (*Cicer arietinum* L.) genotypes. *The Bioscan.* 9(2): 879-82.
55. Kuldeep R, Pandey S, Babbar A, Prakash V. Genetic diversity analysis in Chickpea grown under heat stress conditions of Madhya Pradesh. *Electron J Plant Breed.* 2015;6(4):962-71.
56. Thakur NR, Toprope VN, Phanindra KS. Genetic diversity analysis in chickpea (*Cicer arietinum* L.). *Int J Curr Microbiol Appl Sci.* 2018 [Special Issue];6:904-10.

UNDER PEER REVIEW




Table 1. Clustering pattern of 71 genotypes of chickpea on the basis of genetic divergence

Clusters	No. of genotypes	Genotypes

Comment [B53]: Compulsory for acceptance:
Please add this Table.

UNDER PEER REVIEW

Table 1 | Average intra and inter cluster D² values of 71 genotypes of Chickpea

Comment [B54]: Table 2. Average....

	C1	C2	C3	C4	C5	C6	C7	C8	C9	C10	C11	C12	C13	C14	C15	C16	C17	C18	C19	C20	C21	C22	C23	C24	C25	C26
C1	19.7	33.98	27.79	32.19	34.27	44.10	31.43	29.21	46.93	29.8	49.85	36.77	54.38	29.85	32.51	108.50	27.77	48.85	31.19	26.82	53.34	40.13	38.70	47.66	55.70	97.55
C2		0.00	60.69	6.59	8.71	85.59	29.94	26.55	10.76	21.68	38.37	68.95	87.49	18.02	20.64	107.55	28.13	21.83	51.88	50.23	22.60	60.58	43.87	23.91	36.68	131.52
C3			0.00	68.32	62.24	7.54	42.16	61.51	79.11	43.3	87.63	59.13	40.28	49.40	64.98	120.63	53.84	82.83	57.38	23.00	79.34	59.74	61.72	56.93	52.88	83.58
C4				0.00	6.79	91.29	19.62	12.54	11.67	27.42	22.02	46.74	74.53	23.77	18.99	80.56	18.04	18.43	47.22	57.35	15.32	68.44	46.58	42.22	40.74	106.99
C5					0.00	85.71	24.38	15.77	8.84	18.5	30.98	61.22	75.87	31.99	16.54	85.20	34.05	12.88	62.04	51.46	23.15	72.95	40.88	31.32	25.38	117.28
C6						0.00	54.81	86.08	101.09	61	110.08	78.00	51.56	62.80	92.17	138.61	68.60	105.30	83.44	40.49	93.84	85.99	87.05	86.23	69.63	79.31
C7							0.00	26.63	31.34	47.1	22.58	35.24	45.79	49.68	36.80	60.41	33.56	39.81	52.26	61.84	24.70	83.54	66.62	58.69	35.50	55.85
C8								0.00	34.69	30.47	27.17	22.80	59.57	38.82	21.45	76.95	19.01	20.98	41.21	46.26	37.50	71.29	30.22	59.76	57.61	82.57
C9									0.00	23.84	35.57	88.09	102.54	30.41	17.22	105.36	39.90	27.87	78.94	75.66	18.54	79.26	66.45	41.33	33.93	148.00
C10										0	59.49	74.48	81.83	18.34	16.59	125.33	31.67	38.52	63.46	33.24	48.80	57.65	41.08	35.42	45.38	
C11											22.82	40.88	83.32	62.92	36.67	78.46	42.30	48.16	56.19	92.86	35.18	90.00	74.93	82.27	68.50	
C12												0.00	50.87	71.65	58.43	87.74	35.35	68.08	25.84	59.25	74.78	72.54	52.87	103.04	102.61	53.72
C13													25.02	87.14	98.45	58.88	57.49	97.89	86.84	63.89	74.53	120.64	106.56	112.48	59.30	55.77
C14														0.00	33.63	133.99	15.12	47.53	53.44	41.10	35.84	49.93	51.05	47.87	61.62	138.28
C15															0.00	131.80	34.96	25.85	42.54	45.70	51.72	47.23	31.06	35.52	62.27	147.43
C16																30.50	89.22	119.13	153.40	153.37	60.54	205.81	176.95	167.12	69.49	74.42
C17																	0.00	44.12	42.74	46.01	30.19	67.10	55.07	74.94	67.43	90.11
C18																		0.00	68.14	53.94	47.40	90.50	27.71	42.53	53.06	124.93
C19																			0.00	36.98	92.27	20.83	34.84	60.10	110.01	128.12
C20																				0.00	95.42	36.77	24.93	35.92	67.24	132.75
C21																					0.00	110.12	97.66	77.95	32.31	96.17
C22																						0.00	44.57	49.00	109.57	199.44
C23																							0.00	39.69	94.81	142.49
C24																								0.00	48.38	186.56
C25																									0.00	118.75
C26																										0.00

Comment [B55]: Width?

Table 2 Cluster mean performance for 12 characters of 71 chickpea genotypes

	Days to 50% flowering	Days to maturity	Plant height (cm)	Primary branches per plant	Secondary branches per plant	Pods per plant	Effective pods	Seed per pod	100 seed weight (gm)	Harvest index (%)	Biological yield per plant	Seed yield per plant (gm)
Cluster 1	55.82	115.19	53.48	3.63	8.91	39.49	34.96	1.43	24.16	52.17	35.73	18.56
Cluster 2	53.17	121.33	62.31	4.22	10.65	47.83	42.50	1.40	24.88	59.43	32.38	19.19
Cluster 3	56.50	118.17	53.04	3.68	7.80	40.00	34.83	1.67	18.45	46.69	39.92	18.56
Cluster 4	53.83	116.17	61.70	4.03	9.98	46.83	41.50	1.20	28.24	59.48	34.31	20.40
Cluster 5	57.67	117.50	55.68	3.75	9.47	50.67	45.83	1.40	27.09	60.98	33.64	20.51
Cluster 6	56.83	113.00	56.46	3.42	7.82	39.00	36.17	1.45	17.41	40.59	42.06	17.06
Cluster 7	54.50	114.33	58.35	4.23	9.51	47.50	41.33	1.47	27.48	46.96	41.06	19.22
Cluster 8	56.67	116.50	52.10	3.55	7.95	41.67	36.33	1.32	29.24	63.35	35.36	22.34
Cluster 9	57.50	117.00	66.57	3.85	8.97	52.00	46.50	1.73	27.54	55.63	32.24	17.91
Cluster 10	54.17	115.83	57.84	2.55	6.37	45.83	41.67	1.60	23.28	57.92	31.72	18.34
Cluster 11	55.17	116.78	56.09	3.76	9.95	45.06	40.11	1.44	31.88	49.54	37.73	18.70
Cluster 12	54.50	115.00	45.06	3.62	9.20	32.83	28.33	1.13	29.77	51.82	40.28	20.85
Cluster 13	56.31	114.05	55.03	3.64	8.38	40.98	36.40	1.49	22.64	56.12	45.58	25.59
Cluster 14	54.17	115.83	70.82	3.42	9.31	40.50	35.33	1.25	22.26	57.27	31.21	17.85
Cluster 15	54.17	114.17	53.50	3.42	6.91	45.33	39.33	1.93	28.58	56.25	30.37	17.10
Cluster 16	58.13	117.50	63.42	3.68	8.93	50.17	44.79	1.21	28.74	59.41	48.48	28.71
Cluster 17	52.17	112.17	68.42	3.42	8.59	36.83	32.50	1.40	25.92	60.11	35.60	21.38
Cluster 18	57.83	112.83	52.23	4.22	10.88	48.67	43.17	1.60	27.86	68.57	31.29	21.41
Cluster 19	49.50	115.17	42.99	4.27	10.57	32.17	29.17	1.20	27.24	49.55	34.03	16.82
Cluster 20	53.33	111.83	45.20	3.93	7.83	38.00	35.17	1.42	20.32	60.82	33.75	20.49
Cluster 21	59.17	120.00	75.67	4.25	10.26	48.33	44.00	1.33	26.89	55.17	38.41	21.13
Cluster 22	56.17	117.00	46.11	4.42	9.11	33.50	31.50	1.25	24.28	46.52	29.70	13.82
Cluster 23	57.67	117.17	39.04	4.02	9.24	38.17	33.33	1.40	25.60	63.64	29.14	18.52
Cluster 24	51.83	119.83	48.32	4.87	9.08	52.17	45.17	1.18	21.78	58.07	30.63	17.78
Cluster 25	61.17	118.83	60.90	4.48	9.17	55.50	52.17	1.52	22.27	58.03	39.78	23.08
Cluster 26	57.67	118.83	55.66	3.72	9.60	37.00	33.00	1.30	26.18	45.84	51.07	23.41

Comment [B56]: Table 3.

Comment [B57]: unit?

Table 3 Number percent contribution of characters towards divergence in 71 chickpea genotypes

Comment [B58]: Table 4. Number

S.No.	Characters	Time ranked first	Percent contribution
1	Days to 50 % flowering	19	0.76
2	Days to maturity	14	0.56
3	Plant height (cm)	159	6.4
4	Number of primary branches per plant	44	1.77
5	Number of secondary branches per plant	13	0.52
6	Number of pods per plant	362	14.57
7	Number of effective pods per plant	2	0.08
8	Number of seeds per pod	16	0.64
9	100 seed weight (gm)	696	28.01
10	Harvest Index(gm)	59	2.37
11	Biological yield per plant(gm)	946	38.07
12	Yield per plant(gm)	155	6.25
	Total		100




Fig.1 Dendrogram based on Euclidian distance using Tocher's method in 71 chickpea genotypes

Comment [B59]: Optional correction not compulsory correction for acceptance:

Instead of this Tocher Dendrogram, Please provide 'An unrooted neighbour-joining tree' for better explaining the relationship among the studied genotypes, if possible.

or it is OK.





Fig. 2 Cluster diagram depicting intra and inter-cluster distance based on Mahalanobis Euclidean distance

Comment [B60]: Fig. 2. Cluster ...

Please make it clear if possible!!!
or it is OK.



Comment [B61]: Both 1% is not clear. Make it clear- may be by giving arrow.

Fig. 3 | Percent contribution of characters

Comment [B62]: Fig. 3. Percent ...

UNDER PEE