

## GCMS, FTIR and NMR analysis of chloroform extract of *Azadirachta indica*

### Abstract

*Azadirachta indica*, tree of Meliaceae family contains a flexible source of bioactive compounds for controlling plant diseases. But nowadays, people have become more aware of the dangerous side effects of these chemicals, so using biocontrol agents is becoming more and more important. Researchers have started to focus on using plants and microbes as a biocontrol agent as the introduction of some ecofriendly and secure alternative control tactics for agriculture is currently a key problem in the field of plant pathology. It is well known for its therapeutic and ethnomedicinal relevance since it has a variety of pharmacological effects. The goal of the ongoing study is to use GC-MS, FTIR, and NMR investigation to identify the chemical components of *Azadirachta indica* chloroform extract. The Soxhlet method was used for extraction process. To isolate and identify the antibacterial fraction from *Azadirachta indica* chloroform extract, TLC-bioautography was used. Phenol 3,5-bis (1,1-dimethyl ethyl), Phthalic acid bis (7-methylcyl), Dido decyl phthalates, Oxalic acid, allyl hexadecyl ester, and 2-Piperidinone, N-(4-bromo-n-butyl) were the five primary antibacterial chemicals identified by the GC-MS study. The findings of the FTIR study revealed the presence of functional groups C-H str, C=O str, and C=C str as well as alcohol and the carboxylate ion. While the  $^{13}\text{C}$  NMR data demonstrated the existence of carbonyl, aromatic carbon, quaternary carbon, olefinic carbon, and methyl group, and the  $^1\text{H}$  NMR results revealed the presence of aliphatic OH, methyl, aromatic OH, and olefinic proton. The phytoconstituents found using GC-MS analysis showed a wide range of pharmacological properties, including antioxidant, antibacterial, antifungal, anti-inflammatory, and antimalarial effects. Thus, *Azadirachta indica* plant has a high concentration of medicinal chemicals constituents that can be used to make antimicrobial medications to fight against plant pathogenic bacteria.

**Keywords:** *Azadirachta indica*, TLC-bioautography, GC-MS, FTIR, NMR.

## Introduction

Secondary metabolites are the broad category of organic chemicals that are created by the plants, many of which don't seem to directly contribute to growth and development (Hartmann, 1991). The unique source of supply that these secondary metabolites provide for medications, food additives, flavour, and other industrial materials is crucial (Zhao et al. 2005). These substances are a very diverse set of organic materials produced by a wide range of organisms, including plants, fungi, bacteria, algae, and mammals. The lack of secondary metabolites has no detrimental effects on a plant's ability to survive, but it does have a significant impact on its defence mechanisms (Stamp and Nancy, 2003). It has been estimated that, between 14 and 28 percent of higher plant species are used for medicinal purposes, and from that 74% of plant species are utilised pharmacologically, based on research into the ethnomedical uses of the plants (Ncube et al. 2008). Many therapeutic benefits are demonstrated by the diverse array of phytochemicals that medicinal plants produce, including phenolic acids, flavonoids, tannins, and other substances (El-Najjar et al. 2007). Terpenes, phenolics, and chemicals with nitrogen are the three primary categories of secondary metabolites. In this, phenolics include phenolic acids, coumarins, flavonoids, tannins, and lignin, while nitrogen-containing chemicals include alkaloids and glycosylates. Terpenes include plant volatiles, cardiac glycosides, carotenoids, and sterols.

Many efforts have been made to find novel antimicrobial chemicals from a variety of sources, including microorganisms, animals, and plants. Plants now defend themselves against microbial infection and degeneration because of the discovery to novel, powerful antibacterial chemicals from plants (Cowan, 1999). To prevent plant pathogenic infections, there is increased interest in employing natural antibacterial chemicals, particularly those obtained from plants. As a result of the increased public knowledge of the dangerous side effects of chemicals, the employment of biocontrol agents is becoming increasingly important. Researchers have started to focus on using plants and microbes as a biocontrol agent due to the introduction of some ecofriendly and secure alternative control tactics in agriculture, which is currently a key problem in the field of plant pathology.

The plant *Azadirachta indica*, also known as neem, belongs to the Meliaceae family and has already risen to the top of the list of potential biocontrol agents. Many pharmacological activities, including antibacterial, antifungal, antiulcer, antifeedant, repellent, pesticide, inhibitor, and sterilant, have been linked to the neem leaf, bark, fruit, stem, and flower. The presence of bioactive chemicals makes it commercially viable and

enables it to be employed against a variety of plant diseases historically. Due to the plant's medicinal and bioactive properties, researchers from around the world are interested in studying its bioactive components (Luo et al. 1999). It has been claimed that bioactive substances derived from natural sources can be used to diagnosis a variety of diseases (Hamoburger and Hostettmann, 1991). The general screening of plants to find those with bioactivity against pathogenic organisms serves as the first step in the isolation and characterization of bioactive chemicals (Oyewale et al. 2004).

Alkaloids, tannin, flavonoids, phenolic compounds, dicarboxylic acid, and plasticizer-type chemicals are among the bioactive substances mostly found in *Azadirachta indica* (Hill, 1985). These groups of substances exhibit a variety of pharmacological properties, including antimicrobial and antimalarial activity (Elaiyaraja and Chandramohan, 2016), antioxidant activity (Ajayi et al., 2011), anti-inflammatory activity (Chandrashekar et al., 2015), antifouling activity (Kumari et al., 2012), anti-hypersensitive activity (Mallika Devi et al., 2012), cancer (Lee et al. 2000).

Thus, the aim of this paper is to report on the isolation and partial characterization of a bioactive antimicrobial compound from chloroform extract of *Azadirachta indica* using GCMS, FT-IR and NMR ( $^1\text{H}$  and  $^{13}\text{C}$ ).

## **Materials and methods**

### **Preparation of plant extract**

The fresh leaves of *Azadirachta indica* were collected and thoroughly washed under running tap water to get rid of dirt and other contaminants. The leaves were ground into a powder using a grinder, then stored in an airtight container for later use. The leaves were dried individually under shade with occasional shifting for around 3 to 4 weeks. The Soxhlet's method procedure given by Gupta et al. 2013, was carried out for extraction of leaves. Following extraction, the supernatant was collected in the flask individually by filtering it using Whatman No. 1 filter paper and allowed to evaporate at room temperature. Air dried extracts were weighed separately and kept in little tubes at 5 °C in the refrigerator.

### **Thin Layer Chromatography**

To determine the chemical composition of *Azadirachta indica* leaf chloroform extract, thin layer chromatography was used. The TLC plates were made by combining 25 g of silica gel-G (Hi media) with 50 ml of distilled water, and then using a spreader to evenly spread the resulting slurry across the plates with a thickness of 0.25 mm. The plates were

heated in an oven at 110 ° C for one hour after being allowed to dry at room temperature. With the aid of capillary tubes, a 10- $\mu$ l sample of the *Azadirachta indica* chloroform extract was put on TLC plates at identical distances after being diluted in DMSO. The TLC plate was retained and allowed to run until it reached the 3/4 position in the hexane: ethyl acetate (1:1) solvent system. The produced chromatogram on the TLC plates was examined with visible and ultraviolet light after being allowed to air dry. The distance travelled by the solvent front and the solute front was used to calculate the bands' R<sub>f</sub> values (Relative front).

### **Bioautography**

According to Ahmed and Beg 2001, the bioautography approach was used to isolate the bioactive chemicals. On TLC plates, the zone of inhibition was visible as a transparent spot on a red background. The position of the compound exhibiting antibacterial activity was confirmed using bioautography technique on the TLC plate. Later the antibacterial fraction was scraped out from the silica gel and was thoroughly dissolved in chloroform. It was then centrifuged for 10 minutes at 10,000 rpm. For complete solvent evaporation, the supernatant was evaporated at 60°C for 50 min using a vacuum concentrator.

### **FT-IR, GC-MS, <sup>1</sup>H and <sup>13</sup>C NMR analysis**

Nuclear Magnetic Resonance (NMR), Gas Chromatography-Mass Spectrometry (GC-MS), and Fourier Transform Infrared (FT-IR) spectroscopy (FT-IR) techniques were used to conduct additional study on the purified chemical compound. For this analysis, separate 70 mg active compound was stored in tiny, sterile glass vials. The samples were sent to the Sophisticated Analytical Facility, IIT, Powai, Bombay, for chemical analysis using FT-IR, GC-MS, and <sup>1</sup>H and <sup>13</sup>C NMR. Tables and Figures showed the interpreted data.

## Results

### Extraction yield of *Azadirachta indica*

Extraction yield of *Azadirachta indica* leaves in different solvents are presented in Table 1 and

### Extraction yield of *Azadirachta indica*

Distilled water exhibited (9.08%) maximum extraction from *Azadirachta indica* leaves whereas minimum extraction yield was observed in petroleum ether (2.9%).

### Thin Layer Chromatography (TLC)

Thin layer chromatography was used for separation of different chemical constituents present in chloroform extract of *Azadirachta indica*.

### TLC-Bioautography

To assess the antibacterial activity of isolated compounds against the tested bacterium, bioautography technique was utilized on TLC plates run in hexane: ethyl acetate (1:1). The TLC plate displayed a transparent zone of inhibition against a red background around the band that contained the active ingredient responsible for the antibacterial activity after being sprayed with 2, 3,5-tri phenyl tetrazolium chloride. One compound from *Azadirachta indica* extract demonstrated well-resolved suppression of *Xanthomonas axonopodis* pv. *citri* at Rf- 0.74 while emitting a pink color when illuminated by UV light.

### GCMS analysis of *Azadirachta indica*

Chloroform leaf extract from *Azadirachta indica* revealed the existence of five major components together with their retention times, peak areas, and molecular weights. Phenol 3, 5-bis (1,1-dimethyl ethyl) at retention time 9.88, mol. wt. 206, and peak area 659085.64 was the first compound. Phthalic acid bis (7-methyl octyl) ester was second compound at retention time 15.58, mol. wt. 418, and peak area 1188911.27. The third compound was Dodecyl phthalates with retention time 20.17, mol. wt. 502 and peak area 3732124.99. The fourth and fifth compound are Oxalic acid, allyl hexadecyl ester, and 2-Piperidinone, N-(4-bromo-n-butyl) at retention times of 27.24 and 31.34, mol. wt. 354 and 233, their peak areas were 1528807.42 and 2565012.55, respectively, (Table 2 and Figure 1,2,3,4 and 5).

### **FTIR analysis of *Azadirachta indica***

The following peaks and functional groups were identified using FT-IR analysis. Peak 3427 indicates the existence of a hydroxyl methyl group, while peak 2952 and peak 1734 indicate C-H stretching and carboxylic acid-like C=O stretching, respectively. C-H bending was found at 1089.46 and 801.61 peak, and C=C stretching was seen at 1465.19. (Table 3 and Figure 6).

### **<sup>1</sup>H NMR analysis of *Azadirachta indica***

Different signals for various proton types were discovered by <sup>1</sup>H NMR analysis. The initial signal indicated the existence of an aromatic proton between 6.95 and 7.78. The existence of olefinic proton was indicated by the second signal, 6.67 to 6.81. At 4.51 to 4.72 signal, phenolic hydroxyl group presence was evident. While the 4.21 to 3.31 signal indicated the presence of a hydroxyl group, or a methylene group connected to an electronegative atom. At 3.62 to 3.70 signal, the ester group was present, and at 2.62 signal, the ketone group. And the final signal (0.9-1.48) indicated the existence of a methyl group (Table 4 and Figure 7).

### **<sup>13</sup>C NMR analysis of *Azadirachta indica***

Carbonyl group was present at signals 169.50, 67.1, and 67.3 according to <sup>13</sup>C NMR study. Quaternary carbons were observed at signals 129.7 to 132.4 while aromatic carbons were present at signals 116.7 to 127.57. Likely at 27.85 to 23.79 and 14.52 signals, methyl groups were present respectively, (Table 4a and Figure 8).

## Discussion

### Extraction

Table 1 provides the extraction yields of the various solvents employed in this study. The polarity and capacity of a solvent to extract additional chemical compounds from the *Azadirachta indica* plant determines how extractable it is. *Azadirachta indica* was recognised to yield more bioactive compounds when extracted with distilled water compared with other solvents. The findings were like those of Raja Pandiyan et al. (2011), who found that water extract had the maximum extraction yield (4.5g) due to the presence of highly polar alkaloids, flavones, and sugars. Babu et al. (2016) also previously recorded the highest extraction yield (0.6882g) from *Azadirachta indica* leaves in water extract.

### Thin Layer Chromatography

Several antibacterial fractions or secondary metabolites responsible for antibacterial activity were separated using thin layer chromatography. The findings of the above investigation, revealed the retention factors (R<sub>f</sub>) of ethanol extracts of *Azadirachta indica* in various solvent systems, are consistent with those of Mondali et al. (2014). In hexane: ethyl acetate (1:1) solvent system, the ethanol extracts generated nine fractions with R<sub>f</sub> 0.09, 0.10, 0.19, 0.22, 0.38, 0.48, 0.58, 0.66, and 0.91. In the current study, the same solvent system hexane: ethyl acetate (1:1) produced the maximum band separation using chloroform extract. As a result, the TLC results show that chloroform extracts contain a variety of chemical components.

### Bioautography

The distinct antibacterial fraction of *Azadirachta indicia* chloroform extract eluted on TLC plates in a hexane: ethyl acetate (1:1) solvent system was identified using the bioautography method. The 2,3,5-triphenyl tetrazolium chloride was sprayed onto TLC plates, which displayed a whitish or translucent zone of inhibition against a pink or red background at R<sub>f</sub>-0.74. Kruzseliyi et al. (2016) used High Performance Liquid Chromatography (HPLC) and electrospray ionization mass spectrometry to identify the active chemicals present in *Azadirachta indica* oil and designated them as linoleic and oleic acid. Using TLC-bioautography and spectroscopic analysis, Shubham et al. (2016) also reported the existence of an active chemical tetra nor-triterpenoid limonoid with an R<sub>f</sub>- 0.56 and a retention period of 3.8 minutes in *Azadirachta indica* leaves.

## GCMS analysis

The several bioactive components found in *Azadirachta indica* chloroform extract are identified using Gas Chromatography and Mass Spectrometry (GC-MS) technique. Five main chemicals that are responsible for *Azadirachta indica* therapeutic potential were discovered using GC-MS research. Table 4 lists the recognised compounds along with their retention time, molecular weight, molecular formula, and peak area. These components fall under the categories of alkaloid, phenolic, dicarboxylic, and plasticizer chemicals.

Phenol, 3,5, bis-(1,1-dimethyl ethyl), Dodecyl phthalate, 2-piperidinone, N-(4-bromo-n-butyl), Oxalic acid, allyl hexadecyl ester, and Phthalic acid, bis (7-methyloctyl) ester were the most prevalent of the above-mentioned described compounds. The first compound Phenol, 3,5, bis-(1,1-dimethyl ethyl), was discovered in the leaves of *Indoneesiella echiodes* (Elaiyaraja and Chandramohan, 2016), *Azadirachta indica* (Sandanasamy et al. 2014), *Hibiscus micranthus* (Kumar et al. 2011), *Nerium oleander* (Dey and Chaudhary, 2016), and *Ninbapatradi chooram* (Chandrasekhar et al. 2015). This phytochemical demonstrated a variety of activities, including antimicrobial (Elaiyaraja and Chandramohan 2016, Sandanasamy et al. 2014; Lawal et al. 2016; Wagay and Rothe 2016; Arora et al. 2017; Rai et al. 2016 and Chandrasekhar et al. 2015), antioxidant (Ajayi et al. 2011; Victoria and Samrot 2016; Rai et al (Rai et al. 2016). Lawal et al. 2016 and Rukhsana et al. 2015 and Govindappa et al. 2014), antimalarial (Elaiyaraja and Chandramohan 2016), anti-inflammatory (Chandrasekhar et al. 2015), analgesic, anesthetic, antiseptic, antiviral, cancer preventive and fungicidal (Rai et al. 2016).

Dodecyl phthalate, the second compound, was recognised for its plasticizing properties (Priya and Vijayalakshmi, 2011). These phthalates are present in cosmetics, detergents, lubricating oils, alternatives for polychlorinated biphenyls, carriers in pesticide formulations, solvents, and building materials like flooring, sheeting, and films (George and Prest, 2002). This phytocompound was identified in a variety of plant extracts, including *Mukia maderaspatana* (Mallikadevi et al. 2012), *Sarcostemma secamone* (Kumari et al. 2012), *Blighia sapida* (Ojo et al. 2018), *Viola odorata* (Jasim et al. 2018) flower and *Trigonella foenum* (Priya and Vijayalakshmi, 2011). Additionally, a variety of activities, including antimicrobial and antifouling (Kumari et al. 2012; Priya and Vijayalakshmi 2011; Chandel and Kumar 2015 and Kalaiarasan et al. 2012), anti-hypersensitive, vasodialator,

diuretic, and angiotensin ATZ receptor antagonist, were demonstrated by this compound (Mallikadevi et al. 2012).

The third compound, 2-piperidinone, N-(4-bromo-n-butyl), was reported to be found in a variety of plants, including the leaves extract of *Microcosmus exaspeatus* (Meenakshi et al. 2012), *Asparagus racemosus* (Selvam et al. 2014), sesame seed (Olaleye et al. 2018), the leaves, fruit, and latex of *Croton bonplandianum* (Vennila and Udayakumar, 2015), and in *Aspergillus tamaris* and *Penicillium islandicum* (Hady et al. 2016). This compound showed antibacterial, anti-inflammatory (Vennila and Udayakumar, 2015; Meenakshi et al., 2012), and antioxidant properties (Meenakshi et al. 2012).

The presence of fourth antibacterial compound Oxalic acid, allyl hexadecyl ester was found by various researchers in other plant species like in *Laurencia brandenii* (Manilal et al. 2011), *Aloe vera* plants (Arunkumar et al. 2009), Nigerian rice (Adekoyeni et al. 2018) and in *Pongamia pinnata* (Anuradha and Krishnamoorthy 2012). This compound produced a variety of activities such as antimicrobial (Sathya et al. 2012), acaricide, irritant, pesticidal, renotoxic and varroacidial (Zayed and Samling 2016).

The final compound, Phthalic acid, bis (7-methyloctyl) ester was similarly known for its plasticizing properties (Santhi et al. 2016; Ramakrishnan and Venkataraman 2011). Later, different plant taxa including *Tabebuia argentea* (Melappa et al. 2017), *Aporosa lindleyana* (Ramakrishnan and Venkataraman 2011), *Calotropis gigantea* (Singh and Javed 2015), *Purpura persica* (Santhi et al. 2016), and *Centratherum punctatum* (Sivasubramanian and Brindha, 2013). were shown to contain this compound. According to Ramakrishnan and Venkataraman (2011), the antibacterial compound exhibited antibacterial and antifouling properties as well as tumour-fighting properties against mice sarcoma 180 cell lines (Lee et al. 2000).

### **FTIR analysis**

*Azadirachta indica* chloroform extract was further analysed using Fourier Transform Infrared Spectroscopy to detect the various functional groups that were present. Table 5 presents the findings of the discovered functional groups by FTIR analysis together with their peaks and average range. The results of the FTIR analysis discussed above were consistent with those of Shaikh et al. (2017), who found that O-H str was present at peak 3456.55, C=O str was present at peak 1653.05, and alkene and alkyl halide group were present at peak 675.11. Moreover, the existence of C=O str and C=C str was noted at maxima in 1730 and 1452, respectively (Reshmi et al. 2014). At peaks 873.75 and 721.38, 2922.26

and 2852.72, 1463.97 and 1741.72, C-H bend, C-H str, C=C, and C=O were all detected. (Banerjee *et al.* 2016).

### **NMR analysis**

To determine the types of protons and carbon contained in various compounds from the *Azadirachta indica* plant, the Nuclear Magnetic Resonance technique was used. The first type of NMR is  $^1\text{H}$ , which identifies the proton type, and the second type is  $^{13}\text{C}$ , which identifies the carbon type contained in the corresponding plant extracts. Tables 5 and 6 show the chemical shifts, types of protons, and carbon atoms. Similar findings were made in the study of  $^1\text{H}$  NMR analysis by Kumar *et al.* (2012), who indicated the existence of an aromatic proton at peak 7.2–6.8, a  $\text{CH}_2$  group at peak 3.1–3.8, a  $\text{CH}_3$  group at peak 2.2–2.9, and a methyl group at peak 1.28. Furthermore, the methyl groups at 0.77–0.78 and 1.14–1.33 (Patil *et al.* 2017) and aromatic protons at 6.19–7.2, 7.54, 6.40 and 6.88 peaks were given by Sambadam *et al.* 2016.

The above mentioned FTIR and NMR ( $^1\text{H}$  and  $^{13}\text{C}$ ) data revealed various functional groups, proton types, and carbons, which as a result are responsible for producing various pharmacological actions as demonstrated by the bioactive compounds from the *Azadirachta indica* plant that were found. These chemical constituents and the groups they were placed in the study play a significant part in having various activities against plant pathogens, which will be more crucial in the future for controlling the most serious horticultural and agricultural diseases.

## References

- Arunkumar S, Muthuselvam M (2009) Analysis of phytochemical constituents and antimicrobial activities of *Aloe vera* L. against clinical pathogens. *World J Agri Sci* 5:572-576.
- Cowan MM (1999) Plant products as antimicrobial agents. *Clin Microbial Rev* 12:564-582.
- El-najjar N, Saliba N, Talhouk S and Gali-muhtasib H (2007) *Onopordum cynarocephalum* Induces Apoptosis and protects against 1, 2-dimethylhydrazine Induced Colon Cancer. *Oncology Reports* 17: 1517-1523
- Kumar VS, Navaratnam V, Rajasekaran A, Nair N, Soundaraj D, Matharasi P, Narasimhan S and Subramaniam (2012). Isolation and characterization of glucosamine from *Azadirachta indica* leaves: An evaluation of immunostimulant activity in mice. *Asian Pacific J of Trop Biomed.* 1561-1567.
- Patil V, Dhunjibhoy K and Dasgupta D (2017). Novel antioption bacterium activity of embelin and chebulagic acid on screening of Indian medicinal plants. *Int. Res. J. Pharm*, 8 (5):45-52.
- Ncube RNS, Afolayan AJ and Okoh A (2008). Assessment techniques of antimicrobial properties of natural compounds of plant origin: current methods and future trends. *Afri J Biotech* 7(12): 1797-1806
- Sambadam B, Thiagarajan D, Ayyaswamy A and Raman P Extraction and isolation of flavonoid quercetin from the leaves of *Trigonella foenum-graecum* and their antioxidant activity. *Int J Pharm Pharm Sci*, 8(6): 120-124
- Banerjee K, Thiagarajan N and Thiagarajan P *Azadirachta indica* A. Juss Based Emollient Cream for Potential Dermatological Applications. *Ind J Pharm Sci* 2016;78(3):320-325
- Resmi CR., Sreejamol P and Pillai P Green synthesis of silver nanoparticles using *Azadirachta indica* leaves extract and evaluation of antibacterial activities. *Int J. Adv Bio Res*, 4(3) 2014: 300-303.

Shaikh1 TN and Chaudhari S Characterization of Green Synthesized Silver Nanoparticles Using *Azadirachta Indica* (Neem) Leaf Extract. Int J Adv Res in Sci and Eng 6(9):1127-1138

Elaiyaraja A and Chandramohan G (2016) Comparative phytochemical profile of *Indoneesiella echioides* (L) Nees leaves using GC-MS. J of Pharmaco and Phytochem 5:158-171.

Hamburger H and Hostettmann K (1991). The link between phytochemistry and medicine. Phytochemistry 30:3864-3874.

Hill RA (1985) Terpenoids in Thomson RH, (ed). Chemistry of Natural Products, Blackie Academic and Professional. London :106-134

Govindappa M, Hanabusa R, Sadananda TS, Chandrappa CP and Umashankar T (2014) Identification of bioactive metabolites by GC-MS from an endophytic fungus, *Alternaria alternata* from *Tabebuia argentea* and their *in vitro* cytotoxicity activity. Int J Biolog Pharmac Res 5(6): 527-534

Ramakrishnan S and Venkataraman R (2011) Screening of antioxidant activity, total phenolics and gas chromatography-mass spectrophotometer (GC-MS) study of ethanolic extract of *Aporosa lindleyana* Baill. Afri J Biochem Res 5(14), 360-364.

Kumari TK, Muthukumarasamy, and Mohan VR GC-MS analysis of ethanol extract of *Sarcostemma secamone* (l) bennet (Asclepiadaceae) Sci Res Rep 2(3):187-191.

Sathya S, Lakshmi S and Nakkeeran S Combined effect of biopriming and polymer coating on chemical constituents of root exudation in chilli (*Capsicum annuum* L.) cv. K 2 seedlings J of Appl and Nat Sci 8 (4):2141-2154.

Chandrasekar T, Mudiganti RKR, Vijaya KR, Prabhu K, Nandha KS, and Divya D (2015) GC-MS analysis, antimicrobial, antioxidant activity of an Ayurvedic medicine, Ninbapatradi Chooram. J Chem and Pharma Res 7(8):124-136.

Rukhsana K, Varghese V, Akhilesh VP, Jisha KEK, Baskaran KP, Bindu PU and Sebastian CD (2015) GC-MS determination of chemical components in the bioactive secretion of *Anoplodesmus saussurii* (Humbert, 1865). Int J Pharma Sci and Res 6(4):650-653.

Mallikadevi T, Paulsamy S, Jamuna S and Karthika K Analysis for phytochemicals and bioinformatics approach for the evaluation of therapeutic properties of whole plant methanolic extract of *Mukia maderaspatana* (L.)– A traditional medicinal plant in

- western districts of Tamilnadu, INDIA. Asian J Pharm Clin Res 5(4): 2012, 163-168.
- Kumar KA, Shetty SR and Narsu ML GC-MS Analysis of n-Hexane Extracts of Hibiscus micranthus Linn. Asian J of Chem 23(2) (2011) 561-565.
- Lawal RA, Odesanmi OS, Ozaslan MD, Ebuehi OAT, Karagoz ID, Kilic IH, Uyar C and Badmus IA. Gas Chromatography-Mass Spectrometry and Cytotoxicity of *Securidaca long epedunculata* (polygalaceae) Root Bark Extract. Fountain Journal of Natural and Applied Sciences: 2016; 5(1): 19 – 24.
- Rai DK, Sharma V, Pal K and Gupta RK. Comparative phytochemical analysis of *Cuscuta reflexa* Roxb. Parasite grown on north India by GC-MS (2016) Trop Plant Res 3(2): 428–433.
- Melappa G, Shree SCB, Basava C and Prakash B. In vitro antimutagenic, antiproliferative and GC-MS studies on the methanolic extract of endophytic fungi, penicillium species of *Tabebuia argentea* bur & k. Sch Farmacia 2017 65(2) 301-309.
- Ajayi GO, Olagunju JA, Ademuyiwa O and Martins OC (2011) Gas chromatography-mass spectrometry analysis and phytochemical screening of ethanolic root extract of *Plumbago zeylanica*, Linn. J Med Plan Res 5(9):1756-176.
- Arora M, Mahajan AM and Sembi JK (2017) Essential oils analysis of pseudobulbs of *Crepidium acuminatum* (D.DON) SZLACH by GC-MS. Asian Pac J Health Sci 4(3):198-204.
- Victoria DT and Samrot AV (2016) Identification of antioxidant activity of bark of *Aegle Marmelos*. Der Pharma Chemical 8(18):359-363.
- Kumar DR, Sharma V, Pal K and Kumar RG (2012) Comparative phytochemical analysis of *Cuscuta reflexa* Roxb. Parasite grown on north India by GC-MS. Trop Plant Res 3(2): 428–433.
- Dey P and Chaudhuri TK (2016) Comparative phytochemical profiling and effects of *Nerium oleander* extracts on the activities of murine peritoneal macrophages. Arch Biol Sci 68(3):515-531.
- Adekoyeni OO, Adegoke AF and Sogunle KA (2018) Volatile aromatic components of two varieties of parboiled Nigerian rice. Life Journal of Science 20(1).
- Manilal A, Sujith S, Sabarathnam B, George SK, Selvin J, Shakir C, Aaron P and Lipton (2011) Biological activity of the red alga *Laurencia brandenii*. Acta Bot Croat 70(1): 81–90.

- George C and Prest H (2002) Determination of Phthalate Esters by Positive Chemical Ionization MS with Retention-Time Locked GC. NORTH AMERICA, 20(2).
- Kalaiarasan A, Kumar P, Ahmed JS (2012) Biochemical Investigation of *Bulbophyllum kaitense* RECHIB. Root By GC-MS. Eastern ghats of India. Nature and Science 10 (2):29-31.
- Ojo OA, Ajiboye BO, Imiere OD, Adeyonu O, Olayide I and Fadaka A (2018) Antioxidative Properties of *Blighia sapida* K.D. Koenig Stem Bark Extract and Inhibitory Effects on Carbohydrate Hydrolyzing Enzymes Associated with Non-Insulin Dependent Diabetes Mellitus. Pharmacogn J 10(2):376-383.
- Jasim SF, Baqer NN and Alraheem E (2018) Detection of Phytochemical Constituent In Flowers Of *Viola Odorata* By Gas Chromatography-Mass Spectrometry. Asian J Pharm Clin Res 11(5): 262-269.
- Chandel E and Kumar B (2015) Antimicrobial Activity and Phytochemical Analysis Of *Cynodon dactylon*: A Review. World J Pharm and Pharmaceu Sci 4(11):515- 530
- Priya V, Jananie RK and Vijayalakshmi K (2011) GC/MS determination of bioactive components of *Trigonella foenum grecum*. J Chem Pharm Res 3(5):35-40.
- Santhi V, Sivakumar V, Jayalakshmi S, Thilaga RD and Mukilarasi Isolating M (2016) Bioactive Compound from Marine Prosobranch *Purpura persica* from Tuticorin Coast. Int J Env Protection and Policy 4(3): 64-76.
- Hady HA, Abdel-Wareth MTA, El-Wakil EA and Helmy EA (2016) Identification and evaluation of antimicrobial and cytotoxic activities of *Penicillium islandicum* and *Aspergillus tamarii* ethyl acetate extracts. World J of Pharm and Pharmaceu Sci 5(9):2021-2039.
- Meenakshi VK, Gomathy S, Senthamarai S, Paripooranaselvi M and Chamundeswari KP (2012) GC-MS determination of the bioactive components of *Microcosmus exasperates*. J Curr Chem Pharm Sci 2(4):271-276.
- Selvam PR, Srinivasam V, Gunasekaran S and Palani S (2014) Phytochemical and GCMS analysis of ethanolic extracts of *Asparagus racemosus*. J Curr Chem Pharm Sci 2(4):271-276.
- Sandanasamy JN, Hamid A, Tajuddin, Nizam S and Hamid NA (2014) Chemical Characterization and Biological Study of *Azadirachta indica* Extracts. Euro J of Acad Essa 1:9-16.

- Olaleye OO, Kukwa RE, Eke MO and Aondo TO (2018) Extraction, Physicochemical and Phytochemical Characterization of Oil from Sesame Seed. *Asian Food Sci Jou* 1(4): 1-12.
- Vennila V and Udayakumar R (2015) GC-MS Analysis of Leaf, Fruits, and Latex of *Croton bonplandianum* Baill. *Int J of Biochem Res & Rev* 5(3): 187-197.
- Govindappa CM, Chandrappa CP and Sadananda TS (2014) *In Vitro* Antidiabetic Activity of Three Fractions of Methanol Extracts of *Loranthus Micranthus*, Identification of Phytoconstituents by GC-MS and Possible Mechanism Identified by GEMDOCK Method. *Asian J of Biomed and Pharmaceu Sci* 4(34):34-41.
- Zhaoa JT, Davisb LC and Verpoortec R (2005) Elicitor signal transduction leading to production of plant secondary metabolites. *Biotech Adv* 23:283–333.
- Stamp and Nancy (2003) Out of the quagmire of plant defense hypotheses. *The Quar Rev of Bio* 78(1): 23 55.
- Hartmann T (1991) Alkaloids. In herbivores; their interaction with secondary plant metabolites, The chemical participants, 2nd ed, Rosenthal GA and Berenbaum MR eds Academic press, San Diego. 1: 33-85.
- Oyewale AO, Audu PA and Amupitan JO (2004) A Survey of the Chemical Constituents and Biological Activities of Some Medicinal Plants. *Chem Class J* 162-165.
- Lee SM, Ha CS, Cho WJ (2000) Antitumor and Antiangiogenic Activities of Phthalic Acid Derivative Polymers with Medium Molecular-Weight. *Mol Crys and Liquid Crys Sci and Tech* 354:287-301.
- Luo XD, Ma YB, Wu SH, Wu DG (1999) Two novels azadirachtin derivatives from *Azadirachta indica*. *J Nat Prod* 62(7):1022–1024.
- Shubham, Bhardwaj U, Sharma N and Mathur A (2016) Evaluation of potent hydro-alcoholic extract of leaves of *Azadirachta Indica* for isolation and identification of anti-helminthic compound. *Int J Med Res Health Sci* 5(5): 88-95.
- Singh M and Javed K (2015) Comparative study of chemical composition of *Calotropis gigantea* flower, leaf, and fruit essential oil. *Euro Chem Bull* 4:477-480.
- Sivasubramanian R and Brindha P (2013) In-vitro cytotoxic, antioxidant and GC-MS studies on *Centratherum punctatum*. *Int J Pharm Pharm Sci* 5:364-367.

- Rajender B, Saikumar A and Venkatesham A (2015) Comparative antibacterial activities of combined crude leaf extracts of *Eucalyptus globules*, *Azadirachta indica* and *Ocimum scantum*. J Pharm Res 4(4):164-166.
- Rajapandiyan K, Shanthi S, Murugan AM, Muthu GA, and Singh AJR (2011) *Azadirachta indica* - cow urine extract, a novel controlling agent towards Clinically significant Multi Drug Resistant Pathogens. J App Pharm Sci 1 (10):107-113.
- Mondall NK, Mojumdar A, Chatterje SK, Banarjee JK and Gupt AS (2009) Antifungal activities and chemical characterization of Neem leaf extracts on the growth of some selected fungal species in vitro culture medium. J Appl Sci Environ Manage 13(1):49–53.
- Kruzselyi D, Nagy R, Ott PG and Móricz AM (2016) Rapid, Bioassay-Guided Process for the Detection and Identification of Antibacterial Neem Oil Compounds. J Chrom Sci 54(7):1084–1089.
- Babu SK, Naik VKM, Latha J and Ramanjaneyulu K (2016) Extraction, Isolation and Phytochemical Investigation of Natural Products by Using Chromatographic (TLC) Method. Int J Pharm Pharma Res 7(10):380-393.
- Gupta AK, Ahirwar NK, Shinde N, Choudhary M, Rajput YS and Singh A (2013) Phytochemical Screening and Antimicrobial Assessment of Leaves of *Adhatoda vasica*, *Azadirachta indica* and *Datura stramonium*. U K J Pharma Biosci 1(1): 42-47.
- Zayed MZ and Samling B (2016) Phytochemical constituents of the leaves of *Leucaena eucocephala* from Malaysia. Int J Pharm Pharma Sci 8:174-179.

## Tables

**Table 1. Effect of different solvents on per cent extraction yield from dry weight of leaves**

Plant	Solvent	Yield in %
<i>Azadirachta indica</i>	Petroleum ether	2.9
	Chloroform	6.7
	Dichloromethane	4.64
	Distilled water	9.08

**Table 2. Identification of compounds from chloroform extract of *Azadirachta indica* leaves by GCMS**

Sr. no.	Name of compound	Formula	MW	Retention time	Peak area
1	Phenol 3,5-bis (1,1-dimethyl ethyl)	C <sub>14</sub> H <sub>22</sub> O	206	9.88	659085.64
2	Phthalic acid bis (7-methyloctyl) ester	C <sub>25</sub> H <sub>42</sub> O <sub>4</sub>	418	15.58	1188911.27
3	Didodecyl phthalates	C <sub>32</sub> H <sub>54</sub> O <sub>4</sub>	502	20.17	3732124.99
4	Oxalic acid, allyl hexadecyl ester	C <sub>21</sub> H <sub>38</sub> O <sub>4</sub>	354	27.24	1528807.42
5	2-Piperidinone, N-(4-bromo-n-butyl)	C <sub>9</sub> H <sub>16</sub> BrN <sub>0</sub>	233	31.34	2565012.55

**Table 3. Identification of functional group from chloroform extract *Azadirachta indica* leaves by FTIR analysis**

Sr.no.	Peak	Functional group	Average range
1	3427	R-CH <sub>2</sub> OH, R <sub>2</sub> -CHOH R <sub>3</sub> -C-OH	3400-3600
2	2952	C-H str. Hydrocarbons aliphatic aromatic	2850-3000
3	1734	C=O str. Carbonyl group	1650-1800
4	1465.19	C=C str. Aromatic compounds	1450-1600
5	1089.46 and 801.61	C-H bending hydrocarbons, aliphatic aromatic	650-1000


**Table 4. Identification of types of protons from *Azadirachta indica* leaves by Proton NMR analysis (deuterated methanol)**

Sr. no.	( $\delta$ ) Chemical shift	Type of proton
1	6.95-7.78	Aromatic proton
2	6.67-6.81	Olefinic proton
3	4.51-4.72	Phenolic OH
4	4.21-4.31	CH <sub>2</sub> or CH attached to electronegative atom
5	3.62-3.70	CH <sub>2</sub> or CH or Ar-O=CH <sub>3</sub> or R-O-C-CH <sub>3</sub> (ester)
6	2.62	Ar-CH <sub>3</sub> or O=C-CH <sub>3</sub> (ketone)
7	0.9-1.48	R-CH <sub>3</sub>

**Table 4a. Identification of types of carbon from *Azadirachta indica* leaves by <sup>13</sup>C NMR analysis (deuterated methanol)**

Sr. no,	( $\delta$ ) Chemical shift	Type of carbon
1	169.50	C=O
2	129.7-132.4	Quaternary carbon aromatic
3	116.7-127.57	Aromatic carbon Olefinic carbon
4	31.9-49.67	CH or CH <sub>2</sub> attached to electronegative atom, O-CH <sub>3</sub>
5	27.85 and 23.79	R-CH or R-CH <sub>2</sub>
6	14.52	R-CH <sub>3</sub>

UNDER PEER REVIEW



**Fig. 1** Mass spectra of compound Phenol 3, 5, bis (1,1-dimethyl ethyl) from chloroform extract of *Azadirachta indica* leaves







Fig. 2 Mass spectra of compound Phthalic acid bis (7-methyloctyl) ester from chloroform extract of *Azadirachta indica* leaves




**Fig. 3** Mass spectra of compound Didodecyl phthalate from chloroform extract of *Azadirachta indica* leaves




**Fig. 4** Mass spectra of compound Oxalic acid, allyl hexadecyl ester from chloroform extract of *Azadirachta indica* leaves




**Fig. 5** Mass spectra of compound 2-piperidinone, N- (4-bromo-n-butyl) from chloroform extract of *Azadirachta indica* leaves



**Fig. 6 FTIR spectra of compound isolated from chloroform extract of *Azadirachta indica* leaves**



**Fig. 7**  $^1\text{H}$  NMR spectra of compound isolated from *Azadirachta indica* leaves (deuterated methanol)



**Fig. 8**  $^{13}\text{C}$  NMR spectra of compound isolated from *Azadirachta indica* leaves (deuterated methanol).