

Survey of species cultivated in Brazil and their biological applications: A review

ABSTRACT

Medicinal plants are the basis for the development of numerous drugs, ensuring their effectiveness through pharmaceutical properties, from antimicrobial action to the treatment of infectious diseases. Thus, the present study aimed to carry out a survey of the species cultivated in the municipal garden of Sobral - CE, Brazil, in order to verify in the literature, the medicinal potential and their biological properties. For this, the survey was carried out between January and February 2022, while the bibliographic review followed through the PubMed, SciELO and LILACS platforms. The survey identified 70 species, divided into gardening, fruit, medicinal, native and arboreal groups. Of these, about 90.9% had some therapeutic properties, whether antimicrobial, antiparasitic, antioxidant, anti-inflammatory or antinociceptive. Some of these even exhibited all these properties, as is the case of *Moringa oleifera*, becoming a model for studies. Thus, the raised species have potential for the development of new therapeutic alternatives, whether in the development of a new drug from extracts and essential oils, or even through the enhancement of existing drugs.

Keywords: Antimicrobial action; Antioxidant action; Anti-inflammatory action; Medicinal plants.

1. INTRODUCTION

The biodiversity of plants present in northeastern Brazil represents a large portion nationwide, being used as ornamental, fruitful or even as herbal medicine for the treatment of a multitude of diseases [1]. These species combined with ethnic and cultural diversity resulted in a rich traditional medicine, becoming the basis for the development of knowledge and innovative scientific technologies capable of transforming the pharmaceutical market [2].

In recent years, there has been an increase in the use of medicinal plants in the public health system as a complement to some treatments [3]. According to the World Health Organization (WHO), the use of this Traditional and Complementary Medicine is defined as a set of practices, products and knowledge grouped together and not belonging to conventional medicine [3]. In this line, the WHO is implementing the strategy of traditional medicine from 2014 to 2023, having as guidelines the quality and safety of herbal medicines, methodologies for the clinical study of traditional medicine and definition of terminology and classification of species and possible treatments [4].

Numerous studies point to the action of these plant species, from antimicrobial, antiviral, antiparasitic potential, to analgesic, anti-inflammatory action and in the treatment of minor diseases [5]. This activity is related to the active principles that are responsible for provoking responses in the body that consumes it [5]. Such principles are substances synthesized from light energy and nutrients that the plant can extract from the soil [6]. According to the chemical structure, these substances can be organic acids, alkaloids, anthraquinones, phenolic compounds, coumarins, steroids, flavonoids, cardioactive or cardiotoxic glycosides, heterosides, mucilages, essential oils, among others [6].

Within this perspective, Gardens of Medicinal Plants represent great importance, contributing significantly to the preservation of species, and consequently, knowledge and tradition in the use of these plants [7]. Located 242 kilometers from the capital Fortaleza - CE, Brazil, the garden in the Municipality of Sobral - CE, Brazil, produces about 100,000 species per year, meeting the demands for planting in public areas, as well as making the population available for their own planting.

Thus, the study aimed to carry out a survey of the species cultivated in the municipal garden of Sobral - CE, Brazil, in order to verify in the literature their medicinal potential and their antimicrobial properties. Therefore, the study seeks to characterize these plants in order to scientifically base their use for therapeutic purposes, as well as being the basis for applied studies.

2. MATERIALS AND METHODS

The survey of species was carried out between the months of January and February 2022 in the Horto of the Municipality of Sobral - CE, Brazil, in which it was verified which species were most requested, both for afforestation in the city and for local residents. With the objective of describing the possible potential of these plants, a study of the current literature was carried out, pointing out antimicrobial, antiparasitic, antioxidant, anti-inflammatory and antinociceptive action. The literature review was performed using PubMed, SciELO and LILACS platforms.

3. DEVELOPMENT

The survey resulted in 70 species organized into groups, such as those for gardening, fruit, medicinal, native and afforestation plants (Table 1).

Table 1 – Species cultivated in the municipal garden of Sobral – CE, Brazil.

GARDENING			
POPULAR NAME	SCIENTIFIC NAME		
money	<i>Callisia repens</i>	bilberry	<i>Plectranthus ornatos</i>
Me and you	<i>Euphorbia milii</i>	lemon balm	<i>Melissa officinalis</i>
Horse face	<i>Philodendron</i>	Holy grass	<i>Cymbopogon citratus</i>
	<i>Panduriforme</i>	NATIVE	
garden flamboyant	<i>Delonix regia</i>	inga	<i>Inga edulis</i>
palm tree hawaii	<i>Veitchia merrillii</i>	white stick	<i>Picconia azorica</i>
gold drop	<i>Duranta erecta</i>	white ipe	<i>Tabebuia roseo-alba</i>
white grass	<i>Chloris polydactyla</i>	yellow ipe	<i>Handroanthus albus</i>
Zabumba	<i>Brugmansia suaveolens</i>	pink ipe	<i>Handroanthus</i>
purple vine	<i>Allamanda blanchetti</i>		<i>heptaphyllus</i>
Spear of Saint George	<i>Brugmansia suaveolens</i>	fake white dick	<i>Auxemma oncocalyx</i>
Croton (painted)	*	Mungulu	<i>Erythrina verna</i>
croton (petra)	<i>Codiaeum variegatum</i>	imburana	<i>Commiphora leptophloeos</i>
pitaya	*	Pereiro	<i>Aspidosperma pyrifolium</i>
Roxinha	*	cow foot	<i>Aspidosperma pyrifolium</i>
Cambric	*	Ipe perobinha	*
Juicy	*	Purple IPE	<i>Handroanthus</i>
mini seal	*		<i>impetiginosus</i>
Eleven o'clock	<i>Portulaca grandiflora</i>	Timbaúba	<i>Enterolobium</i>
crawl	*		<i>contortisiliquum</i>
white jasmine	*	eighth	<i>Licania tomentosa</i>
Mango	<i>Mangifera indica</i>	Pajeú	<i>Triplaris gardneriana</i>
cashew	<i>Spondias mombin</i>	Munguba	<i>Pachira aquatica</i>
Acerola	<i>Malpighia emarginata</i>	Jucá	<i>Libidibia ferrea</i>
Cherry	<i>Prunus subg. Cerasus</i>	Marizeira	<i>Calliandra spinosa</i>
cashew tree	<i>Anacardium occidentale</i>	big belly	<i>Ceiba glaziovii</i>
earl	*	trapia	<i>Crataeva tapia L.</i>
Guava	<i>Psidium guajava</i>	ipê do cerrado	<i>Tabebuia ochracea</i>
		Craibeira	<i>Tabebuia aurea</i>
		You knew	<i>Mimosa caesalpinhiifolia</i>
		Canafistula	<i>Peltophorum dubium</i>
		Coité	<i>Crescentia cujete</i>
MEDICINAL		AFFORESTATION	
anador	<i>Justicia pectoralis</i>	Acacia	<i>Acacia spp.</i>
Artemisia	<i>Artemisia vulgaris</i>	moringa	<i>Moringa oleifera</i>
mauve	<i>Plectranthus amboinicus</i>	Jatoba	<i>Hymenaea courbaril</i>
Mint	<i>Mentha piperita</i>	mango jasmine	<i>Plumeria rubra</i>
Saffron	<i>Curcuma longa</i>	Cola	*
Meracillin	<i>Alternanthera brasiliiana</i>	gliricidia	<i>Gliricidia sepium</i>
Courama (Malva)	<i>Bryophyllum pinnatum</i>	flamboyant	<i>Delonix regia</i>
coruda)			

*No scientific identification.

It is possible to notice that 38.57% of the plants present in the municipal garden of Sobral-CE, Brazil, are part of the plants used for gardening, 14.28% are medicinal plants, 35.71% are native plants of Brazil, and that 11.43% are used for afforestation.

3.1 Antimicrobial activity

It is known that some plants, mainly extracts and essential oils, exhibit antimicrobial action against a variety of pathogens, effectively contributing to the development of new drugs [8]. Among the plants raised in the study, *Euphorbia milii*, *Allamanda blanchetti* and *Codiaeum variegatum* from the gardening group, have antimicrobial activity [8]. The first, *E. milii*, also popularly known as the “eu e tu” plant, has antimicrobial potential against *Klebsiella pneumonia* and *Staphylococcus epidermidis*, mainly related to the presence of cardiac glycosides, steroids/phytosterols, anthocyanins, proteins, terpenoids, flavonoids and tannins in its extract [8]. The extract of the species *A. blanchetti* presents action against the bacteria *Bacillus subtilis*, *Staphylococcus aureus*, *Sarcina lutea*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Salmonella paratyphi*, *Shigella dysenteriae*, *Vibrio parahaemolyticus* and the fungi *Saccharomyces cerevisiae* and *Aspergillus niger* [9]. Finally, the species *C. variegatum* demonstrates to have antibacterial action against bacteria *Escherichia coli*, *P. aeruginosa*, *Bacillus subtilis* and *S. aureus*, antifungal action against *Alternaria alternata* and *Fusarium oxysporum*, as well as, antiviral action against influenza A viruses [10-11].

In the group of fruit plants, *Mangifera indica* is of great importance, exhibiting antimicrobial action against *E. coli*, *S. aureus*, *Candida albicans* and *Mycobacterium smegmatis*, in addition to action against oral microorganisms [12-13]. In addition to this, the extract of the species *Spondias mombin*, routinely used against gastrointestinal infections, demonstrates action against Gram-negative bacteria, and synergistic effect with some antibiotics (gentamicin and imipenem against *S. aureus* and *E. coli* and norfloxacin against *P. aeruginosa*) [14]. Another study showed the antiviral action of the extract of this species against the Herpes Simplex virus type-1, related to the interaction of the geraniin molecule with the gB viral surface glycoprotein, which is responsible for adsorption [15]. As for the fruit species *Malpighia emarginata*, studies have shown the antifungal action of its saline extract against some species of the genus *Candida*, also demonstrating its ability to interfere with the formation of the biofilm of these fungi [16].

Along the same lines, studies have shown that fractions of the species *Prunus subg. Cerasus* also has antimicrobial action, inhibiting the growth in the planktonic form and biofilm of *Candida albicans*, *Streptococcus mutans* and *Fusobacterium nucleatum*, also functioning as an epithelial barrier preventing infection by these pathogens [17]. With regard to the species *Anacardium occidentale*, studies demonstrate the action of extracts and fractions of its leaves against *Listeria monocytogenes* [18], *S. aureus*, *B. subtilis*, *Salmonella enterica*, *Shigella*, *E. coli* [19], *P. aeruginosa*, *Proteus mirabilis*, *K. pneumoniae*, *C. albicans* [20] and stem bark extract against *Streptococcus mitis*, *S. mutans*, *S. sanguis* and *S. sobrinus* [21], among others. Finally, among the fruit species raised in the current study, the extract of *Psidium guajava* also demonstrates great antimicrobial potential against oral bacteria, mainly related to the presence of the compounds β -selinene, α -humulene and β -caryophyllene [22-23], in addition to this action, the essential oil of this species also demonstrated activity against *Curvularia lunata* and *Fusarium chlamydosporum* [24].

Among the medicinal plants surveyed, the oily fraction of *Artemisia vulgaris* demonstrates great antimicrobial action against the fungus *C. albicans* and *A. niger*, in addition to the bacteria *E. coli*, *Salmonella enteritidis*, *P. aeruginosa*, *K. pneumoniae*, *S. aureus* and *S. mutans*, probably associated with the presence of essential oils, 1,8-cineol, α -thujone and camphene [25-26]. In addition, *Plectranthus amboinicus* exhibits antimicrobial action against planktonic cells and biofilm of *S. aureus* resistant to vancomycin and oxacillin, in addition to *E. coli* and *Salmonella typhimurium*, this activity is related to the compound cracavol [27]. It also exhibits antifungal activity against *A. flavus*, *A. niger*, *A. ochraceus*, *A. oryzae*, *C. versatilis*, *Fusarium* sp., *Penicillium* sp., *S. cerevisiae*, *C. albicans*, *C. tropicalis*, *C. krusei* and *C. stellatoidea* [28]. Another species that has antimicrobial activity is *Mentha piperita*, with essential oil demonstrating action against *Klebsiella* spp., *S. aureus*, *S. epidermidis*, *E. coli*, *S. enteritidis*, *Micrococcus flavus*, among others [29], in addition to the fungal strains *Fusarium moniliforme*, *A. niger*, *A. fumigatus* and *C. albicans* [30].

In addition, the extract of the species *Curcuma longa* has been shown to be effective against bacteria that cause periodontitis, in addition to inhibiting the planktonic form and biofilm of *S. aureus* and *P. aeruginosa* [31-32]. Antifungal action was also evidenced against *Trichophyton mentagrophytes*, *T. rubrum*, *Epidermophyton floccosum* and *Microsporium gypseum* [33]. With regard to the *Alternanthera brasiliana* species, studies point to its effectiveness when used in combination with the antibiotic gentamicin, exhibiting a synergistic effect against *S. aureus*, *E. coli* and *P. aeruginosa*, related to the compound luteolin, the majority in its extract [34]. As for the species *Bryophyllum pinnatum*,

studies indicate that its extract demonstrates effectiveness in inhibiting the bacterium *Helicobacter pylori* in an *in vivo* model, with a significant reduction of this in the gastric mucosa [35]. In this line, the dichloromethane extract and the essential oil of the *Plectranthus ornatos* species also demonstrate antibacterial action against Gram-positive bacteria in their planktonic form, as well as in biofilm [36]. In addition, the *Melissa officinalis* species demonstrates activity against Gram-positive and Gram-negative bacteria, as well as the fungi *C. albicans* and *S. cerevisiae* [37]. Finally, the essential oil of the *Cymbopogon citratus* species demonstrates antibacterial action against Gram-positive and Gram-negative bacteria related to the synergism of its compounds [38].

In the native group, *Inga edulis* stands out for its antifungal action against species of the genus *Candida* [39]. In addition, *Auxemma onocalyx* demonstrates action against *S. aureus*, both in its planktonic form and in biofilm, related to the benzoquinone molecule [40]. Another species with this action is *Commiphora leptophloeos*, exhibiting antimicrobial potential against methicillin-resistant *S. aureus*, *Mycobacterium smegmatis*, *M. tuberculosis*, in addition to antifungal action against *Candida* species in planktonic and biofilm forms [41-42].

In addition, *Handroanthus impetiginosus* also exhibits antibacterial action against *S. aureus*, *Listeria monocytogenes* and *Streptococcus agalactiae*, this action is associated with coumarins, tannins, flavonoids and steroids present in the ethanolic extract of flowers of this species [43]. *Licania tomentosa* also demonstrates antimicrobial action against *S. aureus*, *Bacillus cereus*, *E. coli* and *K. pneumoniae* strains, related to the synergism present in the extract [44]. With regard to the *Triplaris gardneriana* species, studies have shown quercetin as one of the main compounds, with antimicrobial action against Gram-positive and Gram-negative bacteria [45]. Also, *Libidibia ferrea* demonstrates action against the planktonic form and biofilm of *S. mutans*, exhibiting anticariogenic action [46]. *Mimosa caesalpiniiifolia* also has antimicrobial action, mainly against *S. aureus* and *C. albicans* with a mechanism aimed at inhibiting the efflux pump [47]. In addition to these, *Peltophorum dubium* demonstrates antifungal action against *A. flavus*, with a mechanism aimed at the disruption of hyphae [48]. Finally, *Crescentia cujete* has antibacterial action against *S. aureus*, *E. coli* and *Vibrio harveyi* with a mechanism aimed at disturbing the microbial membrane [49-50].

In addition, species destined for afforestation were also surveyed, in this sense the ethanol, chloroform and ethyl acetate *extreacts* of *Moringa oleifera* leaves stand out with action against Gram-positive and Gram-negative bacteria, in addition to antifungal activity against *A. flavus*, *A. niger*, *Trichoderma* sp and *Candida* spp. [51].

2.2 Antiparasitic activity

The antiparasitic activity has also been very evident among botanical species [52]. The current research identified the species *Euphorbia milii*, *Veitchia merrillii* and *Codiaeum variegatum* with this potential within the gardening group [52]. Studies have shown that the hydroalcoholic extract of the species *Euphorbia milii* has ovicidal action against *Ancylostoma* spp., with significantly low toxicity against *Artemia salina* [52]. In this line, the ethanol extract of the species *Veitchia merrillii* is capable of inducing surface changes in adult worms of *Ascaridia galli* [53]. While *in vitro* studies show the antiamoebic action of the species *Codiaeum variegatum* [10].

Among the species of the fruit group, *Mangifera indica* also has antiparasitic action, preventing the development of *Haemonchus contortus* larvae [54]. In addition, the extract of the species *Spondias mombin* has demonstrated antiparasitic action against *Haemonchus placei* related to the compound Phaeophorbide-a, however, it does not present cytotoxicity for cell lines H460, Caco-2, MC3T3-E1 [55]. In this line, some constituents of the *Anacardium occidentale* species have demonstrated effective action against *Trypanosoma cruzi*, mainly cardol, acting as a sirtuin inhibitor, these are involved in the deacetylation of lysine residues of proteins acting as regulators of DNA repair, in the morphology of the parasite and other crucial aspects for its survival [56]. In addition, a study carried out by Alvarenga et al. (2016) demonstrated the action of the compound 2-methylcardol diene from *A. occidentale* against adult worms of *Schistosoma mansoni*, causing damage to the interstitial tissue and mitochondria [57]. Also, among fruit species, *Psidium guajava* demonstrates great antiparasitic action against *Leishmania infantum* related to two main compounds, jacumaric acid and corosolic acid [58]. Furthermore, in a study carried out by Silva et al (2019), the antiparasitic action of this species was demonstrated from the pro-oxidative activity, caused by the increase in total proteins, intracellular H₂O₂ and lipid peroxidation products, in addition to the increase in the level of enzymes glutathione S-transferase and superoxide dismutase [59].

In the group of medicinal plants raised in the current study, *Artemisia vulgaris* shows action against *Plasmodium yoelii* and *P. berghei* and *Trichinella spiralis* [60-61]. The species *Curcuma longa* also demonstrates an antiparasitic effect, with nanoformulations based on curcumin, with effect against *S. mansoni* demonstrating damage at the molecular level [62]. In addition to these, the *Bryophyllum pinnatum* extract demonstrates action against *Plasmodium falciparum*, one of the species that causes malaria in humans [63]. For the *Melissa officinalis* species, studies point to the action of the essential

oil against *Leishmania amazonenses*, mainly associated with its neral and citral compounds [64]. In this line, the species *Cymbopogon citratus*, studies point to its effectiveness against *P. chabaudi* and *P. berghei*, as well as, anthelmintic action against *Haemonchus contortus* [65-66].

With regard to the group of native plants, *Aspidosperma pyrifolium* stands out, demonstrating great *in vitro* potential of its extract and fractions against *P. berghei*, without showing cytotoxicity [67]. Furthermore, fractions of the leaves of *Tabebuia aurea* showed antileishmania action related to the compounds oleanolic acid and ursolic acid [68].

Among the species intended for afforestation, *Moringa oleifera* has a high antiparasitic action, studies indicate that the hydroalcoholic extract of its leaves is capable of reducing the egg count of *Trichuris* ssp. and *Ostertagia* sp., also contributing as a food additive for some animals [69]. In addition, the ethanolic extract of the leaves of *Gliricidia sepium* demonstrates high activity against *H. contortus*, exhibiting approximately 91.2% inhibition [70].

2.3 Antioxidant activity

The antioxidant action of these plants has also been discussed [71]. In the gardening group, the species *Callisia repens*, *Euphorbia milii*, *Delonix regia*, *Duranta erecta* and *Allamanda blanchetti* stand out. In this line, the species *Callisia repens* has as main components total flavonoids, total anthocyanins and sugars, having its high antioxidant action related to the synergism of these molecules, with elimination of DPPH·, and scavenging of free radical ·OH, in addition to a low content of heavy metals [71]. The *Euphorbia mili* species also has an antioxidant action, especially when chloroform fractions are used, demonstrating a great elimination of DPPH radicals [8]. The *Delonix regia* chloroform fractions also has antioxidant action, especially when are used, demonstrating a great elimination of DPPH radicals [8]. The *Delonix regia* species, on the other hand, demonstrates potent antioxidant action related to three flavonoids: rutin, isoquercitrin and myricetin, still presenting low toxicity to healthy cells [72]. Leaf and fruit extracts of *Duranta erecta*, has an antioxidant action mainly associated with several photochemical compounds including flavonoids [73-74-75]. Finally, *Allamanda blanchetti*, demonstrates the high elimination of free radicals of DPPH, with soluble fraction of carbon tetrachloride [9].

In the second group, fruit trees, *Mangifera indica* is highlighted, with studies demonstrating its antioxidant potential [76]. Study carried out by Banerjee et al. (2020). Also, in another study, the antioxidant potential phenolic compounds obtained from the skin of the fruit of this species was demonstrated using the for TPC, DPPH and ORAC methods [77]. The *Spondias mombin* species also demonstrates antioxidant action related to the compounds found in its seeds, namely dodecanoic acid, tetradecanoic acid, n-hexadecanoic acid, capsaicin and dihydrocapsaicin phenolamides [78]. As for the *Malpighia emarginata* species, its antioxidant capacity is supported by the phenolic compounds, ascorbic acid and malic acid [79-80]. In addition, a study carried out by Barros and Melo (2019) verified that the extract of *Malpighia emarginata* is capable of increasing the scavenging of free radicals (38.59%) promoting a greater reduction of ferric ions [81]. With regard to the species *Prunus subg. Cerasus*, Garcia et al. (2015) demonstrated the obtaining of peptides with antioxidant potential with the use of three enzymes (Alcalase, Thermolysin and Flavourzyme) from its seeds, evaluating the ability to scavenge radicals and reduce lipid peroxidation [82]. The results showed that those peptides obtained with Flavourzyme were the ones with the lowest antioxidant action compared to the other two groups [82]. Another plant from the fruit group is *Anacardium occidentale*, whose leaf extract exhibits antioxidant action in the treatment of RAW 264.7 macrophage cells, with a significant reduction in oxidative damage, in addition to reversing the oxidative damage of cells induced by LPS [83]. In addition to these, leaf and fruit extracts of the *Psidium guajava* species show great antioxidant action using the DPPH, CUPRAC and FRAP methods related to the content of phenolic compounds, more precisely: quercetin, quercetin-3-O- α -D-arabinopyranoside, quercetin-3-O- α -D-ribofuranoside, quercetin-3-O- β -D-galactopyranoside, quercetin-3-O- α -D-glucopyranoside, and quercetin-3-O- α -D-xylfuranoside [85-86].

For the group of medicinal plants, *Artemisia vulgaris* has proven antioxidant action by different methods: DPPH, lipid peroxidation, protein glycation, xanthine oxidases, ABTS, hydroxyl superoxide, nitric oxide, ferric reducing power activity and inhibition of lipid peroxidation by assays of reactive species of thiobarbituric acid, with the capacity to increase the level of ascorbic acid and glutathione [87]. This action is related to flavonoid compounds, flavonols, phenolic acids [87]. In this sense, *Plectranthus amboinicus* also demonstrates oxidizing action, significantly inhibiting the formation of free radicals and DPPH hydroxyl radicals, with action related to carvocrol and thymol [88]. Another species with this action is *Mentha piperita*, with essential oil showing DPPH radical scavenging [89]. The *Curcuma longa* species also has antioxidant effects, studies indicate that its extract reduces the levels of malondialdehyde and nitric oxide, increasing the levels of thiol, superoxide dismutase and catalase under conditions of oxidative stress [90]. With regard to the species *Alternanthera brasiliana*, a study carried out by Marchete et al. (2021) demonstrated its antioxidant action *in vitro* and *in vivo* in a murine model, stimulating angiogenesis and tissue healing [91]. Studies indicate that the extract of *Bryophyllum pinnatum* has a great antioxidant effect, mainly associated with the synergism of the compounds luteolin-7-glucoside, carlinoside and quercetin [92]. The *Melissa officinalis* species, on the other hand, presents high antioxidant activity, through its chemical compounds, including high amounts of flavonoids, rosmarinic acid, gallic acid and phenolic content

[93]. Finally, the *Cymbopogon citratus* species exhibits antioxidant action in kidney lineage cells, increasing cell viability and proliferation and decreasing oxidative stress caused by rotenone [94].

Another group raised was that of native plants, in which *Inga edulis* has a high antioxidant action related to flavonoids compounds [95]. In addition, *Commiphora leptophloeos* demonstrates antioxidant action in DPPH and superoxide radical scavenging assays, associated with the presence of polyphenols such as rutin, vitexin, and quercetin diglycosides [96]. With regard to the *Pachira aquatic* species, studies indicate that the phenolic fractions of seed have great antioxidant potential, as measured by oxygen radical absorption capacity (ORAC) and trolox equivalent antioxidant capacity (TEAC) assays [97]. Furthermore, the extract fractions of the fruit of *Libidibia ferrea* have shown antioxidant efficacy, increasing glutathione levels and reducing MDA levels [98].

For the group of plants intended for afforestation, *Moringa oleifera* has a great antioxidant action, studies with some fractions of leaves of this species have shown that this activity is related to the compounds kaempferol 3-O-rutinoside, quercetin 3-O-(6"-malonyl- glycoside), kaempferol 3-O-glycoside and quercetin derivative [99]. The *Hymenaea courbaril* species also exhibits antioxidant action with the hydroethanolic extract of its seeds, with flavones as the major compound and possibly responsible for this action [100]. In addition, the methanolic extract of flowers of the *Plumeria rubra* species has a high antioxidant action, eliminating DPPH, FRAP, metal chelating, hydrogen peroxide, and superoxide nitric oxide radicals [101]. Finally, the fruit extract of the *Adansonia digitata* species due to phenolic compounds, inhibiting α -glucosidase [102].

2.4 anti-inflammatory activity

With regard to the anti-inflammatory action of these plants, great potential is evidenced, highlighting the species *Delonix regia* from the gardening group [103]. This, when significant anti-inflammatory activity of *Delonix regia* was observed using the ethanolic extract of its leaves in an *in vivo* model, with a reduction in the edema developed for experimentation [103].

The *Mangifera indica* species of the fruit group also has a great anti-inflammatory action, studies indicate that this effect has been shown to prevent or mitigate inflammation and other symptoms associated with chronic intestinal diseases, colon cancer, leaky gut and constipation through the suppression of pro cytokines, in addition to improving intestinal health [104]. Furthermore, a study carried out by Oliveira et al. (2017) demonstrates that the essential oil of this species is able to reduce edema in a murine model [105]. The species *Spondias mombin*, on the other hand, has shown great action in combating inflammation in murine models [105]. A study carried out by Abiodun, Nnoruka and Tijani (2020) showed the extract of the seeds of this species as a reducer of inflammation [78]. Still, Gomes et al. (2020) demonstrated the anti-inflammatory potential of the hydroethanolic extract with a significant reduction of SOD, MDA, IL-1 β and TNF- α [106]. Another species of the fruit group with great anti-inflammatory power is *Malpighia emarginata*. Dias et al. (2014) demonstrated in a study with mice that only the ingestion of fruit juice of this species was able to reduce triglyceride levels, increasing the IL-10/TNF- α ratio in adipose tissue, in addition to acting to reduce levels of JNK (c-Jun N-terminal kinase), responsible for regulating the cell self-destruction process [107]. The species *Prunus subg. Cerasus* also exhibits great anti-inflammatory ability due to cyanidin-3-glucoside present in the fruit and linoleic acid in the seeds, these compounds were able to protect the gastric mucosa against lesions induced by HCl/EtOH and reduce pro-inflammatory cytokines TNF-alpha and IL-6 and increase levels of the anti-inflammatory cytokine IL-10 [108]. In this sense, the species *Anacardium occidentale* also exhibits anti-inflammatory properties, in a murine colon inflammation model, the seeds of this species were able to increase the levels of anti-inflammatory cytokines, in addition to activating the nuclear factor (NF) pathway. - κ B, in addition, in an *in vitro* model, the leaf extract of this species was able to inhibit the release of TNF - α and IL-1 β in cells stimulated by LPS [83-84].

With regard to the group of medicinal plants, the anti-inflammatory action of *Artemisia vulgaris* is linked to the normalization of the serum lipid profile, increased activity of paraoxonase-1 and decreased serum levels of malondialdehyde, nitric oxide and tumor necrosis factor- α . proven by lipoxigenase inhibitory activity assay (LOX) and "Cotton Pellets Granuloma Method" [109]. Furthermore, the species *Plectranthus amboinicus* demonstrates inhibition of DNA-binding activities, binding of AP-1 to its consensus DNA sequence, decrease in carrageenan-induced paw edema, and significant increase in IgG, IgM, and lysozyme activity in rats, related to the presence of rosmarinic acid, shimobashiric acid, alvianolic acid L, rutin, thymoquinone, quercetin [110]. Another species that demonstrates anti-inflammatory action is *Mentha piperita*, exhibiting significant results in reducing edema in a murine model, without inducing cytotoxicity [111]. Anti-inflammatory effects were also observed for the extract of the *Curcuma longa* species, demonstrating a reduction in the number of leukocytes, neutrophils and eosinophils, a protective effect related to phospholipase A2 and total protein in several inflammatory disorders, in addition to reducing the rates of IgE, IL- 4, IL-17 and interferon gamma [90]. Still, the *Alternanthera brasiliana* species also demonstrates anti-inflammatory action, reducing nitric oxide and the production of pro-inflammatory cytokines, decreasing levels of inflammatory infiltrate when evaluated in an *in vivo* model [91]. The anti-inflammatory action was also observed for the species *Bryophyllum pinnatum*,

promoting the reduction of the regulation of the Toll-like and kappa receptor, in addition to the expression of the nuclear factor B p65 gene, reducing pro-inflammatory and oxidative mediators, chemokines and molecules of cell adhesion [112].

Some species of the group of native plants presents proven anti-inflammatory action, a polysaccharide in the leaf extract of *Handroanthus albus*, capable of reducing the leukocyte infiltration induced by acetic acid in the peritoneal cavity and showed activity antiedematogenic, decreasing mechanical allodynia and myeloperoxidase activity in the carrageenan-induced paw edema model [79]. Another species with this potential is *Commiphora leptophloeos*, through the extract of its leaves it was evidenced anti-inflammatory action related to the modulation of the cytokine level, decrease of TNF- α , increase of IL-10 *in vivo* and also the inhibition of the production of nitric oxide RAW 264.7 activated by LPS [113]. Furthermore, the species *Aspidosperma pyrifolium* has been highlighted as a possible anti-inflammatory, in a study carried out by Lima et al. (2017) the action of the extract of this species was verified in a murine model of envenoming, the results showed a reduction in cell migration to the peritoneal cavity, and likewise the envenomed animals also showed a reduction in edema, infiltration of inflammatory cells and vasodilation in the lungs [95]. In addition, the *Triplaris gardneriana* seed extract also has anti-inflammatory action acting as a modulator in the degranulation of human neutrophils and in the activity of scavenging free radicals, related to the presence of flavonoids [114]. *Libidibia ferrea* also exhibits anti-inflammatory action through the extract and fractions of its fruit, acting in the reduction of leukocyte migration [98]. Another species with this action is *Mimosa caesalpinifolia*, acting in the modulation of COX-2 and TNF- α expression, attenuating the induced lesions in a murine model [115].

The group of plants for afforestation also has some species with anti-inflammatory characteristics, as is the case of *Moringa oleifera* [116]. Studies indicate that the seeds of this species have a polysaccharide called MRP-1 that exhibits significant effects on inflammatory processes, inhibiting the production of nitric oxide and TNF- α induced by LPS [116]. In addition, the *Gliricidia sepium* species demonstrates anti-inflammatory action, reducing the number of inflammatory cells, and the expression of IL-1 β and IL-6. This activity is related to flavonoids, saponins and tannins present in the leaves [117].

4. CONCLUSION

The present study demonstrates the great medicinal potential of the species present in the Horto Municipal de Sobral - CE, Brazil, showing the antimicrobial, antiparasitic, antioxidant, anti-inflammatory and antinociceptive action. Some of the species like *Moringa oleifera* exhibit all these properties, becoming a model for studies. Thus, the raised species have potential for the development of new therapeutic alternatives, whether in the development of a new drug from extracts and essential oils, or even through the enhancement of existing drugs.

REFERENCES

- [1]. SILVA, G. A., ISHIKAWA, T., & DA SILVA, M. A. (2011). Implementation project for the medicinal plant garden at the Faculty of Pharmaceutical Sciences. Federal University of Alfenas. Faculty of Pharmaceutical Sciences. Department of Food and Medicine. Alfenas/MG.
- [2]. BRAZIL. Ministry of Health. Secretariat for Science, Technology and Strategic Inputs. Department of Pharmaceutical Assistance. National policy on medicinal plants and herbal medicines / Ministry of Health, Secretariat for Science, Technology and Strategic Inputs, Department of Pharmaceutical Assistance. – Brasilia: Ministry of Health, 2006. 60 p. – (Series B. Basic Health Texts).
- [3]. Sousa, I. M. C. D., & Tesser, C. D. (2017). Traditional and Complementary Medicine in Brazil: insertion in the Unified Health System and integration with primary care. *Public Health Books.*, 33, e00150215.
- [4]. World Health Organization. Implementation of the WHO Traditional Medicine Strategy 2014-2023. 2022.
- [5]. HECK, R. M., Ribeiro, M. V., BARBIERI, R. L., RITA MARIA HECK, U. F. P. E. L., & MÁRCIA VAZ RIBEIRO, U. F. P. E. L. (2017). Medicinal plants from the pampa biome in health care.
- [6]. Garlet, T. M. B., Conti, P. N., & Vargas, T. G. C. D. (2022). Native medicinal plants of popular use in Rio Grande do Sul.
- [7]. SILVA, G. A.; ISHIKAWA, T; SILVA, M. A. Implementation project of the medicinal plant garden at the Faculty of Pharmaceutical Sciences. Federal University of Alfenas, Alfenas. 2011.

- [8]. Rauf, A., Khan, A., Uddin, N., Akram, M., Arfan, M., Uddin, G., & Qaisar, M. (2014). Preliminary phytochemical screening, antimicrobial and antioxidant activities of *Euphorbia milli*. *Pakistan journal of pharmaceutical sciences*, 27(4).
- [9]. Sharmin, Tasnuva, et al. "Investigation of biological activities of *Allamanda blanchetii*, the violet Allamanda." *journal of pharmacy research* 6.7 (2013): 761-764.
- [10]. Nojoya, E. M., Fewou, P. M., & Niedermeyer, T. H. (2021). *Codiaeum variegatum* (L.) Rumph. ex A. Juss.(Euphorbiaceae): An overview of its botanical diversity, traditional uses, phytochemistry, pharmacological effects and perspectives towards developing its plant-based products. *Journal of Ethnopharmacology*, 277, 114244.
- [11]. Forero, J. E., Avila, L., Taborda, N., Tabares, P., López, A., Torres, F., ... & Echeverri, F. (2008). *In vitro* anti-influenza screening of several Euphorbiaceae species: Structure of a bioactive Cyanoglucoside from *Codiaeum variegatum*. *Phytochemistry*, 69(16), 2815-2819.
- [12]. Anand, G., Ravinathan, M., Basaviah, R., & Shetty, A. V. (2015). *In vitro* antimicrobial and cytotoxic effects of *Anacardium occidentale* and *Mangifera indica* in oral care. *Journal of pharmacy & bioallied sciences*, 7(1), 69.
- [13]. El-Gied, A. A. A., Abdelkareem, A. M., & Hamedelniei, E. I. (2015). Investigation of cream and ointment on antimicrobial activity of *Mangifera indica* extract. *Journal of advanced pharmaceutical technology & research*, 6(2), 53.
- [14]. Freitas, M. A., da Cruz, R. P., Dos Santos, A. T. L., Almeida-Bezerra, J. W., Machado, A. J. T., Dos Santos, J. F. S., ... & Morais-Braga, M. F. B. (2022). HPLC–DAD analysis and antimicrobial activities of *Spondias mombin* L. (Anacardiaceae). *3 Biotech*, 12(3), 61.
- [15]. Siqueira, E. M. D. S., Lima, T. L., Boff, L., Lima, S. G., Lourenço, E. M., Ferreira, É. G., ... & Zucolotto, S. M. (2020). Antiviral potential of *Spondias mombin* L. leaves extract against herpes simplex virus type-1 replication using *in vitro* and *in silico* approaches. *Planta medica*, 86(07), 505-515.
- [16]. Barros, B. R., Barboza, B. R., Ramos, B. A., MOURA, M. C., Coelho, L. C., Napoleao, T. H., ... & MELO, C. M. D. (2019). Saline extract from *Malpighia emarginata* DC leaves showed higher polyphenol presence, antioxidant and antifungal activity and promoted cell proliferation in mice splenocytes. *Annals of the Brazilian Academy of Sciences*, 91.
- [17]. Lagha, A., LeBel, G., & Grenier, D. (2020). Tart cherry (*Prunus cerasus* L.) fractions inhibit biofilm formation and adherence properties of oral pathogens and enhance oral epithelial barrier function. *Phytotherapy research*, 34(4), 886-895.
- [18]. Sousa, J. M. S., de Abreu, F. A. P., Ruiz, A. L. T. G., da Silva, G. G., Machado, S. L., Garcia, C. P. G., ... & Dionísio, A. P. (2021). Cashew apple (*Anacardium occidentale* L.) extract from a by-product of juice processing: assessment of its toxicity, antiproliferative and antimicrobial activities. *Journal of Food Science and Technology*, 58, 764-776.
- [19]. Aguilar, Y., Soto Rodríguez, F., Almeida Saavedra, M., Hermosilla Espinosa, R., & Martínez Yero, O. (2012). Metabolitos secundarios y actividad antibacteriana *in vitro* de extractos de hojas de *Anacardium occidentale* L.(marañón). *Revista Cubana de Plantas Medicinales*, 17(4), 320-329.
- [20]. Ajileye, O. O., Obuotor, E. M., Akinkunmi, E. O., & Aderogba, M. A. (2015). Isolation and characterization of antioxidant and antimicrobial compounds from *Anacardium occidentale* L. (Anacardiaceae) leaf extract. *Journal of King Saud University-Science*, 27(3), 244-252.
- [21]. Silva, N. L., Bezerra, R. A., Costa, F. N. D., Rocha, M. M. D. N. P., & Pereira, S. L. D. S. (2013). Evaluation of the effect of cashew bark extract on subgingival biofilm microorganisms. *In vitro* experimental study. *Periodontics*, 26-30.
- [22]. Shetty, S., Shetty, R. M., Rahman, B., Vannala, V., Desai, V., & Shetty, S. R. (2020). Efficacy of *Psidium guajava* and *Allium sativum* extracts as antimicrobial agents against periodontal pathogens. *Journal of Pharmacy & Bioallied Sciences*, 12(Suppl 1), S589.
- [23]. Souza, W. F. C., de Lucena, F. A., de Castro, R. J. S., de Oliveira, C. P., Quirino, M. R., & Martins, L. P. (2021). Exploiting the chemical composition of essential oils from *Psidium cattleianum* and *Psidium guajava* and its antimicrobial and antioxidant properties. *Journal of food science*, 86(10), 4637-4649.

- [24]. Chaturvedi, T., Singh, S., Nishad, I., Kumar, A., Tiwari, N., Tandon, S., ... & Verma, R. S. (2021). Chemical composition and antimicrobial activity of the essential oil of senescent leaves of guava (*Psidium guajava* L.). *Natural product research*, 35(8), 1393-1397.
- [25]. Malik, S., de Mesquita, L. S. S., Silva, C. R., de Mesquita, J. W. C., de Sá Rocha, E., Bose, J., ... & Costa-Júnior, L. M. (2019). Chemical profile and biological activities of essential oil from *Artemisia vulgaris* L. cultivated in Brazil. *Pharmaceuticals*, 12(2), 49.
- [26]. Obistoiu, D., Cristina, R. T., Schmerold, I., Chizzola, R., Stolze, K., Nichita, I., & Chiurciu, V. (2014). Chemical characterization by GC-MS and *in vitro* activity against *Candida albicans* of volatile fractions prepared from *Artemisia dracunculus*, *Artemisia abrotanum*, *Artemisia absinthium* and *Artemisia vulgaris*. *Chemistry Central Journal*, 8(1), 1-11.
- [27]. Vasconcelos, S. E. C. B., Melo, H. M., Cavalcante, T. T. A., Júnior, F. E. A. C., de Carvalho, M. G., Menezes, F. G. R., ... & Costa, R. A. (2017). *Plectranthus amboinicus* essential oil and carvacrol bioactive against planktonic and biofilm of oxacillin-and vancomycin-resistant *Staphylococcus aureus*. *BMC complementary and alternative medicine*, 17, 1-9.
- [28]. Murthy, P. S., Ramalakshmi, K., & Srinivas, P. (2009). Fungitoxic activity of Indian borage (*Plectranthus amboinicus*) volatiles. *Food Chemistry*, 114(3), 1014-1018.
- [29]. Zaidi, S., & Dahiya, P. (2015). *In vitro* antimicrobial activity, phytochemical analysis and total phenolic content of essential oil from *Mentha spicata* and *Mentha piperita*. *International Food Research Journal*, 22(6), 2440.
- [30]. Nilo, M. C. S., Riachi, L. G., Simas, D. L. R., Coelho, G. C., da Silva, A. J. R., Costa, D. C. M., ... & De Maria, C. A. B. (2017). Chemical composition and antioxidant and antifungal properties of *Mentha x piperita* L.(peppermint) and *Mentha arvensis* L.(cornmint) samples.
- [31]. Suwal, N., Subba, R. K., Paudyal, P., Khanal, D. P., Panthi, M., Suwal, N., ... & Koirala, N. (2021). Antimicrobial and antibiofilm potential of *Curcuma longa* Linn. Rhizome extract against biofilm producing *Staphylococcus aureus* and *Pseudomonas aeruginosa* isolates. *Cellular and Molecular Biology*, 67(1), 17-23.
- [32]. Jalaluddin, M., Jayanti, I., Gowdar, I. M., Roshan, R., Varkey, R. R., & Thirutheri, A. (2019). Antimicrobial activity of *Curcuma longa* L. extract on periodontal pathogens. *Journal of pharmacy & bioallied sciences*, 11(Suppl 2), S203.
- [33]. Jankasem, M., Wuthi-Udomlert, M., & Gritsanapan, W. (2013). Antidermatophytic properties of ar-turmerone, turmeric oil, and *Curcuma longa* preparations. *International Scholarly Research Notices*, 2013.
- [34]. Coutinho, H. D. M., de Moraes Oliveira-Tintino, C. D., Tintino, S. R., Pereira, R. L. S., de Freitas, T. S., da Silva, M. A. P., ... & Dos Santos, J. F. S. (2018). Toxicity against *Drosophila melanogaster* and antiedematogenic and antimicrobial activities of *Alternanthera brasiliana* (L.) Kuntze (Amaranthaceae). *Environmental Science and Pollution Research*, 25, 10353-10361.
- [35]. Mabeku, L. B., Eyoum Bille, B., Tchouangueu, T. F., Nguépi, E., & Leundji, H. (2017). Treatment of *Helicobacter pylori* infected mice with *Bryophyllum pinnatum*, a medicinal plant with antioxidant and antimicrobial properties, reduces bacterial load. *Pharmaceutical biology*, 55(1), 603-610.
- [36]. Nascimento, F. R., Albuquerque, K. R., Oliveira, M. R., Pizziolo, V. R., Brasileiro, B. G., Diaz, G., & Diaz, M. A. (2017). Antibiotic activity of *Plectranthus ornatus* Codd., a traditional medicinal plant. *Annals of the Brazilian Academy of Sciences*, 89, 2461-2469.
- [37]. Abdellatif, F., Akram, M., Begaa, S., Messaoudi, M., Benarfa, A., Egbuna, C., ... & Simal-Gandara, J. (2021). Minerals, essential oils, and biological properties of *Melissa officinalis* L. *Plants*, 10(6), 1066.
- [38]. Boeira, C. P., Piovesan, N., Flores, D. C. B., Soquetta, M. B., Lucas, B. N., Heck, R. T., ... & Terra, N. N. (2020). Phytochemical characterization and antimicrobial activity of *Cymbopogon citratus* extract for application as natural antioxidant in fresh sausage. *Food chemistry*, 319, 126553.
- [39]. Dib, H. X., de Oliveira, D. G. L., de Oliveira, C. F. R., Taveira, G. B., de Oliveira Mello, E., Verbisk, N. V., ... & Macedo, M. L. R. (2019). Biochemical characterization of a Kunitz inhibitor from *Inga edulis* seeds with antifungal activity against *Candida* spp. *Archives of microbiology*, 201, 223-233.

- [40]. Silva, R. E., Ribeiro, F. D. O. S., de Carvalho, A. M. A., Daboit, T. C., Marinho-Filho, J. D. B., Matos, T. S., ... & dos Santos Soares, M. J. (2020). Antimicrobial and antibiofilm activity of the benzoquinone oncocalyxone A. *Microbial Pathogenesis*, 149, 104513.
- [41]. Dantas-Medeiros, R., Zanatta, A. C., de Souza, L. B. F. C., Fernandes, J. M., Amorim-Carmo, B., Torres-Rêgo, M., ... & Zucolotto, S. M. (2021). Antifungal and Antibiofilm Activities of B-Type Oligomeric Procyanidins From *Commiphora leptophloeos* Used Alone or in Combination With Fluconazole Against *Candida* spp. *Frontiers in microbiology*, 12, 613155.
- [42]. Souza Pereira, J. J., Pereira, A. D. P., Jandú, J. J., Da Paz, J. A., Crovella, S., dos Santos Correia, M. T., & de Azevêdo Silva, J. (2017). *Commiphora leptophloeos* phytochemical and antimicrobial characterization. *Frontiers in microbiology*, 8, 52.
- [43]. Morais, M. G., de Oliveira Junior, A. S., Aguiar, E. L. C. C., Ferreira, M. V. G., Martins, M. F., de Fátima Oliveira, P. P., ... & dos Santos Lima, L. A. R. (2020). PHYTOCHEMICAL SCREENING AND EVALUATION OF THE ANTIBACTERIAL ACTIVITY OF FLOWERS OF *Handroanthus impetiginosus*. *Biodiversidade*, 19(2).
- [44]. Silva, J. B. N. F., Menezes, I. R. A., Coutinho, H. D. M., Rodrigues, F. F. G., Costa, J. G. M., & Felipe, C. F. B. (2012). Antibacterial and antioxidant activities of *Licania tomentosa* (Benth.) Fritsch (Crhysobalanaceae). *Archives of Biological Sciences*, 64(2), 459-464.
- [45]. Macêdo, S. K., Almeida, T. S., Alencar Filho, J. M. T., Lima, K. S. B., Libório, R. C., Costa, M. M., ... & Nunes, X. P. (2021). Phytochemical identification and quantification of quercetin in *Triplaris gardneriana* wedd. leaves by HPLC-DAD with evaluation of antibacterial activity. *Natural product research*, 35(18), 3083-3088.
- [46]. Passos, M. R., Almeida, R. S., Lima, B. O., de Souza Rodrigues, J. Z., de Macêdo Neres, N. S., Pita, L. S., ... & Yatsuda, R. (2021). Anticariogenic activities of *Libidibia ferrea*, gallic acid and ethyl gallate against *Streptococcus mutans* in biofilm model. *Journal of Ethnopharmacology*, 274, 114059.
- [47]. Silva, S. W. C., Monção, N. B. N., Araújo, B. Q., Arcanjo, D. D. R., Ferreira, J. H. L., Lima Neto, J. S., ... & Barreto, H. M. (2019). Antimicrobial activity of *Mimosa caesalpinifolia* Benth and its interaction with antibiotics against *Staphylococcus aureus* strains overexpressing efflux pump genes. *Letters in applied microbiology*, 69(1), 57-63.
- [48]. Di Ciaccio, L. S., Catalano, A. V., López, P. G., Rojas, D., Cristos, D., Fortunato, R. H., & Salvat, A. E. (2020). *In vitro* antifungal activity of *Peltophorum dubium* (Spreng.) taub. extracts against *Aspergillus flavus*. *Plants*, 9(4), 438.
- [49]. Rahmaningsih, S., & Pujiastutik, H. (2019). An *in vitro* and in silico evaluation of the antibacterial activity of the bioactive compounds in Majapahit (*Crescentia cujete* L.) fruit. *Veterinary World*, 12(12), 1959.
- [50]. Parvin, M. S., Das, N., Jahan, N., Akhter, M. A., Nahar, L., & Islam, M. E. (2015). Evaluation of *in vitro* anti-inflammatory and antibacterial potential of *Crescentia cujete* leaves and stem bark. *BMC research notes*, 8, 1-7.
- [51]. Oladeji, O. S., Odelade, K. A., & Oloke, J. K. (2020). Phytochemical screening and antimicrobial investigation of *Moringa oleifera* leaf extracts. *African Journal of Science, Technology, Innovation and Development*, 12(1), 79-84.
- [52]. Coêlho, M. D. G., Maciel, L. T. R., de Fátima Kieko Ozaki, T., Silva, M. E. G., Bozo, L. S. O., Consoli, Y. A., ... & da Silva-Coêlho, F. A. (2021). Ovicidal and toxicological effect of hydroalcoholic extracts of *Euphorbia milli* var *splendens*, *Synadenium carinatum* Boiss and *Tagetes minuta* L. against *Ancylostoma* spp.: *In vitro* study. *Journal of Parasitic Diseases*, 45, 252-257.
- [53]. Balqis, U., Hambal, M., Athaillah, F., & Vanda, H. (2017). Cuticular surface damage of *Ascaridia galli* adult worms treated with *Veitchia merrillii* betel nuts extract *in vitro*. *Veterinary World*, 10(7), 732.
- [54]. Macedo, K. M. (2017). *Mangifera indica* leaves as a nutritional alternative and for the control of *Haemonchus contortus* in sheep.
- [55]. Ogedengbe-Olowofoyeku, A. N., Ademola, I. O., Wright, C. W., Idowu, S. O., & Fatokun, A. A. (2021). Anthelmintic activity and non-cytotoxicity of phaeophorbide-a isolated from the leaf of *Spondias mombin* L. *Journal of Ethnopharmacology*, 280, 114392.

- [56]. Matutino Bastos, T., Mannocho Russo, H., Silvio Moretti, N., Schenkman, S., Marcourt, L., Gupta, M. P., ... & Botelho Pereira Soares, M. (2019). Chemical constituents of *Anacardium occidentale* as inhibitors of *Trypanosoma cruzi* Sirtuins. *Molecules*, 24(7), 1299.
- [57]. Alvarenga, T. A., de Oliveira, P. F., de Souza, J. M., Tavares, D. C., Andrade e Silva, M. L., Cunha, W. R., ... & Pauletti, P. M. (2016). Schistosomicidal activity of alkyl-phenols from the Cashew *Anacardium occidentale* against *Schistosoma mansoni* adult worms. *Journal of agricultural and food chemistry*, 64(46), 8821-8827.
- [58]. Phakeovilay, C., Bourgeade-Delmas, S., Perio, P., Valentin, A., Chassagne, F., Deharo, E., ... & Marti, G. (2019). Antileishmanial compounds isolated from *Psidium guajava* L. using a metabolomic approach. *Molecules*, 24(24), 4536.
- [59]. Silva, L. P., Debiage, R. R., Bronzel-Junior, J. L., Silva, R. M. D., & PEIXOTO, E. C. M. (2020). *In vitro* anthelmintic activity of *Psidium guajava* hydroalcoholic extract against gastro-intestinal sheep nematodes. *Annals of the Brazilian Academy of Sciences*, 92.
- [60]. Bamunuarachchi, G. S., Ratnasooriya, W. D., Premakumara, S., & Udagama, P. V. (2013). Antimalarial properties of *Artemisia vulgaris* L. ethanolic leaf extract in a *Plasmodium berghei* murine malaria model.
- [61]. Caner, A., Döşkaya, M., Değirmenci, A., Can, H., Baykan, Ş., Üner, A., ... & Gürüz, Y. (2008). Comparison of the effects of *Artemisia vulgaris* and *Artemisia absinthium* growing in western Anatolia against trichinellosis (*Trichinella spiralis*) in rats. *Experimental parasitology*, 119(1), 173-179.
- [62]. Allam, G. (2009). Immunomodulatory effects of curcumin treatment on murine schistosomiasis mansoni. *Immunobiology*, 214(8), 712-727.
- [63]. Singh, N., Kaushik, N. K., Mohanakrishnan, D., Tiwari, S. K., & Sahal, D. (2015). Antiplasmodial activity of medicinal plants from Chhotanagpur plateau, Jharkhand, India. *Journal of Ethnopharmacology*, 165, 152-162.
- [64]. Andrade, M. A., Azevedo, C. D. S., Motta, F. N., Santos, M. L. D., Silva, C. L., Santana, J. M. D., & Bastos, I. M. (2016). Essential oils: *in vitro* activity against *Leishmania amazonensis*, cytotoxicity and chemical composition. *BMC Complementary and Alternative Medicine*, 16, 1-8.
- [65]. Macedo, I. T. F., Oliveira, L. M. B. D., Ribeiro, W. L. C., Santos, J. M. L. D., Silva, K. D. C., Araújo Filho, J. V. D., ... & Bevilaqua, C. M. L. (2015). Anthelmintic activity of *Cymbopogon citratus* against *Haemonchus contortus*. *Revista Brasileira de Parasitologia Veterinária*, 24, 268-275.
- [66]. Chukwuocha, U. M., Fernández-Rivera, O., & Legorreta-Herrera, M. (2016). Exploring the antimalarial potential of whole *Cymbopogon citratus* plant therapy. *Journal of ethnopharmacology*, 193, 517-523.
- [67]. Ceravolo, I. P., Zani, C. L., Figueiredo, F. J., Kohlhoff, M., Santana, A. E., & Krettli, A. U. (2018). *Aspidosperma pyriforme*, a medicinal plant from the Brazilian caatinga, displays a high antiplasmodial activity and low cytotoxicity. *Malaria journal*, 17(1), 1-11.
- [68]. Mahmoud, A. H., Mahmoud, B. K., Samy, M. N., Fouad, M. A., Kamel, M. S., & Matsunami, K. (2022). Cytotoxic and antileishmanial triterpenes of *Tabebuia aurea* (Silva Manso) leaves. *Natural Product Research*, 36(23), 6181-6185.
- [69]. Pedraza-Hernández, J., Elghandour, M. M., Khusro, A., Salem, M. Z., Camacho-Diaz, L. M., Barbabosa-Pliego, A., & Salem, A. Z. (2021). Assessment on bioactive role of *Moringa oleifera* leaves as anthelmintic agent and improved growth performance in goats. *Tropical Animal Health and Production*, 53(2), 318.
- [70]. Romero, N., Areche, C., Cubides-Cárdenas, J., Escobar, N., García-Beltrán, O., Simirgiotis, M. J., & Céspedes, Á. (2020). *In vitro* anthelmintic evaluation of *Gliricidia sepium*, *Leucaena leucocephala*, and *Pithecellobium dulce*: Fingerprint analysis of extracts by UHPLC-orbitrap mass spectrometry. *Molecules*, 25(13), 3002.
- [71]. Leng, B., Cai, S., Li, X., Huang, J., Xue, Y., Lin, Z., ... & Li, C. (2019). Main Components and Antioxidant Activity of *Callisia repens* Extracts. *Asian Agricultural Research*, 11(1812-2020-063), 76-79.

- [72]. Khongkaew, P., Wattanaarsakit, P., Papadopoulos, K. I., & Chaemsawang, W. (2021). Antioxidant Effects and *in vitro* Cytotoxicity on Human Cancer Cell Lines of Flavonoid-Rich *Flamboyant* (*Delonix regia* (Bojer) Raf.) Flower Extract. *Current Pharmaceutical Biotechnology*, 22(13), 1821-1831.
- [73]. Donkor, S., Larbie, C., Komlaga, G., & Emikpe, B. O. (2019). Phytochemical, antimicrobial, and antioxidant profiles of *Duranta erecta* L. parts. *Biochemistry research international*, 2019.
- [74]. Donkor, S., Larbie, C., Komlaga, G., & Emikpe, B. O. (2020). Evaluation of the acute hepatoprotective potential of hydroethanolic extract of *Duranta erecta* L. Parts. *Journal of oxicology*, 2020, 1-10.
- [75]. Eke, I. G., & Okpara, G. C. (2021). Anti-hyperglycemic and anti-dyslipidemic activities of methanol ripe fruit extract of *Duranta erecta* L (Verbenaceae) in normoglycemic and hyperglycemic rats. *Journal of Traditional and Complementary Medicine*, 11(3), 209.
- [76]. Banerjee, P., Jana, S., Mukherjee, S., Bera, K., Majee, S. K., Ali, I., ... & Ray, S. (2020). The heteropolysaccharide of *Mangifera indica* fruit: Isolation, chemical profile, complexation with β -lactoglobulin and antioxidant activity. *International Journal of Biological Macromolecules*, 165, 93-99.
- [77]. Navarro, M., Arnaez, E., Moreira, I., Quesada, S., Azofeifa, G., Wilhelm, K., ... & Chen, P. (2019). Polyphenolic characterization, antioxidant, and cytotoxic activities of *Mangifera indica* cultivars from Costa Rica. *Foods*, 8(9), 384.
- [78]. Abiodun, O. O., Nnoruka, M. E., & Tijani, R. O. (2020). Phytochemical constituents, antioxidant activity, and toxicity assessment of the seed of *Spondias mombin* L.(Anacardiaceae). *Turkish Journal of Pharmaceutical Sciences*, 17(3), 343.
- [79]. Carneiro Ferreira, I., Pereira da Silva, V., Vilvert, J. C., de França Souza, F., de Freitas, S. T., & dos Santos Lima, M. (2021). Brazilian varieties of acerola (*Malpighia emarginata* DC.) produced under tropical semi-arid conditions: Bioactive phenolic compounds, sugars, organic acids, and antioxidant capacity. *Journal of Food Biochemistry*, 45(8), e13829.
- [80]. Gualberto, N. C., de Oliveira, C. S., Nogueira, J. P., de Jesus, M. S., Araujo, H. C. S., Rajan, M., ... & Narain, N. (2021). Bioactive compounds and antioxidant activities in the agro-industrial residues of acerola (*Malpighia emarginata* L.), guava (*Psidium guajava* L.), genipap (*Genipa americana* L.) and umbu (*Spondias tuberosa* L.) fruits assisted by ultrasonic or shaker extraction. *Food Research International*, 147, 110538.
- [81]. Barros, B. R., Barboza, B. R., Ramos, B. A., MOURA, M. C., Coelho, L. C., Napoleao, T. H., ... & MELO, C. M. D. (2019). Saline extract from *Malpighia emarginata* DC leaves showed higher polyphenol presence, antioxidant and antifungal activity and promoted cell proliferation in mice splenocytes. *Annals of the Brazilian Academy of Sciences*, 91.
- [82]. García, M. C., Endermann, J., Gonzalez-Garcia, E., & Marina, M. L. (2015). HPLC-Q-TOF-MS identification of antioxidant and antihypertensive peptides recovered from cherry (*Prunus cerasus* L.) subproducts. *Journal of agricultural and food chemistry*, 63(5), 1514-1520.
- [83]. Souza, N. C., de Oliveira, J. M., Morrone, M. D. S., Albanus, R. D. O., Amarante, M. D. S. M., Camillo, C. D. S., ... & Pasquali, M. A. D. B. (2017). Antioxidant and anti-inflammatory properties of *Anacardium occidentale* leaf extract. *Evidence-Based Complementary and Alternative Medicine*, 2017.
- [84]. Siracusa, R., Fusco, R., Peritore, A. F., Cordaro, M., D'Amico, R., Genovese, T., ... & Impellizzeri, D. (2020). The antioxidant and anti-inflammatory properties of *Anacardium occidentale* L. cashew nuts in a mouse model of colitis. *Nutrients*, 12(3), 834.
- [85]. Hartati, R., Nadifan, H. I., & Fidrianny, I. (2020). Crystal guava (*Psidium guajava* L."Crystal"): Evaluation of *in vitro* antioxidant capacities and phytochemical content. *The Scientific World Journal*, 2020.
- [86]. Zhao, Y. J., Li, J. K., Zhang, X., & Gao, J. P. (2018). Study on antioxidant activity of flavonoids from leaves of *Psidium guajava*. *Zhongguo Zhong yao za zhi= Zhongguo Zhongyao Zazhi= China Journal of Chinese Materia Medica*, 43(4), 760-765.
- [87]. Melguizo-Melguizo, D., Diaz-de-Cerio, E., Quirantes-Piné, R., Švarc-Gajić, J., & Segura-Carretero, A. (2014). The potential of *Artemisia vulgaris* leaves as a source of antioxidant phenolic compounds. *Journal of functional foods*, 10, 192-200.

- [88]. Praveena, B., & Pradeep S, N. (2012). Antioxidant and antibacterial activities in the leaf extracts of Indian borage (*Plectranthus amboinicus*). *Food and Nutrition Sciences*, 2012.
- [89]. Mallick, B., Sinha, S., & Roy, D. (2016). Evaluation of antioxidative potential of field grown and tissue culture derived *Mentha piperita* L. plants. *Int. J. Curr. Microbiol. App. Sci*, 5(3), 382-391.
- [90]. Memarzia, A., Khazdair, M. R., Behrouz, S., Gholamnezhad, Z., Jafarnejhad, M., Saadat, S., & Boskabady, M. H. (2021). Experimental and clinical reports on anti-inflammatory, antioxidant, and immunomodulatory effects of *Curcuma longa* and curcumin, an updated and comprehensive review. *BioFactors*, 47(3), 311-350.
- [91]. Marchete, R., Oliveira, S., Bagne, L., Silva, J. I. D. S., Valverde, A. P., Aro, A. A. D., ... & Andrade, T. A. M. (2021). Anti-inflammatory and antioxidant properties of *Alternanthera brasiliana* improve cutaneous wound healing in rats. *Inflammopharmacology*, 29(5), 1443-1458.
- [92]. Ogidigo, J. O., Anosike, C. A., Joshua, P. E., Ibeji, C. U., Ekpo, D. E., Nwanguma, B. C., & Nwodo, O. F. C. (2021). UPLC-PDA-ESI-QTOF-MS/MS fingerprint of purified flavonoid enriched fraction of *Bryophyllum pinnatum*; antioxidant properties, anticholinesterase activity and in silico studies. *Pharmaceutical Biology*, 59(1), 442-454.
- [93]. Miraj, S., Rafieian-Kopaei, & Kiani, S. (2017). *Melissa officinalis* L: A Review study with an antioxidant prospective. *Journal of evidence-based complementary & alternative medicine*, 22(3), 385-394.
- [94]. Pan, D., Machado, L., Bica, C. G., Machado, A. K., Steffani, J. A., & Cadoná, F. C. (2022). *In vitro* evaluation of antioxidant and anticancer activity of lemongrass (*Cymbopogon citratus* (DC) Stapf). *Nutrition and Cancer*, 74(4), 1474-1488.
- [95]. Souza Lima, M. C. J. D., Oliveira Bitencourt, M. A., Furtado, A. A., Torres-Rêgo, M., Siqueira, E. M. D. S., Oliveira, R. M., ... & Fernandes-Pedrosa, M. D. F. (2017). *Aspidosperma pyrifolium* has anti-inflammatory properties: an experimental study in mice with peritonitis induced by *Tityus serrulatus* venom or carrageenan. *International Journal of Molecular Sciences*, 18(11), 2248.
- [96]. Cordeiro, M. L. D. S., Ribeiro, A. R., Melo, L. F. M. D., Silva, L. F. D., Fidelis, G. P., Silva, L. M. P., ... & Scortecchi, K. C. (2021). Antioxidant activities of *Commiphora leptophloeos* (Mart.) JB Gillett (Burseraceae) leaf extracts using *in vitro* and *in vivo* assays. *Oxidative Medicine and Cellular Longevity*, 2021, 1-11.
- [97]. Rodrigues, A. P., Pereira, G. A., Tomé, P. H. F., Arruda, H. S., Eberlin, M. N., & Pastore, G. M. (2019). Chemical composition and antioxidant activity of Monguba (*Pachira aquatica*) seeds. *Food research international*, 121, 880-887.
- [98]. Falcão, T. R., Araújo, A. A. D., Soares, L. A. L., Farias, I. B. D., Silva, W. A. V. D., Ferreira, M. R. A., ... & Guerra, G. C. B. (2019). *Libidibia ferrea* fruit crude extract and fractions show anti-inflammatory, antioxidant, and antinociceptive effect *in vivo* and increase cell viability *in vitro*. *Evidence-Based Complementary and Alternative Medicine*, 2019.
- [99]. Xu, Y., Chen, G., & Guo, M. (2021). Correlations between phytochemical fingerprints of *Moringa oleifera* leaf extracts and their antioxidant activities revealed by chemometric analysis. *Phytochemical analysis*, 32(5), 698-709.
- [100]. Spera, K. D., Figueiredo, P. A., Santos, P. C., Barbosa, F. C., Alves, C. P., Dokkedal, A. L., ... & Silva, R. M. D. (2019). Genotoxicity, anti-melanoma and antioxidant activities of *Hymenaea courbaril* L. seed extract. *Annals of the Brazilian Academy of Sciences*, 91.
- [101]. Mohamed Isa, S. S. P., Ablat, A., & Mohamad, J. (2018). The antioxidant and xanthine oxidase inhibitory activity of *Plumeria rubra* flowers. *Molecules*, 23(2), 400.
- [102]. Braca, A., Sinisgalli, C., De Leo, M., Muscatello, B., Cioni, P. L., Milella, L., ... & Sanogo, R. (2018). Phytochemical profile, antioxidant and antidiabetic activities of *Adansonia digitata* L. (Baobab) from Mali, as a source of health-promoting compounds. *Molecules*, 23(12), 3104.
- [103]. Shewale, V. D., Deshmukh, T. A., Patil, L. S., & Patil, V. R. (2012). Anti-inflammatory activity of *delonix regia* (boj. Ex. Hook). *Advances in Pharmacological and Pharmaceutical Sciences*, 2012.

- [104]. Kim, H., Castellon-Chicas, M. J., Arbizu, S., Talcott, S. T., Drury, N. L., Smith, S., & Mertens-Talcott, S. U. (2021). Mango (*Mangifera indica* L.) polyphenols: Anti-inflammatory intestinal microbial health benefits, and associated mechanisms of actions. *Molecules*, 26(9), 2732.
- [105]. Oliveira, R. M., Dutra, T. S., Simionatto, E., Ré, N., Kassuya, C. A. L., & Cardoso, C. A. L. (2017). Anti-inflammatory effects of essential oils from *Mangifera indica*. *Genet. Mol. Res*, 16(1), 1-9.
- [106]. Gomes, M. S., Lins, R. D. A. U., Langassner, S. M. Z., da Silveira, É. J. D., de Carvalho, T. G., de Sousa Lopes, M. L. D., ... & de Araujo, A. A. (2020). Anti-inflammatory and antioxidant activity of hydroethanolic extract of *Spondias mombin* leaf in an oral mucositis experimental model. *Archives of Oral Biology*, 111, 104664.
- [107]. Dias, F. M., Leffa, D. D., Daumann, F., de Oliveira Marques, S., Luciano, T. F., Possato, J. C., ... & de Lira, F. S. (2014). Acerola (*Malpighia emarginata* DC.) juice intake protects against alterations to proteins involved in inflammatory and lipolysis pathways in the adipose tissue of obese mice fed a cafeteria diet. *Lipids in health and disease*, 13(1), 1-9.
- [108]. Raafat, K., El-Darra, N., & Saleh, F. A. (2020). Gastroprotective and anti-inflammatory effects of *Prunus cerasus* phytochemicals and their possible mechanisms of action. *Journal of Traditional and Complementary Medicine*, 10(4), 345-353.
- [109]. Ben Nasr, S., Aazza, S., Mnif, W., & Miguel, M. (2020). In-vitro antioxidant and anti-inflammatory activities of *Pituranthos chloranthus* and *Artemisia vulgaris* from Tunisia. *International Journal of Applied Pharmaceutical Sciences and Research*, 11(2), 605-614.
- [110]. Silitonga, M., Ilyas, S., Hutahaeen, S., & Sipahutar, H. (2014). Levels of apigenin and immunostimulatory activity of leaf extracts of bangunbangun (*Plectranthus amboinicus* Lour). *International Journal of Biology*, 7(1), 46-53.
- [111]. Mahendran, G., & Rahman, L. U. (2020). Ethnomedicinal, phytochemical and pharmacological updates on Peppermint (*Mentha piperita* L.)—A review. *Phytotherapy Research*, 34(9), 2088-2139.
- [112]. Andrade, A. W. L., Guerra, G. C. B., de Souza Araújo, D. F., de Araújo Júnior, R. F., de Araújo, A. A., de Carvalho, T. G., ... & Zucolotto, S. M. (2020). Anti-inflammatory and chemopreventive effects of *Bryophyllum pinnatum* (Lamarck) leaf extract in experimental colitis models in rodents. *Frontiers in pharmacology*, 11, 998.
- [113]. Dantas-Medeiros, R., Furtado, A. A., Zanatta, A. C., Torres-Rêgo, M., Lourenço, E. M. G., Alves, J. S. F., ... & Zucolotto, S. M. (2021). Mass spectrometry characterization of *Commiphora leptophloeos* leaf extract and preclinical evaluation of toxicity and anti-inflammatory potential effect. *Journal of Ethnopharmacology*, 264, 113229.
- [114]. Neto, J. J. L., de Almeida, T. S., Almeida Filho, L. C. P., Rocha, T. M., Nogara, P. A., Nogara, K. F., ... & Carvalho, A. F. U. (2020). *Triplaris gardneriana* seeds extract exhibits *in vitro* anti-inflammatory properties in human neutrophils after oxidative treatment. *Journal of ethnopharmacology*, 250, 112474.
- [115]. Silva, M. J. D., Vilegas, W., Da Silva, M. A., Paiotti, A. P. R., Pastrelo, M. M., Ruiz, P. L. M., ... & Ribeiro, D. A. (2018). The anti-inflammatory potential of *Mimosa caesalpiniiifolia* following experimental colitis: role of COX-2 and TNF-alpha expression. *Drug Research*, 68(04), 196-204.
- [116]. Cui, C., Chen, S., Wang, X., Yuan, G., Jiang, F., Chen, X., & Wang, L. (2019). Characterization of *Moringa oleifera* roots polysaccharide MRP-1 with anti-inflammatory effect. *International journal of biological macromolecules*, 132, 844-851.
- [117]. Aulanni'am, A. A., Ora, K. M., Ariandini, N. A., Wuragil, D. K., Permata, F. S., Riawan, W., & Beltran, M. A. G. (2021). Wound healing properties of *Gliricidia sepium* leaves from Indonesia and the Philippines in rats (*Rattus norvegicus*). *Veterinary World*, 14(3).