

## **Vitamin, Bioactive compound Contents and Antioxidant Activity of *Ficus capensis* Fruit.**

### **Abstract**

*Ficus capensis* is a tropical, evergreen tree with more than 800 species. Virtually all the parts are locally used for the treatment of various illnesses. This study investigated the Vitamin, Bioactive compound contents and the Antioxidant activity of the fruits of *Ficus capensis*. The analyses were conducted using standard biochemical methods. Various concentrations of 20, 40, 60, 80 and 100 µg/ml of the sample were subjected to 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging, Reducing Power and inhibition of lipid peroxidation in-vitro antioxidant tests and compared with standard antioxidant (Butylated hydroxyl anisole (BHA)). From the results of the vitamin analysis, the fruit contained Vitamins A ( $1.15 \pm 0.18$  mg/g), B<sub>1</sub> ( $0.14 \pm 0.00$  µg/g), B<sub>2</sub> ( $0.12 \pm 0.01$  µg/g), B<sub>3</sub> ( $0.26 \pm 0.02$  µg/g), C ( $8.12 \pm 0.16$  mg/100g), E ( $1.44 \pm 0.45$  mg/g) and Folic acid ( $81.23 \pm 6.29$  µg/g). The bioactive compound evaluated were Total Phenol ( $13.97 \pm 1.40$  mgGAE/100g), Flavonoids ( $11.03 \pm 0.16$  mgCE/100g), β-carotene ( $0.34 \pm 0.001$ mg/100g) and Lycopene ( $0.14 \pm 0.002$ ). The antioxidant activity assays showed that the fruits of *Ficus capensis* exhibited scavenging activity on DPPH with EC<sub>50</sub> of 30 µg/ml while that of the standard was 28µg/ml. It inhibited lipid peroxidation with EC<sub>50</sub> of 80 µg/ml while that of the standard was 62 µg/ml and exhibited reducing power activity with OD<sub>0.5</sub> of 38 µg/ml while that of the standard was 34 µg/ml. Results from this study revealed that fruits of *Ficus capensis* is a potential antioxidant agent due to its free radical scavenging and reducing power activities which were comparable with that of the standard antioxidant. The presence of antioxidant vitamins and bioactive compounds makes it a natural source of antioxidants.

Keywords: *Ficus capensis*, fruits, vitamin, bioactive, antioxidant

### **Introduction**

Greater percentage of the world's population still depend on herbal medicines for their health needs which includes management, cure and treatment of diseases. Herbal medicines usually play roles in first-line and basic health services, both to people living in remote areas where it is the only available health service and to people living in poor areas where it offers the only affordable remedy (Adesina, 2014). Even in areas where modern medicine is available, the interest in herbal medicines and their utilization have been increasing rapidly in recent years. Herbal medicines account for a significant percentage of the pharmaceutical market (Adesina, 2014). Plants which include fruits and vegetables contain many bioactive compounds. They serve as herbal medicines and used in traditional medicine. Bioactive compounds are present in small quantities in foods, mainly in fruits, vegetables and whole grains and provide health benefits beyond the basic nutritional value (Gökmen, 2016). Bioactive compounds are

molecules that can present therapeutic potential with influence on energy intake, while reducing pro-inflammatory state, oxidative stress, and metabolic disorders (Siriwardhana et al., 2013). Studies have shown that high consumption of food rich in bioactive compounds such as flavonoids and carotenoid, has a positive effect on human health and could diminish the risk of numerous diseases such as cancer, heart diseases, stroke, Alzheimer's, diabetes, cataracts and age related functional decadence(Hassimotto et al.,2015).Plants are rich in natural antioxidants of which the best known are carotenoids, tocopherols, vitamin C, flavonoids, and different other phenolic compounds (Iqbal and Bhanger, 2006). In recent times, flavonoids, amongst other natural antioxidants have received increasing attention. In comparison to vitamin C and E, dietary flavonoids are considered to be more powerful antioxidants (Sultana and Anwar, 2008).Flavonoids are known antioxidants which are highly effective in scavenging oxygen radicals. They exhibit anti-cancer, anti-ageing, hypolipidemic, and anti-inflammatory activities (Braca et al., 2002). More so, the protective effects of flavonoids in biological systems are attributed to their capacity to scavenge free radicals, chelate metal catalysts, activate antioxidant enzymes, reduce alpha-tocopherol radicals, and inhibit oxidases (Heim et al., 2002).

Antioxidants provide protection against degenerative diseases including cancer, coronary heart, and Alzheimer's diseases (Iqbal and Bhanger, 2006). Reactive Oxygen Species (ROS), contribute to cellular aging, mutagenesis, carcinogenesis, and coronary heart disease, likely through destabilization of membranes, DNA and protein damage, and oxidation of low-density lipoprotein (LDL) (2). Mechanism of action of antioxidants includes the suppression of ROS formation, the inhibition of enzymes or chelating of elements involved in free-radical production. Furthermore, antioxidants scavenge reactive species, and upregulate antioxidant defences (Montoro et al., 2005).

The plants of *Ficus* species are well known in the field of traditional medicine. *Ficus* species have been found to be rich source of phenolic acid and flavonoids making them capable of protecting against disorders of oxidative stress (Sirisha et al, 2010). *Ficus capensis* commonly known as bush fig is a fast growing deciduous or evergreen tree(Onyeleke et al., 2008). The fruits are often hairy, soft and edible but watery and tasteless, having many seeds. Traditionally, *Ficus capensis* has been used in gastrointestinal tract relaxation (Ayinde and Owolabi, 2009; Ayinde et al., 2013) as antioxidant (Sirisha et al., 2010), antimicrobial (Adebayo-Tayo and Odeniyi, 2012) immune booster(Daikwo et al., 2012).

## **Materials and methods**

### **Collection and identification of plants material**

Fresh fruits of *Ficus capensis* were collected from department of Applied Biology and Biotechnology garden of Enugu State University of Science and Technology and was authenticated in the Department of Applied Biology and Biotechnology, Enugu State University of Science and Technology, Enugu, Nigeria.

### **Sample Preparation**

The fresh fruits of *Ficus capensis* were rinsed thoroughly with distilled water and were cut into halves. The fruits were ground with mortar and pestle.

### **Vitamin Analysis**

Vitamins A and E were determined according to the method of Rutkoski *et al.* (2006), Vitamin C was determined according to the method of Klein and Perry (1982), Vitamins B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub> were determined using the method of Kirk and Sawyer (1991) while Vitamin B<sub>9</sub> was determined according to the method of Padmarajaiah *et al.* (2002).

### Bioactive Compound Analysis.

Total phenol, flavonoids,  $\beta$ -carotene and Lycopene contents were determined by using the slightly modified colorimetry method described by Barros *et al.* (2007).

### Antioxidant Assay

The stable 2,2-diphenyl-1-picryl hydrazyl radical (DPPH) was used for the determination of free radical scavenging activity of the sample. This was assayed using the method of Ebrahimzadem, *et al.*, (2009). The reducing power and inhibition of lipid peroxidation were determined according to the method of Barros *et al.*, (2007).

### Statistical Analysis

Data was analyzed using Excel Statistical Package and presented as mean  $\pm$  standard deviation of duplicate analysis

### Results

#### Vitamin Composition

The result of the vitamin analysis is presented in Table 1. From the result, vitamin B<sub>9</sub> (81.23  $\pm$  6.29  $\mu$ g/g) was the highest in composition while vitamin B<sub>2</sub> (0.12  $\pm$  0.01  $\mu$ g/g) was the least.

Table 1: Vitamin Composition of fruit of *Ficus capensis*

Vitamins	Composition
A(mg/g)	1.15 $\pm$ 0.18
B <sub>1</sub> ( $\mu$ g/g)	0.14 $\pm$ 0.00
B <sub>2</sub> ( $\mu$ g/g)	0.12 $\pm$ 0.01
B <sub>3</sub> ( $\mu$ g/g)	0.26 $\pm$ 0.02
C (mg/100g)	8.12 $\pm$ 0.16
E (mg/g)	1.44 $\pm$ 0.15
B <sub>9</sub> ( $\mu$ g/g)	81.23 $\pm$ 6.29

Values are mean  $\pm$  standard deviation (n=2)

#### Bioactive Compound Content

Table 2 shows the result of the bioactive compound composition of fruit of *Ficus capensis*. Flavonoid content was higher (13.97 mgGAE/100) than other bioactive compounds evaluated.

Table 2: Bioactive Compound Composition of fruit of *Ficus capensis*.

Bioactive Compounds	Concentration
Total Phenol (mgGAE/100g)	13.97 ± 1.39
Flavonoids (mgCE/100g)	11.03 ± 0.16
Beta Carotene (mg/100g)	0.34 ± 0.001
Lycopene (mg/100g)	0.14 ± 0.002

Values are mean ± standard deviation (n=2)

## Antioxidant Activity

### DPPH Radical Scavenging Activity

The DPPH radical scavenging activity of the fruit of *Ficus capensis* was dose-dependent. The highest activity was observed at the highest concentration tested (100 µg/ml) and gave 21.19% activity while the standard antioxidant (BHA) gave 89.96% activity as shown in Fig. 1. The effective concentration at 50% (EC<sub>50</sub>) was 30 µg/ml and 28 µg/ml for the fruit of *Ficus capensis* and BHA respectively (Fig 2).




Fig. 1: Percentage DPPH Radical Scavenging Activity (RSA) of fruit of *Ficus capensis* and BHA




Fig. 2: The EC<sub>50</sub> of fruit of *Ficus capensis* and BHA

### Reducing Power Activity

The reducing power activity of fruit of *Ficus capensis* and BHA is shown in Fig. 3. As the concentration increased, the reducing power activity increased shown by the increase in the absorbance. At 100 µg/ml, the absorbance of fruit of *Ficus capensis* was 0.623 while that of BHA was 1.352. Fig. 4 shows the OD<sub>0.5</sub> of 38µg/ml and 34 µg/ml for fruit of *Ficus capensis* and BHA respectively.




Fig. 3: Reducing Power Activity of fruit of *Ficus capensis* and BHA




Fig. 4: OD<sub>0.5</sub> of *Ficus capensis* fruit and BHA

### Inhibition of Lipid Peroxidation

The percentage inhibition of lipid peroxidation activity of fruit of *Ficus capensis* and BHA is presented in Fig.5. The activities were concentration-dependent with BHA exhibiting higher activity than *Ficus capensis* fruit. At 100 µg/ml, the activities were 29.294% and 72.703% for the fruit of *Ficus capensis* and BHA respectively. The EC<sub>50</sub> for the fruit of *Ficus capensis* was 80 µg/ml while that of BHA was 62 µg/ml (Fig. 6).




Fig. 5: Percentage Inhibitory ratio of fruit of *Ficus capensis* and BHA




Fig 6:  $EC_{50}$  of fruit of *Ficus capensis* and BHA.

## Discussion

Plants have proven to be effective in the treatment and management of diseases. Hence, there is increase in efforts at identifying and exploiting potentially beneficial plants. Other parts of *Ficus capensis* has been reported to possess several potential health benefits, little is known about health benefits of the fruits. In view of this, this work intends to unravel the vitamin and bioactive compound contents of the fruits as well as the *in-vitro* antioxidant mechanism.

## Vitamin Composition

The result of the vitamin composition showed the presence of vitamins A, B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>9</sub>, C and E (Table 1). Vitamin B<sub>9</sub> was the highest in concentration (81.23  $\mu\text{g/g}$ ). These vitamins are molecules required in small amount and play important roles in cellular metabolism where they act as coenzymes as well as in health and well-being. Their deficiencies have been linked to some diseases (Traber and Stevens, 2011). The B vitamins majorly function as coenzymes in metabolic pathways of carbohydrate, proteins and amino acids. Vitamin C and E are well-known antioxidants that play vital roles in combating oxidative stress in cells such as in cancer cells (Risvi et al., 2014; Montecinos et al., 2014). Vitamin C is involved in many body functions including formation of collagen, absorption of iron, the proper functioning of the immune system, wound healing and maintenance of cartilage, bones and teeth (Haworth et al., 1993). Derivatives of vitamin A such as all-trans- retinoic acid are used in the management of acute promyelocytic leukaemia (Sanz et al., 2010). Some of these vitamins and their derivatives have been suggested as chemo-preventive agents (Freemantle et al., 2003). Therefore, the discovery of more sources of these vitamins and their derivatives like *Ficus capensis* fruit can be a way to manage chronic diseases.

## Bioactive Compounds Composition

Bioactive compounds are active medicinal chemical constituents. The presence of bioactive phytochemical constituents present in different plants parts gives them their medicinal value and biological activities (Thilakarathna and Rupasinghe, 2013). Important bioactive compounds such as phenol, flavonoids, beta carotene and lycopene were present in the fruit of *Ficus capensis* (Table 2). *Ficus capensis* fruits showed reasonable amount of phenol at the concentration of  $13.97 \pm 1.39$  mgGAE/100g which have been reported to possess some medicinal properties. It has a defense role against free radicals and exhibits several properties beneficial to humans and its antioxidant property are important in determining their role as protecting agents against free radical-mediated diseases process (Cai et al., 2006). The presence of phenol may indicate that fruit of *Ficus capensis* have antioxidant activity. Flavonoids was present in the concentration of  $11.03 \pm 0.16$  mg/g. Flavonoids possess many medicinal benefits such as anti-cancer, anti-oxidant, antiviral properties and anti-inflammatory as well as cardio-protective and neuroprotective effects (Zhao et al., 2019; Zhao et al., 2018; Camero et al., 2018; Patel et al., 2018; Mazidi et al., 2019). The concentration of lycopene ( $0.14 \pm 0.002$  mg/100g) was very low. Lycopene is a carotenoid that naturally gives foods their pink/red colour. It is said to be good for bone, heart and vascular health (Russo et al., 2020; Li and Xu, 2014) and, as a potent antioxidant especially effective at scavenging singlet oxygen (Sies and Stahl, 1995). Beta carotene was slightly present and was found in the concentration of ( $0.34 \pm 0.001$  mg/g).  $\beta$ -Carotene is regarded as the major carotenoid present in human diet (Johnson, 2002) and the main source of vitamin A in humans. It plays a crucial role in prevention of degenerative diseases such as cardiovascular diseases, diabetes and several types of cancers (Singh and Goyal, 2008).

## Antioxidant Activity

### DPPH scavenging activity:

One of the in-vitro antioxidant parameters used for testing the potency of plants is their ability to scavenge free radicals. DPPH radical scavenging assay is the most common method used in the study of antioxidant activity of plant extracts. This mechanism involves the protonation of the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radicals turning it to stable free radical which is visually noticeable as a decoloration from purple to golden yellow and detected by spectrophotometric technique. Decrease in absorbance shows the more efficient antioxidant activity of the extract in terms of hydrogen atom donating capacity. *Ficus capensis* fruit demonstrated radical scavenging activity in a dose-dependent manner. Antioxidant activity of *Ficus capensis* fruit as measured by ability to scavenge (DPPH) free radicals was compared with the standard antioxidant; butylated hydroxyl toluene (BHT). The standard however gave a higher activity. The presence of phenolic compounds in the fruit of *Ficus capensis* may contribute directly to the observed antioxidant effect as phenolic compounds have been reported to have antioxidant activities (Afari et al., 2014).

The effective concentration of the extract at half maximum DPPH scavenging activity ( $EC_{50}$ ) was calculated from graph of percentage scavenging activity against concentration of the extract was  $30 \mu\text{g/ml}$  and that of BHA was  $28 \mu\text{g/ml}$ . The  $EC_{50}$  was used to evaluate the antioxidant effectiveness of the sample. The lower the  $EC_{50}$  value, the higher is the

scavenging potential. This shows that BHA was the stronger DPPH scavenger having the lower EC<sub>50</sub> value.

### **Reducing Power Assay**

In reducing power assay, the yellow colour of the **test solution** changes to different shades of green depending on the reducing power of the test specimen. The presence of the reductants in the solution causes the reduction of the Fe<sup>3+</sup>/ferricyanide complex to the ferrous form. Therefore, Fe<sup>2+</sup> can be monitored by absorbance measurement at 700 nm. Previous studies have reported and suggested that the reducing properties have been shown to exert antioxidant action by donating a hydrogen atom to break the free radical chain (Gordon, 1990). In this study, the observed increasing absorbance at 700 nm as the concentration of the extract increased indicates an increase in reducing ability. The antioxidants present in the fruit of *Ficus capensis* caused the reduction of Fe<sup>3+</sup>/ferricyanide complex to the ferrous form, and thus proved the reducing power.

### **Inhibition of lipid peroxidation**

The thiobarbituric acid reactive substances (TBARS) method has been used widely in determination of the extent of lipid peroxidation via formation of malondialdehyde (MDA) compound from lipid peroxidation of polyunsaturated fatty acids. Using brain homogenate as a medium of peroxidation in this study, percentage lipid peroxidation inhibition by the extract of fruit of *Ficus capensis* increased with increase in concentration. Antioxidant potential of extract correlates directly to the scavenging of hydroxyl radical and consequently the inhibition of lipid peroxidation (Orabi et al., 2001). Lipid hydroperoxides generation can cause damage to molecules of the bio-system and have the ability to bind with DNA leading to strand breaks, mutation and carcinogenesis (Sharififaret et al., 2009). Intrinsically, erythrocytes are highly susceptible to peroxidation due to the heavy buildup of polyunsaturated fatty acids and haemoglobin. During respiration erythrocytes are frequently exposed to high tension of oxygen, which can induce oxidative damage (Babu et al., 2001).

In this study, fruit of *Ficus capensis* exhibited scavenging of radicals indicating the presence of primary antioxidants which possess anti-lipid peroxidation potential. Phenolic compounds have been reported to delay oxidative degradation of lipids as well as enhance the quality and nutritional value of food (Kähkönen et al., 1994).

### **Conclusion**

Plant sources with antioxidant abilities is of great interest to industries and researchers because of their medicinal properties in the supportive and definitive care of debilitating diseases. This study investigated the vitamin content, bioactive compound composition and antioxidant properties of fruit of *Ficus capensis*. The study showed that vitamins and bioactive compounds were present in varying concentrations. The antioxidant activity exhibited in different *in-vitro* antioxidant assays carried out in this study may be linked to the antioxidant constituents such as phenolic, flavonoid, ascorbic acid, lycopene and beta-carotene found in the fruit of *Ficus capensis*. These results suggest that fruit of *Ficus capensis* may be a potential source of natural antioxidants.

## References

1. Adesina SK. Traditional medical care in Nigeria. Online Nigeria Daily News; 2014.
2. Gökmen V. Acrylamide in food: Analysis, content and potential health effects; Elsevier. 2015.
3. Siriwardhana N, Kalupahana NS, Cekanova M, LeMieux M, Greer B, Moustaid-Moussa N. Modulation of adipose tissue inflammation by bioactive food compounds. *J Nutr Biochem.* 2013; 24(4): 613-623.
4. Hassimotto NM, Genovese MI, Lajola FM. Antioxidant activity of dietary fruits, vegetables and commercial frozen fruit pulps. *J Agri Food Chem.* 2005; 53 (8): 2928-2935.
5. Iqbal S, Bhangar M. Effect of season and production location on antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. *J Food Comp Anal.* 2006; 19:544-551
6. Sultana B, Anwar F. Flavonols (kaempferol, quercetin, myricetin) contents of selected fruits, vegetables and medicinal plants. *Food Chem.* 2008; 108:879-884.
7. Braca A, Sortino C, Politi M, Morelli I, Mendez J. Antioxidant activity of flavonoids from *Licanialicaniaeflora*. *J Ethnopharmacol.* 2002; 79:379-381
8. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J Nutr Biochem.* 2002; 13:572- 584
9. Montoro P, Braca A, Pizza C, De Tommasi N. Structure-antioxidant activity relationships of flavonoids isolated from different plant species. *Food Chem.* 2005; 92:349-355.
10. Sirisha N, Screenivasulu M, Sangeeta K, Madhusudhana C. Antioxidant properties of ficus species a review. *Intern J Pharm. Tech. Res.* 2010;2(4):2174-2182
11. Onyeleke SB, Dauda SB, Boye OA. Antibacterial activity of *Ficus capensis* Thunb. Africa. *Journal Biotechnology.* 2008; 7(10): 1414-1417
12. Ayinde BA, Owolabi OJ. Effects of the aqueous extract of *Ficus capensis* Thunb. (Moraceae) leaf on gastrointestinal motility. *J Pharmacog. Phyto.* 2009; 1(3):031-035.
13. Ayinde BA, Owolabi OJ, Jesuoroba RI. Active ileum relaxant fraction from the leaves of *Ficus capensis*, Thunb (Moraaceae). *Nig J Pharm.* 2013; 10(1):1-10.
14. Adebayo-Tayo BC, Odeniyi AO. Phytochemical screening and microbial inhibitory activities of *Ficus capensis*. *Afri J Biomed. Res.* 2012; 15:35- 40.
15. Daikwo OA, Tende JA, Okey SM, Eze ED, Isa AS. The effect of aqueous extract of leaf of *Ficus capensis* Thunb (Moraceae) on in Vivo leukocyte mobilization in wistar rats. *Bri. J Pharmacol. Toxicol.* 2012; 3(3): 110-114
16. Rutkowski M, Grzegorzcyk K, Gendek E, Kedziora J. Laboratory convenient modification of Bessey method for vitamin A determination in blood plasma *J. Physiol. Pharm.* 2006; 57 (2):221
17. Klein BP, Perry AK. Ascorbic acid and vitamin A activity in selected vegetables from different geographical areas of the United States. *J Food Sci.* 1982; 47: 941-945.
18. Kirk R, Sawyer R. Perason's composition and analysis of foods, 9th edition. Longman Scientific and Technical. 1991; p. 647-648
19. Padmarajaiah N, Ramanathapura AV, Hemmige SY. Spectrophotometric determination of folic acid in pharmaceutical preparations by coupling reactions with iminodibenzyl or 3-aminophenol or sodium molybdate-pyrocatechol. *Anal Biochem.* 2002; 307(2):316-321.

20. Barros L, Ferreira MJ, Queiros B, Ferreira ICFR, Baptista P. Total phenols, ascorbic acid,  $\beta$ -carotene and lycopene in portuguese wild edible mushrooms and their antioxidant activities. *Food chemistry*. 2007; 103(2): 413-419.
21. Ebrahimzadeh MA, Seyed MN, Seyed FN, Fatemeh B, Ahmad RB. Antioxidant and free radical scavenging activity of *H. officinalis*, *L. angustifolius*, *V. odorata*, *B. hyrcana* and *C. speciosum*. *Pak. Journal of Pharm Sci*. 2009; 23(1):29-34.
22. Traber MG, Stevens JF. Vitamins C and E: Beneficial effects from a mechanistic perspective. *Free Radical Biology and Medicine*. 2011; 51(5):1000-13
23. Rizvi S, Raza ST, Ahmed F, Ahmad A, Abbas S, Mahdi F. The Role of Vitamin E in Human Health and Some Diseases. *Sultan Qaboos University Medical Journal*. 2014; 14(2):e157-65.
24. Montecinos V, Guzmán P, Barra V, Villagrán M, Muñoz-Montesino C, et al. Vitamin C is an Essential Antioxidant That Enhances Survival of Oxidatively Stressed Human Vascular Endothelial Cells in the Presence of a Vast Molar Excess of Glutathione. *The Journal of Biological Chemistry*. 2014; 282(21):15506- 15.
25. Haworth WN, Hirt El. Synthesis of ascorbic acid. *Journal society of Chemical industry (london)*. 1993; 52:645-647.
26. Sanz MA, Montesinos P, Rayón C, Holowiecka A, Serna JDL, Milone G, et al. (on behalf of the PETHEMA and HOVON Groups). Risk-adapted treatment of acute promyelocytic leukaemia based on all-trans retinoic acid and anthracycline and addition of cytarabine in consolidation therapy for high-risk patients: further improvement in treatment outcome. *Blood*. 2010; 115:5137-46.
27. Freemantle SJ, Spinella MJ, Dmitrovsky E. Retinoids in cancer therapy and chemoprevention: promise meets resistance. *Oncogene*. 2003; 22(47):7305-15.
28. Thilakarathna S, Rupasinghe H. Flavonoid Bioavailability and Attempts for Bioavailability Enhancement. *Nutrients*. 2013; 5, 3367–3387
29. Cai YZ, Sun M, Xing J, Luo Q, Corke H. Structure–radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Sci*. 2006;78:2872-2888.
30. Zhao L, Yuan X, Wang J, Feng Y, Ji F, Li Z, Bian J. A review on flavones targeting serine/threonine protein kinases for potential anticancer drugs. *Bioorganic Med. Chem*. 2019; 27, 677–685.
31. Zhao K, Yuan Y, Lin B, Miao Z, Li Z, Guo Q, Lu N. LW-215, a newly synthesized flavonoid, exhibits potent anti-angiogenic activity in vitro and in vivo. *Gene*. 2018; 642, 533–541.
32. Camero CM, Germanò MP, Rapisarda A, D'Angelo V, Amira S, Benchikh F, Braca A, De Leo M. Anti-angiogenic activity of iridoids from *Galiumtunetanum*. *Rev. Bras. de Farmacogn*. 2018; 28, 374–377.
33. Patel K, Kumar V, Rahman M, Verma A, Patel DK. New insights into the medicinal importance, physiological functions and bioanalytical aspects of an important bioactive compound of foods 'Hyperin': Health benefits of the past, the present, the future. *Beni-Suef Univ. J. Basic Appl. Sci*. 2018; 7, 31–42.
34. Mazidi M, Katsiki N, Banach M. A higher flavonoid intake is associated with less likelihood of nonalcoholic fatty liver disease: Results from a multiethnic study. *J. Nutr. Biochem*. 2019; 65, 66–71.

35. Russo C, Ferro Y, Maurotti S, et al., Lycopene and bone: an in-vitro investigation and pilot prospective clinical study. *J Transl Med.* 2020; 18(1):43.
36. Li X, Xu J. dietary and circulating lycopene and stroke risk: a meta-analysis of prospective studies. *Scientific Reports.* 2014; 4(1).
37. Sies H, Stahl W. vitamin E and Vitamin C, beta-carotene, and other carotenoids as antioxidants. *Am J clin Nutr,* 1995; 62(6): 1315S-21S.
38. Johnson EJ. The role of carotenoids in human health. *Nutrition clinic care.* 2002; (2):56-65.
39. Singh P, Goyal GK. Dietary Lycopene: Its properties and anticarcinogenic effects. *Comprehensive Reviews in Food Science and Food safety.* 7(3): 255-270.
40. Afari S, Saeidnia S, Abdollahi M. Role of natural phenolic compounds in cancer chemoprevention via regulation of the cell cycle. *Current Pharmaceutical Biotechnology.* 2014; 15(4):409-21
41. Gordon MH. The mechanism of antioxidant action in vitro. In *Food antioxidants.* Edited by Hudson BJ. London: Elsevier Applied Science; 1990: 1–18.
42. Orabi KY, Al-Qasoumi SI, EI-Qlemy MM, Mossa JS, Muhammad I: (Dihydroagarofuran alkaloid and triterpenes from *Maytenus heterophylla* and *Maytenus arbutifolia*. *Phytochemistry.* 2001; 58:475–480.
43. Sharififar F, Dehghan-nudeh GH, Mirtajaldini M: Major flavonoids with antioxidant activity from *Teucrium polium* L. *Food Chem.* 2009; 112:885–888.
44. Babu BH, Shylesh BS, Padikkala J: Antioxidant and hepatoprotective effect of *Alanthusicifocus*. *Fitoterapia.* 2001; 72:272–27
45. Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M: Antioxidant activity of plant extracts containing phenolic compounds. *J Agri Food Chem.* 1999; 1999(47):3954–3962