

Screening of different genotypes of field pea against major insect pests.

Abstract:

In the rabi season of 2020-21, a field trial was executed at the Breeder Seed Production Farm, JNKVV, Jabalpur, M.P., aimed at assessing the resistance of 71 field pea genotypes against Aphid, Leaf hopper, and Gram pod Borer. Through weekly observations and subsequent statistical analysis, genotypes were categorized into distinct resistance groups. Prominent findings revealed that JP-180, Matar Rangpur, and Aman 1-206 exhibited high resistance to aphids and pod borers. Additionally, Kashi Samriddhi, JP-180, and Matar Rangpur demonstrated resistance to leafhoppers. Conversely, B-22, DDR-39, and KPMR-485 displayed high susceptibility to Aphids, while Batripatiram, VL-3, and VL-3 were highly susceptible to Gram pod borer. KPMR-485, VL-3, and VL-1 were found to be susceptible to both Aphid and Gram pod borer. The study further revealed that the genotype KPMR-485 achieved the highest pea yield, closely followed by Jayanti and IPF-99-25. In contrast, Safed Batra Gudda, Matar Rangpur, and Aman 1-206 exhibited the minimum yield. These findings provide valuable insights into the resistance patterns of field pea genotypes, aiding in the selection of cultivars with enhanced resistance and productivity for sustainable agricultural practices.

Keywords: Screening, Resistance, Susceptible, Yield, Aphid, Podborer, Leafhopper.

Introduction:

Field pea, a diploid crop that undergoes self-pollination, holds significant agricultural importance as a high-quality protein source. This annual crop is predominantly cultivated in cooler temperate zones and tropical highlands globally. While versatile in adapting to various soil types, ranging from light sandy loam to heavy clays, it is sensitive to both saline and waterlogged conditions.

Field peas boast a rich content of high-quality vegetable protein, encompassing all essential amino acids. While sulfur-containing amino acids like cysteine and methionine are relatively scarce, field peas are abundant in lysine and other essential amino acids. With a protein content ranging from 23% to 25%, field

peas serve as a valuable source of essential amino acids, providing substantial nutritional benefits, particularly for low-income families (Nawab et al., 2008).

Moreover, peas are a plentiful source of essential minerals, including calcium, phosphorus, and iron—nutrients that are often deficient in cereals (Haque et al., 2015). Globally, peas serve as a significant export and cash crop, contributing to approximately 40% of the total world pulse trade. Notably, peas have the unique ability to fix atmospheric nitrogen, making it available for the crop, further enhancing their agricultural value.

Field peas are commonly integrated into forage crop mixtures alongside small grains. The protein content in field pea forage typically falls within the range of 18-20%. When inter-seeded at a rate of 60-100 pounds per acre with a small grain like oats, field peas can enhance the protein concentration of the mixed forage by 2-4% points, thereby elevating the relative feed value by 20 points.

Insect pests pose significant biotic challenges, leading to substantial global losses. Notably, pests such as pea leaf miner (*Chromatomyia horticola* Goureau), pea aphid (*Acyrtosiphon pisum*), pod borer complex (*Helicoverpa armigera* (H), *Lampides boeticus* (L), and *Etiellazinkenella* Tr.), and thrips (*Caliothrips indicus* Bagnall) are prominent among these constraints, causing considerable damage to the crop.

The American bollworm, *H. armigera*, is a widely distributed pest with a global presence in Africa, Asia, Europe, and Australia. Among its host plants is the field pea, where the larvae inflict damage by creating circular holes in pea pods, directly feeding on the seeds. This feeding behavior often leads to substantial seed damage.

The pea aphid (*Acyrtosiphon pisum*), commonly found in field pea crops, is a small insect measuring approximately 3 millimeters (1/8 inches) in length. It has a light green color and long legs, and individuals may either be wingless or possess prominent, translucent wings. By extracting sap from the plant, the pea aphid weakens the field pea directly. When aphids feed on peas during the early pod stage, it can result in reduced yields due to diminished seed formation and smaller seed size. Although protein content and other quality aspects do not seem to be affected, the overall health and productivity of the plant are compromised.

Materials And Methods

The experiment was carried out at the Experimental Field of the Department of Plant Breeding, Seed Breeding Farm, JNKVV Jabalpur (MP) during the Rabi season of 2020-21. A Randomized Block Design (RBD) was employed, with a plot size of 10x10 and three replications, incorporating 71 genotypes. The spacing maintained was 30x10 cm. Throughout the growing season, no pesticidal treatment was administered to the crop. Regular observations were conducted during various weather weeks to document the density of major insect pests affecting field pea, namely Pea Aphids (*Acyrtosiphonpissum*), Pod Borer (*Helicoverpaarmigera* H.), and Leaf Hopper (*Empoasca fabae*).

The sample unit for recording the density of sucking pests, such as aphids, was defined as the top 10 cm apical twig of the sample plant. Five plants were randomly selected from each plot, and observations were recorded weekly. Pest population assessments were conducted at different crop stages during varying weather weeks. Weekly observations were made to record borer density, sucking insect pests like aphids, pod damage (%), and pea yields under different treatments. The collected data were tabulated into transformed values and subjected to statistical analysis using the Analysis of Variance technique at a 5% level of significance.

Result And Discussion:

71 genotypes of pea, were screened for relative resistance to major sucking insect pests during the year, 2020-21. The infestation was recorded at weekly interval on five randomly selected and tagged plants just after initiation of sucking insect pests to disappearance.

Aphid

The first observation recorded on 30 DAS. The mean aphid population ranged from 8.08 (B-22) to 16.35 (IPF-99-25(Local check) /10 cm apical twigs/ plant. The minimum infestation was observed on genotype B-22 followed by KPMR-420, IPF-99-25, P-3, Batripatiram, VL-3, VL-1, KPMR-485, Jayanti, DDR-39 which were statistically at par with each other. The maximum infestation was observed on genotype IPF-99-25 (Local check) followed by Safed Batra gudda, Demo-JP-180, JP-885 (local check), Matar Rangpur, Double Branching, Aman 1-206, Kashi Samriddhi, JPMR-402 and NDVP-20. These were statistically significant in degree of infestation with IPF-99-25 (Local check).

Based on overall mean population of the season on different genotypes of pea, it was minimum on genotype B-22 (19.57 aphid/ **10 cm apical twigs**) followed by DDR-39, KPMR-485, JFP-99-25, VL-3, Jayanti, KPMR-420, Shikha, P-3, Batripatiram, VL-1, these were found at par with each other (except B-22). The maximum population of aphid was recorded in the genotype Double branching (31.15 aphid/ **10 cm apical twigs**) followed by Demo-JP-180, Safed Batra Gudda, Double branching, Matar Rangpur, NDVP-20, IPF-99-25 (local check), Aman 1-206 and KPMR-402, which were found at par with each other.

Present findings also supported with the findings of Wale (2002) Identified *Acyrtosiphonpissum*(Harris), a pea aphid that was once a minor pest in Ethiopia, has recently emerged as a major pest of field pea (*Pisumsativum* L.) in the northwestern region. Field experiments were performed in Adet and Zema in northwestern Ethiopia in 1994, 1995, and 1996 to determine the pest's population dynamics. In May, June, and July, planting dates were spaced 15 days apart. In August, aphid colonies at Adet were at their peak. In both 1994 and 1995, weekly counts of pea aphid numbers peaked in late August, and in mid- to late-August in

1996. In 1995, a delay in sowing date resulted in a large increase in pea aphid numbers, and a nearly identical increase in 1996.

Table 1.1: Categorization of pea genotypes into degree of susceptibility against

S. No.	Mean aphid population/ 10 cm apical twigs	Name of genotypes	Category
1	19.57-24.02	B-22, DDR-39, KPMR-485, JFP-99-25, VL-3, Jayanti, KPMR-420, Shikha, P-3, BatriPatiram and VL-1	Least susceptible

3	>25.18	Triple branching, Demo-JP-180, Safed Batra Gudda, Double branching, Kashi samriddhi, Matar Rangpur, NDVP-20, IPF-99-25 (Local check), Aman 1-206, KPMR-402 and JP-885 (local check)	Highly susceptible
---	--------	---	--------------------

Aphid

2	24.16-25.18	IPF-99-25, SPS-2, KPMR-30, NDVP-4, PP-14, DhanoliBatri, Kala matar, HUVP-2, DDR-43, DDR-27, KPMR-503, PP-155, NDVP-20, Chhoti safed Anju, Atrumatar, LEP-260, PP-96, JFP-27, PP-96, KFP-151, PP-86, HUVP-12, DDP 94-14, BatanaMoolchand, Kashmiri samriddhi, DDR-44, Gol Batra Tenduna, DDR-23, DDR-54, Late sown, Dhan Batri, Rachna, HUP-2, HFP-94-12, KPMR-327, KPMR-504, RP-3, JM-91-01, KPMR-402, DDR-52, KPMR-302, Kali Batri, KPMR-400, KPMR-486 and HFP-94-13	Moderately susceptible
---	-------------	---	------------------------

UNDER PEER REVIEW

Table .1: Screening of different genotypes of field pea against aphid of pea

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
IPF-99-25	0.00	10.25	28.15	37.21	56.21	51.28	26.01	20.77	18.69	5.45	4.52	2.39	23.09
	(0.71)	(3.28)	(5.35)	(6.14)	(7.53)	(7.20)	(5.15)	(4.61)	(4.38)	(2.44)	(2.24)	(1.70)	(4.86)
B-22	0.00	8.08	26.10	34.25	50.25	48.62	20.05	14.72	12.55	3.32	2.44	0.36	19.57
	(0.71)	(2.93)	(5.16)	(5.89)	(7.12)	(7.01)	(4.53)	(3.90)	(3.61)	(1.95)	(1.71)	(0.93)	(4.48)
DDR-27	0.00	10.25	28.24	37.24	57.24	52.31	27.06	21.76	19.92	6.68	5.79	3.66	23.87
	(0.71)	(3.28)	(5.36)	(6.14)	(7.60)	(7.27)	(5.25)	(4.72)	(4.52)	(2.68)	(2.51)	(2.04)	(4.94)
PP-155	0.00	10.35	28.39	37.39	57.39	52.46	27.19	21.92	19.81	6.57	5.60	3.47	23.91
	(0.71)	(3.29)	(5.37)	(6.16)	(7.61)	(7.28)	(5.26)	(4.73)	(4.51)	(2.66)	(2.47)	(1.99)	(4.94)
Kashmiri samriddhi	0.00	11.00	29.15	38.15	58.15	53.22	27.99	22.75	20.55	7.31	6.40	4.21	24.66
	(0.71)	(3.39)	(5.45)	(6.22)	(7.66)	(7.33)	(5.34)	(4.82)	(4.59)	(2.79)	(2.63)	(2.17)	(5.02)
DDR-43	0.00	10.35	28.35	37.35	57.31	52.38	27.11	21.82	19.71	6.47	5.58	3.49	23.86
	(0.71)	(3.29)	(5.37)	(6.15)	(7.60)	(7.27)	(5.25)	(4.72)	(4.50)	(2.64)	(2.47)	(2.00)	(4.94)
DDR-44	0.00	11.10	29.16	38.16	58.11	53.18	27.92	22.61	20.55	7.33	6.42	4.33	24.67
	(0.71)	(3.41)	(5.45)	(6.22)	(7.66)	(7.33)	(5.33)	(4.81)	(4.59)	(2.80)	(2.63)	(2.20)	(5.02)
DDR-39	0.00	8.12	26.08	34.08	54.18	47.25	23.98	18.59	16.49	3.65	2.70	0.54	20.81
	(0.71)	(2.94)	(5.16)	(5.88)	(7.39)	(6.91)	(4.95)	(4.37)	(4.12)	(2.04)	(1.79)	(1.02)	(4.62)
Late sown	0.00	11.14	29.29	38.29	58.33	53.40	28.16	22.71	20.55	7.31	6.42	4.29	24.76
	(0.71)	(3.41)	(5.46)	(6.23)	(7.67)	(7.34)	(5.35)	(4.82)	(4.59)	(2.79)	(2.63)	(2.19)	(5.03)

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
DDR-23	0.00	11.21	29.22	38.22	58.26	53.33	28.14	22.69	20.56	7.32	6.42	4.29	24.74
	(0.71)	(3.42)	(5.45)	(6.22)	(7.67)	(7.34)	(5.35)	(4.82)	(4.59)	(2.80)	(2.63)	(2.19)	(5.02)
HUP-2	0.00	10.95	28.99	37.99	58.62	53.69	28.40	23.16	21.03	7.79	6.88	4.75	24.94
	(0.71)	(3.38)	(5.43)	(6.20)	(7.69)	(7.36)	(5.38)	(4.86)	(4.64)	(2.88)	(2.72)	(2.29)	(5.04)
DDP94-14	0.00	10.99	29.05	38.10	58.10	53.17	27.88	22.59	20.49	7.25	6.36	4.23	24.60
	(0.71)	(3.39)	(5.44)	(6.21)	(7.66)	(7.33)	(5.33)	(4.81)	(4.58)	(2.78)	(2.62)	(2.17)	(5.01)
HUVP-2	0.00	10.11	28.26	37.29	57.29	52.36	27.07	21.78	19.69	6.45	5.56	3.43	23.79
	(0.71)	(3.26)	(5.36)	(6.15)	(7.60)	(7.27)	(5.25)	(4.72)	(4.49)	(2.64)	(2.46)	(1.98)	(4.93)
KPMR-402	0.00	14.21	33.39	42.39	63.69	57.62	33.47	29.18	27.11	10.28	9.39	6.26	28.93
	(0.71)	(3.84)	(5.82)	(6.55)	(8.01)	(7.62)	(5.83)	(5.45)	(5.25)	(3.28)	(3.14)	(2.60)	(5.43)
NDVP-20	0.00	13.25	34.29	43.31	65.36	59.34	35.14	30.85	28.72	9.48	8.57	5.44	29.42
	(0.71)	(3.71)	(5.90)	(6.62)	(8.12)	(7.74)	(5.97)	(5.60)	(5.41)	(3.16)	(3.01)	(2.44)	(5.47)
Atrumatar	0.00	10.25	28.29	37.33	57.33	52.40	27.11	22.82	20.69	6.59	5.67	3.54	24.02
	(0.71)	(3.28)	(5.37)	(6.15)	(7.60)	(7.27)	(5.25)	(4.83)	(4.60)	(2.66)	(2.48)	(2.01)	(4.95)
Triple branching	0.00	14.82	34.98	43.98	66.98	62.05	39.77	34.48	32.35	9.62	8.70	5.24	31.15
	(0.71)	(3.91)	(5.96)	(6.67)	(8.21)	(7.91)	(6.35)	(5.91)	(5.73)	(3.18)	(3.03)	(2.40)	(5.63)
Double Branching	0.00	14.96	33.98	42.98	67.28	62.35	37.11	31.82	29.95	8.71	7.79	4.66	30.21
	(0.71)	(3.93)	(5.87)	(6.59)	(8.23)	(7.93)	(6.13)	(5.69)	(5.52)	(3.03)	(2.88)	(2.27)	(5.54)
SPS-2	0.00	10.50	28.52	37.52	57.00	52.07	26.81	21.52	19.39	5.16	4.24	2.11	23.45
	(0.71)	(3.32)	(5.39)	(6.17)	(7.58)	(7.25)	(5.23)	(4.69)	(4.46)	(2.38)	(2.18)	(1.62)	(4.89)
JM-91-01	0.00	11.75	29.75	38.75	58.75	53.92	28.53	23.24	21.11	7.87	6.95	3.82	25.18

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
	(0.71)	(3.50)	(5.50)	(6.26)	(7.70)	(7.38)	(5.39)	(4.87)	(4.65)	(2.89)	(2.73)	(2.08)	(5.07)
P-3	0.00	8.88	26.88	34.88	54.88	48.95	24.66	19.12	16.99	3.79	2.80	0.62	21.44
	(0.71)	(3.06)	(5.23)	(5.95)	(7.44)	(7.03)	(5.02)	(4.43)	(4.18)	(2.07)	(1.82)	(1.06)	(4.68)
RP-3	0.00	10.75	28.75	37.75	57.75	52.82	27.53	24.62	22.49	9.25	8.33	5.12	25.16
	(0.71)	(3.35)	(5.41)	(6.18)	(7.63)	(7.30)	(5.29)	(5.01)	(4.79)	(3.12)	(2.97)	(2.37)	(5.07)
VL-3	0.00	8.65	26.65	34.65	54.65	48.54	24.43	18.14	16.01	3.65	2.79	0.69	21.13
	(0.71)	(3.02)	(5.21)	(5.93)	(7.43)	(7.00)	(4.99)	(4.32)	(4.06)	(2.04)	(1.81)	(1.09)	(4.65)
JFP-27	0.00	10.76	28.76	37.76	57.76	52.83	27.54	22.25	20.12	6.88	5.96	3.84	24.27
	(0.71)	(3.36)	(5.41)	(6.19)	(7.63)	(7.30)	(5.30)	(4.77)	(4.54)	(2.72)	(2.54)	(2.08)	(4.98)
PP-86	0.00	11.01	29.01	38.01	58.01	53.08	27.79	22.50	20.37	7.13	6.21	4.11	24.52
	(0.71)	(3.39)	(5.43)	(6.21)	(7.65)	(7.32)	(5.32)	(4.80)	(4.57)	(2.76)	(2.59)	(2.15)	(5.00)
NDVP-4	0.00	10.05	28.09	37.09	57.09	52.16	26.87	21.58	19.45	6.21	5.29	3.16	23.59
	(0.71)	(3.25)	(5.35)	(6.13)	(7.59)	(7.26)	(5.23)	(4.70)	(4.47)	(2.59)	(2.41)	(1.91)	(4.91)
NDVP-20	0.00	9.65	27.69	37.69	57.65	52.77	27.43	22.14	20.14	6.90	5.98	3.85	23.97
	(0.71)	(3.19)	(5.31)	(6.18)	(7.63)	(7.30)	(5.28)	(4.76)	(4.54)	(2.72)	(2.55)	(2.09)	(4.95)
KFP-151	0.00	9.99	27.99	37.99	58.22	53.36	28.00	22.69	20.59	7.35	6.43	4.30	24.41
	(0.71)	(3.24)	(5.34)	(6.20)	(7.66)	(7.34)	(5.34)	(4.82)	(4.59)	(2.80)	(2.63)	(2.19)	(4.99)
HUVP-12	0.00	10.14	28.15	38.15	58.34	53.41	28.12	22.83	20.73	7.49	6.57	4.44	24.54
	(0.71)	(3.26)	(5.35)	(6.22)	(7.67)	(7.34)	(5.35)	(4.83)	(4.61)	(2.83)	(2.66)	(2.22)	(5.00)
LEP-260	0.00	10.62	28.71	37.71	57.61	52.68	27.39	22.10	19.98	6.79	5.87	3.74	24.16
	(0.71)	(3.33)	(5.40)	(6.18)	(7.62)	(7.29)	(5.28)	(4.75)	(4.53)	(2.70)	(2.52)	(2.06)	(4.97)

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
DhanoliBatri	0.00	9.35	27.33	37.33	57.39	52.46	27.17	21.95	19.82	6.59	5.66	3.53	23.66
	(0.71)	(3.14)	(5.28)	(6.15)	(7.61)	(7.28)	(5.26)	(4.74)	(4.51)	(2.66)	(2.48)	(2.01)	(4.92)
Gol Batra Tenduna	0.00	9.66	27.69	38.69	58.62	53.69	28.64	24.00	21.87	7.16	6.29	4.16	24.68
	(0.71)	(3.19)	(5.31)	(6.26)	(7.69)	(7.36)	(5.40)	(4.95)	(4.73)	(2.77)	(2.61)	(2.16)	(5.02)
Matar Rangpur	0.00	15.35	34.05	43.05	65.05	60.05	34.88	30.59	28.49	10.25	8.33	5.10	29.69
	(0.71)	(3.98)	(5.88)	(6.60)	(8.10)	(7.78)	(5.95)	(5.58)	(5.38)	(3.28)	(2.97)	(2.37)	(5.49)
Kashi samridhi	0.00	14.22	33.85	42.85	66.81	61.57	36.61	31.32	29.35	10.16	8.64	5.29	30.15
	(0.71)	(3.84)	(5.86)	(6.58)	(8.20)	(7.88)	(6.09)	(5.64)	(5.46)	(3.26)	(3.02)	(2.41)	(5.54)
Kala matar	0.00	10.20	28.25	37.25	57.25	52.32	27.03	21.74	19.61	6.37	5.45	3.32	23.75
	(0.71)	(3.27)	(5.36)	(6.14)	(7.60)	(7.27)	(5.25)	(4.72)	(4.48)	(2.62)	(2.44)	(1.95)	(4.92)
KPMR-503	0.00	10.36	28.39	37.39	57.39	52.46	27.17	21.88	19.75	6.51	5.59	3.46	23.89
	(0.71)	(3.30)	(5.37)	(6.16)	(7.61)	(7.28)	(5.26)	(4.73)	(4.50)	(2.65)	(2.47)	(1.99)	(4.94)
DDR-52	0.00	11.71	29.75	38.75	58.75	53.82	28.53	23.24	21.11	7.87	6.95	4.82	25.25
	(0.71)	(3.49)	(5.50)	(6.26)	(7.70)	(7.37)	(5.39)	(4.87)	(4.65)	(2.89)	(2.73)	(2.31)	(5.07)
DDR-54	0.00	11.24	29.25	38.25	58.25	53.32	28.03	22.74	20.61	7.37	6.45	4.32	24.76
	(0.71)	(3.43)	(5.45)	(6.22)	(7.66)	(7.34)	(5.34)	(4.82)	(4.59)	(2.81)	(2.64)	(2.20)	(5.03)
PP-96	0.00	10.75	28.74	37.74	57.74	52.81	27.52	22.23	20.10	6.86	5.94	3.81	24.25
	(0.71)	(3.35)	(5.41)	(6.18)	(7.63)	(7.30)	(5.29)	(4.77)	(4.54)	(2.71)	(2.54)	(2.08)	(4.97)
KPMR-30	0.00	9.92	27.96	36.96	56.96	52.03	26.74	21.45	19.32	6.08	5.16	3.10	23.47
	(0.71)	(3.23)	(5.33)	(6.12)	(7.58)	(7.25)	(5.22)	(4.69)	(4.45)	(2.57)	(2.38)	(1.90)	(4.90)
JFP-99-25	0.00	8.98	26.68	34.68	53.68	48.69	23.46	18.22	16.95	3.69	2.77	0.63	21.12

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
	(0.71)	(3.08)	(5.21)	(5.93)	(7.36)	(7.01)	(4.89)	(4.33)	(4.18)	(2.05)	(1.81)	(1.06)	(4.65)
PP-96	0.00	10.35	28.35	37.35	57.35	52.42	27.13	21.91	21.51	8.27	7.35	4.29	24.39
	(0.71)	(3.29)	(5.37)	(6.15)	(7.61)	(7.27)	(5.26)	(4.73)	(4.69)	(2.96)	(2.80)	(2.19)	(4.99)
PP-14	0.00	9.67	27.67	36.67	56.67	51.74	26.45	21.35	20.26	7.02	6.15	4.02	23.61
	(0.71)	(3.19)	(5.31)	(6.10)	(7.56)	(7.23)	(5.19)	(4.67)	(4.56)	(2.74)	(2.58)	(2.13)	(4.91)
Aman1-206	0.00	14.25	33.25	42.25	63.65	58.71	34.55	30.26	28.13	9.89	8.97	5.47	29.14
	(0.71)	(3.84)	(5.81)	(6.54)	(8.01)	(7.69)	(5.92)	(5.55)	(5.35)	(3.22)	(3.08)	(2.44)	(5.44)
Demo-JP-180	0.00	15.66	35.66	44.66	66.64	61.46	37.42	33.13	31.00	10.24	9.32	5.62	31.04
	(0.71)	(4.02)	(6.01)	(6.72)	(8.19)	(7.87)	(6.16)	(5.80)	(5.61)	(3.28)	(3.13)	(2.47)	(5.62)
Chhoti safed Anju	0.00	10.32	28.32	37.32	57.32	52.39	27.10	22.18	20.05	6.81	5.89	3.76	23.98
	(0.71)	(3.29)	(5.37)	(6.15)	(7.60)	(7.27)	(5.25)	(4.76)	(4.53)	(2.70)	(2.53)	(2.06)	(4.95)
BatanaMoolchand	0.00	10.28	28.28	37.28	57.28	52.35	27.06	23.77	21.64	8.40	7.48	5.35	24.62
	(0.71)	(3.28)	(5.36)	(6.15)	(7.60)	(7.27)	(5.25)	(4.93)	(4.71)	(2.98)	(2.82)	(2.42)	(5.01)
Batripatiram	0.00	8.75	26.75	35.75	54.67	49.71	24.45	19.16	17.03	3.56	2.69	0.53	21.48
	(0.71)	(3.04)	(5.22)	(6.02)	(7.43)	(7.09)	(4.99)	(4.43)	(4.19)	(2.01)	(1.79)	(1.01)	(4.69)
Rachna	0.00	10.62	28.69	37.69	57.69	52.76	27.47	24.18	23.50	8.26	7.34	4.29	24.93
	(0.71)	(3.33)	(5.40)	(6.18)	(7.63)	(7.30)	(5.29)	(4.97)	(4.90)	(2.96)	(2.80)	(2.19)	(5.04)
Shikha	0.00	8.65	26.71	35.71	54.71	49.72	24.15	18.86	16.73	3.49	2.69	0.56	21.39
	(0.71)	(3.02)	(5.22)	(6.02)	(7.43)	(7.09)	(4.96)	(4.40)	(4.15)	(2.00)	(1.79)	(1.03)	(4.68)
KPMR-420	0.00	9.10	27.15	36.15	55.15	48.65	23.93	18.64	16.51	3.27	2.35	0.22	21.35

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
	(0.71)	(3.10)	(5.26)	(6.05)	(7.46)	(7.01)	(4.94)	(4.37)	(4.12)	(1.94)	(1.69)	(0.85)	(4.67)
KPMR-402	0.00	10.35	28.14	37.14	57.14	52.21	26.92	26.63	24.50	9.14	8.22	5.68	25.20
	(0.71)	(3.29)	(5.35)	(6.14)	(7.59)	(7.26)	(5.24)	(5.21)	(5.00)	(3.10)	(2.95)	(2.49)	(5.07)
KPMR-327	0.00	10.39	28.39	37.40	57.40	52.47	27.18	25.84	23.88	8.69	7.77	5.92	25.14
	(0.71)	(3.30)	(5.37)	(6.16)	(7.61)	(7.28)	(5.26)	(5.13)	(4.94)	(3.03)	(2.88)	(2.53)	(5.06)
KPMR-302	0.00	11.00	29.00	38.05	58.05	53.12	27.83	23.54	22.65	9.41	8.49	5.95	25.34
	(0.71)	(3.39)	(5.43)	(6.21)	(7.65)	(7.32)	(5.32)	(4.90)	(4.81)	(3.15)	(3.00)	(2.54)	(5.08)
KPMR-485	0.00	8.35	26.33	35.15	54.16	48.22	23.99	18.70	16.57	3.33	2.42	0.29	20.99
	(0.71)	(2.97)	(5.18)	(5.97)	(7.39)	(6.98)	(4.95)	(4.38)	(4.13)	(1.96)	(1.71)	(0.89)	(4.64)
Kali Batri	0.00	10.21	29.35	38.35	58.35	53.42	28.13	24.82	24.62	11.38	9.62	6.10	25.89
	(0.71)	(3.27)	(5.46)	(6.23)	(7.67)	(7.34)	(5.35)	(5.03)	(5.01)	(3.45)	(3.18)	(2.57)	(5.14)
Safed Batra Gudda	0.00	15.75	34.04	43.14	66.35	61.34	36.13	30.84	28.71	11.62	9.71	5.98	30.45
	(0.71)	(4.03)	(5.88)	(6.61)	(8.18)	(7.86)	(6.05)	(5.60)	(5.40)	(3.48)	(3.20)	(2.55)	(5.56)
Dhan Batri	0.00	9.68	29.66	38.69	58.69	53.76	28.47	23.39	21.26	8.02	7.10	4.97	24.86
	(0.71)	(3.19)	(5.49)	(6.26)	(7.69)	(7.37)	(5.38)	(4.89)	(4.66)	(2.92)	(2.76)	(2.34)	(5.04)
Jayanti	0.00	8.26	26.26	35.29	54.29	48.35	24.17	18.88	16.75	4.01	3.09	0.96	21.21
	(0.71)	(2.96)	(5.17)	(5.98)	(7.40)	(6.99)	(4.97)	(4.40)	(4.15)	(2.12)	(1.89)	(1.21)	(4.66)
VL-1	0.00	8.55	26.51	35.26	55.16	49.22	24.81	19.50	17.37	4.13	3.21	1.08	21.61
	(0.71)	(3.01)	(5.20)	(5.98)	(7.46)	(7.05)	(5.03)	(4.47)	(4.23)	(2.15)	(1.93)	(1.26)	(4.70)
KPMR-504	0.00	10.35	29.35	38.35	58.35	53.42	28.23	23.94	21.81	8.57	7.65	5.52	25.16
	(0.71)	(3.29)	(5.46)	(6.23)	(7.67)	(7.34)	(5.36)	(4.94)	(4.72)	(3.01)	(2.85)	(2.45)	(5.07)

Genotypes	Population of aphid/ 10 cm apical twigs/plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS**	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
KPMR-400	0.00	11.00	30.01	39.01	60.01	55.08	29.91	25.00	22.87	9.63	8.71	5.58	26.15
	(0.71)	(3.39)	(5.52)	(6.29)	(7.78)	(7.46)	(5.51)	(5.05)	(4.83)	(3.18)	(3.03)	(2.47)	(5.16)
KPMR-486	0.00	11.35	30.31	39.31	60.31	55.38	30.19	25.15	23.02	9.78	8.86	5.73	26.40
	(0.71)	(3.44)	(5.55)	(6.31)	(7.80)	(7.48)	(5.54)	(5.06)	(4.85)	(3.21)	(3.06)	(2.50)	(5.19)
HFP-94-13	0.00	11.36	32.25	41.25	61.25	56.32	31.06	25.78	23.66	9.42	8.52	5.39	26.97
	(0.71)	(3.44)	(5.72)	(6.46)	(7.86)	(7.54)	(5.62)	(5.13)	(4.92)	(3.15)	(3.00)	(2.43)	(5.24)
HFP-94-12	0.00	11.71	29.69	38.69	58.69	53.76	28.52	23.29	21.00	7.76	6.89	3.76	25.12
	(0.71)	(3.49)	(5.49)	(6.26)	(7.69)	(7.37)	(5.39)	(4.88)	(4.64)	(2.87)	(2.72)	(2.06)	(5.06)
JP-885 (Local check)	0.00	15.65	33.63	42.66	62.58	57.66	32.39	27.15	25.02	9.89	8.74	5.61	28.55
	(0.71)	(4.02)	(5.84)	(6.57)	(7.94)	(7.63)	(5.73)	(5.26)	(5.05)	(3.22)	(3.04)	(2.47)	(5.39)
IPF-99-25 (Local check)	0.00	16.35	34.35	43.39	63.64	58.65	33.42	28.19	26.16	9.92	8.95	5.48	29.24
	(0.71)	(4.10)	(5.90)	(6.62)	(8.01)	(7.69)	(5.82)	(5.36)	(5.16)	(3.23)	(3.07)	(2.45)	(5.45)
SEm_±		0.14	0.27	0.34	0.27	0.25	0.31	0.24	0.22	0.13	0.11	0.09	0.28
CD (p=0.05)		(0.40)	(0.76)	(0.94)	(0.76)	(0.71)	(0.86)	(0.66)	(0.60)	(0.38)	(0.32)	(0.25)	(0.79)

* Mean of three replications

** Peak population of aphid

Figures in the parentheses are $\sqrt{X + 0.5}$ values

4.2.2 Leaf hopper

The first observation was taken on 24th December. The mean leaf hopper population ranged from 0.22 (KPMR-420) to 2.62 (Demo-JP-180) leaf hopper/six leaves/ plant. The minimum infestation was observed on genotype, KPMR-420 followed by KPMR-485, VL-1, B-22, DDR-39, Batripatiram, IPF-99-25, Jayanti, Shikha, P-3, VL-3, and these were statistically at par with each other. The maximum infestation was observed on genotype Demo-JP-180 followed by Kashi samriddhi, Double branching, Aman 1-206, Matar rangpur, NDVP-20, Safed Batra Gudda and KPMR-402 were found at par with each other.

Based on overall mean population of the season on different genotypes of pea, it was minimum on genotypes KPMR-485 (1.73 leaf hopper/six leaves) followed by VL-3, KPMR-420, VL-1, Shikha, Jayanti, B-22, Batripatiram, DDR-39, JFP-99-25 and P-3, these were found at par with each other. The maximum population of leaf hopper was recorded in the genotypes Kashi samriddhi (4.13 leaf hopper/six leaves) followed by Demo-JP-180, Double branching, Matar Rangpur, Triple branching, Aman 1-206, NDVP-20, Safed Batra Gudda and KPMR-402, which were found at par with each other. The average population data of leafhoppers revealed that all genotypes differed from one another in terms of population leafhopper/six leaves/plant. Among the screened genotypes the highest population of leaf hopper was recorded in genotype Kashi samriddhi followed by JP-180, matar rangpur while minimum was recorded in genotype KPMR-485 followed by VL-3, VL-1. The genotype KPMR-485 was superior amongst all in respect to lowest leafhopper population.

Table 2.1: Categorization of pea genotypes into degree of susceptibility against leafhopper

S. No.	Mean leaf hopper population/ six leaves	Name of genotypes	Category
1	2.97-1.77	KPMR-485, VL-3, KPMR-420, VL-1, Shikha, jayanti, B-22, BatriPatiram, DDR-39JFP-99-25 and P-3	Least susceptible
2	1.77-3.28	PP-14, KPMR-30, Kali Batri, Atrumatar, SPS-2, PP-86, JFP-27, RP-3, Gol Batra Tenduna, NDVP-4, Kala matar, DDP 94-14, DDR-44, PP-96, DDR-54, KFP-151, BatanaMoolchand, HFP-94-13, JPF-99-25, Kashmiri samriddhi, DDR-43, NDVP-20, Dhan Batri, KPMR-503, HUVP-12, DDR-23, Lat sown, IPF-99-25 (local check), KPMR-504, DhanoliBatri, KPMR-302, Rachna, JP-885 (Local check), KPMR-400, KPMR-327, DDR-27, KPMR-402, HUVP-2, PP-96, KPMR-486, Chhoti safed Anju, LEP-260, HFP-94-12, DDR-52, PP-155, JM-91-01 and HUP-2	Moderately susceptible
3	>3.28	Kashi samriddhi, Demo-JP-180, Double branching, Matar Rangpur, Triple branching, Aman 1-206, NDVP-20, Safed Batra Gudda and KPMR-402	Highly susceptible

Table .2: Screening of different genotypes of pea against leafhopper of pea

Genotypes	Population of leaf hopper/ six leaves/ plant												Mean
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	
IPF-99-25	0.00	1.39	4.39	4.52	5.61	5.3	3.99	2.66	2.01	1.35	1.05	0.00	2.97
	(0.71)	(1.37)	(2.21)	(2.24)	(2.47)	(2.41)	(2.12)	(1.78)	(1.58)	(1.36)	(1.24)	(0.71)	(1.86)
B-22	0.00	0.36	3.09	3.18	4.27	3.96	2.45	1.15	0.55	0.42	0.1	0.00	1.8
	(0.71)	(0.93)	(1.89)	(1.92)	(2.18)	(2.11)	(1.72)	(1.28)	(1.02)	(0.96)	(0.77)	(0.71)	(1.52)
DDR-27	0.00	1.69	4.71	4.88	5.97	5.66	4.14	2.84	2.26	1.75	1.32	0.00	3.25
	(0.71)	(1.48)	(2.28)	(2.32)	(2.54)	(2.48)	(2.15)	(1.83)	(1.66)	(1.50)	(1.35)	(0.71)	(1.94)
PP-155	0.00	1.92	4.94	5.01	6.1	5.79	4.32	3.04	2.31	1.82	1.1	0.00	3.36
	(0.71)	(1.56)	(2.33)	(2.35)	(2.57)	(2.51)	(2.20)	(1.88)	(1.68)	(1.52)	(1.26)	(0.71)	(1.96)
Kashmiri samriddhi	0.00	2.21	4.23	4.33	5.42	5.11	3.82	2.48	1.86	1.98	1.15	0.00	2.99
	(0.71)	(1.65)	(2.17)	(2.20)	(2.43)	(2.37)	(2.08)	(1.73)	(1.54)	(1.57)	(1.28)	(0.71)	(1.87)
DDR-43	0.00	1.49	4.43	4.51	5.6	5.29	3.99	2.69	2.06	1.49	1.16	0.00	3.01
	(0.71)	(1.41)	(2.22)	(2.24)	(2.47)	(2.41)	(2.12)	(1.79)	(1.60)	(1.41)	(1.29)	(0.71)	(1.87)
DDR-44	0.00	1.33	4.31	4.38	5.47	5.16	3.86	2.52	1.88	1.33	1	0.00	2.88
	(0.71)	(1.35)	(2.19)	(2.21)	(2.44)	(2.38)	(2.09)	(1.74)	(1.54)	(1.35)	(1.22)	(0.71)	(1.84)
DDR-39	0.00	0.36	3.08	3.2	4.29	3.98	2.68	1.39	0.76	0.44	0.11	0.00	1.87
	(0.71)	(0.93)	(1.89)	(1.92)	(2.19)	(2.12)	(1.78)	(1.37)	(1.12)	(0.97)	(0.78)	(0.71)	(1.54)
Late sown	0.00	2.26	4.28	4.44	5.53	5.22	3.92	2.66	2.03	1.95	1.21	0.00	3.07
	(0.71)	(1.66)	(2.19)	(2.22)	(2.46)	(2.39)	(2.10)	(1.78)	(1.59)	(1.57)	(1.31)	(0.71)	(1.89)

Genotypes	Population of leaf hopper/ six leaves/ plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
DDR-23	0.00	2.21	4.23	4.49	5.58	5.27	3.95	2.62	1.99	1.88	1.25	0.00	3.06
	(0.71)	(1.65)	(2.17)	(2.23)	(2.47)	(2.4)	(2.11)	(1.77)	(1.58)	(1.54)	(1.32)	(0.71)	(1.89)
HUP-2	0.00	1.95	4.98	5.11	6.2	5.89	4.57	3.26	2.63	1.95	1.29	0.00	3.49
	(0.71)	(1.57)	(2.34)	(2.37)	(2.59)	(2.53)	(2.25)	(1.94)	(1.77)	(1.57)	(1.34)	(0.71)	(2.00)
DDP94-14	0.00	1.98	4	4.1	5.19	4.88	3.56	2.25	1.62	1.98	1.33	0.00	2.83
	(0.71)	(1.57)	(2.12)	(2.14)	(2.39)	(2.32)	(2.01)	(1.66)	(1.46)	(1.57)	(1.35)	(0.71)	(1.82)
HUVP-2	0.00	1.68	4.7	4.82	5.91	5.6	4.28	2.97	2.34	1.68	1.35	0.00	3.25
	(0.71)	(1.48)	(2.28)	(2.31)	(2.53)	(2.47)	(2.19)	(1.86)	(1.69)	(1.48)	(1.36)	(0.71)	(1.94)
KPMR-402	0.00	2.26	5.34	5.45	6.54	6.23	4.91	3.6	2.97	2.4	1.95	0.00	3.83
	(0.71)	(1.66)	(2.42)	(2.44)	(2.65)	(2.59)	(2.33)	(2.02)	(1.86)	(1.70)	(1.57)	(0.71)	(2.08)
NDVP-20	0.00	2.45	5.49	5.6	6.69	6.38	5.06	3.75	3.12	2.49	1.88	0.00	3.95
	(0.71)	(1.72)	(2.45)	(2.47)	(2.68)	(2.62)	(2.36)	(2.06)	(1.90)	(1.73)	(1.54)	(0.71)	(2.11)
Atrumatar	0.00	1.54	3.99	4.1	5.19	4.88	3.5	2.19	1.56	1.54	1.21	0.00	2.73
	(0.71)	(1.43)	(2.12)	(2.14)	(2.39)	(2.32)	(2.00)	(1.64)	(1.44)	(1.43)	(1.31)	(0.71)	(1.80)
Triple branching	0.00	2.51	5.53	5.64	6.73	6.42	5.1	3.79	3.16	2.51	1.98	0.00	3.99
	(0.71)	(1.73)	(2.46)	(2.48)	(2.69)	(2.63)	(2.37)	(2.07)	(1.91)	(1.73)	(1.57)	(0.71)	(2.12)
Double Branching	0.00	2.55	5.57	5.68	6.77	6.46	5.14	3.83	3.2	2.55	2	0.00	4.03
	(0.71)	(1.75)	(2.46)	(2.49)	(2.70)	(2.64)	(2.37)	(2.08)	(1.92)	(1.75)	(1.58)	(0.71)	(2.13)
SPS-2	0.00	1.11	4.13	4.24	5.33	5.02	3.7	2.39	1.76	1.11	0.78	0.00	2.73
	(0.71)	(1.27)	(2.15)	(2.18)	(2.41)	(2.35)	(2.05)	(1.70)	(1.50)	(1.27)	(1.13)	(0.71)	(1.80)
JM-91-01	0.00	1.82	4.84	4.95	6.04	5.73	4.42	3.11	2.48	1.82	1.39	0.00	3.37
	(0.71)	(1.52)	(2.31)	(2.33)	(2.56)	(2.50)	(2.22)	(1.90)	(1.73)	(1.52)	(1.37)	(0.71)	(1.97)
P-3	0.00	0.62	3.15	3.21	4.3	3.99	2.67	1.32	0.62	0.62	0.18	0.00	1.9
	(0.71)	(1.06)	(1.91)	(1.93)	(2.19)	(2.12)	(1.78)	(1.35)	(1.06)	(1.06)	(0.82)	(0.71)	(1.55)

Genotypes	Population of leaf hopper/ six leaves/ plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
RP-3	0.00	1.12	4.14	4.25	5.34	5.03	3.71	2.4	1.77	1.12	0.79	0.00	2.74
	(0.71)	(1.27)	(2.15)	(2.18)	(2.42)	(2.35)	(2.05)	(1.70)	(1.51)	(1.27)	(1.14)	(0.71)	(1.80)
VL-3	0.00	0.62	2.94	2.95	4.04	3.73	2.44	1.19	0.56	0.62	0.15	0.00	1.77
	(0.71)	(1.06)	(1.85)	(1.86)	(2.13)	(2.06)	(1.71)	(1.30)	(1.03)	(1.06)	(0.81)	(0.71)	(1.51)
JFP-27	0.00	1.84	3.86	3.97	5.06	4.75	3.43	2.12	1.52	1.84	1.51	0.00	2.73
	(0.71)	(1.53)	(2.09)	(2.11)	(2.36)	(2.29)	(1.98)	(1.62)	(1.42)	(1.53)	(1.42)	(0.71)	(1.80)
PP-86	0.00	1.11	4.13	4.24	5.33	5.02	3.70	2.39	1.77	1.11	0.78	0.00	2.73
	(0.71)	(1.27)	(2.15)	(2.18)	(2.41)	(2.35)	(2.05)	(1.70)	(1.51)	(1.27)	(1.13)	(0.71)	(1.80)
NDVP-4	0.00	1.16	4.18	4.29	5.38	5.07	3.75	2.44	1.79	1.16	0.83	0.00	2.77
	(0.71)	(1.29)	(2.16)	(2.19)	(2.42)	(2.36)	(2.06)	(1.71)	(1.51)	(1.29)	(1.15)	(0.71)	(1.81)
NDVP-20	0.00	1.85	4.34	4.45	5.54	5.23	3.91	2.6	1.98	1.85	1.12	0.00	3.02
	(0.71)	(1.53)	(2.20)	(2.22)	(2.46)	(2.39)	(2.10)	(1.76)	(1.57)	(1.53)	(1.27)	(0.71)	(1.88)
KFP-151	0.00	1.3	4.32	4.43	5.52	5.21	3.89	2.58	1.95	1.3	0.97	0.00	2.9
	(0.71)	(1.34)	(2.20)	(2.22)	(2.45)	(2.39)	(2.10)	(1.75)	(1.57)	(1.34)	(1.21)	(0.71)	(1.84)
HUVP-12	0.00	1.44	4.46	4.57	5.66	5.35	4.11	2.8	2.17	1.44	1.11	0.00	3.05
	(0.71)	(1.39)	(2.23)	(2.25)	(2.48)	(2.42)	(2.15)	(1.82)	(1.63)	(1.39)	(1.27)	(0.71)	(1.88)
LEP-260	0.00	1.74	4.76	4.87	5.96	5.65	4.36	3.05	2.42	1.76	1.13	0.00	3.29
	(0.71)	(1.50)	(2.29)	(2.32)	(2.54)	(2.48)	(2.20)	(1.88)	(1.71)	(1.50)	(1.28)	(0.71)	(1.95)
DhanoliBatri	0.00	1.53	4.55	4.66	5.75	5.44	4.12	2.81	2.18	1.55	1.15	0.00	3.11
	(0.71)	(1.42)	(2.25)	(2.27)	(2.50)	(2.44)	(2.15)	(1.82)	(1.64)	(1.43)	(1.28)	(0.71)	(1.90)
Gol Batra Tenduna	0.00	1.16	4.18	4.29	5.38	5.07	3.75	2.42	1.79	1.16	0.83	0.00	2.77
	(0.71)	(1.29)	(2.16)	(2.19)	(2.42)	(2.36)	(2.06)	(1.71)	(1.51)	(1.29)	(1.15)	(0.71)	(1.81)
Matar Rangpur	0.00	2.46	5.51	5.66	6.75	6.44	5.12	3.81	3.18	2.46	2.01	0.00	3.99
	(0.71)	(1.72)	(2.45)	(2.48)	(2.69)	(2.63)	(2.37)	(2.08)	(1.92)	(1.72)	(1.58)	(0.71)	(2.12)

Genotypes	Population of leaf hopper/ six leaves/ plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
Kashi samriddhi	0.00	2.62	5.68	5.81	6.9	6.59	5.31	4.00	3.37	2.62	1.99	0.00	4.13
	(0.71)	(1.77)	(2.49)	(2.51)	(2.72)	(2.66)	(2.41)	(2.12)	(1.97)	(1.77)	(1.58)	(0.71)	(2.15)
Kala matar	0.00	1.32	4.34	4.45	5.54	5.23	3.42	2.11	1.48	1.32	0.99	0.00	2.8
	(0.71)	(1.35)	(2.20)	(2.22)	(2.46)	(2.39)	(1.98)	(1.62)	(1.41)	(1.35)	(1.22)	(0.71)	(1.82)
KPMR-503	0.00	1.46	4.48	4.59	5.68	5.37	4.05	2.74	2.11	1.46	1.13	0.00	3.05
	(0.71)	(1.40)	(2.23)	(2.26)	(2.49)	(2.42)	(2.13)	(1.80)	(1.62)	(1.40)	(1.28)	(0.71)	(1.88)
DDR-52	0.00	1.76	4.78	4.89	5.98	5.67	4.35	3.06	2.31	1.76	1.33	0.00	3.31
	(0.71)	(1.50)	(2.30)	(2.32)	(2.55)	(2.48)	(2.20)	(1.89)	(1.68)	(1.50)	(1.35)	(0.71)	(1.95)
DDR-54	0.00	1.31	4.33	4.44	5.53	5.22	3.90	2.59	1.96	1.23	0.90	0.00	2.90
	(0.71)	(1.35)	(2.20)	(2.22)	(2.46)	(2.39)	(2.10)	(1.76)	(1.57)	(1.32)	(1.18)	(0.71)	(1.84)
PP-96	0.00	1.81	4.83	4.94	6.03	5.72	4.14	2.83	2.18	1.62	1.29	0.00	3.27
	(0.71)	(1.52)	(2.31)	(2.33)	(2.56)	(2.49)	(2.15)	(1.82)	(1.64)	(1.46)	(1.34)	(0.71)	(1.94)
KPMR-30	0.00	1.1	4.12	4.23	5.32	5.00	3.68	2.37	1.74	1.10	0.77	0.00	2.71
	(0.71)	(1.26)	(2.15)	(2.17)	(2.41)	(2.35)	(2.04)	(1.69)	(1.50)	(1.26)	(1.13)	(0.71)	(1.79)
JFP-99-25	0.00	0.44	3.16	3.22	4.31	3.99	2.67	1.39	0.76	0.44	0.11	0.00	1.89
	(0.71)	(0.97)	(1.91)	(1.93)	(2.19)	(2.12)	(1.78)	(1.37)	(1.12)	(0.97)	(0.78)	(0.71)	(1.55)
PP-96	0.00	1.29	4.31	4.42	5.51	5.19	3.87	2.56	1.93	1.29	0.96	0.00	2.89
	(0.71)	(1.34)	(2.19)	(2.22)	(2.45)	(2.39)	(2.09)	(1.75)	(1.56)	(1.34)	(1.21)	(0.71)	(1.84)
PP-14	0.00	1.02	4.04	4.15	5.24	4.92	3.60	2.29	1.66	1.02	0.69	0.00	2.64
	(0.71)	(1.23)	(2.13)	(2.16)	(2.40)	(2.33)	(2.02)	(1.67)	(1.47)	(1.23)	(1.09)	(0.71)	(1.77)
Aman1-206	0.00	2.47	5.52	5.63	6.72	6.40	5.08	3.79	3.19	2.31	1.98	0.00	3.97
	(0.71)	(1.72)	(2.45)	(2.48)	(2.69)	(2.63)	(2.36)	(2.07)	(1.92)	(1.68)	(1.57)	(0.71)	(2.11)
Demo-JP-180	0.00	2.62	5.68	5.79	6.88	6.56	5.24	3.96	3.33	2.12	1.79	0.00	4.06
	(0.71)	(1.77)	(2.49)	(2.51)	(2.72)	(2.66)	(2.40)	(2.11)	(1.96)	(1.62)	(1.51)	(0.71)	(2.13)

Genotypes	Population of leaf hopper/ six leaves/ plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
Chhoti safed Anju	0.00	1.76	4.78	4.89	5.98	5.66	4.34	3.03	2.4	1.55	1.22	0.00	3.28
	(0.71)	(1.50)	(2.30)	(2.32)	(2.55)	(2.48)	(2.20)	(1.88)	(1.70)	(1.43)	(1.31)	(0.71)	(1.95)
BatanaMoolchand	0.00	1.35	4.37	4.48	5.57	5.25	3.93	2.62	1.99	1.35	1.02	0.00	2.94
	(0.71)	(1.36)	(2.21)	(2.23)	(2.46)	(2.40)	(2.10)	(1.77)	(1.58)	(1.36)	(1.23)	(0.71)	(1.86)
Batripatiram	0.00	0.42	3.2	3.26	4.35	4.03	2.34	1.06	0.43	0.42	0.09	0.00	1.82
	(0.71)	(0.96)	(1.92)	(1.94)	(2.20)	(2.13)	(1.69)	(1.25)	(0.96)	(0.96)	(0.77)	(0.71)	(1.52)
Rachna	0.00	1.58	4.6	4.71	5.80	5.48	4.16	2.85	2.22	1.58	1.25	0.00	3.15
	(0.71)	(1.44)	(2.26)	(2.28)	(2.51)	(2.45)	(2.16)	(1.83)	(1.65)	(1.44)	(1.32)	(0.71)	(1.91)
Shikha	0.00	0.56	2.94	3.11	4.20	3.88	2.36	1.08	0.45	0.56	0.23	0.00	1.78
	(0.71)	(1.03)	(1.85)	(1.90)	(2.17)	(2.09)	(1.69)	(1.26)	(0.97)	(1.03)	(0.85)	(0.71)	(1.51)
KPMR-420	0.00	0.22	2.97	3.1	4.19	3.87	2.51	1.26	0.63	0.32	0.14	0.00	1.77
	(0.71)	(0.85)	(1.86)	(1.90)	(2.17)	(2.09)	(1.73)	(1.33)	(1.06)	(0.91)	(0.80)	(0.71)	(1.51)
KPMR-402	0.00	1.68	4.7	4.81	5.9	5.58	4.26	2.95	2.32	1.72	1.39	0.00	3.25
	(0.71)	(1.48)	(2.28)	(2.30)	(2.53)	(2.47)	(2.18)	(1.86)	(1.68)	(1.49)	(1.37)	(0.71)	(1.94)
KPMR-327	0.00	1.62	4.64	4.75	5.84	5.52	4.2	2.89	2.26	1.62	1.29	0.00	3.19
	(0.71)	(1.46)	(2.27)	(2.29)	(2.52)	(2.45)	(2.17)	(1.84)	(1.66)	(1.46)	(1.34)	(0.71)	(1.92)
KPMR-302	0.00	1.54	4.56	4.67	5.76	5.44	4.12	2.81	2.18	1.54	1.21	0.00	3.12
	(0.71)	(1.43)	(2.25)	(2.27)	(2.50)	(2.44)	(2.15)	(1.82)	(1.64)	(1.43)	(1.31)	(0.71)	(1.90)
KPMR-485	0.00	0.29	3.01	3.01	4.1	3.78	2.42	1.14	0.52	0.29	0.15	0.00	1.73
	(0.71)	(0.89)	(1.87)	(1.87)	(2.14)	(2.07)	(1.71)	(1.28)	(1.01)	(0.89)	(0.81)	(0.71)	(1.49)
Kali Batri	0.00	1.10	4.12	4.23	5.32	5.00	3.69	2.38	1.77	1.1	0.77	0.00	2.72
	(0.71)	(1.26)	(2.15)	(2.17)	(2.41)	(2.35)	(2.05)	(1.70)	(1.51)	(1.26)	(1.13)	(0.71)	(1.79)
Safed Batra Gudda	0.00	2.39	5.41	5.52	6.61	6.29	4.97	3.66	3.06	2.39	2.06	0.00	3.9
	(0.71)	(1.70)	(2.43)	(2.45)	(2.67)	(2.61)	(2.34)	(2.04)	(1.89)	(1.70)	(1.60)	(0.71)	(2.10)

Genotypes	Population of leaf hopper/ six leaves/ plant												
	30 DAS	38 DAS	47 DAS	54 DAS	61 DAS* *	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
Dhan Batri	0.00	1.46	4.48	4.59	5.68	5.36	4.04	2.73	2.12	1.46	1.13	0.00	3.05
	(0.71)	(1.40)	(2.23)	(2.26)	(2.49)	(2.42)	(2.13)	(1.80)	(1.62)	(1.40)	(1.28)	(0.71)	(1.88)
Jayanti	0.00	0.46	3.18	3.18	4.27	3.95	2.13	1.14	0.51	0.46	0.13	0.00	1.8
	(0.71)	(0.98)	(1.92)	(1.92)	(2.18)	(2.11)	(1.62)	(1.28)	(1.00)	(0.98)	(0.79)	(0.71)	(1.52)
VL-1	0.00	0.32	3.12	3.22	4.31	3.99	2.18	1.07	0.44	0.32	0.10	0.00	1.77
	(0.71)	(0.91)	(1.90)	(1.93)	(2.19)	(2.12)	(1.64)	(1.25)	(0.97)	(0.91)	(0.77)	(0.71)	(1.51)
KPMR-504	0.00	1.52	4.54	4.65	5.74	5.42	4.10	2.79	2.16	1.52	1.19	0.00	3.1
	(0.71)	(1.42)	(2.24)	(2.27)	(2.50)	(2.43)	(2.14)	(1.81)	(1.63)	(1.42)	(1.30)	(0.71)	(1.90)
KPMR-400	0.00	1.58	4.64	4.75	5.84	5.52	4.20	2.89	2.26	1.58	1.25	0.00	3.18
	(0.71)	(1.44)	(2.27)	(2.29)	(2.52)	(2.45)	(2.17)	(1.84)	(1.66)	(1.44)	(1.32)	(0.71)	(1.92)
KPMR-486	0.00	1.73	4.76	4.82	5.91	5.59	4.27	2.96	2.33	1.73	1.38	0.00	3.27
	(0.71)	(1.49)	(2.29)	(2.31)	(2.53)	(2.47)	(2.18)	(1.86)	(1.68)	(1.49)	(1.37)	(0.71)	(1.94)
HFP-94-13	0.00	1.39	4.41	4.48	5.57	5.25	3.94	2.65	2.02	1.39	1.06	0.00	2.97
	(0.71)	(1.37)	(2.22)	(2.23)	(2.46)	(2.40)	(2.11)	(1.77)	(1.59)	(1.37)	(1.25)	(0.71)	(1.86)
HFP-94-12	0.00	1.75	4.77	4.86	5.95	5.63	4.31	3.01	2.38	1.75	1.42	0.00	3.30
	(0.71)	(1.50)	(2.30)	(2.32)	(2.54)	(2.48)	(2.19)	(1.87)	(1.70)	(1.50)	(1.39)	(0.71)	(1.95)
JP-885(Local check)	0.00	1.54	4.58	4.70	5.79	5.47	4.21	2.92	2.36	1.54	1.21	0.00	3.16
	(0.71)	(1.43)	(2.25)	(2.28)	(2.51)	(2.44)	(2.17)	(1.85)	(1.69)	(1.43)	(1.31)	(0.71)	(1.91)
IPF-99-25 (Local check)	0.00	1.48	4.54	4.68	5.77	5.45	4.18	2.84	2.33	1.32	0.99	0.00	3.09
	(0.71)	(1.41)	(2.24)	(2.28)	(2.50)	(2.44)	(2.16)	(1.83)	(1.68)	(1.35)	(1.22)	(0.71)	(1.90)
SEm+		0.05	0.11	0.11	0.14	0.13	0.09	0.08	0.08	0.05	0.04	0.00	0.07
CD (p=0.05)		0.145	0.304	0.318	0.379	0.351	0.25	0.234	0.227	0.13	0.122	0.00	0.202

** Peak population of leafhopper,
 Figures in the parentheses are $\sqrt{X + 0.5}$ values

4.2.3 Pod borer, *Helicoverpa armigera*

The first observation recorded on 17th December, 2021. The mean *Helicoverpa* population ranged from 0.31 (KPMR-485) to 2.55 (Double branching) / plant. The minimum infestation was observed on genotype KPMR-485 followed by KPMR-420, VL-1, Batripatiram, JFP-99-25, Jayanti, B-22, DDR-39, P-3, Shikha, and VL-3 which was statistically at par with each other. The maximum infestation was observed on genotype Double branching followed by Triple branching, NDVP-20, KPMR-402, Matar Rangpur, Safed Batra Gudda, Aman 1-206, Kashi samriddhi and Demo-JP-180. These were statistically at par with each other in degree of infestation.

Based on overall mean population of the season on different genotypes of pea, it was minimum on genotypes Batripatiram (0.14 *Helicoverpa* /plant) followed by KPMR-485, Jayanti, KPMR-420, JFP-99-25, B-22, DDR-39, Shikha, P-3, VL-3, these were found at par with each other. The maximum population of *helicoverpa* was recorded in the genotypes Matar Rangpur (1.21 *Helicoverpa*/ plant) followed by Double branching, Triple branching, Aman 1-206, Kashi samriddhi, Safed Batra Gudda, Demo-JP-180, NDVP-20 and KPMR-402, which were found at par with each other.

The average population data of pod borer revealed that all genotypes differed from one another in terms of population of aphid/10cm apical twigs/plant. Among the screened genotypes the highest population of pod borer was recorded in genotype Matar rangpur followed by double branching, Aman1-206 while minimum was recorded in genotype Batripatiram followed by VL-3, VL-3.

Genotypes	Population of pod borer, <i>helicoverpaarmigera</i>												
	30 DAS	38 DAS	47** DAS	54 DAS	61 DAS	68 DAS	75 DAS	81 DAS	89 DAS	95 DAS	103 DAS	107 DAS	Mean
	(1.42)	(1.31)	(1.45)	(1.32)	(1.28)	(1.06)	(0.91)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.12)
KPMR-400	1.58	1.27	1.66	1.29	1.21	0.68	0.38	0.00	0.00	0.00	0.00	0.00	0.79
	(1.44)	(1.33)	(1.47)	(1.34)	(1.31)	(1.09)	(0.94)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.13)
KPMR-486	1.73	1.12	1.52	1.14	1.06	0.66	0.36	0.00	0.00	0.00	0.00	0.00	0.73
	(1.49)	(1.27)	(1.42)	(1.28)	(1.25)	(1.08)	(0.93)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.11)
HFP-94-13	1.39	1.08	1.48	1.10	1.02	0.71	0.33	0.00	0.00	0.00	0.00	0.00	0.69
	(1.37)	(1.26)	(1.41)	(1.26)	(1.23)	(1.10)	(0.91)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.09)
HFP-94-12	1.77	1.19	1.59	1.15	1.07	0.72	0.40	0.00	0.00	0.00	0.00	0.00	0.77
	(1.51)	(1.30)	(1.45)	(1.28)	(1.25)	(1.10)	(0.95)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.12)
JP-885 (Local check)	1.53	1.22	1.60	1.16	1.00	0.68	0.39	0.00	0.00	0.00	0.00	0.00	0.74
	(1.42)	(1.31)	(1.45)	(1.29)	(1.22)	(1.09)	(0.94)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.11)
IPF-99-25 (Local check)	1.35	1.00	1.48	0.98	0.85	0.55	0.35	0.00	0.00	0.00	0.00	0.00	0.64
	(1.36)	(1.22)	(1.41)	(1.22)	(1.16)	(1.02)	(0.92)	(0.71)	(0.71)	(0.71)	(0.71)	(0.71)	(1.07)
SEm+	0.05	0.04	0.05	0.03	0.03	0.03	0.02	-	-	-	-	-	0.03
CD (p=0.05)	0.137	0.119	0.132	0.097	0.089	0.073	0.057	-	-	--	-	--	0.088

Table 3.1: Categorization of pea genotypes into degree of susceptibility against pod borer

S. No.	Mean pod borer/ plant	Name of genotypes	Category
1	0.63-0.68	Batripatiram, VL-1, KPMR-485, Jayanti, KPMR-420, JFP-99-25, B-22, DDR-39, Shikha, P-3, VL-3	Least susceptible
2	0.78-1.06	PP-14, KPMR-30, Kali Batri, Atrumatar, SPS-2, PP-86, JFP-27, RP-3, Gol Batra Tenduna, NDVP-4, Kalamatar, DDP 94-14, DDR-44, PP-96, DDR-54, KFP-151, BatanaMoolchand, HFP-94-13, JPF-99-25, Kashmiri samridhi, DDR-43, NDVP-20, Dhan Batri, KPMR-503, HUV-12, DDR-23, Lat sown, IPF-99-25 (local check), KPMR-504, DhanoliBatri, KPMR-302, Rachna, JP-885 (Local check), KPMR-400, KPMR-327, DDR-27, KPMR-402, HUV-2, HUP-2 KPMR-486, Chhoti safed Anju, LEP-260, HFP-94-12, DDR-52, PP-155, JM-91-01 and PP-96	Moderately susceptible
3	>1.06	KPMR-402, NDVP-20, Demo-JP-180, Safed Batra Gudda, Kashi samridhi, Aman 1-206, Triple branching, Double branching and Matar Rangpur	Highly susceptible

Pod damage (%)

The minimum pod damage was recorded in genotypes KPMR-485 (2.24%) followed by genotypes B-22, KPMR-420, VL-3, DDR-39, P-3, IPF-99-25, Batripatiram, Shikha, Jayanti and VL-1, these treatments observed at par each other (except Jayanti and VL-1) as well as best in reducing the percentage of pod damage due to pod borer. The maximum infestation of by-pod borer in genotype Matar Rangpur (8.14%) followed by KPMR-402, Kashi samriddhi, NDVP-20, Aman 1-206, Triple branching, Safed Batra Gudda, Demo-JP-180, Double branching in pod damage, these treatments had remained at par with each other.

Seed yield of pea

The seed yield of different genotypes of field pea were observed in ranged from 1023 kg/ha (Safed Batra Gudda) to 1699 kg/ha (KPMR-485). Based on mean yield of pea on different genotypes of pea, it was maximum seed yield 1699(kg/ha) on genotype (KPMR-485) followed by Batripatiram (1620 kg/ha), Jayanti (1611 kg/ha), IPF-99-25 (1285 kg/ha), Shikha (1599 kg/ha), KPMR-420 (1597 kg/ha), VL-1 (1588 kg/ha), VL-3 (1568 kg/ha), P-3 (1532kg/ha), DDR-39 (1514 kg/ha), and B-22 (1412kg/ha) these were found at par with each other (except KPMR-485). The minimum yield of pea was recorded in the genotypes Safed Batra Gudda (1023kg/ha) followed by Aman 1-206 (1065 kg/ha), Kashi samriddhi (1310 kg/ha), Double branching (1095 kg/ha), Matar Rangpur (1095 kg/ha), Demo-JP-180 (1100 kg/ha), Triple branching (1119 kg/ha), KPMR-402 (1165 kg/ha) and NDVP-20 (1169 kg/ha), which were found at par with each other (except KPMR-402).

Table .4: Yield and pod damage in different genotypes of pea

S. No	Genotypes	Pod damage (%)	Yield of pea (kg/ha)
1	IPF-99-25	6.95(15.29)	1285
2	B-22	2.25(8.63)	1412
3	DDR-27	6.35(14.60)	1291
4	PP-155	6.33(14.57)	1299
5	Kashmiri samriddhi	5.39(13.42)	1301
6	DDR-43	5.21(13.19)	1285
7	DDR-44	5.29(13.30)	1287
8	DDR-39	2.55(9.19)	1514
9	Late sown	5.35(13.37)	1265
10	DDR-23	5.39(13.42)	1271
11	HUP-2	5.10(13.05)	1275
12	HUVP-2	5.32(13.34)	1275
13	KPMR-402	8.15(16.59)	1165
14	NDVP-20	58.35(16.80)	1169
15	Atrumatar	5.95(14.12)	1312
16	Triple branching	8.36(16.81)	1119
17	Double Branching	8.84(17.30)	1095
18	SPS-2	5.25(13.25)	1295
19	NDVP-20	5.24(13.25)	1335
20	KFP-151	5.29(13.30)	1268
21	HUVP-12	5.46(13.51)	1275
23	DhanoliBatri	5.85(14.00)	1278
24	Gol Batra Tenduna	5.68(13.79)	1225
25	Matar Rangpur	8.14(16.58)	1095
26	Kashi samriddhi	8.19(16.39)	1080
27	Kala matar	5.58(13.66)	1267
28	KPMR-503	5.68(13.79)	1278
29	DDR-52	5.66(13.76)	1245
30	DDR-54	5.26(13.26)	1225
31	PP-96	5.45(13.50)	1300
32	KPMR-30	5.55(13.63)	1325
33	IPF-99-25	2.72(9.49)	1610
34	PP-96	5.58(13.66)	1315
35	PP-14	5.95(14.12)	1311
36	Aman1-206	8.35(16.80)	1065
37	Demo-JP-180	8.68(17.13)	1100
38	Chhoti safed Anju	5.65(13.75)	1350

S. No	Genotypes	Pod damage (%)	Yield of pea (kg/ha)
39	BatanaMoolchand	5.69(13.80)	1322
40	Batripatiram	2.75(9.55)	1620
41	Rachna	5.69(13.80)	1325
42	Shikha	2.79(9.62)	1599
43	KPMR-420	2.26(8.65)	1597
44	KPMR-402	5.26(13.26)	1312
45	KPMR-327	5.49(13.55)	1327
46	KPMR-302	5.56(13.64)	1295
47	KPMR-485	2.24(8.61)	1699
48	Kali Batri	5.15(13.12)	1323
49	Safed Batra Gudda	8.64(17.09)	1023
51	Dhan Batri	5.32(13.34)	1351
52	Jayanti	3.56(10.88)	1611
53	VL-1	3.59(10.92)	1588
54	KPMR-504	5.11(13.06)	1314
55	KPMR-400	5.69(13.80)	1295
56	KPMR-486	5.78(13.91)	1313
57	HFP-94-13	5.79(13.92)	1269
58	HFP-94-12	5.69(13.80)	1299
59	JP-885 (Local check)	5.84(13.98)	1310
60	IPF-99-25 (Local check)	5.94(14.11)	1324
61	Batripatiram	2.75(9.55)	1620
62	Rachna	5.69(13.80)	1325
63	Shikha	2.79(9.62)	1599
64	KPMR-420	2.26(8.65)	1597
65	KPMR-402	5.26(13.26)	1312
66	KPMR-327	5.49(13.55)	1327
67	KPMR-302	5.56(13.64)	1295
68	KPMR-485	2.24(8.61)	1699
69	Kali batri	5.15(13.12)	1323
70	Aman1-206	8.35(16.80)	1065
71	Demo-JP-180	8.68(17.13)	1100
		SEm±0.11	74
		CD (p=0.05)	208

* Mean of three replications

Figures in the parentheses are angular transformation value

Conclusion

The field pea crop encountered the presence of four significant insect pests, with aphids and borers emerging at the early stages, and leafhoppers making an appearance in the first week of January. These insect pests played a pivotal role in reducing the overall yield of the field pea, establishing them as significant contributors to the yield decline. In terms of aphid populations, the genotype Double Branching exhibited the highest count, followed by JP-180, Matar Rangpur, and Aman 1-206. Conversely, the minimum aphid population was observed in the B-22 genotype, followed by DDR-39 and KPMR-485. Pod borer populations were most pronounced in the Matar Rangpur genotype, followed by Double Branching and Aman 1-206, while the minimum pod borer population was recorded in the BatriPatiram genotype, followed by VL-3 and VL-3. Leafhopper populations peaked in the Kashi Samriddhi genotype, followed by JP-180 and Matar Rangpur, while the minimum count was observed in the KPMR-485 genotype, followed by VL-3 and VL-1.

The lowest percentage of pod damage was noted in the KPMR-485 genotype, closely followed by B-22. Genotypes with high susceptibility were identified as Double Branching, whereas the least susceptible genotype was found to be B-22. The highest pea yield was achieved by the KPMR-485 genotype, followed by Jayanti and IPF-99-25. Conversely, the lowest yield was recorded in the Safed Batra Gudda genotype, followed by Matar Rangpur and Aman 1-206.

References:

- Chatterjee Puja, Mondal Shanowly and Das Ayan. 2019. Screening of Different Genotypes of okra (*Abelmoschus esculentus* L.) against Leafhopper (*Amrascabi guttulabiguttula* L) and Whitefly (*Bemisia tabaci* G.) under New Gangetic Alluvial Zone of West Bengal., Int. J. Curr. Microbiol. App. Sci (2019), 8(3): 1087-1095.
- Dixon, A.FG. and Harrington, R. 2000. Causes of regional and early variation in pea aphid numbers in Eastern England. Rev. Agril. Ent, 88: 281.
- Haque SR, Akter N, Khan MAH, Kabir K, Islam MM. 2015. Yield potential of garden pea varieties at varied harvesting dates, Bangladesh Agronomy Journal ,17(2):21-28.
- Hooda V S, Dhankhar B S and Singh Ram. 2011. Evaluation of okra cultivars for field resistance to the leafhopper *Amrascabi guttulabiguttula* (Ishida), International Journal of Tropical Insect Science, Volume 17, Issue 3-4, December 199, pp. 323 – 327.
- Honda, K. 1989. Seasonal changes in population density of the pea aphid *Acyrtosiphon pisum* Harris in pea fields and environmental conditions for the emergence of alate virginoparae. Annual Report of the Society of Plant Protection of North Japan, (40): 110-111.

- Krishna H, Singh AK, Kumar P and Kumar S.2019. Germplasm screening against major insect pests in field pea. *Journal of Entomology and Zoology Studies* 2019, 7(3): 270-272.
- Kushwaha, K. 2002. Succession of insect pests of pea. M.Sc. (Ag.) Thesis, submitted to J.N.K.V.V, Jabalpur, M.P.
- Makasheva RK. 1983. The pea. Amerind Publ Co Pvt Ltd, New Delhi, India:267.
- Mandloi Ishwar. 2020.To study the effect of different dates of sowing on the incidence of major pests of niger.M.sc.Thesis, JNKVV, Jabalpur 1-75p.
- Manivannan A, Philip R S, Karthikeyan S and Pillai A K. 2017. Screening of cotton genotypes against leafhopper, *Amrascabiguttulabiguttula* (Ishida) (Homoptera: Cicadellidae), *Journal of Entomology and Zoology Studies* 2017, 5(6): 1305-1310
- Melesse T and Singh SK.2012. Effect of climatic factors on pea aphid, *Acyrtosiphonpisum* population and its management through planting dates and biopesticides in field pea. *Journal of Agricultural Technology*, 8(1): 125-132. Mittal V and Ram U.2007.
- Nawab NN, Subhani GM, Mahmood K, Shakil Q, Saeed A. 2008. Genetic variability, correlation, and path analysis studies in garden pea (*Pisum sativum L.*). *Journal of Agricultural Research*, 46(4):333-34.
- Chauhan Rohit, Singh AK, Sharma Ravi Kamal, and Ali Amjad. 2018.Screening of mungbean (*Vigna radiata L.*) germplasm against major sucking pest. *Journal of Pharmacognosy and Phytochemistry*.2018,7(5): 1784-1787
- Snedecor GW and Cochran WG.1967. Statistical methods. 6th ed. Ames: Iowa state university press, 3(12): 66-67.
- Ceyhan E and Avci MA. 2005. Combining ability and heterosis for grain yield and some yield components in pea (*Pisum sativum L.*). *Pakistan Journal of Biological Sciences* 8(10):1447-1452. *Sci. Agri*, 16 (3), 2016: 98-103.
- Saleem Muhammad, Hassan Waqar Muhammad and Jamil Moazzam. 2017.Screening of sunflower (*Helianthus annus L.*) varieties against insect pests, predator's populations, and their yield comparison under semi-arid climatic conditions. *Journal of Entomology and Zoology Studies* 2017, 5(3): 1219-1225
- Singh N, Panday DK and Dikshit HK .2002. Status of germplasm its management and utilization in pulse crop. *Farmers Forum*, 2(4):23-27. 42
- Teshome A, Mendesil E, Geleta M, Andargie D, Anderson P, Mert B, Seyoum E, Hillbur Y, Dagne K and Bryngelsson T 2015. Screening the primary gene pool of Field pean (*Pisum sativum L. subsp. sativum*) in Ethopia for resistance against pea weevil (*Bruchus pisorum L.*). *Genet Resoure Crop Evol*, 62:525–538
- Abhilasha CR Shekharappa. 2017. Field screening of pea, (*Pisum sativum L.*) Varieties for resistance against major insect pests An international journal of life sciences, The Bioscan, 12(2):815-818.