

Physicochemical properties of sweet potato chips (*Ipomoea batatas*)

ABSTRACT

Aims: The study aimed to produce and characterize sweet potato-based chips, focusing on pH, titratable acidity, moisture, ash, fat, protein, carbohydrates, and calorific value. Sensory quality was assessed using acceptance tests.

Study Design: The research followed an experimental design with four formulations of sweet potato chips.

Methodology: Four different formulations of sweet potato chips were prepared, varying in the addition of ingredients such as sodium chloride and spice seasoning. Various chemical analyses were conducted to determine the nutritional content of the chips, and sensory evaluations were performed.

Results: The study found differences in pH, moisture, carbohydrates, proteins, lipids, ash, titratable acidity, and calorific value among the different chip formulations. However, there were no significant differences in terms of moisture, protein, and titratable acidity. Formulations (B) and (C) achieved a 70 percent acceptance rate in sensory evaluations.

Conclusion: Sweet potato chips offer an alternative way to enjoy the nutritional benefits of sweet potatoes. The study provides insights into the composition and sensory acceptability of different chip formulations, which can guide future product development.

Keywords: Food diversification, food quality, sensory analysis.

1. INTRODUCTION

The consumption of industrialized food is directly related to the way of life that characterizes today's societies. The demand for quality food has been growing, from the raw material to the finished product, in order to reduce harm and guarantee its quality [26].

The group of processed foods includes sweet potato chips, which are currently well received by consumers, with low levels of cholesterol, trans fats, low sugar levels and are beneficial because they reduce the rate of diseases such as diabetes, high blood pressure, cancer and lower blood levels in the body [4].

Chips are a food produced from potatoes cut into thin slices, fried in vegetable oil or baked in the oven and salted. Various condiments and flavourings can be added at the end of production to improve their sensory and physicochemical characteristics [28].

The chips must have basic characteristics such as high dry matter content (minimum 18%), low reducing sugar content (maximum 0.5%), low fat absorption [39]

The production of chips using sweet potatoes has been gaining ground in an attempt to take advantage of their nutritional and medicinal quality. They contain levels of carbohydrates, vitamins, minerals, fiber and beta-carotene [6].

According to [24], sweet potatoes help strengthen the human immune system, act as antioxidants that reduce the onset of chronic diseases and help reduce premature ageing.

Given the post-harvest losses of sweet potatoes, the implication for producers and consumers is a major problem that contributes to the poor utilization of this crop [50]. In this context, this research aimed to produce and evaluate the physicochemical and organoleptic qualities of sweet potato-based chips.

2. MATERIALS AND METHODS

2.1 Study area

The study was conducted at the laboratory of Higher Polytechnic Institute of Gaza (ISPG), Lionde administrative post. According to [37], Chókwè district is located in the south of the Gaza province, in the middle of Limpopo River, bordered to the north by Limpopo River, on south by district of Bilene, Mazimuchope and Xai-Xai, to the east by districts of Bilene and Chibuto and to the west by districts of Magude and Massingir.

Chókwè district contributes around 27.6% of sweet potato production, 58% of maize, 27.6% of rice, 31% of cassava, 3% of tomatoes and 6.9% of beans, making it important and fertile in the context of developing new technologies for the conservation and use of this tuber [1].

2.2 Acquisition of raw materials

2kg of sweet potato samples, variety "Irene", absent of cracks, with a firm consistency, and no cuts on the surface of the peel, were purchased from the local market, along with sodium chloride (Impala), potato seasoning (spice and paprika), and vegetable oil (Dona), packed in high-density polythene bags and taken to the laboratory for production and analyses.

2.3 Production of sweet potato chips

2.3.1. Weighing and washing

The material with the characteristics described in point 2.2 was weighed using an analytical balance. The sweet potato was washed by immersing it in running water to remove impurities and the microbial load on the surface of the product.

2.3.2. Peeling, sanitising

Peeling was carried out manually using a stainless steel knife, eliminating any defects that affected the pulp. The material was sanitized by placing it in a solution of chlorinated water at 10 ppm active chlorine for 10 minutes.

2.3.3 Moulding and margining

Moulding was carried out using a potato peeler (Riastvy brand) with which uniform longitudinal cuts of ± 2 cm in thickness were made. 5kg of this material was then placed in 3

liters of brine previously prepared with citric acid and water in a ratio of 1:9 and left to marinate for 30 minutes. The material was then removed and left to rest on cotton cloths for 30 minutes.

2.3.4 Mixing the ingredients

The ingredients listed in Table 1 were mixed by hand and left to stand for 10 minutes. This was followed by frying using vegetable oil (Dona). To do this, 1.3kg of the moulds were fried in 250mL of vegetable oil placed in a stainless steel pan. The heat source used was an ANVIL COA3004 gas cooker with a temperature of 180°C for ± 9 minutes, after which the chips were removed with the aid of a grid-shaped foamier.

Table 1. Sweet potato chip formulations.

Ingredients (%)	Formulations			
	A	B	C	D
Sweet potatoes	100	98	90	95
Salt	0	2	0	0
Spice Seasoning	0	0	10	0
Paprika powder	0	0	0	5

2.4 PHYSICOCHEMICAL ANALYSES

Based on [3], pH, total titratable acidity, carbohydrates, calorific value, proteins, lipids, ash and moisture were determined.

2.4.1 Moisture

Moisture was determined using the gravimetric method. For this purpose, 5g of chips were previously crushed in a mortar, added to a petri dish previously weighed on a 0.0001g precision Adam analytical balance, dissected in an ecotherm oven with circulating air at 105°C for 2 hours, then cooled to room temperature (±25°C). Equation 1 was used to determine moisture.

$$\% \text{ moisture content} = \frac{(P_i - P_f)}{P_i} * 100$$

Where:

P_i- Initial weight of the sample;

P_f- Final weight of the sample.

2.4.2 Fat determination

The fat content was determined using the Goldfish method. 5g of the sample was crushed in a mortar and subjected to direct extraction of fat with petroleum ether, extracting (55°C) for 4 hours in a Goldfish apparatus. Expression 2 was used to determine the fat percentage.

$$\% \text{ Fat} = \frac{(\text{capsule weight} + \text{fat}) - (\text{capsule weight})}{\text{capsule weight}} * 100$$

2.4.3 Determination of titratable total acidity (ATT)

In triplicate, 5g of sample was weighed using a 0.0001g ADAM analytical balance and diluted in a 200 mL beaker containing 95 mL of water, homogenizing manually for 5 minutes.

Subsequently, 0.3 mL of phenolphthalein solution was added and titrated with sodium hydroxide solution (0.1M NaOH), homogenizing until the turning point was reached. Equation 3 was used to determine treatable acidity.

$$\% \text{ acidity} = \frac{(V * F * M)}{p} * 100$$

Where:

V -Number of ml of sodium hydroxide solution used in the titration;

F -Correction factor for the sodium hydroxide solution;

p -Sample mass in g or pipetted volume in mL;

M -Molarity of the sodium hydroxide solution.

2.4.4 Ash

On an ADAM analytical balance with 0.0001g precision, 5g of sample was weighed into porcelain crucibles and placed in an OPTIL IVYMEN muffle furnace at 550°C until the incineration was verified. The crucibles were then transferred to an oven at 105°C for 30 minutes with the emphasis on lowering the temperature, followed by weighing them with the incinerated sample in inorganic matter. The percentage of ash was determined using equation 4:

$$\% \text{ ash} = \frac{(m2 - m)}{(m1 - m)} * 100$$

Where:

m - Crucible weight;

m1 - weight of crucible with wet sample;

m2 - weight of crucible with ash;

2.4.5 Carbohydrates

They were determined by difference. To do this, the percentages of moisture, fat, protein and ash in 100g of the sample were added and subtracted from the maximum percentage (100%). Equation 5 was used to determine the carbohydrate content.

$$\text{Carbohydrates \%} = (\% \text{ moisture} + \% \text{ ash} + \% \text{ protein} + \% \text{ fat}) - 100 \quad [5]$$

2.4.6 Calorific or energy value

The calorific or energy value of the samples was determined using the method of calculating the sum of macronutrients that supply energy to the human body, using the coefficients and based on the equation:

$$\text{Calorific Value (Kcal)} = ((\% \text{ protein} * 4 + \% \text{ carbohydrates} * 4) + \% \text{ lipids} * 9) \quad [6]$$

2.4.7 Determining pH

In triplicate, 10g of sample was weighed into an erlenmeyer flask and diluted in 90ml of water. The contents were shaken to remove the suspended particles from the sample, then the pH was read after emptying the electrode of the HANNA model pH meter (HI 2212 pH/ORP Meter), previously calibrated with buffer solutions 7 and 10.

2.4.8 Proteins

The protein content was determined using the Biuret method, where 300 μL of each extract (prepared in the ratio: 10g of Chips and 90 mL of water) was mixed with 2000 μL of Biuret reagent and left in a dark place for 30 minutes to give a purple complex color, and then the absorbance was read at 540 nm on a brand spectrophotometer (YOKE) previously zeroed with distilled water. The protein content of samples was determined by extrapolation using a calibration curve made up of casein n in proportions of 0 to 10 mg/ml.

2.5 Sensory analysis

The analysis continued with the application of affective methods. The test was carried out by a panel made up of (50) untrained tasters of both sexes aged between 18 and 45, including teachers, students and staff selected at random. The following attributes were evaluated: appearance, color, crunchiness, aroma, flavor, aftertaste and overall assessment, using a 9-point hedonic scale ranging from "1" extremely disliked to "9" extremely liked". The acceptance index (AI) was calculated using equation 7:

$$AI = \frac{(A * 100)}{B}$$

Where:

A - Represents the average score obtained for the product;
B - Maximum score given to the product.

2.6 Statistical analysis

The analysis of variance (ANOVA) was carried out using the general linear model (GLM) using the RStudio statistical package 4.2.1. A completely randomized design with 12 experimental units was used to analyze the physicochemical parameters and a randomized block design with 50 experimental units was used for the sensory analysis. In the event of significant effects, the difference between the means of experimental units was evaluated using Tukey test at 5% level.

3. Results and discussion

3.1 Physicochemical composition of sweet potato chips

The centesimal composition of chip samples evaluated is shown in Table 2.

Table 2. Physicochemical composition of chips.

Parameters	Formulations			
	A	B	C	D
Moisture (%)	2,6 \pm 0,0 ^a	2,5 \pm 0,8 ^a	2,2 \pm 0,3 ^a	2,4 \pm 1,0 ^a
Carbohydrates (%)	77,3 \pm 0,5 ^b	82,1 \pm 3,4 ^a	81,2 \pm 0,8 ^{ab}	79,0 \pm 0,9 ^{ab}
Protein (%)	8,0 \pm 1,8 ^a	5,9 \pm 1,9 ^a	6,3 \pm 0,7 ^a	4,8 \pm 2,1 ^a
Lipids (%)	9,8 \pm 0,9 ^{ab}	7,9 \pm 1,2 ^b	8,7 \pm 0,7 ^{ab}	11,3 \pm 1,7 ^a
Ash (%)	1,5 \pm 0,2 ^{ab}	1,6 \pm 0,0 ^{ab}	1,3 \pm 0,0 ^b	1,9 \pm 0,1 ^a
Calorific value (Kcal)	430,8 \pm 1,8 ^{ab}	423,6 \pm 4,9 ^b	428,7 \pm 5,2 ^{ab}	447,0 \pm 4,4 ^a

Others	pH	5,5±0,1 ^a	5,2±0,0 ^b	5,1±0 ^b	5,0±0,1 ^b
	Acidity (%)	3,2±0,2 ^a	2,9±0,1 ^a	3,1±0,2 ^a	3,2±0 ^a

Means ± standard deviation followed by the same letter on the same line do not show significant differences at Tukey's 5% level. A, Sweet potato pulp; B, 98% sweet potato and 2% salt; C, 90% sweet potato and 10% chips spice seasoning; and D, 95% sweet potato and 5% paprika powder.

3.1.1 Moisture

The results showed that the formulations ranged from 2.2±0.3 to 2.6±0.0%. The highest moisture content (2.6%) was observed in formulation A. On the other hand, the lowest content (2.2%) was obtained in formulation C. This result is probably linked to the moulding, rinsing and frying stages which, according to [38] in their study on production of sweet potato chips with the replacement of sodium chloride by potassium chloride, provide partial dehydration of the moulds and evaporation of free water to the extent influenced by high temperatures (180°C). No significant differences were found between formulations.

Higher results than those obtained in the present study were reported by [40], when evaluating chips formulated with *Discoreabulbifera*, in which the moisture content was 49.70%, by [38], when producing sweet potato chips with sodium chloride substitution, who obtained moisture ranging from 7.3% to 8.3%, by [17], studying physicochemical and sensory characterization of sweet potato chips, mentioned moisture contents around 7% and 8%, and by [14], when evaluating jicama potato chips made by dehydration and drying, who obtained moisture contents of 8.0 to 10%. This difference in moisture content is possibly linked to the physical process each product was subjected to, considering that [77] related that quickly evaporation of volatile compounds is promoted by heating the vegetable oil, directly influencing the desorption of moisture.

3.1.2 Carbohydrates

There were significant differences ($p < 0.05$) between formulations A (77.3± 0.5%) and B (82.1± 3.4%), and the high levels of carbohydrates may be associated with the Irene variety, which according to [69], is considered a complex carbohydrate with up to 90% sugars in its dry matter, being higher than the varieties (white and yellow).

Lower results were reported by [38], when evaluating the physicochemical characterization of sweet potato chips with sodium chloride substitution, indicating carbohydrates ranging from 73.9 to 76.7%, [75], in their research on the preparation of cassava-based chips, reported carbohydrates of 56%, and by [18], evaluating ipomoea batata chips, found carbohydrates of 56.24%. Unlike the authors mentioned above, the high level of carbohydrates found in this study can be explained by the characteristics of Irene variety which, according to [20], has a higher carbohydrate content around 80 to 81 percent for 100g. Similar results (81 and 82% carbohydrates) were reported by [21], when evaluating production of roasted *D. bulbifera* chips, who found 60% and 81% of carbohydrate content, and by [13], in their study about preparation and characterization of chips, who obtained 82.6% of carbohydrates.

3.1.3 Proteins

The samples evaluated showed protein values ranging from 4.8 to 8.0%, with no significant differences ($p < 0.05$) between them.

Similar results to those obtained in this research were reported by [62] in his study about production of dehydrated potato chips, who found 11.4% of protein content. Results that were not in agreement were described by [23] with aim of developing cassava-based chips, reported protein content of 0.92%, [51] studying the centesimal composition of *Soibamdioscorea* spp. chips, obtained protein of 1.2%, and by [12], when developing dehydrated *D. rotundata* cv. chips, found a protein content of 2.1%. For their part, [70], when producing chip formulations with *D. bulbifera*, revealed protein content ranging from 3.7 to 4.1%, and [67] around 4.12% of protein, when developing baked chips, and [64], when studying the effect of osmo-convective dehydration on physical and physicochemical characteristics of chips, obtained protein content ranging from 1.95 to 2.04%, and [11] found protein content of 0.63% in fried cassava chips. The fact that the results of this study do not agree can be explained by [74], who mention the high lipoprotein content in soya oil, which possibly added greater protein value to the chips.

3.1.4 Lipids

Statistical differences were evident in formulations B and D, and this difference may be associated with the frying temperature and time, since it had an effect on the degradation of lipoproteins and fatty acids in the oil as a function of the time of use.

Different results were reported by [57] who, in their study on the characterization of cassava chips, obtained 34.8 to 44.2% of lipids and by [18] who, when studying the production and physicochemical and sensory evaluation of cassava chips subjected to pre-treatment, obtained of 36.5% lipids. The difference between the results of these authors and this study may be associated with the amount of oil used and the cooling stage, which may have led to the low levels of lipids.

Lower values than those obtained in this research were reported by [38], studying *Dioscorea bulbifera* chips as an innovative alternative for the technological processing of unconventional food plants from Amazon, who obtained lipid content ranging from 0.39% to 3.41%, and by [52] when developing chips based on *Lupinus albus* L, who found 3.65% of protein. The difference in lipid content may be related to the cooking processes and/or the processing methods (frying and baking) used to produce the chips.

Reports by [69], in their study of production and centesimal characterization of chips based on *Ipomeia batata*, obtained levels of 6.75%, close to those found in this study.

3.1.5 Ashes

The results for fixed mineral residue ranged from 1.3 to 1.9%. The highest value (1.9%) was observed in formulation D, which is probably related to the properties of added potato spice, which contains a high mineral content and concentrated dry matter. Formulations A and B were not statistically different from each other. On the other hand, significant differences were seen in formulations C and D, possibly due to the concentration of dry matter and mineral content present in the potato spice compared to paprika powder.

In the assessment carried out by [38] in his characterization study physicochemical action of sweet potato chips with replacement of sodium chloride by potassium chloride, obtained ash around 3.1 ± 0.1 to 3.9 ± 0.2 %. Higher results than those obtained in the present study, where the differentiation of the raw material and concentration of potassium ions may have influenced these results.

[23], developing moulded and fried cassava chips, obtained ash content of 2.83%, [16], working on sweet potato snacks, obtained ash content of 3.40%, [36], studying the stability of chips in different packaging systems, found ash content of 3.35%, [65] reported ash content of 2.41% when evaluating Dioscorea chips obtained by oven drying, [7] found 2.40% of fixed mineral residue in their study about production and sensory characterization of Arracacia xanthorrhiza Bancroft chips. These results are above on levels found in this study, where the cooking time and type of raw material may have had an influence.

3.1.6 Calorific value

There was a statistical difference ($p < 0.05$) in formulations B and D with levels around 423.65 ± 4.9 kcal and 447.02 ± 4.4 kcal, and these differences, according to [31] and [46], probably correlate with the variation of lipids concentration, glucose, dietary fibre and lipoproteins in the product.

In the study carried out by [38], with aim of producing chip formulations using potato, lupin and pea flour, they reported energy value ranging from 319.52 to 346.29 Kcal/100g, lower than the results found in this research, which may be directly linked to the constitution of the raw material used in chip production. Results reported by [39], analysing *Dioscorea bulbifera* chips as an innovative alternative for the technological processing of unconventional food plants from Amazon, obtained values ranging from 284.84 kcal to 371.26 Kcal/100g, which is lower than results of this research, and possibly the cooking method, time and temperature used contributed to these results.

3.1.7 Acidity

The highest level of acidity (3.2%) were observed in formulations A and D, with no significant differences. Lower results than those found in this study were reported by [54] studying physicochemical analysis of *D. bulbifera* chips, who obtained 0.25% of acidity, and by [70], 0.27%, in their study about production of *Mussa spp.* chips. According to [71] the differentiation of acidity levels in the application of multidimensional techniques and the nutritional composition of sweet potato chips may be linked to the type of raw material, as well as the cooking stage and, according to [76] the adhesion of the oil to the surface of the chips may have contributed to the increase in acid levels such as lauric and palmitic, considering the sweet potato genotype.

3.1.8 pH

The pH values ranged from 5.0 to 5.5, where formulation A differed significantly from the others, which may be related to concentration of citric acid (90mL of bioactive) added and the ingredients (sodium chloride "salt", spice seasoning and paprika powder) used as seasonings in production. Values in line with those obtained in this study were revealed by [68], who produced sweet potato chips and obtained a pH of 5, within the levels found in this study.

Results close to those obtained in this research were described by [60] when evaluating *Musa spp* chips, which obtained a pH of 5.89. [30], when drying *D. esculenta* chips, obtained a pH of 6.56, which is higher than the results obtained in this study, which may be linked to the drying stage.

3.2 Sensory analysis

Figure 1 shows the results of sensory analysis of sweet potato chip formulations.




Fig 1. Level of acceptance of the sweet potato chips.

Means ± standard deviation followed by the same letter on the same line do not show significant differences at Tukey's 5% level. (A), sweet potato pulp; (B), 98% sweet potato and 2% salt; (C), 90% sweet potato and 10% spice seasoning chips; and (D), 95% sweet potato and 5% paprika powder.

Source: Authors.

3.2.1 Appearance

Formulation D (6.0 ± 2) differed significantly ($p < 0.05$) from the others, probably due to the brownish color of the paprika which may have given it a different appearance. [64] studying sensory analysis of yacon chips, obtained a score of 6.5, a result in line with that found in formulation D of the present research, which is in line with that found by [55], making yacon chips, who reported value of 6.5 for the appearance attribute. Similarly, [38], studying the production of chips with partial replacement of sodium chloride with potassium chloride, obtained average of 7.5, which is also in line with the results of this research.

3.2.2 Color

The averages ranged from 7.14 to 7.5, with statistical differences in the formulations produced. These differences were caused by the addition of spice, the ingredient that affected the intensification of the color and the caramelization process. Similar results to those obtained in this study were reported by [47] studying the frying temperature of chips and sensory acceptance, who obtained a score of 7, and by [19] in his study related with sensory analysis of chips stored at low temperatures, obtained a score of 7. Different values were described by [17] when carrying out a sensory analysis of potato chips with fish flavouring flavoured with lemon, obtaining 8.5, and by [55] when evaluating the quality of the color of chips, who obtained a score on a scale of 6.

3.2.3 Flavor

The flavor attribute ranged from 5.8 to 7.5. The highest score (7.5) was given to formulation B, which was statistically different from the others. This variation may be linked to the different ingredients used to make the chips. Similar results (6, 7, 7.4 and 7.46) to those found in this research were obtained by [64] at around (7.4), when studying the dehydration of chips, [10] in the range of (6 to 7), seeking to evaluate the sensory analysis for launching a product on the market, and by [59] when carrying out their study about physical and sensory quality of dehydrated Yacon chips, they obtained values of 7.46.

3.2.4 Aftertaste

The highest score was observed in formulation C (7.4), followed by sample B with (6.7). Significant differences ($p < 0.05$) were observed in formulations A and D compared to formulations B and C. This variation in aftertaste preference is possibly due to its acidic and spicy properties. [22], in his sensory analysis of chips, revealed aftertaste scored 6 on hedonic scale, while [44], seeking to study the sensory acceptance of chips made with yams, obtained scores ranging from 6.7 to 7.4, results in line with those obtained in the present study. Higher results than those obtained in this research were referenced by [39] studying on *Dioscorea bulbifera* chips as an innovative alternative for the technological processing of unconventional food plants from Amazon, who obtained a score on the scale of 8.1.

3.2.5 Crunchiness

There were no significant differences ($p > 0.05$) between the formulations. This finding may be associated with the thickness of the chips, moisture content and frying temperature. [38] in his research seeking to produce chips with sodium chloride substitution, obtained average of 7.7, [28] producing cassava chips, who obtained a score of 7.6, results similar to those found in this research. Different results to those found in this research were reported by [17] evaluating the sensory quality of cassava chips, who obtained score in range of 6.5 to 8.3, and by [41] in the range of (5 to 8.45), when processing potatoes of different genotypes and frying them, taking into account [11], this finding may be linked to the dry matter of the food, inferring an increase in crispiness levels due to dehydration and removal of the sweet potato's moisture content.

3.2.6 Flavor

Flavor scores ranged from 5.5 ± 2.2 to 7.3 ± 1.8 . The highest score (7.3) was found in formulation C, which was statistically different ($p < 0.05$) from the others. This variation is linked to composition of the spice seasoning. Similar results to those obtained in this research were obtained by [34] studying about sensory analysis of sweet potatoes subjected to different cooking methods, obtained score of 6.8, [5] seeking to develop cassava chips, who obtained score of 6.8, and by [44], around 6.8, evaluating chip consumers in relation to

o [39] when carrying out their study on *Dioscorea bulbifera* chips as an innovative alternative for the technological processing of unconventional food plants from Amazon, obtained score of 7.8.

3.2.7 Overall assessment

No statistical differences were observed between the samples, with values ranging from 5.5 ± 1.8 to 6.3 ± 2.0 . This variation was correlated to dietary habits and the conditions in which the analyses were carried out. [33] in their study about production and physicochemical characterization of crisps, found a score of 6.3, a result in agreement with the present study. Higher results were reported by Santos, [59], who analyzed the sensory characteristics of sweet potato chips using different oils and obtained an overall score of 6.8, by [67] in their studies on chips, who obtained a score of 8.2, and also by [72] studying yam chips, who obtained a score of 8.

3.3 Purchase intention

The results of buy intention of Chips are shown in figure 2.




Fig 2. Chip purchase intention test.

(A), sweet potato pulp; (B), 98% sweet potato and 2% salt; (C), 90% sweet potato and 10% spice seasoning chips; and (D), 95% sweet potato and 5% paprika powder.

Source: Authors.

Formulation C obtained the best purchase intention index (70%), which may be associated with eating habits and the composition of formulations, which tended towards commercial chips, and the taste and appearance of product were associated with these characteristics. [44], studying about production and sensory approval of sweet potato chips, revealed 70% of purchase intention, in agreement with the results obtained in the present study. [25], in his research aimed to study sensory quality of cassava chips of different genotypes and subjected to different frying temperatures, obtained 74.22% of purchase intention, [73], when carrying out sensory analysis of potato chips, obtained 44%, and [58] reported 38% in their studies evaluating the purchase intention of potato chips flavored with fish.

3.4 Acceptability index

Figure 3 shows the results of the acceptability index for (*Ipomeabatatas*) chips.




Fig 3. Acceptability index of sweet potato chip formulations.

(A), sweet potato pulp; (B), 98% sweet potato and 2% salt; (C), 90% sweet potato and 10% spice seasoning chips; and (D), 95% sweet potato and 5% paprika powder.

Source: Authors.

The acceptability index of the sweet potato chips showed that formulations B and C had the best with 70%.

Consistent results were reported by [49] in his study about sensory acceptance, who obtained results of 70% acceptability, [53], when evaluating the acceptability of banana chips from the tip of the bunch, in the range of 70% to 75%, and [2] when evaluating the development of potato chips, who obtained acceptance of 82%, which is within the recommendations for product acceptance. On the other hand, [58], when producing fish-flavored chips, obtained 38% product acceptance, which is lower than the result observed in this study.

4. Conclusion

No significant differences were found in terms of moisture, protein and titratable acidity levels. Differences were found in lipid content, carbohydrates, ash, calorific value and pH. Sensorially, the chips in formulations A and B, with 97.5% sweet potato and 2.5% spice seasoning and 99.2% sweet potato and 0.8% salt, had satisfactory acceptance ratings around 70%. The sweet potato proved to be technologically viable for the production of chips, helping to guarantee food and nutritional security as well as reducing waste.

ACKNOWLEDGEMENTS

We express with love and affection, our gratitude to all those who directly or indirectly provided their preponderant help, whether moral or material, for the development of this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

References

1. Amade, A. Survey of Rainwater Collection and Conservation Techniques in Agricultural Production in the District of Chókwe, 2006.
2. Ambrósio CLB, Campos FACES, Faro ZP. Carotenoids as an alternative against hypovitaminosis A. *Nutritional Journal*, Campinas. 2006;19 (2):233 -243.
3. AOAC - Official Methods of the Association of the Agriculture Chemists. Guidelines for standard method performance requirements, 2016.
4. Augustin MA, Riley M, Stockmann R, Bennett L, Kahl A, Lockett T, Osmond M, Sanguansri P, Stonehouse W, Zajac I. Food processing and nutritional safety rules, 2016.
5. Barbosa, AF, Lopes FJ, Silva VRO, Silva MHL, Minim VPR, Rita CSNS. Sensory acceptance of peach-flavoured yoghurt with different concentrations of flavouring and pulp using the preference map technique. *Journal of the "CândidoTostes" Dairy Institute*, 2016;(68):52-58.
6. Bernardes IB. Breaking dormancy in botanical sweet potato seeds using different chemical compounds, 2019.
7. Bessa LAS, Jardim, FBB, Dias LCFC, Costa LL. Physico-chemical and sensory evaluation of manioc chips - *parsley. roots and starchTropical magazine*. 2016; 12(1):83-95.
8. Bobbio PA, Bobbio FO. Chemistry of food processing, 2001.
9. Borges JTS, Paula CD, Pirozi M. Physico-chemical composition, physical and sensory quality of cassava chips. *Evolution and engineering*. 2023;1(2):59-69.
10. Borges JTS, Pirozi MR, Paula CD, Vidigal JG, Sousa e Silva NA, Caliman FRB. Yacon in human nutrition: nutritional and functional aspects, utilisation and toxicity. *Scientia Amazonia*. 2012;1(3): 3-16.
11. Borges J, De Paula C, Pirozi M. Physicochemical composition, physical and sensory quality of cassava chips. *journal of evolution and engineering*. 2013;1(2):59-69.

12. Bovi DCML, Correia AFK, Gutierrez EMR, Bovi VML, Harder MNC. Determination of betalain content and centesimal composition of beetroot in natura and chips. *Brasileira Agro-industrial Products*. 2019;17(1):17-23.
13. Brito CM, Boitrago WJ, Almeida ES, Melo CMT, Queiroz CRAA. Preparation and characterisation of Jicama potato chips (*Pachyrhizuserosus*). 2020.
14. Brito CM, Boitrago WJ, Almeida ES, Melo CMT, Queiroz CRAA. Rural Extension: practices and research for strengthening family farming - Preparation and characterisation of jicama potato chips (*Pachyrhizuserosus*). 2021; 12(1):282-297.
15. Caetano PK, Mariano-Nasser FAC, Mendonça VZ, Furlaneto KA, Daiuto EG, Vieites RL. Physicochemical and sensory characteristics of sweet potato chips produced using different cooking methods. *Food Science and Technology*. 2018;38(3):434-440.
16. Campos VR, Calliari CM. Elaboration of Sweet Potato Snack (*Ipomoea Batatas*). In: Oliveira AF, Storto LJ. *Topics in Food Science and Technology: results of academic research, São Paulo*. 2016;(2).263-284. <https://doi.org/10.21897/23460466.774>
17. Caroline BT, John DP, Robert C, Jin-Shei L, Berend T, David C, Pip G, Lidwine B. Mokkink. A conceptual clarification and systematic review of MIC estimates of PROMIS measures. *Quality of Life Research*. 2021.2729-2754.
18. Carvalho AV, Seccadio LL, Ferreira T. *Production and physico-chemical and sensory evaluation of pre-treated cassava chips, 2010.*
19. Celestino SMC. *Principles of Food Drying*, 2010.
20. Cosmo BMN, Galeriani TM. Vitamins in animal feed. *Revista de agronomia brasileira*. 2020;(4)1-9. 10.29372/rab20 2020. Available at: <http://rsdjornal.org>> Accessed: 02/05/2023.
21. Djeri B, Tchobo PF, Adjrah Y, Karou DS, Ameyapoh Y, Soumanou MM. (Reuters) - M. E Souza, C. (in English). Nutritional potential of yam chips (*Dioscoreacayenensis* and *DioscorearotundataPoir*) obtained using two production methods in Togo. *African Journal of Food Science*. 2015;9(5):278- 284.
22. Dutcosky SD. *Sensory Analysis of Food*. 5. ed. Curitiba: Champagnat – Pucpress. 2019. 540.
23. Elaine MF. *Development of cassava chips, moulds and fries*, 2011.
24. Fernandes AM, Soratto RP, Evangelista RM, Nardin I. Physico-chemical and frying quality of potato cultivars in the winter harvest, *Brazilian Horticulture*. 2010;28: 299 - 304.
25. Ferrarezzo EM. *Development of cassava chips, moulded and fried*. Thesis (PhD) -- Faculty of Animal Science and Food Engineering - University of São Paulo. Department of Basic Sciences. 2011:131-172.
26. Galeriani TM, Cosmo NMB, Ceccon PPA, Marchi SC, Mundt TT. Cultivars of cortex roses. *Revisit agronomic brasielira*. 2020:1-4.

27. Geisa PAGM. Sweet potato chips with replacement of sodium chloride by potassium chloride. Rio Verde. 2019:22.
28. Grizotto KR, Reuters R, Menezes HC. Evaluation of the acceptance of cassava chips. Food Science and Technology. 2003;23:79-86.
29. Grizotto, RK. Cassava Chips a technology for utilising cassava (ManihotesculentaCrantz), 2010.
30. Hilman A, Harmayani E, Cahyanto M. Extraction of Inulin and Characterisation of Gembili (Dioscoreaesculenta) Extract by Ultrasound Assisted Extraction Proceed of the International Conference on Science, Technology, Engineering, Environment and Branching Research. 2018;1(1):47-53.
31. Kayode MO, Buhari OJ, Otutu LO, Ajibola TB, Oyeyinka SA, Opaleke DO, Akeem SA. Physicochemical Properties of Processed Aerial Yam (Dioscoreabulbifera) and Sensory Properties of Paste (Amala) Prepared with Cassava Flour. The Journal of Agricultural Sciences. 2017;12(2):84-94,.
32. Kita A. Influence of crispy texture on the chemical composition of potatoes, 2002.
33. Laurie SM, Faber M, Calitz FJ, Moelich EI, Nina M, Labuschagne MT. Use of sensory attributes, sugar, instrumental analyses and consumer acceptability in the selection of sweet potato varieties. Journal of the Science of Food and Agriculture. 2013;93(7):1610-1619.
34. Liana CFLV. Evaluation of nutrient retention, sensory and microbiological aspects of sweet potatoes (Ipomoea batatas (L.) Lam.) subjected to different cooking methods. 112f, ilcolour. Federal University of Ceará, 2016.
35. Low JT, Walker RKR, Hijmans DRP, Ewell YDZ. The potential impact of beta-carotene-rich sweet potatoes on vitamin A intake in Sub-Saharan Africa. 2003.
36. Luís MJ, Biofortified sweet potato chips: product development and stability study in different packaging systems. 2017.
37. MAE, Ministry of State Administration Chókwè district profile. Gaza Province. 2014. Available at: <http://www.govnet.gov.mz>. Accessed: 12/10/2022.
38. Maha IKA, Esraa AM, Mousa NAA. Hassan. Production of Healthy Chips Ready to Eat Using Potato, Green Pea and Lupine Flour for Malnourished Children. International Journal of Food Science and Biotechnology. 2019:1-9.
39. Maia AG, Melo NGM, Dantas LOO, Souza RP, Moreno MN, Marinho NMM, Reuters MV, Martim SR, Reuters R. Chips from Dioscoreabulbifera: an innovative alternative for the technological processing of unconventional food plants from the Amazon. Research, Society and Development, [S. I.]. 2021;(10)15.
40. Mapiemfu-Lamare D, Josiane M, Ngome A, Dingwan M, Eliane E. The quality of fried chips varies with cassava (ManihotesculentaCrantz) varieties and maceration regime. 2018.

41. Márcia ML, Caroline DB, Marina VS, Daniela SA, Flávia FP, Ana CRK. Physicochemical and sensory characteristics of potato chips made from blanched potatoes of the cultivar BRS Ana and coated with methylcellulose. *Brazilian Journal of Food Technology*. 2015;1- 9.
42. Mesquita AS. Yam and taro: international, Brazilian and Bahian market scenarios. 2002.
43. Michels I. Technological aspects of the minimum processing of yacon tubers (*polymniasonchifolia*) stored in modified atmosphere packaging. (Master's thesis). Federal University of Paraná, Curitiba. 2005.
44. Minim VPR. *Sensory Analysis: Consumer Studies*. 4. ed. Viçosa: Editora UFV. 2018:362.
45. Mogrovejo Sanchez & Diego Isique Turpo, José Carlos. Obtaining and determining bioactive compounds (phenolic compounds, anthocyanins and polyphenols) from pigmented native potatoes from the High Andean region. 2021. Accessed at: <http://repositorio.ucsm.edu.pe/bitstream/handle/20.500.12920/10537/41.0308.AG.pdf?sequence=1&isAllowed=y>, 2023.
46. Moraes DP, Lozano-Sánchez J, Machado ML, Vizzotto M, Lazzaretti M, Leyva-Jimenez FJJ, Barcia MT. Characterisation of a new blackberry cultivar BRS Xingu: Chemical composition, phenolic compounds, and antioxidant capacity in vitro and in vivo. *Food Chemistry*. 2020; 322: 126783.
47. Neves VA, Silva MA. Polyphenol oxidase from yacon (*Smallanthussonchifolius*) roots. *Journal of Agricultural and Food Chemistry*. 2007;55(6):2424-2430.
48. Neves, V. A., & Silva, M. A. (2007). Polyphenol oxidase from yacon roots (*Smallanthussonchifolius*). *Journal of Agricultural and Food Chemistry*, 55 (6), 2424-2430.
49. Noronha JF. *Notes on Sensory Analysis*. Coimbra: ESAC. 2003.75p.
50. Oliveira MC et al. Nutritional composition and fatty acid profile of potato chips and extruded snacks (Master's thesis). **State UniversityPaulista**, São José do Rio Preto, 2013.
51. Omohimi CI, Piccirillo C, Roriz M, Ferraro V, Vasconcelos MW, Sanni LO, Tomlins K, Pintado MM, Abayomi LA. A study of the centesimal and mineral composition of different Nigerian yam chips, flakes and flours. *Journal of Food Science and Technology*. 2018;55(1): 42-51.
52. Ozcan M.M., Ipek D., Ghafoor K., Al Juhaimi F., Uslu N., Babiker E.E., Ahmed I.A.M., Alsawmahi O.N. Physico-chemical and sensory properties of chips produced using different lupin (*Lupinusalbus* L.) flour formulations and cooking methods. *Int. J. Food Sci. Technol.* 2021;56:2780-2788.
53. Paula Aia et al. Evaluation of the acceptability of banana chips from the tip of the bunch. 2012. Available at: <http://propi.ifto.edu.br/ocs/index.php/connepi/vii/paper/viewFile/2244/2882> .

54. Pedro IS, Suelma FO, Deyzi SG, Mércia MM, Rebeca LD, Cláudia GR. Effects of osmo-convective dehydration on the physical and physicochemical characteristics of savoury sweet potato chips. 2019.

55. Rabelo NM, Silva LMR, Canuto KM, Vieira JMM. Elaboration of yacon chips by the process of osmotic dehydration and drying. *Brazilian Journal of Food Research*. 2017;8(2):145-154.

Rib

1. eiro JA. Chemical and biochemical studies of yacon (*Smallanthus sonchifolius*) in natura and processed and influence of its consumption on glycaemic levels and faecal lipids in rats. 2008.

57. Rogério WF, Leonel M. Effect of slice thickness and pre-cooking on the quality of fried snacks (Chips) made from tropical tubers. 2004.

58. Sales PVG, Sales VHG, Oliveira EMO. Sensory evaluation of two fish burger formulations. *Brazilian Magazine of Agroindustrial Products, Campina Grande*. 2015;17(1):17 - 23.

59. Santos CSP, Cunha SC, Casal S. Deep or air frying? A comparative study with different vegetable oils. *European Journal of Lipid Science and Technology*. 2017;6:119.

60. Santos MRL, Souza PB. Physicochemical, microbiological and sensory evaluation of green banana chips (*Musa spp.*). *Multidisciplinary Science Journal*. 2020;7(1): 1 - 7.

61. Severo J, Dos Santos FN, Samborski T, Rosane R, Mello AFS. Biofortified sweet potatoes as a tool to combat vitamin A deficiency: Effect of food processing in carotenoid content. *Chilean Journal of Nutrition*. 2021;48(3):414-424. Available at: <<http://dx.doi.org/10.4067/S0717-75182021000300414>>.

62. Silva EL. Preparation of dehydrated potato chips. Ponta grossa. 2011.

63. Silva PIS, Oriente SF, Gouveia DS, Mota MMA, Dantas RL, Rodrigues CG. Effect of osmo-convective dehydration on the physical and physico-chemical characteristics of chips. *Magazine Agropecuária Técnica, Areia-PB*. 2018;39(3):264-269.

64. Silva AF, Minim VPR, Ribeiro MM. Sensory analysis of different commercial brands of organic coffee (*Coffea arabica* L.). *Science and Agroecology, Lavras*. 2006;29:1224-1230.

65. Silva LSC, Martim SR, Souza RAT, Machado ARG, Teixeira LS, Sousa LB, Teixeira MFS. Extraction and characterisation of starch from *Dioscorea* species cultivated in Amazonia. *Bulletin of the Museu Paraense Emílio Goeldi-Natural Sciences*. 2020;14(3): 439-452.

66. Silva V, Pereira S, Vilela A, Bacelar E, Guedes F, Ribeiro C, Gonçalves B. Preliminary insights in sensory profile of sweet cherries. *Foods*, 2021;10(3):612.

67. Silvia V. Lopez, Rosana G. Moreira. Increased Phenolic Compounds in Potato Chips Vacuum Impregnated with Green Tea. *Food Engineering, Materials Science, & Nanotechnology*. 2018:807 - 817.

68. Smith TP, Stoddard S, Shankle M, Schultheis J. Sweet potato production in the United States of America. 14 Science + Business Media B.V, 2009.
69. Soibam H, Ayam V, Mitra S. Effect of heat treatment on chemical composition, microbiology and sensory evaluation of yam chips during storage. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(5):1705-1711.
70. Soibam H, Ayam V, MITRA, S. Effect of temperature treatment on the chemical composition, microbiology and sensory evaluation of Yam chips during storage, 2017.
71. Suárez MH, Hernández AM, Galdón BR, Rodríguez LH, Cabrera CEM, Mesa DR, Rodríguez-Rodríguez EM, Romero CD. Application of multidimensional scaling technique to differentiate sweet potato (*Ipomoea batatas* (L.) Lam) cultivars according to their chemical composition. *Journal of Food Composition and Analysis*. 2016;46:43-49. Available at: <<https://doi.org/10.1016/j.jfca.2015.10.008>>.
72. Taciano P. Osmotic dehydration and frying of yam sticks in coconut and soya oils. *Campina Grande-PB*. 2015:1-173.
73. Teferi D, Geremew C. Effect of Blanching and Frying Time on the Sensory Quality of Fried Sweet Potato Chips. *Food and Nutrition Science - An International Journal*. 2019: 1-17.
74. Vital ANS, Messias CMBO. *Beauregard Sweet Potato: Literature Review*, 2020.
75. Vitrac O, Dufour D, Trystram G, Raoult-Wack AL. Characterisation of heat and mass transfer during frying and its effect on the quality of cassava chips. 2002.
76. Vizzotto M et al. Mineral composition in sweet potato genotypes with coloured pulps and adequacy of consumption for risk groups. *Brazilian Journal of Food Technology*. 2018;(21).1-8.
77. Zambrano M, Dutta B, Mercer D, Maclean H, Touchie M. Evaluation of moisture content measurement methods for dry food products in small-scale operations in developing countries: A review. *Trends in Food Science & Technology*. 2019 (88). 484-496