

Study the impact of abiotic factors on the population fluctuation of major insect pest of field Pea

Abstract :-

The present investigation entitled Study the impact of abiotic factors on the population fluctuation of major insect pest of field peawas conducted at Experiment field of Department of Plant breeding, seed Breeding farm, JNKVV Jabalpur (MP), Madhya Pradesh during Rabi season 2020-21. Positive correlation of *Acrythosiphonpissum*, pod borer and leafhopper with Maximum Temperature (0.581, 0.581, 0.582) and *Acrythosiphonpissum* with morning vapour pressure (0.453) may be given due weightage in formulating forewarning modules in the pest. Similarly positive correlation was found between Relative humidity and aphid ($r = -0.304$) and was found significant. Max temperature showed significant negative correlation ($r = -0.581$) with aphid, Pod borer and leafhopper's population where minimum temperature and rainfall showed negative correlation ($r = -0.453$ and $r = -0.153$) with aphid population. Whereas min temperature showed positive non-significant correlation ($r = 0.413$). Max temperature showed significant negative correlation ($r = -0.581$) with pod borer population. Morning vapour pressure shows significant negative correlation ($r = -0.453$) with aphid population.

Keywords:- Correlation, Fieldpea, leafhopper, Pod borer, *Acrythosiphonpissum*.

Introduction:

Field pea is the self-pollinated diploid crop and it is the most important annual season crop that is valued as a high-quality protein food. It is commonly cultivated in the cooler temperate zones and tropical highlands around the world. The crop can be grown in variety of soil types, from light sandy loam to hard clays but it does not tolerate salty or wet conditions.

Field peas are rich in high quality vegetable protein and does contain all the essential amino acid. The seeds of peas are prized for their nutritional value. Sulphur containing amino acids including cysteine and methionine are scarce in pea, but lysine

and other essential amino acids are plentiful. It is good source of essential amino acids in the form of protein (23-25%) which have high nutritional benefit to low-income families (Nawab *et al.* 2008). Furthermore, pea contains ample amounts of essential minerals such as calcium, phosphorus and iron which are deficient in cereals (Haque *et al.* 2015). Peas have a starch content (20-25%), sugar content (4-10%), fat content (0.6-1.5%), mineral content (2-4%) (Makasheva 1983). It is mainly an export and cash crop for the world, accounting for around 40% of total world pulse trade.

The nitrogen fixing capacity of this crop help in restoring soil fertility (Singh *et al.*, 2002). Due to high market price in the neighboring states of Maharashtra, Pea production for vegetable purpose in the Jabalpur district has been a huge success. Although its area has increase but the average production is modest.

Field pea is frequently used in forage crop mixtures with small grains. Protein content in field pea forage ranges from 18-20%. Pea interseeded at 60-100 pounds per acre with a small grain like oat will increase the protein concentration of the mixed forage by 2-4% points and relative feed value by 20 points.

The crop is destroyed by 17 insect pests which are lowering down the yield of crop both qualitatively and quantitatively. Prasad *et al.* (1983) reported 19 species of insect pest of which 8 could be considered of major significance. Kushwaha (2002) recorded 9 insect pests on pea crop at different stages of its growth and estimated losses ranged 13 to 17 percent at Jabalpur Madhya Pradesh, were as under.

Insect pests are among the main biotic constraints which causes huge losses worldwide. Among the biotic constraints, *i.e.*, insect pests like pea leaf miner (*Chromatomyia horticola* Goureau), pea aphid (*Acrythosiphon pisum*), Pod borer complex (*Helicoverpa armigera* (H), *Lampides boeticus* (L) and *Etiellazinkenella* Tr.) and thrips (*Caliothrips indicus* Bagnall) often causes substantial losses to the crop.

A large number of insect pests damage field pea at different phases of crop growth. Both abiotic (temperature, humidity and rainfall) and biotic factors (parasites and predators) are recognized as important factors for the multiplication of insect's pests influencing their population. Scanty information is available on the pest's status, crop losses and management of crop pests of pea. Therefore, to update this information present study is proposed.

Materials and Methods:-

The mean pest populations during different weather weeks were subjected sample to correlation studies considering studies considering weather parameters like that mean minimum and maximum temperature (°c) and minimum and maximum relative humidity (%) as the independent variables and the density of insect pests as the dependent variable.

Regression equations were worked out for insect populations showing correlation with weather parameters. Correlation and regression of the biotic factors on major insect pest populations were worked out by using the following formula as suggested by Snedecor and Cochran (1967).

$$\text{Correlation 'r'} = \frac{\sum xy - \frac{\sum x \cdot \sum y}{n}}{\sqrt{\left\{ \sum x^2 - \frac{(\sum x)^2}{n} \right\} \left\{ \sum y^2 - \frac{(\sum y)^2}{N} \right\}}}$$

The following formula were used for various estimation, population were worked out by using the following formula.

Regression $Y = a + bX$

Where, a=Intercept

b=Regression coefficient

R²=Coefficient of multiple determination

Test of significance of correlation coefficient

$$t = \frac{r}{\sqrt{1-r^2}} \sqrt{n-2}$$

Regression $Y = a + bX$

b = Regression coefficient

R² = Coefficient of multiple determination

Result and Discussion:

Incidence of aphid

The data presented in table .1 revealed that, the population of aphid appeared on crop in the fourth week of December (three week after sowing) and built up in

population started from 52nd standard week means fourth week of December and increased up to the third week of January and declined gradually till the crop was matured in March. The population of aphid ranged from 1.25 to 65.69 /10 cm apical twigs. The peak activity of aphid (65.69/10 cm apical twigs) was recorded in the 4th standard week or third week of January.

The incidence of aphid started when maximum and minimum temperature was 23.8°C and 7.2°C and morning relative humidity was 83% and evening relative humidity was 43%. The aphid population increased gradually and reached to its peak (65.69 aphid/10 cm apical twigs) at 24.6°C maximum, 8.6°C minimum temperature and 86 per cent morning relative humidity and 49% Evening relative humidity. The data presented in table 4.2 revealed that the maximum temperature showed negative significant correlation ($r = -0.581$) and morning vapour pressure shows significant negative correlation ($r = -0.453$) with the aphid population, whereas, minimum temperature and rainfall showed negative correlation ($r = -0.453$ and $r = -0.153$) with the aphid population.

Biswal and Patel (2015) identified aphid, whitefly and thrip's were the first to infiltrate and colonize the field crop, followed by leaf miner, and remained active until the crop was harvested.

Dixon and Harrington (2000) found that the number of aphids fluctuated in lockstep with the temperature from January to July. Cool temperature in January to February results in large number of aphids population. Their conclusion are consistent with the current studies, which shows that peak populations were observed at a minimum temperature of 9.7°C.

Population of *Acrythosiphonpissum* indicated more affinity with evening relative humidity with r value of 0.398 and regression equation obtained was $y = 0.872x - 7.445$. Whereas Wale (2011) observed negative correlation with aphid population and relative humidity.

Melease and Singh (2012) reported negative correlation of rainfall with aphid population ($r = -0.98$) means when rainfall decreased, the aphid population increased at $P < 0.05$. Negative correlation was also found between aphid population and minimum temperature ($r = -0.453$) but it was non-significant. Significant negative correlation was found between maximum temperature and aphid population.

Correlation was observed between aphid population and wind speed, morning relative humidity, evening relative humidity and it was positively correlated but was found non-significant. Correlation was observed between aphid population and morning vapour pressure so it was negatively correlated and found significant.

Bhadoria (1993) Chakraborty & Dutta (1999), Kushwaha (2002), Mittal & Ram (2007), and Singh & Mishra (2013) have reported the incidence of pod borers in green peas. However, under Jabalpur conditions the borer's incidence in green peas was almost absent.

Tomar *et al.* (2004) reported the most favourable period of insect pests from first week of December to the second week of January for green pea. Relative humidity and rainfall were favourable for the development of almost all the pest (except *spodopteralitura*). The maximum and minimum temperatures were negatively correlated with the population of all the insect pests.

Positive correlation was found between minimum temperature and percentage pod damage by pod borer. Negative correlation was found between percentage pod damage by pod borer and Maximum temperature, rainfall, relative humidity and was found to be significant. The peak population of gram pod borer occurred in December and January and the temp. And R.H. showed positive correlation with pest population while rainfall was negatively correlated.




Fig. 1: Impact of Maximum Temperature on aphid




Fig. 2: Impact of morning vapour pressure on incidence of aphid

Incidence of leaf hopper

The observations on population of leafhopper on pea were recorded three weeks after sowing and till maturity of the crop. The data presented in the table.1 revealed that the leaf hopper remained on pea crop throughout the crop season and population varied from 0.23 to 4.96 /six leaves/plants. In the starting the population was low but it increased gradually and reached to its peak in 4th standard weeks *i.e.*, third week of January and then declined.

The incidence of leaf hopper started when maximum and minimum temperature was 23.8°C and 7.2°C and morning relative humidity was 83% and evening relative humidity was 43%. The leaf hopper population increased gradually and reached to its peak (4.96 leaf hopper/six leaves/plants) at 24.6°C maximum, 8.6°C minimum temperature, 86% morning relative humidity and 49% evening relative humidity and rainfall was none and when the morning vapour pressure was 8.4 and evening vapour pressure was 9.4. The data presented in table .2 revealed that the maximum temperature showed negative significant correlation ($r = -0.582$) with the population, whereas, minimum temperature and rainfall showed negative non-significant correlation ($r = -0.363$ and $r = -0.176$) and minimum temperature showed positive non-significant correlation ($r = 0.413$) with leafhopper population.




Fig. 3: Impact of maximum temperature on incidence of leafhopper

Findings of Darandaleet *al.* (2015) are partially supported the present findings they reported that the incidence of leaf hopper was started from the fourth week of sowing *i.e.*, second week of December (0.27 jassid/leaf). Initially the population was low, increased slowly and reached the peak level (8.52 jassids/ leaf) at 12th week after sowing coinciding with 5th week of January.

Incidence of *Helicoverpaarmigera*

The data presented in table.1 revealed that the incidence of *Helicoverpaarmigera* was observed in the third week of December (one month after sowing). The population increased gradually with slight up and down and reached to its peak (0.65 *Helicoverpaarmigera*/plant) in the second week of January (2nd standard meteorological week) thereafter, it was decreased gradually. The population of *Helicoverpaarmigera* on pea crop ranged from 0.05 to 0.65 *Helicoverpaarmigera* /plant throughout the crop period

The incidence of *Helicoverpaarmigera* started when maximum and minimum temperature was 23.2°C and 5.5°C, morning relative humidity was 74% and evening relative humidity was 31% and rainfall was none. The *Helicoverpaarmigera* population increased gradually and reached to its peak (0.65 *Helicoverpaarmigera*/ plants) at 25.0°C maximum, 11.8°C minimum temperature, 86% morning relative humidity and 49% evening relative humidity whereas rainfall was none. The data presented in table.2 revealed that the maximum temperature showed significant negative correlation ($r = -0.581$) with the population, whereas minimum temperature and rainfall showed negative non-significant correlation ($r = -0.367$ and $r = -0.324$) with *Helicoverpaarmigera* population.




Fig. 4: Impact of maximum temperature on incidence on pod bore

Table .1: Seasonal incidence of major insect pests of field pea

| SMW | Meteorological parameters | | | | | | | | | | |
|-----|---------------------------------|------------------|------------------------------|------------------|---------|----------------------|---------------------|----------------------|---------|-----------------------|---------|
| | Insect pests | | | Temperature (°C) | | Wind speed (km/h r.) | Total rainfall (mm) | Vapour pressure (mm) | | Relative humidity (%) | |
| | Aphid/ 10cm apical twigs/ plant | Pod borer/ plant | Leaf hopper/ 6 leaves/ plant | Maximum | Minimum | | | Morning | Evening | Morning | Evening |
| 51 | 0 | 0 | 0.6 | 23.2 | 5.5 | 1.6 | 0.0 | 7.1 | 6.5 | 74 | 31 |
| 52 | 8.08 | 0.6 | 0.25 | 23.8 | 7.2 | 2.2 | 0.0 | 7.6 | 8.9 | 83 | 43 |
| 1 | 18.8 | 2.75 | 0.36 | 26.5 | 12.4 | 2.1 | 0.5 | 11.4 | 12.9 | 87 | 50 |
| 2 | 32.68 | 3.69 | 0.65 | 25 | 11.8 | 3.2 | 0.4 | 10.9 | 11.3 | 86 | 49 |
| 3 | 40.96 | 3.78 | 0.3 | 25.4 | 8 | 2.6 | 0.0 | 7.7 | 7.9 | 75 | 31 |
| 4 | 65.69 | 4.96 | 0.24 | 24.6 | 8.6 | 3.2 | 0.0 | 8.4 | 9.4 | 86 | 49 |
| 5 | 60.15 | 4.61 | 0.1 | 21.4 | 4.8 | 3.7 | 0.0 | 5.7 | 6.1 | 73 | 31 |
| 6 | 31.25 | 3.25 | 0.05 | 26.4 | 8.9 | 3.2 | 0.0 | 7.7 | 8.1 | 72 | 34 |
| 7 | 20.84 | 1.95 | 0 | 27.7 | 11.8 | 2.7 | 12.6 | 10.0 | 10.8 | 83 | 42 |
| 8 | 18.64 | 1.15 | 0 | 28.6 | 10.6 | 1.9 | 0.0 | 9.2 | 8.2 | 79 | 28 |
| 9 | 4.19 | 0.65 | 0 | 32.7 | 12.4 | 2.9 | 0.0 | 9.8 | 8.9 | 74 | 25 |
| 10 | 3.69 | 0.23 | 0 | 34.8 | 13 | 2.8 | 0.0 | 10.3 | 7.8 | 74 | 20 |
| 11 | 1.25 | 0 | 0 | 32.3 | 15.3 | 3.6 | 6.2 | 12.2 | 10.1 | 78 | 29 |

Correlation coefficient

Table.2 Correlation coefficient between the pest population and weather parameters of field pea

| Insect pests | Rainfall | Temperature | | Relative humidity | | Wind speed | Vapour Pressure | |
|----------------|----------------|----------------|-----------------|-------------------|---------------|----------------|-----------------|----------------|
| | | Max. | Min. | Mor. | Eve. | | Mor. | Eve. |
| Aphid | -0.175 (NS) | -0.581* (S) | - 0.453 (NS) | 0.166 (NS) | 0.398 (NS) | 0.465 (NS) | -0.453* (S) | -0.104 (NS) |
| Gram pod borer | -0.324 (NS) | -0.581* (S) | 0.367 (NS) | 0.309 (NS) | 0.477 (NS) | -0.362 (NS) | -0.127 (NS) | 0.102 (NS) |
| Leaf hopper | -0.176 (NS) | -0.582* (S) | -0.363 (NS) | 0.242 (NS) | 0.515 (NS) | 0.440 (NS) | -0.333 (NS) | 0.075 (NS) |

NS = Non significant

S = Significant

* = Mean of 3 replications

Conclusion

Field pea crop was found to be affected by four major insect pest, aphid and borer appear in early stage while leafhopper appears in first week of January. These insect pests major reason for reducing yield of crop. Seasonal activity of insect's pest indicated high population of *Acrythosiphonpissum* during SMW # 4, high incidence or gram pod borer at SMW #52 and high incidence of leafhopper during SMW # 4. Hence the last week of January was the most crucial period for application of insecticidal measure.

References:

- Chaukikar K, Marabi RS, Das SB, and Bhowmick AK. 2016. Pest succession and population dynamics of major insect pests of Niger (*Guizotia abyssinica* (L.F.) CASS). Progressive research - An international journal, Volume11(Special II): pp 986-989.
- Chakraborty A and Dutta SK.1999. Population build-up of thrips *M. distalis* in peacrop. J. Agric. Sci. Soc. North East India, 12(1): 46-51.
- Darandale SA, Pandya HV, Patel SD, and Patel SM. 2015. Population dynamics of major sucking pests infesting niger and their relation to weather parameters, International Journal of Plant Protection, 8(1), 61-64.
- Dixon, A.FG. and Harrington, R. 2000. Causes of regional and early variation in pea aphid numbers in Eastern England. Rev. Agril. Ent, 88: 281.
- Honda, K. 1989. Seasonal changes in population density of the pea aphid *Acyrtosiphon pisum* Harris in pea fields and environmental conditions for the emergence of alate virginoparae. Annual Report of the Society of Plant Protection of North Japan, (40): 110-111.
- Janu Arun, Ombir, Dahiya KK, Jakhar Pritish. 2017. Population dynamics of leafhopper, *Amrascabigutullabigutulla* (Homoptera: Cicadellidae) in upland cotton (*Gossypium hirsutum* L.), Journal of Cotton Research and Development 2017, Vol.31 No.2 pp.298-304.
- Kushwaha, K. 2002. Succession of insect pests of pea. M.Sc. (Ag.) Thesis, submitted to J.N.K.V.V, Jabalpur, M.P.
- Kushwaha, K. 2002. Succession of insect pests of pea. M.Sc. Thesis, JNKVV, Jabalpur. Martinovich .1993. A study of population dynamics of the gram pod borer *Helicoverpa armigera* (Hub.) in pea seed production. Bulletin Zoldseger Mesztesi Kutata Intezet ,16:115-130.
- Melesse T and Singh SK.2012. Effect of climatic factors on pea aphid, *Acyrtosiphon pisum* population and its management through planting dates and biopesticides in field pea. Journal of Agricultural Technology, 8(1): 125-132. Mittal V and Ram U.2007.
- Mittal Vishal and Ram Ujagir .2007. Succession of insect pests associated with pea crop (*Pisum sativum* Linnaeus) at Pantnagar, India. En vi. Eco, 25 (4): 1030-1035.
- Panday A K, Bisen R, Jain S and Sahu R. 2019. Effect of climatic factors on population intensity of insect pests damaging niger crop. XIX International Plant Protection Congress IPPC2019 Hyderabad, Telangana. P. 278
- Prasad, D., Singh, K.M. and Katiyar, R.N. 1983. Succession of insect pest in early maturing high yielding variety of pea (*Pisum sativum* Linn.). Ind. J. Ent, 45(4): 451-455.

- Scnedecor GW and Cochran WG.1967. Statistical methods. 6th ed. Ames: Iowa state university press, 3(12): 66-67.
- Singh N, Panday DK and Dikshit HK .2002. Status of germplasm its management and utilization in pulse crop. Farmers Forum, 2(4):23-27. 42
- Tomar SPS, Dubey OP and Tomar R. 2004. Succession of insect pests on green pea. JNKVV Research Journal, 38(1):82-85.
- Vaishampayan, S.M. and Veda, O.P. 1980. Population dynamics of gram pod borer [*H. armigera* (Hub)] and its outbreak situation on gram [*Cicer arietinum* (L)] at Jabalpur. Ind. J. Ent. 42 (3): 453- 459.
- Wale M.2011. Population dynamics of the pea aphid, *Acyrtosiphonpissum* (Homoptera: Aphididae) on field pea in Northwestern Ethiopia. [http://dx. doi. org/10.1017/S174275840001521](http://dx.doi.org/10.1017/S174275840001521)

UNDER PEER REVIEW

UNDER PEER REVIEW