

## Review Article

# Biostimulatory Impacts of Seed Priming through Botanical Extracts on Crop Production-A Critical Review

## ABSTRACT

Rapid human population increase, food scarcity, and climate change are some of the global challenges in the current scenario. Among these, climate change adversely impacts crop production by creating abiotic stresses such as drought, salinity, and metal toxicity. Seed emergence, the critical stage in crop development, is highly vulnerable to these abiotic stresses. Therefore, short-term approaches to increase the sprouting of seeds and the initial growth of plants were investigated in advance as a solution. Seed priming is a pre-sowing treatment of seeds which is considered a low-cost, eco-friendly, and sustainable technique to promote seed germination and the initial growth of plants. Though various priming methods are available, this review discusses only botanical extracts (both seaweed and plant extracts) used as priming agents in seed priming. Among the seaweed extracts, green and brown algae species such as *Ascophyllum nodosum*, *Sargassum spp.*, and *Ulva spp.* enhanced the crop performances even under stress conditions. In addition, leaf extracts derived from *Moringa oleifera* and *Azadirachta indica* widely used to increase the productivity of vegetable and field crops under stress and non-stress environments. Furthermore, in this review we discussed about priming mechanism, different extraction methods and bioextracts behavior under diverse agroecosystems are discussed in detail.

**Keywords:** *Abiotic stresses, antioxidant, bioextracts, biostimulant, seed priming*

## 1. INTRODUCTION

The challenge of a rising population affects the world. Due to population expansion, more individuals would require assistance with food security and supply. However, climate change, environmental pollution and abiotic stressors significantly impact on seed germination, emergence, seedling vigour, and crop productivity [1]. These undesirable impacts on crop production and performance significantly influence the food and agricultural systems. Several approaches, such as conventional breeding and advanced technologies (e.g., genetic engineering, mutation breeding, and polyploidy breeding) are being assessed to enhance crop production under environmental challenges. However, these techniques have significant drawbacks, including sizeable human resource requirements, biosafety apprehensions, and ethical concerns [1]. Thus, it is necessary to develop simple, efficient, and sustainable technology to solve these issues [2]. In this regard, seed priming is an alternative approach to getting around these constraints and increasing the capacity of plants to withstand stress [3,4].

Seed priming is a simple and an efficient method that would enhance plants' physiological and biochemical responses to abiotic stresses; thus, regulates hydration of seeds that permits pre-germination of metabolic activities without causing the radicle to emerge [5]. Basically, seed priming improves the seed germination or seedling growth rate through entrusting uniformity of seedling establishment [6]. Besides, priming process ensures early emergence, efficient water usage, mitigation of excessive fertilizer usage, promote roots growth, initiate the growth of reproductive organs, early flowering and maturity, resistance to abiotic stresses and soil-borne pathogens [2,6]. According to the way of conducting priming procedure, seed priming methods can be classified as hydro-priming, halo-priming/chemo-priming, osmo-priming, osmo-hardening, on-farm priming, matrix priming, nutri-priming, nano-priming, bio-priming, thermo-priming, UV priming and hormone priming [2,7]. Hydro-priming is achieved by soaking seeds in clean water and re-drying them to their initial moisture content before planting [8]. In this regard, hydro-priming shortens the lag phase of the germination process and facilitates rapid germination for vigorous development [9,10]. In osmo-priming, a low water potential osmotic solution is used to soak seeds rather than pure water [1]. Numerous materials are used in osmo-priming, including mannitol, sorbitol, glycerol, and inorganic salts [7].

Osmo-hardening includes re-drying of seeds after soaking them in tap water for 24 hours, then hardening them with  $\text{CaCl}_2$  and  $\text{KCl}$  solutions [2]. In solid matrix priming a solid or semi-solid media is employed instead of a liquid medium. This method involves combining seeds with a solid or semi-solid substrate with certain quantity of water [9]. Solid matrix priming would regulate water absorption by properly adjusting of moisture content [7]. In nutrient priming, seeds are exposed to a particular concentration of macro or micronutrients for a predetermined duration before planting. One of the novel seed priming techniques is nano-priming which uses nanoparticles (NPs), such as zinc oxide, iron oxide, titanium dioxide, silver nano-particles etc. [2]. In bio-priming, specific microorganisms or bioactive molecules are used for seed treatment [7]. Thermo-priming involves treating seeds at various temperatures while minimizing the influence on their critical physiological functions [7]. In hormonal priming plant growth regulators like auxins, gibberellins, kinetin, salicylic acid, abscisic acid, ethylene, and ascorbic acids are widely utilized [9]. Apart from these methods, in chemical priming seeds are treated with either organic (e.g., organic acids, botanical extracts, chitosan, polyamines, mannose, and trehalose) or inorganic (e.g., sodium nitroprusside, and sodium hypochlorite) substances [11]. As organic substances, botanical extracts are significant for seed priming. The main types of botanical extracts are being obtained from seaweed and plant extracts. Hence, in this review we would discuss the numerous botanical extracts, extraction methods, and their significance under abiotic stress conditions.

## **2. SEAWEED EXTRACTS (SWEs) USED IN SEED PRIMING**

Seaweed, often known as marine macroalgae, comprises a large portion of the world's marine life resources [12]. Most of their distribution recorded in shallow coastal waters, estuaries, and backwaters; but relatively small quantity has been found on solid surfaces like rocks, dead coral, and pebbles [12]. Approximately 10,000 species of macroalgae exist in globally which account for 10% of marine productivity [13]. According to presence of pigments, marine macroalgae would divide into three categories namely; (a) Chlorophyta or green algae, (b) Rhodophyta or red algae, and (c) Phaeophyta or brown algae respectively [14]. Brown algae mainly encompasses chlorophyll a and c with carotenoids and fucoxanthin while green algae contain pigments that are identical to the chlorophyll a, b and carotenoids which found in terrestrial plants [14]. Red algae contains both chlorophyll a, b, and phycoerythrin pigment which is responsible for the color of the red algae [15, 16].

SWEs have been used as a seed treatment to improve the seed growth and development [17]. These are widely recognized to include a number of bioactive compounds [18]. A variety of biologically active components including polyphenols (e.g., phenolic acids, flavonoids, cinnamic acid, isoflavones, benzoic acid, lignans, quercetin), polysaccharides (e.g., galactan, fucoidan, alginate, laminarin), proteins (e.g., lectins), polyamines, pigments, free amino acids, vitamins, micronutrients (e.g., B, Co, Cu, Fe, Mn, Mo, Ni, Si, Zn), macronutrients (e.g. Ca, K, Mg, Na, P, S), and natural phytohormones (e.g., cytokinins, auxins, gibberellins, abscisic acid), antioxidants, osmo-protectants betains, antimicrobial compounds are consist in SWEs [16, 18, 19]. Furthermore, seaweeds have elicitor chemical precursors that aid for seed germination [20]. Several SWEs used for seed priming are summarized in table 1.

**Table 1:**Seed priming through seaweed extract

No	Seaweed extract type	Crop type	Priming condition	Impact of seed priming	Country of study	Stress condition	Reference
1	<i>Ascophyllum nodosum</i>	Spinach	Seeds were soaked in solutions ranging from 0.15% to 1.2%	A concentration of 0.3% treatment showed the highest germination percentage and seedling	Brazil	Thermal stress	[21]
		Moth bean	Seeds were soaked with various concentrations (0%, 0.01%, 0.05%, 0.10%, 0.50% and 1.0%) for different periods ranging from 0 to 24 hours	Improved the seed germination, shoot and root length, fresh and dry weight, free radical scavenging, whereas alpha-glucosidase inhibitory action at low concentrations	India	Non-stress	[14]
		Onion	Seeds were soaked for 6 hours in different concentrations ranging from 0ppm to 7500ppm	Improved the growth of onion seedlings	India	Non-stress	[22]
		Pepper	Seeds were primed for 1 to 3	Increased total germination rate	Turkey	Non-stress	[23]

			days in an incubator at 20°C using concentrations ranging from 1:1 to 1:1000 dilutions of seaweed extract and water	and decreased the mean germination time			
	<i>Colpomeniasinuosa</i> <i>Sargassum vulgare</i>	Fenugreek	Seeds were soaked for 12 hours with various concentrations ranging from 5% to 25%	Enhanced the seed germination, biochemical constituents under low concentration	Egypt	Non-stress	[24]
	<i>Codium taylorii</i> <i>Pterocladia capillacea</i>	Radish	Sterilized seeds Were soaked for 2 hours water	Enhanced radish water content and increased plant length significantly	Egypt	Salinity Stress	[25]
	<i>Codium tomentosum</i>	Wheat	Seeds were soaked in Different concentrations ranging from 10% to 50% at 12 hours	Increased shoot length, root length, fresh and dry weights of seedlings	Egypt	Non-stress	[26]
2	<i>Cystoseira compressa</i>	Cowpea/ Maize	Seeds were soaked in a concentration of 20 g L <sup>-1</sup> for 2 hours at 25 °C	Increased the germination rate of both crops, vigorous coleoptile elongation of maize, increased biomass weight and guaiacol peroxidase activity	Egypt	Salinity stress	[27]
3	<i>Ecklonia maxima</i>	Okra	Seeds were immersed for 24 hours in Kelpak® solution under 1:20, 1:40 and 1:100 v/v) levels	Seed germination parameters were increased from 1:100 solution			[28]

		White foxglove	Seeds were treated with different biostimulants for 48 hours at 25 °C	Enhanced the seed germination and seedling growth	South Africa	Heat stress	[29]
		Maize	Soaked seeds in aqueous solution of seaweed extracts for 24 hours	Induced seed germination and shoot weight	Poland	Non-stress	[30]
4	<i>Ecklonia maxima</i> <i>Jania adhaerens</i>	Tomato	Seeds were immersed in concentrations of 2.5, 5.0, and 10.0 mg mL <sup>-1</sup> overnight at room temperature	Extracts provide a sound protection from controlling of <i>Rhizoctonia solani</i> on tomato plants	Italy	Non-stress	[31]
	<i>Gracilariatextorii</i> J. Agardh and <i>Hypneamuscififormis</i> (Turner)	Brinjal  Chilly and Tomato	Seeds were soaked in the various ratios (1:2, 1:4, 1:6, 1:8, 1:10, 1:20,1:30) for 12 hours	Increased seed germination and crop yield		Non-stress	[32]
5	<i>Gracilaria edulis</i>	Chilli	Seeds were primed for various time periods (1:25 for 48 hours, 1:5 for 24 hours and 1:5 for 72 hours)	Improved the vigor index and seedling weight	India	Non-stress	[33]
6	<i>Jania rubens</i>	Maize	N/A	Germination rate, seedling length and weight, and number of lateral radicles were increased	Morocco	Non-stress	[34]
7	<i>Kappaphycusalvarezii</i>	Rice	Seaweed extracts were added to rice seeds at concentrations	Rice seed germination and seedling vigor were increased at	India	Non-stress	[35]

	<i>/Gracilaria edulis</i>		of 2.5%, 5%, 7.5%, 10% and 15% for 24 hours	2.5% and 5% concentrations			
8	<i>Laurencia obtusa</i>	Cowpea/ Maize	Seeds were soaked for 2 hours at 25 °C in a concentration of 20 g L <sup>-1</sup>	Higher germination rate vigorous coleoptile elongation, higher biomass weight and guaiacol peroxidase activity were observed in both crops	Egypt	Salinity stress	[27]
	<i>Padina gymnospora</i>	Tomato	Seeds were treated with different concentrations (0.2, 0.4 and 1.0 %) for 24 hours	Enhanced germination, rate of germination, lower mean germination time, high germination index, greater seedling vigor, greater plumule and radicle length	Mexico	Non-stress	[36]
9	<i>Solieria spp.</i>	Kale	Seeds were soaked for 22 hours at 20°C	Seed primed with seaweed extract has not interfered with seed health	Brazil	Non-stress	[37]
	<i>Sargassum polycystum</i>	Red gram	Seeds were treated with different concentrations of water extracted and alcohol extracted seaweed (0.1, 0.25, 0.50, 0.75, 1.0 and 1.5%) for every 12 hours	The lowest concentration showed higher seed germination, shoot length, root length, fresh weight and dry weight	India	Non-stress	[38]

<i>Sargassum tenerrimum</i>	Tomato	Seeds were immersed in a variety seaweed aqueous extract concentration, including 0.2%, 0.4%, 0.6%, 0.8%, and 1.0%	Seed germination rate was high at 0.8%	India	Non-stress	[39]
<i>Sargassum vulgare</i>	Wheat	Seeds were soaked in different concentrations ranging from 10% to 50% for 12 hours	Increased seed germination, growth and yield	Egypt	Non-stress	[26]
	Tomato	Seeds are treated with 0.2% and 0.5% concentration of seaweed extract and then exposed for 14 days photoperiod	Increased germination percentage and the growth of the radicle	Morocco	Salt stress	[40]
	Wheat	Seeds were treated at different concentrations (0.2, 0.5, 25 and 50 %) and kept under photoperiod for 14 days	Enhance the germination rate with lower mean germination time, greater seedling vigor and greater radicle length	Tunisia	Salt stress	[41]
<i>Sargassum vulgare</i> , <i>Colpomenia sinuosa</i> , <i>Pandia pavonica</i>	Pea	Sterilized seeds were soaked in seaweed extracts or distilled water for two hours	Increased the production of photosynthetic pigments, new peptides, and improved germination, seedling length, and fresh weight		Salt stress	[42]
<i>Sargassum wightii</i> Grev.	Green gram	Seeds were soaked with	In comparison to the high		Non-stress	[43]

		concentrations of 0.5%, 1.0%, and 2.0%, for 6, 12 and 24 hours	concentration (2.0%), 6 hours of soaking in the 0.5% concentration led to higher seed germination and earlier growth			
	<i>Sargassum wightii</i>	Wheat	Seeds were soaked with different concentrations ranging from 5% to 50% for 12 hours	Increased the germination rate, number of lateral roots, shoot length, number of kernels per plant, and kernel length	India	Non-stress [44]
	<i>Thalassiahemprichii</i>	Green gram	Seeds were treated with 50 ppm and 100 ppm of seagrass extract	In comparison to the control treatment, the root: shoot ratio has risen for all of the treatments	India	Non-stress [45]
	<i>Ulva</i> , <i>Polysiphonia</i> , <i>Cladophora</i>	Garden cress	Seeds were treated with concentration of 0.5, 2.5 and 10%	Chlorophyll content was higher compared to non-treated sample		Non-stress [19]
	<i>Ulva fasciata</i> <i>Padina gymnospora</i> <i>Gracilaria edulis</i>	Bell pepper	for 24 hours, seeds were soaked in seaweed extract at various concentrations (2, 4, 6, 8 and 10%).	The phytochemical content and germination were enhanced at 8% concentration	India	Non-stress [46]
12	<i>Ulva fasciata</i> , <i>Padina gymnospora</i> , <i>Gracilaria edulis</i>	Chilli	Seeds were treated in various concentrations range from 2% to 10% for 24 hours	Phytochemical content and germination were enhanced by <i>Padina gymnospora</i> at 8% concentration level	India	Non-stress [15]

Onions	Solid matrix priming using a 2:1:3 mixture of seed, vermiculite, and seaweed extract were carried out at 15 °C for 2 days, seaweed extract was applied at 5% concentration	Enhanced the seed germination and seedling quality		Drought and salinity stresses	[47]
Carrot	N/A	Enhanced the germination fresh and dry weight of seedling	Turkey	Salinity stress	[48]
Tomato	Seeds were soaked for 24 hours at 25°C in a 1 mg mL <sup>-1</sup> solution	Increased the tomato fresh weight, glycine betaine, total phenols and soluble sugars concentration, significantly decreased the hydrogen peroxide (H <sub>2</sub> O <sub>2</sub> ) concentration	Morocco	Salinity stress	[49]
Cowpea and Maize	Seeds were soaked for 2 hours at 25 °C in a concentration of 20 g L <sup>-1</sup>	Observed higher germination rate of both crops, vigorous coleoptile elongation of maize, increased biomass weight and guaiacol peroxidase activity	Egypt	Salinity stress	[27]
Tomato	Soaked for 24 hours	Enhanced germination, rate of germination, lower mean germination time, high	Mexico	Non-stress	[36]

	<i>Ulva lactuca</i> , <i>Caulerpa scalpelliformis</i> <i>Sargassum plagiophyllum</i> <i>Turbinariaconoides</i> <i>Padina Tetrastromatica</i> <i>Dictyotadichotama</i>	Green gram ( <i>Vigna radiata</i> ) seaweed extracts	Seeds were soaked for 12 hours with concentrations ranging from 0.1% to 0.5% seaweed extracts	germination index, greater seedling vigor, greater plumule, radicle length  The 0.3% concentration has the lowest germination percentage and maximum germination (100%) has 0.4% <i>Caulerpa scalpelliformis</i> and 0.1% <i>Padina tetrastromatica</i> extractions	India	Non-stress	[50]
14	<i>Ulva linza</i> <i>Corallina ofcinalis</i>	Wheat	Seeds were soaked for 12 hours in aqueous extract at different concentrations range from 5% to 30%	Increased seedling growth at low concentration	Egypt	Non-stress	[51]

UNDER PEER REVIEW

The commercial SWEs sector employs various exclusive extraction methods to rupture the cells and release beneficial components into the extract. These extraction techniques are further classified as enzyme-assisted extraction, microwave-assisted extraction, pressurized liquid extraction, supercritical fluid extraction, ultrasound-assisted extraction, soxhlet extraction, alkali extraction, acid extraction, and cell bust extraction [13, 18, 19]. In this regard, a variety of solvents, including ethanol, acetone, methanol-toluene, methanol, petroleum ether, ethyl acetate, dichloromethane, and butanol can be utilized for an effective extraction process. However, these procedures are sometimes considered costly and hazardous [19]. Hence, boiling and soaking extraction techniques with distilled water are used to overcome conventional extraction techniques' drawbacks; which are eco-friendly and require no organic solvents [19].

### **3. PLANT EXTRACTS (PEs) USE IN SEED PRIMING**

Another effective method to promote germination is soak the seeds in PEs before planting [52]. Numerous chemical compounds identified in various plant species that improve the growth, development, and productivity of crops grown under adverse conditions. PEs modify the physiological and biochemical processes such as stomatal conductance, phytohormone metabolism, absorption of water and nutrients, antioxidant defensive systems, and photosynthesis [53]. Different PEs used for seed priming are summarized in table 2.

**Table 2:** Seed priming through plant extracts

No	Plant extract type	Crop type	Priming condition	Impact of seed priming	Stress Condition	Country of study	Reference	
1	Moringa leaf extract (MLE)	Maize	Seeds were soaked (0.25, 0.5 and 1.00 g) at the ratio of 1:10 concentrate for 12 to 20 hours	Improved the germination and seedling growth	Non-stress	Egypt	[52]	
		Rice	Seeds were soaked in an aerated solution of MLE at 3% for 8 hours	Improved the speed and spread of emergence of seedlings (e.g., Time to start emergence, emergence index, mean emergence time, final emergence percentage)	Drought stress	Pakistan	[53]	
				Seeds were soaked in the MLE (seed weight: MLE, 1:5) at room temperature in dark conditions under continuous shaking, for 24 hours	Improved seed germination and the plant growth	Arsenic stress	India	[58]
		Onion	Seeds were soaked in 12.5% and 25% of MLE for 6 and 12 hours respectively	Seeds soaked in MLE for 6 hours gave higher seed emergence	Non-stress	Nigeria	[60]	
		Wheat	Seeds were soaked in MLE 1:10 concentration for 12 hours	Improved seed germination and seedling growth	Salinity stress	Libya	[61]	

Raddish	Seeds soaked in 3 and 5% for 6 and 9 hours under continuous aeration	Increase growth with improved biochemical and antioxidant attributes	Non-stress	Pakistan	[62]
Cucumber	Aerated fresh moringa leaf extract for 18 hours	Improved germination and seedling growth	Non-stress	Nigeria	[63]
Pea seeds	Seeds were primed with 3% moringa leaf extract and in combination of magnetized water and moringa leaf extract for 12 hours	Improved germination capacity and seedling vigor	Non-stress	Pakistan	[64]
Cowpea	Primed with 5% and 2% moringa leaf extract for 12 hours	Increase germination percentage with better root and shoot length	Non-stress	Pakistan	[65]
Linola	Primed with 3.3% moringa leaf extract for 12 hours	Improve crop growth and yield	Non-stress	Pakistan	[66]
<i>Sesamum indicum</i> L.	Primed with 2, 4 and 6% solutions of fresh moringa leaf extract for 6 hours	Improve the seedling emergence and growth	Non-stress	Pakistan	[67]
Spring maize	Priming 3% moringa leaf extract for 18 hours	Increased root and shoot lengths, the ratio of root:shoot, length and maximum seedling fresh and dry weights	Under cool condition	Pakistan	[68]
Hybrid maize	Primed with 1:30 and 1:40 moringa leaf extract for 18 hours	Higher emergence rate and better early seedling	Non-stress	Pakistan	[69]

				growth			
2	Neem leaf extract (NSE)	Bread Wheat	Seeds were soaked for 12 hours	Enhanced plant growth and reduced nematode infestation	Non-stress	Pakistan	[57]
		Rice	Seeds were primed with NSE concentrations 25%, 50% and 100% for 24, 48, and 72 hours respectively	Increased the seed germination	Non-stress	Sri Lanka	[70]
		Chickpea	Seeds were primed in 5% and 10% NSEs for 10 hours	10% NSE significantly affected for the emergence percentage, primary branches per plant, pods/plant, seeds per pod, and seed yield per plant	Non-stress	India	[59]
		Lentil	Seeds were primed in 50% NSE solution for overnight	Observed higher germination percentage, number of branches per plant, seeds per pod, 1000 seeds weight, and yield	Non-stress	India	[55]
3	Garlic extract	Brinjal	Seeds were primed in 100, 200, and 300 $\mu\text{g ml}^{-1}$ for 4, 8, and 12 hours respectively	Enhanced the seed germination and initial growth	Non-stress	China	[54]
4	Sorghum water extract	Camelina	Seeds primed in 5% solution for 24 hours at 25°C	Increased the emergence percentage and ionic homeostasis. Enhanced seedling growth, biomass	Salinity stress	Pakistan	[56]

		Sunflower/ Maize	Seeds were primed with 1%, 1.5%,2%,2.5% and 3% concentrations for 8 and 14 hours, respectively	production, and chlorophyll content Primed seeds were showed higher seedling development	Non-stress	Pakistan	[71]
6	Arappu extract	Okra	Seeds were kept in 3% solution for 12 hours	Increased the germination and seedling vigor	Non-stress	India	[72]
7	Sugar beet extract	Wheat	Seeds were primed at different concentrations (10%-50%) for 12 hours	Increased biomass, photosynthetic pigments, and antioxidant defense mechanism	Drought stress	Pakistan	[73]
	<i>Eclipta alba</i>	Sorghum	Seeds soaked in the extract for 6 hours	Increase of yield, emergence and suppression of pathogenic fungi	Non-stress	Denmark	[74]
	Prosopis ( <i>Prosopis juliflora</i> )	Black gram	Seeds were soaked in leaf extracts at 1% solution four hours	Enhance the seed quality characters	Non-stress	India	[75]
		Pigeon pea	Priming with 2% leaf extract for one hour soaking	Enhanced the seed and seedling quality characters	Non-stress	India	[76]
	Marigold	Pepper ( <i>Capsicum baccatum</i> var. <i>pendulum</i> )	18 ml of priming solution at 25 °C for 24 hours in the dark	Improved the emergence, and in the repair of undeveloped seeds	Non-stress	Turkey	[77]

	<i>Willd.</i> )					
White musale, Periwinkle, Neem, Wood apple, Lantana and White cedar	Tomato	Seeds soaked at room temperature 2% and 4% solution four hours	Improvement in seed and seedling growth parameters in presence of pathogen	Non-stress	India	[78]
<i>Acacia nilotica</i> (L.) and <i>Sapindus mukorossi</i> (L.)	Peanut, chickpea, sunflower and okra	Seeds primed with for 5, 10, 20 and 40 minutes	Control of root rot fungi and growth and productivity	Non-stress	Pakistan	[79]
<i>Cyperus esculentus</i> (Della) <i>Axonopus compressus</i> (Itsit), <i>Convolvulus arvensis</i> (Lehli) and <i>Parthenium</i>	Rice	Seeds soaked in the weed extract for 30 minutes	increased germination rate and germination percentage	Non-stress	Pakistan	[80]

*hysterop  
horus*  
(Partheni  
um)

Aloe  
vera,  
*Moringa  
olifera* or  
sugar  
beet  
aqueous  
extracts

Lentil (*Lens  
culinaris*Medi  
k)

Seeds soaked in aerated for 14  
h using five concentrations (1%,  
2%, 3%, 4% and 5%)

Improved seedling growth

Chilling stress

Pakistan

[81]

pungam  
(*Pongami  
a  
pinnata*)  
and  
arappu  
(*Albizia  
amara*)

Cluster bean

Soaked in 1and 2% for 3 hours  
and 6 hours

Enhanced germination,  
vigour and field  
emergence

Non-stress

India

[82]

9 *Cyperus  
rotundus  
L.*

Drumstick

Seeds were soaked in four  
aqueous extracts of *Cyperus  
rotundus* concentrations (0%,  
25%, 50% and 100%)

Seed priming with 50%  
concentartion resulted in  
an increase in initial  
shoots length and  
photosynthetic pigments  
accumulation

Non-stress

Brazil

[83]

10 Tomato  
leaf

Sugar beet

Seeds were primed for 2 days at  
25 °C

Increased seed  
germination

Non-stress

Turkey

[84]

	extract						
11	Carrot leaf extract	Pea	Seeds were primed 75 ml L <sup>-1</sup> concentration for 10 hours	Observed higher plant height	Drought stress	Egypt	[85]
12	Chicory, clerodendron, noni and calotropis leaf extract	Maize	Seeds were soaked in 5%, 10% and 15% chicory extract and 1, 2 and 3 % of clerodendron, noni and calotropis for 12 hours	Increased the seedling vigor and yield parameters	Non-stress	India	[86]

---

According to the literature, several PEs are widely used in crop cultivation such as *Allium sativum* (Garlic), *Zingiber officinale* (Ginger), *Azadirachta indica* (Neem), *Codiaeum variegatum*, (Crouton), *Moringa oleifera* (Moringa), *Datura stramonium* (Jimsonweed), *Aloe vera* (Aloe), *Ricinus communis* (Castor), *Allium cepa* (Onion), *Helianthus annuus* (Sunflower), *Sorghum bicolor* (Sorghum) etc. [54,55,56,57,58]. PEs are rich in macro and micronutrients, antioxidants, amino acids (e.g., alanine, glycine, leucine and proline), ascorbate, zeatin, growth hormones, vitamins (e.g., B-complex, C,  $\beta$ -carotene,  $\alpha$ -tocopherol), allelochemicals, enzymes (e.g., amylase, catalase, lipase, oxidase, superoxide dismutase) and other organic compounds (e.g., triglycerides, triterpenoid, gibberellin, potassium sorbate and salicylic acid) [52,53,59]. However, PEs would have a variety of compositions depending on the plant species.

PEs are commonly prepared as fresh or dry basis [52, 53]. For instance, Khan et al. [53] prepared the plant extract using fresh mature leaves. According to the described method, the extract was extracted from overnight frozen leaves using locally built equipment, and predetermined concentration was prepared by mixing it with distilled water. Ahmed and El-mahdy [52] described the procedure to prepare the extract by air-drying fresh moringa leaves and to store the dried powders at room temperature. The powdered moringa leaves were immersed in distilled water on a weight/volume basis for 24 hours with periodic shaking. The liquid was filtered through four cheesecloth layers and Whatman filter paper to remove fibre debris. Moreover, in some experiments centrifuging process was also carried out as an additional step [57].

#### 4. MECHANISMS OF SEED PRIMING

There are three main phases in the seed priming procedure namely; seed imbibition (phase I), activation phase (phase II) and growth and cell elongation phase (phase III). In phase I, water uptake occurs according to the water potential gradient [11]. In activation phase, most metabolic and repairing actions occur in the cytosol (e.g., protein synthesis, DNA repairing and stimulation of enzymes and antioxidants). During this phase, water imbibition is also continued and numerous cellular activities such as production of ATP, essential lipids and antioxidants take place [87]. Among these activities, DNA repairing is an essential process which protects the cell from oxidative injuries during seed germination [88]. According to latest literature, germination-related enzymes are activated at a higher rate in this phase. For instance,  $\alpha$ -amylase activity amplified in wheat seeds after the seed was primed with sorghum water extract [88]. Moreover, seed priming boosted protein synthesis by improving rRNA synthesis and ribosome integrity [11]. Simultaneously, antioxidant enzymes such as peroxidase (POD), superoxide dismutase (SOD), ascorbate peroxidase (APX), glutathione reductase (GR) and catalase (CAT) ensure cellular protection from reactive oxygen species (ROS) [8]. Stage III consists of rapid water uptake, and the radicle's protrusion indicates that the germination process has advanced to the development and cell elongation phases [87]. Figure 1 illustrates the seed priming mechanism.




Figure 1: Seed priming mechanism

#### 5. SEED PRIMING THROUGH BOTANICAL EXTRACTS UNDER NON-STRESS CONDITION

Seed germination is the most crucial step in crop cultivation which is considered as an internal process aided by a series of enzymatic reactions, mainly governed by  $\alpha$ -amylase [28]. Furthermore,  $\alpha$ -amylase

gelatinizes the starch in the endosperm, thus supplying ATP energy for the growing embryo [88]. However, presently, commercial crop growers might face numerous consequences related to poor performances in the seed germination process due to non-synchronized germination, an uneven growth pattern, and insufficient seedling emergence [28]. Therefore, to mitigate these constraints, seed priming with numerous composites has been introduced as a viable approach [88].

As a biostimulant, SWEs can be incorporated into the seed coat through priming, stimulating the seed germination rate and initial growth [28]. SWEs promote seed germination mainly due to the presence of various compounds in the solution. For instance, at different concentrations, SWEs containing various phytohormones would result the germination and initial seedling development process [16]. However, Nonogaki et al. [89] further reported that auxins had not affected seed germination directly, even though they enabled gibberellic acid biosynthesis. Thus, they promoted seed sprouting through activating of  $\alpha$ -amylase [89]. Concurrently, primed seed germination enhancement might be impacted by solution-retention effects in the pre-germination phase [20]. Furthermore, some mineral nutrients in the seaweed extract might positively affect the plant's germination rate and vigor index [33]. Seed priming through seaweed extraction directly impacts the alternation of seed biochemical properties which accelerates seed germination [33]. According to recent findings, the cell membrane permeability has been reduced in seaweed extract-primed chilli seeds. In contrast, electric conductivity is also reduced in the cytosol, resulting from optimum cell functions avoiding damage or solute leakage from membranes [33]. Furthermore, numerous macro and microelements in the SWEs, such as N, P, K, Ca, Mg and Zn, promote initial plant growth and development [25]. These elements act as building blocks of proteins and other essential substances in the cytoplasm, which play a critical role in osmotic adjustment [25]. Similarly,  $\text{Ca}^{2+}$  and  $\text{K}^{+}$  ions included in SWEs catalyze some enzymatic reactions related to seed germination and growth, thus, ultimately enhancing the final yield [36]. The impacts of SWEs on seed priming have been investigated through several experiments (table 2). In addition, SWE-primed seeds contained a higher concentration of antioxidants; thus, enhancing the metabolic activities and mitigating ROS production would affect seed growth and development [33].

Similar like to SWEs, plant extracts also contain numerous bioactive compounds. [52, 56]. However, the plant type and area where these plants are being grown altered the bioactive compounds' concentration and composition [53]. Seed priming through leaf extractions enhanced plant growth under normal and stress conditions [53]. For example, moringa leaf extraction (MLE) consisted of a mixture of mineral nutrients including K, Ca, Mg, vitamin C, and plant growth regulators (e.g., cytokinin), which expedited the cell division, delayed leaf senescence, expanded the leaf area and increased chlorophyll pigments [56]. Furthermore, plants' biochemical attributes would increase after seed priming through MLE due to various allelochemicals, including secondary composites such as phenolic and ascorbic acid [53]. Simultaneously, with the enhancement of chlorophyll content, it has been observed that mineral nutrients in MLE accumulate in the seed embryo during the seed priming process and facilitate the growth and development of plants [28, 53].

## 6. SEED PRIMING WITH BIOEXTRACTS UNDER STRESS CONDITIONS

As a result of extreme environmental factors such as drought, salinity, temperature and trace metal (loid) s physiochemical reactions in the plant cells are highly impacted [90]. For instance, Siddiqui al. [90] reported that salinity stress enhanced leaf chlorosis by accelerating chlorophyllase activity which depreciates pigment proteins or enhances the enzymatic chlorophyll degradation process. When plants are exposed to stress conditions, they would stimulate to react to the stress by producing reactive oxygen species (ROS) such as singlet oxygen molecules, hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), hydroxyl radicals ( $\text{OH}^\cdot$ ), superoxide anion ( $\text{O}_2^\cdot$ );[33,42; Figure 2).




Figure 2: Different cellular substances present in the cytosol after seed priming under stress condition

These ROS are involved in the degradation of the thylakoid membrane and hinder the chlorophyll synthesis process [90]. Furthermore, Szafrńska et al. [91] observed poor expression of several Mg-chelatase subunit-encoding genes, resulting in low chlorophyll synthesis. In addition, under the salinity stress, water absorption, stomatal conductance nutrient translocation and assimilation [56] were hindered, and this leads to the malfunctioning of cellular activities. In addition, higher salinity level has altered the plant's morphological characteristics by reducing leaf size, developing thicker leaves, and dwarf plants [56]. ROS substances down-regulated gibberellin genes, thus, resulting from low gibberellic acid production and reduction of  $\alpha$ -amylase activity [91]. Overall, seed germination and initial plant growth would retard drastically. Apart from ROS, Malondialdehyde (MDA) synthesis is also induced due to lipid peroxidation, and higher accumulation of MDA in plant cells eventually leads to malfunctioning cell functions [92].

However, to combat with ROS compounds plants produce other osmo-regulatory molecules such as proline and glycine betaine [42]. Since proline is an essential amino acid, it makes a defense against oxidative damage that occurs from ROS [93]. Hence, proline accelerates protein hydrolysis and ensures osmotic balance [42]. Further, glycine betaine production in the cytosol also increased and mitigated the most adverse effects of salinity stress [93]. For example, glycine betaine protects cell membranes and proteins, triggers coenzyme activation, and removes excess ROS compounds from the cell [93]. Interestingly, with elevated saline levels, plants would enhance carotenoid production [53]. Hence, salinity stress triggers overexpression of genes relevant to carotenoid accumulation under stress conditions, directly involved in reducing free radicals in the cytosol [92].

Seaweed and plant extracts contain unique bioactive compounds (e.g., growth regulators, micronutrients, osmo-protectants, and amino acids) that are essential to accelerate defense mechanisms under abiotic stress environments [56]. The protection mainly occurs due to numerous minerals, amino acids, proline and glutathione induce antioxidant enzyme production; thus, reducing ROS levels in stressed plants [93]. Moreover, the exogenous application of SWEs and PEs further increases the antioxidant concentration in cells, which mitigates ROS production [53]. In this regard, the most common antioxidant enzymes, namely, POD, SOD, APX, GR and CAT concentration, have increased after the priming, which protects the plants from oxidative stress by scavenging ROS [8]. In addition, previous research details provide evidence that even under abiotic stress circumstances, seed priming with SWEs would mitigate the harmful effects of those adverse conditions [29]. For instance, seed priming with MLE boosted the initial maize growth due to the presence of phenols and ascorbates [69]. Moreover, Tahira et al. [94] further noticed that MDA production had been reduced after the seed priming.

Abiotic stress conditions widely affect the protein synthesis process in plants. For example, salinity stress-induced *Vigna mungo* varieties showed different protein profiles, known as "new stress-proteins" which are highly important to manipulate osmotic potential [93]. The elevated ROS production in cell resulted DNA alternations and produced DNA fragments, resulting in genome instability [95]. The presence of bioactive substances in SWEs would upregulate the polypeptide synthesis by activating specific enzymes [42]. In addition, salinity stress might involve in occurring of mutations, chromosomal rearrangements, and alternations in DNA base arrangement that leads to genotoxic impairment and structural modifications [95]. Hence, seed priming through SWEs ensures DNA, protein, and RNA repair and synthesis throughout germination phase II [42].

## 7. CONCLUSION AND FUTURE PERSPECTIVES

Climate change and other numerous challenging factors negatively influence on the agricultural production. The sustainability of crop production in a changing environment depends on improved seed quality through using environmentally friendly method like seed priming with botanical extracts. Hence, without endangering the environment or human health, it may be possible to promote crop growth and development, boosting of yield and protecting against numerous abiotic stressors. In addition, crop cultivation under stress and non-stressed conditions, seed priming would become more effective tool.

Although seed priming with bio-extracts would have given positive results, further researches are needed to understand the physiological and biochemical impact of bioextracts at seed germination and seedling growth stages. In this regard, future experiments should target on molecular mechanisms

engage in seed priming process. Furthermore, studies on undiscovered mechanisms of bioextracts on growth promoting processes would be essential to increase usage of these extracts in crop cultivation. Future researches on direct role of botanical extracts in mobilizing seed reserves and the partitioning of biomass in seedlings is also required. Moreover, SWEs differ from one another even processing the same raw material using different extraction methods. Hence, further crop-specific experiments are necessary to maximize the application of botanical extracts to achieve optimum results. The intensity of this impact varies depending on the species and crop/plant treated. However, the bioextract source varies according to these several factors such as maturity, leaves or shoots contained, location etc. Thus, these factors should be more considered in future experiments. Comprehensive studies should be needed to investigate the biological actions of botanical extracts during seed germination, including the control of ROS, their scavenging by antioxidant enzymes. Hence, more studies are required in future to assess the effectiveness of botanical extracts in minimizing the adverse effects of soil salinization on plants, as well as the ideal dose of application.

## REFERENCES

- [1] Jisha KC, Vijayakumari K and Puthur JT. Seed priming for abiotic stress tolerance: an overview. *Acta Physiol Plant*. 2013;35(5): 1381–1396.
- [2] Pawar V and Laware S. Seed Priming-A Critical Review. *International Journal of Scientific Research in Biological Sciences*. 2018;5: 94–101.
- [3] Sen A and Puthur JT. Influence of different seed priming techniques on oxidative and antioxidative responses during the germination of *Oryza sativa* varieties. *Physiol Mol Biol Plants*. 2020;26(3): 551–565.
- [4] Anwar A, Yu X and Li Y. Seed priming as a promising technique to improve growth, chlorophyll, photosynthesis and nutrient contents in cucumber seedlings. *Not Bot Horti Agrobo*. 2020;48(1): 116–127.
- [5] Shah T, Latif S, Saeed F, Ali I, Ullah S, Abdullah AA, Jan S and Ahmad P. Seed priming with titanium dioxide nanoparticles enhances seed vigor, leaf water status, and antioxidant enzyme activities in maize (*Zea mays* L.) under salinity stress. *Journal of King Saud University Science*. 2021;33(1): 101207.
- [6] Galahitigama GAH and Wathugala DL. Effects of presowing seed treatments on seed germination and salinity tolerance of Rice (*Oryza sativa* L.) seedlings. *Int. J. Agron. Agric*. 2016;9(6): 112–116.
- [7] Singh VK, Singh R, Tripathi S, Devi RS, Srivastava P, Singh P, Kumar A and Bhadouria R. Seed priming: state of the art and new perspectives in the era of climate change. In *Climate Change and Soil Interactions*. 2020;143–70.
- [8] Johnson R and Puthur JT. Seed priming as a cost-effective technique for developing plants with cross tolerance to salinity stress. *Plant Physiology and Biochemistry*. 2021;162: 247–57.
- [9] Arun MN, Hebbar SS, Bhanuprakash, Senthivel T, Nair AK, Padmavathi G, Pandey P and Singh A. Seed Priming: The Way Forward to Mitigate Abiotic Stress in Crops. *Physiology*. 2022.
- [10] Ahammad K, Rahman M and Ali M. Effect of hydropriming method on maize (*Zea mays*) seedling emergence. *Bangladesh J Agric Res*. 2014;39(1): 143–50.
- [11] Devika OS, Singh S, Sarkar D, Barnwal P, Suman J and Rakshit A. Seed priming: A potential supplement in integrated resource management under fragile intensive ecosystems. *Front Sustain Food Syst*. 2021;5: 654001.
- [12] Thinh NQ and Sundareswaran S. Effect of seed priming with seaweed extracts on Seed quality parameters in Blackgram (*Vigna mungo* L.) Cv. CO 6. *International Journal of Chemical Studies*. 2019;7(3): 813-817.
- [13] Battacharyya D, Babgohari MZ, Rathor P and Prithiviraj B. Seaweed extracts as biostimulants in horticulture. *Scientia Horticulturae*. 2015;196: 39–48.
- [14] Verma N, Sehrawat KD and Sehrawat AR. Studies on the effect of *Ascophyllum nodosum* extract (ANE) on growth, antioxidant potential and Alpha Glucosidase inhibition activity of *Vigna aconitifolia* (RMO 225). *Int J Curr Microbiol App Sci*. 2017;6(10): 2125–38.

- [15] Rajan MS, Thriunavukkarasu R, Joseph J and Aruni W. Effect of seaweed on seed germination and biochemical constituents of *Capsicum annum*. Biocatalysis and Agricultural Biotechnology. 2020;29: 101761.
- [16] Mahmoud SH, Salama DM, El Tanahy AMM and Abd, and El Samad EH. Utilization of seaweed (*Sargassum vulgare*) extract to enhance growth, yield and nutritional quality of red radish plants. Annals of Agricultural Sciences. 2019;64(2): 167–75.
- [17] Elavarasan E, Padmavathi S, Satheeshkumar P, Suganthi S and Kamaraj A. Role of seaweed extract application on physiological planting value of rice (*Oryza sativa* L.). 2021;6(2): 172–174.
- [18] Nabti E, Jha B and Hartmann A. Impact of seaweeds on agricultural crop production as biofertilizer. Int J Environ Sci Technol. 2017;14(5): 1119–1134.
- [19] Godlewska K, Michalak I, Tuhy Ł and Chojnacka K. Plant growth biostimulants based on different methods of seaweed extraction with water. Bio Med Research International. 2016: 1–11.
- [20] Mohamed HI, Akladios SA and El-Beltagi HS. Mitigation the harmful effect of salt stress on physiological, biochemical and anatomical traits by foliar spray with trehalose on wheat cultivars. Fresenius Environmental Bulletin. 2018;27(10): 7054–7065.
- [21] Antônio Pereira dos Anjos Neto, Fonseca Oliveira, Simone da Costa Mello, Marcio Souza da Silva, Francisco Guilhien Gomes-Junior, Ana Dionisia da Luz Coelho Novembre and Ricardo Antunes Azevedo. Seed priming with seaweed extract mitigate heat stress in spinach: effect on germination, seedling growth and antioxidant capacity. Bragantia. 2020;79(4): 502–11.
- [22] Hidangmayum A and Sharma R. Effect of different concentrations of commercial seaweed liquid extract of *Ascophyllum nodosum* as a plant biostimulant on growth, yield and biochemical constituents of onion (*Allium cepa* L.). International Journal of Science and Research. 2016;391(7): 1488–1491.
- [23] Sivritepe, N and Özkan Sivritepe H. Organic priming with seaweed extract (*Ascophyllum nodosum*) affects viability of pepper seeds. Asian Journal of Chemistry. 2008;20(7): 5689–5694.
- [24] El-Sheekh, MM, Ismail MM and Hamouda MM. Influence of some brown seaweed extracts on germination and cytological responses of *Trigonella foenum-graecum* L. Biotechnol Ind J. 2016;12(6): 104.
- [25] El-Aziz Kasim WA, Saad-Allah KM and Hamouda M. Seed priming with extracts of two seaweeds alleviates the physiological and molecular impacts of salinity stress on radish (*Raphanus sativus*). IJAB. 2016;18(3): 653–60.
- [26] El-Din SM. Utilization of seaweed extracts as bio-fertilizers to stimulate the growth of wheat seedlings. The Egyptian Journal of Experimental Biology. 2015;11: 31-39.
- [27] Hussein MH, Eltanahy E, Al Bakry AF, Elsafty N and Elshamy MM. Seaweed extracts as prospective plant growth bio-stimulant and salinity stress alleviator for *Vigna sinensis* and *Zea mays*. J Appl Phycol. 2021;33(2): 1273–91.
- [28] Makhaye G, Aremu AO, Gerrano AS, Tesfay S, Du Plooy CP, and Amoo SO. Biopriming with seaweed extract and microbial-based commercial biostimulants influences seed germination of five *Abelmoschus esculentus* genotypes. Plants. 2021;10(7): 1327.
- [29] Masondo NA, Kulkarni MG, Finnie JF and Van Staden J. Influence of biostimulants-seed-priming on *Ceratotheca triloba* germination and seedling growth under low temperatures, low osmotic potential and salinity stress. Ecotoxicology and Environmental Safety. 2018;147: 43–48.
- [30] Matysiak K, Kaczmarek S and Krawczyk R. Influence of seaweed extracts and mixture of humic and fulvic acids on germination and growth of *Zea mays* L. Acta Sci. Pol. Agricultura. 2011;10(1): 33-4510.
- [31] Righini H, Francioso O, Di Foggia M, Prodi A, Quintana AM and Roberti R. Tomato seed biopriming with water extracts from *Anabaena minutissima*, *Ecklonia maxima* and *Jania adhaerens* as a new agro-ecological option against *Rhizoctonia solani*. Scientia Horticulturae. 2021;281: 109921.
- [32] Rao GMN and Chatterjee R. Effect of seaweed liquid fertilizer from *Gracilaria Textorii* and *Hypnea Musciformis* on seed germination and productivity of some vegetable crops. Universal Journal of Plant Science. 2014;2(7): 115-120.

- [33] Dutta S, Majzoub A and Agarwal A. Oxidative stress and sperm function: A systematic review on evaluation and management. *Arab Journal of Urology*. 2019; 17(2): 87–97.
- [34] Fayzi L, Dayan M, Cherifi O, Houssine BE and Cherifi K. Biostimulant effect of four moroccan seaweed extracts applied as seed treatment and foliar spray on maize. *Asian Journal of Plant Sciences*. 2020;19(4): 419–428.
- [35] Layek J, Das A, Idapuganti RG, Sarkar D, Ghosh A, Zodape ST, Lal R, Yadav GS, Panwar AS, Ngachan S and Meena RS. Seaweed extract as organic bio-stimulant improves productivity and quality of rice in eastern Himalayas. *J Appl Phycol*. 2018;30(1): 547–58.
- [36] Hernández Herrera RM, Santacruz Ruvalcaba F, Ruiz López MA, Norrie J and Hernández Carmona G. Effect of liquid seaweed extracts on growth of tomato seedlings (*Solanum lycopersicum*L.). *J Appl Phycol*. 2014;26(1): 619-28.
- [37] Lemes TF, Neumann SV, Mendes MP and Tortelli B. Kale seed priming with red seaweed biostimulant. *Adv Hort Sci*. 2021;35(2): 111–8.
- [38] Erulan V, Soundarapandian P, Thirumaran G and Ananthan G. Studies on the effect of *Sargassum polycystum* (C.Agardh, 1824) extract on the growth and biochemical composition of *Cajanus cajan* (L.) Mill sp. *American-Eurasian J. Agric. and Environ. Sci*. 2009;6(4): 392–399.
- [39] Sasikala M, Indumathi E, Radhika S and Sasireka R. Effect of seaweed extract (*Sargassum tenerrimum*) on seed germination and growth of tomato plant (*Solanum lycopersicum*). *International Journal of ChemTech Research*. 2016;9(9): 285–293.
- [40] Mohamed A, Latique S, Chernane H, Hannachi C and Elkaoua M. Effect of Seaweed Extract of *Sargassum Vulgare* on germination behavior of two tomatoes cultivars (*Solanum Lycopersicum* L) under salt stress. *Acta Journal of Environmental Research*. 2014;2(3): 203–210.
- [41] Latique S, Elouaer MA, Chernane H, Hannachi C and Elkaoua M. Effect of seaweed liquid extract of *Sargassum vulgare* on growth of durum wheat seedlings (*Triticum durum* L) under salt stress. *International Journal of Innovation and Applied Studies*. 2014;7(4): 1430-1435.
- [42] Hamouda MM, Badr A, Ali SS, Adham AM, Ahmed HIS and Saad-Allah. Growth, physiological, and molecular responses of three phaeophyte extracts on salt-stressed pea (*Pisum sativum* L.) seedlings. *J Genet Eng Biotechnol*. 2023;21(32).
- [43] Kumar NA, Vanlalzarzova B, Sridhar S and Baluswami, M. Effect of liquid seaweed fertilizer of *Sargassum wightii*greve on the growth and biochemical content of green gram (*Vigna radiata* (L.) R. wilczek). *Recent Research in Science and Technology*. 2012;4(4): 40-45.
- [44] Kumar G and Sahoo D. Effect of seaweed liquid extract on growth and yield of *Triticum aestivum* var. Pusa Gold. *Journal of Applied Phycology*. 2011;23(2): 251-255.
- [45] Dominic Sahaya Rajan R, Stella C, Siva J, and Gokila S. Comparative study on the effect of seed germination activity of seagrass extracts with plant hormones on green gram (*Phaseolus Radiata* L). 2020;7(3): 406–408.
- [46] Shamyia Arokiarajan M, Thriunavukkarasu R, Joseph J and Aruni W. Effect of seaweed on seed germination and biochemical constituents of *Capsicum annum*. *Biocatalysis and Agricultural Biotechnology*. 2020;29(1): 101761.
- [47] Muhie S, Özdamar C, Gökdaş Z, Njie ES, Memiş N and Demir İ. Effect of solid matrix priming with seaweed extract on germination and seedling performance of onion seeds under abiotic stress conditions. *BSJ Agri*. 2020;3(4): 233-238.
- [48] Muhie S, Memiş N, Özdamar C, Gökdaş Z and Demir İ. Biostimulant priming for germination and seedling quality of carrot seeds under drought, salt and high temperature stress conditions. *International Journal of Agriculture Environment and Food Sciences*. 2021;5(3): 352-359.
- [49] El Boukhari MEM, Barakate M, Bouhia Y, and Lyamlouli K. Trends in seaweed extract based biostimulants: manufacturing process and beneficial effect on soil-plant systems. *Plants*. 2020;9(3): 359.
- [50] Kavipriya R, Dhanalakshmi PK., Jayashree S and Thangaraju N. Seaweed extract as a biostimulant for legume crop, green gram. *Journal of Ecobiotechnology*. 2011;3(8): 16–19.
- [51] Hamouda MM, Saad-Allah KM and Gad D. Potential of seaweed extract on growth, physiological, cytological and biochemical parameters of wheat (*Triticum aestivum* L.) seedlings. *J Soil Sci Plant Nutr*. 2022;22(2): 1818–31.

- [52] Ahmed AA and El-Mahdy AA. Improving seed germination and seedling growth of maize (*Zea mays* L.) seed by soaking in water and *Moringa oleifera* leaf extract. *Current Chemistry Letters*. 2022;11: 147–156.
- [53] Khan S, Ibrar D, Bashir S, Rashid N, Hasnain Z, Nawaz M, Al-Ghamdi AA, Elshikh MS, Dvořáčková H and Dvořáček J. Application of moringa leaf extract as a seed priming agent enhances growth and physiological attributes of rice seedlings cultivated under water deficit regime. *Plants*. 2022;11(3): 261.
- [54] Ali M, Hayat S, Ahmad H, Ghani MI, Amin B, Atif MJ and Cheng Z. Priming of *Solanum melongena* L. seeds enhances germination, alters antioxidant enzymes, modulates ROS, and improves early seedling growth: indicating aqueous garlic extract as seed-priming bio-stimulant for eggplant production. *Applied Sciences*. 2019;9(11): 2203.
- [55] Bhatেশwar DC, Prabha D, Jangid D and Salman M. Effect of seed priming with botanicals on plant growth and seed yield of lentil (*Lens culinaris* M.). *Int J CurrMicrobiol App Sci*. 2020;9(7): 3484–99.
- [56] Huang P, He L, Abbas A, Hussain S, Hussain S, Du D, Hafeez MB, Balooch S, Zahra N, Ren X, Rafiq M and Saqib M. Seed priming with sorghum water extract improves the performance of camelina (*Camelina sativa* L. Crantz.) under salt stress. *Plants*. 2021;10(4): 749.
- [57] Arshad U, Jabran M, Ahmed S, Abbas A, Jabbar A, Zahid MS and Amjad M. Seed-priming: a novel approach for improving growth performance and resistance against root-knot nematode (*Meloidogyne incognita*) in Bread Wheat (*Triticum aestivum* L.). *Gesunde Pflanzen*. 2022;74(4): 1041–51.
- [58] Kumar A, Basu S and Kumar G. Evaluating the effect of seed-priming for improving arsenic tolerance in rice. *J Plant Biochem Biotechnol*. 2022;31(1): 197–201.
- [59] Srikanth K and Chaurasia A. Influence of botanicals, coconut water and PGPR treatments on plant growth, nodulation, yield and seed quality parameters of chick pea (*Cicer arietinum* L.). *The Pharma Innovation Journal*. 2021;10(12): 2799–2802.
- [60] Aluko M, Ayodele OJ and Olaleye OE. Seed priming technique as innovation to improve germination in onion (*Allium cepa* L.) *Middle East J. Appl. Sci*. 2020;10(1): 7-17.
- [61] Ehtaiwesh AF and Yarboa LA. Testing of moringa leaf extract (*Moringa oleifera* L.) as seed priming agent in improving wheat germination under salinity. 2020;10(1): 76–90.
- [62] Rizwan A, Bushra S, Saim R, Muhammad M, Munawar I, Arif N, Muhammad A, and Zohaib Z. Fortification of phenolics, antioxidant activities and biochemical attributes of radish root by plant leaf extract seed priming. *Biocatalysis and Agricultural Biotechnology*. 2018;16: 115–120.
- [63] Dunsin O, Ajiboye G and Adeyemo T. Effect of alternative hormones on the root ability of parkiabioglobosa. *Scientia Agriculturae*. 2016;13(2): 113-118.
- [64] Noor MA, Ahmad W, Afzal I, Salamh A, Afzal M, Ahmad A, Ming Z and Wei Ma. Pea seed invigoration by priming with magnetized water and moringa leaf extract. *The Philippine agriculturist*. 2016;99: 171-175.
- [65] Iqbal MA. Improving germination and seedling vigour of cowpea (*Vigna Unguiculata* L.) with different priming techniques. *American-Eurasian Journal of Agricultural and Environmental Sciences*. 2015;15: 265-271.
- [66] Rehman H, Nawaz MQ, Basra SMA, Afzal I, Yasmeen A and Hassan F. Seed priming influence on early crop growth, phenological development and yield performance of linola (*Linum usitatissimum* L.). *Journal of Integrative Agriculture*. 2014;13: 990–996.
- [67] Shabbi I, Ayub M, Tahir M, Bilal M, Tanveer A, Hussain M and Afzal M. Impact of priming techniques on emergence and seedling growth of sesame (*Sesamum indicum* L.) genotypes. *Scientia Agriculture*. 2014;1(3): 92-96.
- [68] Afzal I, Hussain B, Basra SMA and Rehman H. Priming with moringa leaf extract reduces imbibitional chilling injury in spring maize. *Seed Science and Technology*. 2012;40(2): 271–276.
- [69] Basra SMA, Iftikhar MN and Afzal I. Potential of moringa (*Moringa oleifera*) Leaf extract as priming agent for hybrid maize seeds. *Int J Agric Biol*. 2011;13(6): 1006–1010.
- [70] Galappaththi MO, Jayasuriya KMG and Gama- Arachchige NS. Effect of priming with neem seed extract on seeds of four traditional rice varieties of Sri Lanka; Kaluheenati, Kurulurthuda, Madathawalu and Maa-wee. *J Natn Sci Foundation Sri Lanka*. 2021;31;49(4): 525.

- [71] Farooq O, Hussain QM, Sarwar N, Iqbal MM and Shiaz M. Seed Priming with sorghum water extracts and calcium chloride improves the stand establishment and seedling growth of sunflower and maize. *Pakistan Journal of Life and Social Sciences*. 2018;16(2): 97–101.
- [72] Girase IP, Rai PK, Bara BM and Singh BA. Effect of plant extracts on seed germination behaviour and vigour of okra [*Abelmoschus esculentus* (L.) Moench]. *Int J Curr Microbiol App Sci*. 2019;8(8): 830–5.
- [73] Noman A, Ali Q, Naseem J, Javed MT, Kanwal H, Islam W, Aqeel M, Khalid N, Zafar S, Tayyeb M, Iqbal N, Buriro M, Maqsood J and Shahid S. Sugar beet extract acts as a natural bio-stimulant for physio-biochemical attributes in water stressed wheat (*Triticum aestivum* L.). *Acta Physiol Plant*. 2018;40(6): 110.
- [74] Zida PE, Néya BJ, Soalla RW, Sérémé P and Lund OS. Seed-priming of sorghum with antifungal extracts from *Balanites aegyptiaca* and *Eclipta alba* in different agro-ecological zones of Burkina Faso. *African Journal of Agricultural Research*. 2018;13: 2516-2525.
- [75] Gunasekar J, Kamaraj A and Padmavathi S. Effect of botanical seed priming on seed quality characters in Blackgram [*Vigna Mungo* (L.) Hepper] cv. CO 6. *Plant Archives*. 2017;17(2): 1383–1387.
- [76] Sajjan AS, Dhanelappagol MS and Jolli RB. Seed quality enhancement through seed priming in pigeon pea [*Cajanus cajan*(L.) Millsp.]. *Legume Research*. 2017;40(1): 173-177.
- [77] Mavi, K. The Effect of Organic Priming with Marigold Herbal Tea on Seeds Quality in Aji Pepper (*Capsicum baccatum* var. pendulum Willd.). *Journal of Agricultural Faculty of Mustafa Kemal University*. 2016;21(1): 31-39.
- [78] Prabha D, Negi S, Kumari P, Negi YK and Chauhan JS. Effect of seed priming with some plant leaf extract on seedling growth characteristics and root rot disease in tomato. *International Journal of Agriculture System*. 2016;4(1): 46–51.
- [79] Rafi H, Dawar S and Javed Zaki M. Seed priming with extracts of *Acacia nilotica* (L.) Willd. ex Delile and *Sapindus mukorossi* (L.) plant parts in the control of root rot fungi and growth of plants. *Pakistan Journal of Botany*. 2015;47(3): 1129–1135.
- [80] Ashfaq M, Ali A, Haider MS, Ali M, Abdullah M and Mubashar U. A role of weed solvents on seed priming of diverse rice germplasm lines. *Journal of Pure and Applied Microbiology*. 2014;8(3): 2175–2184.
- [81] Imran S, Afzal I, Amjad M, Akram A, Khawar KM and Pretorius S. Seed priming with aqueous plant extracts improved seed germination and seedling growth under chilling stress in Lentil (*Lens culinaris* Medik.). *Acta Advances in Agricultural Sciences*. 2014;11(2): 58-69.
- [82] Jayakumar R, Natarajan N and Srimathi P. Efficacy of Botanicals in Improving the Seeds and Seedling Quality Characteristics of Cluster Bean. *Legume Research*. 2008;3:164-168.
- [83] Costa PS, Ferraz RLS, Dantas-Neto J, Martins VD, Viégas PRA, Meira KS, Ndhala AR, Azevedo CAV and Melo AS. Seed Priming with Light Quality and *Cyperus rotundus* L. Extract Modulate the Germination and Initial Growth of *Moringa oleifera* Lam. seedlings. *Braz J Biol*. 2022; 84.
- [84] Kenanoğlu B, Uşak Üniversitesi and Kor A. Different Effects of Tomato Leaf Extract on Seed Germination. *International Journal of Innovative Approaches in Agricultural Research*. 2018;2(4): 278–84.
- [85] Arafa SA, Attia KA, Niedbala G, Piekutowska M, Alamery S, Abdelaal K, Alateeq TK, A. M. Ali M, Elkelish A and Attallah SY. Seed Priming Boost Adaptation in Pea Plants under Drought Stress. *Plants*. 2021;10(10): 2201.
- [86] Muthuselvan K and Subbaiyan M. Organic Seed Priming and Foliar Nutrition with Medicinal Herbs to Enhance Seedling Vigour and Yield Potential in Maize (*Zea mays* L.). *SP2*. 2019: 693-698.
- [87] Marthandan V, Geetha R, Kumutha K, Renganathan VG, Karthikeyan A and Ramalingam J. Seed Priming: A Feasible Strategy to Enhance Drought Tolerance in Crop Plants. *International Journal of Medical Students*. 2020;21(21): 8258.
- [88] Paparella S, Araújo SS, Rossi G, Wijayasinghe M, Carbonera D and Balestrazzi A. Seed priming: state of the art and new perspectives. *Plant Cell Rep*. 2015;34(8): 1281–93.
- [89] Nonogaki H, Nishiyama E, Ohshima K and Nonogaki M. Ancient Memories of Seeds: ABA-Dependent Growth Arrest and Reserve Accumulation. *Trends in Genetics*. 2020;36(7): 464–73.

- [90] Siddiqui MH, Alamri S, Al-Khaishany MY, Khan MN, Al-Amri A, Ali HM, Alaraidh IA and Alsahli AA. Exogenous Melatonin Counteracts NaCl-Induced Damage by Regulating the Antioxidant System, Proline and Carbohydrates Metabolism in Tomato Seedlings. *Int J Mol Sci.* 2019;20(2): 353.
- [91] Szafrńska K, Reiter RJ and Posmyk MM. Melatonin Improves the Photosynthetic Apparatus in Pea Leaves Stressed by Paraquat via Chlorophyll Breakdown Regulation and Its Accelerated de novo Synthesis. *Front Plant Sci.* 2017;8: 878.
- [92] Liu L, Xia W, Li H, Zeng H, Wei B, Han S and Yin C. Salinity Inhibits Rice Seed Germination by Reducing  $\alpha$ -Amylase Activity via Decreased Bioactive Gibberellin Content. *Front Plant Sci.* 2018;9: 275.
- [93] Wani AS, Ahmad A, Hayat S and Tahir I. Epibrassinolide and proline alleviate the photosynthetic and yield inhibition under salt stress by acting on antioxidant system in mustard. *Plant Physiology and Biochemistry.* 2019;135: 385–94.
- [94] Tahira T, Riaz A, Muhammad F and Basra SMA. Improving salt tolerance in barley by osmopriming and bio priming. *Int. J. Agric. Biol.* 2019;20(11): 2455–2464.
- [95] Win KT and Oo AZ. Salt-stress-induced changes in protein profiles in two blackgram (*Vigna Mungo* L.) varieties differing salinity tolerance. *Adv Plants Agric Res.* 2017; 7(1): 206–210.