

**Antifungal activity of essential oil formulations based on starch and biochar in the postharvest conservation of peanut (*Arachis hypogaea* L.) seeds**

**Abstract**

Poor storage conditions of foodstuffs are responsible for their deterioration by many microorganism species with huge-substantial quantitative and qualitative losses. The antifungal efficacy of essential oil formulations of some plants based on cassava starch and corn-cob biochar against fungi associated with peanut seeds (*Arachis hypogaea* L.) during storage During storage, some plants' antifungal efficacy of essential oil formulations based on cassava starch and corncob biochar against fungi associated with peanut seeds (*Arachis hypogaea* L.) was evaluated. 200-Two hundred peanut seeds were dispersed in plastic bottles; 5 g of essential oil formulation starch-based (3 and 7  $\mu$ /g) were sprinkled on the seeds. For biochar-based essential oils, pieces of biochar were placed in the middle of the seeds in the bottles. Treated and untreated seeds were stored for 70 days in laboratory conditions. Infection rates of fungal were estimated at 10, 20, 40, 60 infections were estimated at 10, 20, 40, 60, and 70 days of storage. From the results obtained, five fungal genera were isolated from peanut seeds, including 4-four species of *Aspergillus*, *Cercospora* sp, *Fusarium* sp, *Phomopsis* sp, and *Rhizopus stolonifer*, with seed infection percentages ranging from 4 to 62%. The antifungal activity of starch-based formulations revealed that at a concentration of 3  $\mu$ /g, *Cymbopogon citratus* essential oil has the highest antifungal activity, with 13% infection of peanut seeds after 70 days of storage. After 70 days of storage, at a dose of 7  $\mu$ /g, *C. sempervirens* essential oil had the best antifungal efficacy with 23.50% of infected seeds for the biochar-based formulation. Based on these results, starch, and biochar can be used to formulate essential oils for the conservation of peanut seeds. Moreover, *Cymbopogon citratus* essential oil formulated with starch is the most effective.

**Keywords:** Essential oils; Starch; Biochar; Formulations; Antifungal activity; Peanut seeds; Conservation.

**Introduction**

Peanut (*Arachis hypogaea* L.) is one of the most important-essential cultivated legumes in sub-Saharan Africa [1-2] (Snapp *et al.*, 2018; Kpatinvoh *et al.*, 2017). Global annual groundnut production is estimated at 47 million tonnes of dry pods, of which nearly 29% is produced in Africa

and 594,000 tonnes from Cameroon [3] (FAOSTAT, 2018). Pulse, in general, and groundnuts—in particular are essential contributors to global food security, health, and poverty eradication [1] (Snapp *et al.*, 2018), especially in developing countries. These legumes are an important source of essential protein, micronutrients, and amino acids [4] (Asif *et al.*, 2013). They can thus contribute to the fight against iron deficiency anemia, one of the most important-critical micronutrient deficiencies observed in Africa, and against protein-energy- deficiency [1] (Snapp *et al.*, 2018).

Poor storage conditions are sometimes observed in rural areas. In fact, high seed moisture content (insufficient drying), inappropriate relative humidity, and temperature of storage structures, thus promoting mould growth [4-5] (Nadjet *et al.*, 2016; Taruvinga *et al.*, 2014). The damage caused by these stock moulds is manifested not only by a significant alteration of the aesthetic quality- and organoleptic and chemical characteristics of- the foodstuffs [4] (Nadjet *et al.*, 2016), but also by the production of mycotoxins such as aflatoxins. The latter are mainly produced by molds of the genus *Aspergillus*. Molds of the genus *Aspergillus* mainly produce the latter, recognized as the major contaminants of peanuts [6] (Torres *et al.*, 2014). Due to their varied toxic effects, mycotoxins can cause acute or chronic poisoning in humans and animals, sometimes fatal, such as carcinogenicity, immune toxicity, neurotoxicity, and hepatotoxicity [7] (Agriopoulou *et al.*, 2020). It has been noticed that the many hazards caused by moulds on agricultural products generally lead to their removal during sorting for the market, resulting in a worldwide loss of food production estimated at between 5 and 10 % [8] (Yiannikouris and Jouany, 2002).

Nowadays, to reduce the incidence of mould in stored commodities, chemical control through the use of synthetic fungicides seems to be the most effective and widespread control method in the world, with a substantial increase in agricultural productivity [9-2] (Aoudou *et al.*, 2010; Kpatinvoh *et al.*, 2017). Nevertheless, synthetic pesticides are currently a the subject of global concern since their adverse effects on human health and the environment have been demonstrated. Indeed, the application of these synthetic chemicals at high concentrations for post-harvest control increases the risk of toxic residues in foodstuffs [10-11] (Adjou *et al.*, 2013; Rosenbaum *et al.*, 2015) and the risk of developing resistant fungal strains [12] (Rapp *et al.*, 2004). The toxic effects of chemical pesticides on human beings are carcinogenic, immunosuppressive, mutagenic, and neurotoxic [13] (Mostafalou and Abdollahi, 2016). [4] Nadjet *et al.* (2016) had shown showed that mycotoxin toxicity could be increased by concomitant human exposure to synthetic pesticides, since the latter inhibits many enzymes, particularly those capable of detoxifying certain mycotoxins.

In view Because of those problems associated with chemical pesticides, the use of using natural products of plant origin for the preservation of to preserve foodstuffs is experiencing significant growth and constitutes a major challenge to be met. Essential oils are a new category of

Field Code Changed

biodegradable plant protection products that have demonstrated their antifungal properties, with the particularity of presenting no toxicity to humans [14-15] (Abd-Alla and Haggag, 2013; Zhu *et al.*, 2016). Many previous studies on essential oils antifungal activity of *Thymus vulgaris* [16-17, 18] (Kritzinger *et al.*, 2002; De Vincenzi *et al.*, 2004; Vitoratos *et al.*, 2013), *Cymbopogon citratus* [19-20, 21] (Yousef, 2013; Aoudou *et al.*, 2017; Premathilake *et al.*, 2018) and *Cupressus sempervirens* [22-23] (Mazari *et al.*, 2010; Ismail *et al.*, 2013) have been demonstrated against some stock fungi such as *Aspergillus niger*, *A. flavus*, *Penicillium sp.*, *Alternaria alternata*, *Fusarium tricinctum*, *Phomopsis*, and *Rhizopus stolonifer*. However, although essential oils do not ~~pose any danger to human health (without overdoses) and environment, some factors limited~~ harm human health (without overdoses) and environment, some factors limit their full adoption. Indeed, essential oils have a short persistence in ~~real-actual~~ storage conditions due to their high volatility and the instability of their constituent molecules [24-25] (Hsieh *et al.*, 2006-; Baptiste Houndou Fokou *et al.*, 2020). ~~In order to ensure their antifungal activity, additional doses are necessary~~ Additional doses are necessary to ensure their antifungal activity, but the extraction yield of essential oils is generally very low. Hence the importance and necessity of developing formulations of essential oils capable of optimizing the effectiveness of low doses of these essential oils as well as improving their persistence during food storage, while maintaining their antifungal properties. With this in mind, the objective of this study is to contribute to improve the conservation of peanuts, by fighting against the development of mold using essential oil formulations.

## Materials and methods

### Plant material and procedure for extracting essential oils

In January 2020, fresh leaves of *Thymus vulgaris*, *Cymbopogon citratus*, and *Cupressus sempervirens* were collected in the locality of Dschang and then air-dried for five days at room temperature ( $25^{\circ}\text{C} \pm 2$ ). ~~The leaves were hydrodistilled with the help of a~~ The leaves were hydrodistilled for about 5 hours at the Research Center for Microbiology and Ontibral Substances of the University of Dschang using a ~~Clevenger-type apparatus~~ Clevenger-type apparatus for about 5 hours at the Research Center for Microbiology and Ontibral Substances of the University of Dschang. Following the recommendations of ~~[26-27]~~ Sessou *et al.* (2012) and Rguez *et al.* (2018), the extracted oils were dried with anhydrous sodium ~~sulphate-sulfate~~ and stored at  $4^{\circ}\text{C}$  away from light until used to prevent spoilage.

Formatted: English (United States)

### **Starch and biochar ~~are~~ made from corn cob.**

Cassava starch (*Manihot esculenta*) was obtained ~~by using~~ the method described by [28] (Maptue *et al.*, 2021). Biochar based on maize cobs was obtained using the methods described ~~by~~ [29] ~~Ioannidou *et al.* (2009), [30] Liu *et al.* (2014)~~ and taken up by [28] Maptue *et al.* (2021). Biochar, in particular corn-cob biochar, has the particularity of having a high relative porosity of nearly 80% [31] (Djousse *et al.*, 2018). This feature has aroused particular interest in its use, as it ~~is able to can~~ retain essential oils in its microporosity and allow increasing volatility over time.

### **Essential oil formulations**

Starch-based essential oil formulations (*Cymbopogon citratus*, *Cupressus sempervirens*, and *Thymus vulgaris*) were made ~~with reference to regarding~~ the previous work of [32] Camara (2009) and the technique described by [28] Maptue *et al.* (2021). For biochar-based essential oil formulations (*Cymbopogon citratus*, *Cupressus sempervirens*), the technique described by Maptue and collaborators in 2021 was applied. Once the formulations were made, they were left to stand for 10 hours to ensure effective diffusion of the volatile elements of the essential oil.

### **Treatment of peanut seeds**

~~200-After removing rotten, cracked, or damaged seeds and waste, two hundred peanut seeds were dispersed in plastic bottles~~ ~~peanut seeds were dispersed in plastic bottles after removing rotten, cracked, or damaged seeds as well as waste.~~ Regarding the starch-based essential oil formulation, 5 g ~~of this formulation~~ was sprinkled on the seeds for each concentration. Each container was ~~then~~ mixed to ensure excellent contact between the powder and the seeds. Each piece of biochar soaked in ~~a~~ ~~dose of~~ essential oil was placed in the middle of the seeds in the bottles. Deltamethrin, a synthetic pesticide for the protection of stored seeds, used at the manufacturer's recommended dose (2g/kg), was ~~the a~~ positive control. Untreated groundnut seeds were used as a negative control. The experiment was performed three (03) times.

### **Effects of ~~essential oil f~~Essential Oil Formulations**

For 70 days, treated and untreated seeds were stored at a laboratory room temperature of 25 ± 2°C. Fungal infection rates were evaluated ~~at~~ 10, 20, 30, 40, 50, 60, and 70 days of storage, using Potato Dextrose Agar- and the ~~blotter~~ blotter technique, following the protocol described by [33] ISTA (2023) with some modifications.

## Seed health

Disinfected peanut seeds on the surface (200 seeds) were placed on PDA (Potato Dextrose Agar) medium to isolate the associated storage fungus. After 7-10 days of incubation at  $25 \pm 2^\circ\text{C}$  and daily observation, fungal colonies produced around the seeds were collected and purified for identification using conventional fungal identification keys as reported in other studies [34-35, 36] (Champion, 1997; Warham *et al.*, 1997; Mathur and Kongsdal, 2003).

## Statistical analysis

The collected data were subjected to analysis of variance and then separation of the means by the Duncan test at the 5% probability threshold using R software version 4.2.2.

## Results and discussion

### Fungi associated with peanut seeds and percentage of infection before treatment

Table 1 shows the fungal species isolated from groundnut seeds and the percentage of infection of each infection percentage. According to this result, groundnut seeds collected in Dschang are infected mainly by eight (08) fungal species of interest. These are: *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, *Aspergillus parasiticus*, *Cercospora sp*, *Fusarium sp*, *Phomopsis sp*, and *Rhizopus stolonifer*. The infection of seeds caused by these different species ranges from 4 to 62%.

**Table 1:** Percentage of peanut seed infection prior to treatment with essential oils by isolated parasites

Identified fungi	Fungal infection rate (%)
	Peanut
<i>A. flavus</i>	39,00
<i>A. fumigatus</i>	11,50
<i>A. Niger</i>	18,00
<i>A. parasiticus</i>	3,50
<i>Cercospora sp</i>	15,50
<i>Fusarium sp</i>	41,50
<i>Phomopsis sp</i>	11,00
<i>Rhizopus stolonifer</i>	22,00

### Antifungal activity of different formulations of essential oils

The *in vivo* antifungal efficacy of essential oil formulations of *C. citratus*, *C. sempervirens*, and *T. vulgaris* based on starch and biochar is presented in Table 2. These results show that all

formulations have an inhibitory effect on fungal infection of peanut seeds. This inhibition varies depending on the type of essential oil, dose, storage period, and formulation support.

**Table 2:** Fungal infection rate of peanuts treated with starch- and biochar-based essential oil formulations as a function of storage period

Infection rate (%)						
Formulations	Doses ( $\mu$ l/g)	10 days	20 days	40 days	60 days	70 days
Starch +	3,00	38.50 <sup>b</sup>	14.50 <sup>a</sup>	22.00 <sup>b</sup>	16.00 <sup>bc</sup>	13.00 <sup>a</sup>
<i>C. citratus</i>	7,00	32.50 <sup>b</sup>	24.50 <sup>b</sup>	19.00 <sup>b</sup>	4.00 <sup>a</sup>	24.16 <sup>bc</sup>
Starch +	3,00	40.50 <sup>b</sup>	30.00 <sup>bc</sup>	18.66 <sup>b</sup>	15.00 <sup>b</sup>	22.50 <sup>bc</sup>
<i>C. sempervirens</i>	7,00	39.00 <sup>b</sup>	41.00 <sup>d</sup>	13.00 <sup>a</sup>	29.00 <sup>d</sup>	28.50 <sup>cd</sup>
Starch +	3,00	66.50 <sup>e</sup>	27.00 <sup>b</sup>	11.50 <sup>a</sup>	14.50 <sup>b</sup>	18.50 <sup>ab</sup>
<i>T. vulgaris</i>	7,00	18.50 <sup>a</sup>	24.50 <sup>b</sup>	9.00 <sup>a</sup>	14.50 <sup>b</sup>	25.00 <sup>bc</sup>
Biochar +	3,00	65.66 <sup>e</sup>	55.66 <sup>e</sup>	44.66 <sup>c</sup>	21.50 <sup>bcd</sup>	34.00 <sup>e</sup>
<i>C. citratus</i>	7,00	54.00 <sup>cd</sup>	54.66 <sup>e</sup>	52.00 <sup>d</sup>	14.66 <sup>b</sup>	57.50 <sup>fg</sup>
Biochar +	3,00	40.16 <sup>b</sup>	34.00 <sup>c</sup>	22.50 <sup>b</sup>	24.66 <sup>d</sup>	38.50 <sup>e</sup>
<i>C. sempervirens</i>	7,00	52.66 <sup>c</sup>	41.16 <sup>d</sup>	20.16 <sup>b</sup>	23.16 <sup>cd</sup>	23.50 <sup>bc</sup>
Control	Control -	62.00 <sup>de</sup>	62.00 <sup>f</sup>	62.00 <sup>th</sup>	62.00 <sup>f</sup>	62.00 <sup>g</sup>
	Control +	50.83 <sup>c</sup>	50.83 <sup>e</sup>	50.83 <sup>d</sup>	50.83 <sup>e</sup>	50.83 <sup>fg</sup>

Means in each column assigned with the same upper letter do not differ significantly according to Duncan's test at the probability threshold  $P = .05$


#### Antifungal activity of starch-based essential oil formulations

Figures 1 (A-B) and 2 (A-B) show, respectively, the percentage of infection as a function of time and the percentage of infection after 70 days of storage of peanut seeds treated with the essential oil formulations of *Cymbopogon citratus*, *Cupressus sempervirens*, and *Thymus vulgaris* starch-based. Based on these results, the inhibitory effects of the formulations were significantly influenced by the storage period, the dose applied, and the type of essential oil. The percentage of fungal infection of seeds treated with essential oils decreased throughout the storage period. After 10 ten days of storage, the percentage of infection of treated seeds the infection rate of treated seeds was high and ranged from 18.50% to 66.50%. After 40 days of storage, the infection percentages of treated seeds were lower than those of positive (58%) and negative (62%) controls; and ranged from 9% to 2.2%. These values are significantly lower ( $P = .05$ ) than those of the control as well as those obtained after 10 ten days of storage. After 60 days of storage, the infection percentages of treated

seeds, which range from 14.50% to 29%, are significantly lower ( $P = .05$ ) than those of the controls, but still higher than the ~~infection percentages~~ obtained after 40 days of storage. These results show that the antifungal activity of different formulations depends on the type of essential oil and its dose. For this purpose, the starch formulation + *Cymbopogon citratus* is the one with the highest antifungal activity with a fungal infection rate of 13.00% at the concentration of 3  $\mu$ l/g after 70 days of storage, followed respectively by starch + *Thymus vulgaris* (18.50%) and starch + *Cupressus sempervirens* (22%) at a concentration of 3  $\mu$ l/g after 60 days of storage.

### Antifungal activity of biochar-based essential oil formulations

Figures 1 (C-D) and 3 (A-B) show the percentages of fungal infection of peanut seeds treated with essential oil formulations of *Cymbopogon citratus* and *Cupressus sempervirens* based on vegetable charcoal. ~~It appears that after 10 ten~~ days of treatment, ~~the percentages of fungal infection of seeds~~ the percentages of fungal infection in seeds range from 97 to 100%, ~~%~~; these average values are statistically higher than those of negative and positive controls, 62.50% and 85.50%, respectively. After 60 days of storage, the percentages of infection of the treated seeds, which vary from 10% to 12%, are significantly lower ( $P = .05$ ) than those of the control, ~~as well as and~~ those obtained after 40 days of storage. ~~It was also noted that~~ Notably, these percentages of fungal infection observed after 60 days were also lower than those recorded after 70 days of storage (9% to 27%). The ~~lowest percentages of fungal infection of the seeds were obtained after 60 days of storage~~ seeds' lowest percentages of fungal infection were obtained after 60 days of storage, and there was no significant difference between the doses applied. However, after 70 days of storage, the biochar + *C. citratus* formulation showed better antifungal activity than that observed at 60 days at 7  $\mu$ l/g, with a 9% seed infection percentage, which is better than that observed at 60 days at 7  $\mu$ l/g (12%).



**Figure 1:** Seed infection rate by substrate type, dose, and storage period. A – B: Formulation at a dose of 3 µg/l. C – D: Formulation at a dose of 7 µg/l.

### Effect of formulation support and dose on the effectiveness of formulated essential oils

Table 2 presents the effect of substrate type on the antifungal activity of essential oil formulations at doses of 3 and 7 µl / g. ~~It is clear that the~~ influence of formulation support on ~~the percentages of fungal infection of seedseeds'~~ fungal infection percentages is not statistically significant ( $P > .05$ ) at a dose of 3 µl/g. Although at 60 days of storage ~~where the lowest percentages of fungal infection were observed (Figure 1), starch-based essential oil formulations were more active,~~ the lowest percentages of fungal infection were observed (Figure 1), starch-based essential oil formulations were more active but with no significant difference from biochar-based formulations.

Looking at the effect of formulation support on the antifungal activity of *Cymbopogon citratus* essential oil formulations at a dose of 7 µl/g, a significant difference ( $P = .05$ ) was observed between starch and biochar formulations at 40 and 70 days of storage. Biochar-based formulations had the most pronounced inhibitory effect on fungal infection of peanut seeds after 70 days of storage.

## Discussion

*In vivo* antifungal tests using the essential oil compositions of *Cymbopogon citratus*, *Cupressus sempervirens*, and *Thymus vulgaris* against fungi associated with peanut seeds revealed a substantial decrease in the rate of fungal infection of treated seeds compared to untreated seeds and positive controls. ~~In the case of starch-based formulations, *Thymus vulgaris* essential oil had the highest antifungal activity.~~ *Thymus vulgaris* essential oil had the highest antifungal activity in starch-based formulations, followed by formulations of *Cupressus sempervirens* and *Cymbopogon citratus* oils at a concentration of 7  $\mu\text{l/g}$ . ~~In the same vein~~ Similarly, for biochar-based formulations, the essential oil of *C. sempervirens* at the concentration of 7  $\mu\text{l/g}$  showed the highest antifungal activity. The ability of these formulations to prevent fungal infections can be explained by the active chemicals ~~contained~~ in essential oils, which can either prevent the development of fungi or stimulate the host plant's defenses, such as pathogenic fungi [37] (Liu *et al.*, 2002). Indeed, numerous studies on the chemical composition and antimicrobial properties of essential oils have shown that they consist of biologically active compounds such as terpenes and terpenoids, especially monoterpenes and sesquiterpenes with their hydrocarbon and oxygenated derivatives [38-39] (Smigielski *et al.*, 2018; Tchoumboungang *et al.*, 2009). These studies have reported the antifungal effects of oxygenate and monoterpene hydrocarbons against a diverse group of phytopathogenic fungi [37-18, 40] (Liu *et al.*, 2002; Vitoratos *et al.*, 2013; Regnier *et al.*, 2014).

Previous studies [41-17] (Thompson *et al.*, 2003; De Vincenzi *et al.*, 2004) showed that *Thymus*-spp essential oil is mainly composed of oxygenated monoterpenes such as carvacrol, thymol, linalool, geraniol and p-cymen. These conclusions justify the results obtained and corroborate the work of [16] Kritzinger *et al.* (2002), which demonstrated that under *in vitro* conditions, *Thymus vulgaris* essential oil significantly inhibited the growth of *Aspergillus flavus*, *A. niger*, *Fusarium oxysporium*, *F. equiseti*, *Penicillium chrysogenum* and *Rhizopus* sp associated with cowpea seeds in storage. Similarly, under *in vivo* conditions, thyme significantly reduced the total incidence of these fungi on naturally infected seeds.

On the other hand, it has been shown that lemongrass essential oil has as its majority compounds citral aldehyde,  $\beta$ -Citral, geraniol, cis-Verbenol, acetal diethyl citral and nerol, which are oxygenated monoterpenes, as well as myrcene and  $\alpha$ -pinene, which are hydrocarbon monoterpenes [42-16] (Shahzadi and Shahzadi, 2017; Premathilake *et al.*, 2018;). Other authors [43-44] (Paranagama *et al.*, 2003; Kakarla and Ganjewala, 2015) explained that citral is the chemical component responsible for the antifungal properties of lemongrass essential oil. The work of [9] Aoudou *et al.* (2010) showed complete inhibition of the growth of *A. parasiticus* mycelia by citral.

Much other research has been conducted on the chemical composition of essential oils of evergreen species [45-46, 47, 22, 23] (Chéraif, 2005; Sacchetti *et al.*, 2005; Emami *et al.*, 2006; Mazari *et al.*, 2010; Ismail *et al.*, 2013). According to these studies, the majority of compounds of this oil are monoterpene hydrocarbons, including  $\alpha$ -pinene,  $\delta$ -3-carene, myrcene, and limonene. They also note the presence of oxygenated monoterpenes in very small proportions. Its higher composition of monoterpene hydrocarbons explains why it has the weakest inhibitory activity against fungi associated with peanut seeds. According to [22] Mazari *et al.* (2010), *C. sempervirens* essential oil moderately inhibited the growth of *A. flavus* and *F. oxysporum*. However, compared to the positive control (amphotericin B), the oil was not significantly active against *Rhizopus stolonifer*.

## Conclusion

Essential oil formulations based on starch and biochar have significantly reduced fungal development in peanut grains during storage. Application of starch-based formulations revealed that *Thymus vulgaris* essential oil almost completely inhibited fungi associated with peanut seeds for up to 60 days of storage at 7  $\mu$ l/g. The biochar-based formulation of *Cymbopogon sempervirens* essential oil had an acceptable effect and inhibitor after 60 days of storage. Based on these results, starch and biochar can be used in the formulation of essential oils for the preservation of peanut seeds. *Thymus vulgaris* essential oil has proven to be the most effective, especially when formulated with starch. It would be fascinating to continue this research to learn more about the durability of these formulations.

## References

- [1] Snapp S, Rahmanian M, Batello C, Calles T. Pulses and sustainable farms in sub-Saharan Africa. FAO, Rome, Italy. 2018; <https://doi.org/10.3/JQUERY-UI.JS>
- [2] Kpatinvoh B, Adjou ES, Dahouenon-Ahoussi E, Konfo TRC, Atrevi B, Soumanou MM, Sohounhloue DCK. Effectiveness of essential oils of three aromatic plants against spoilage mycoflora of cowpea (*Vigna unguiculata* L., Walp) collected in sales stores in South Benin. *J. Apply Biosci.* 2017; 109:10680–10687. <https://doi.org/10.4314/jab.v109i1.12>
- [3] FAOSTAT, 2018. Statistics for the year 2018 [WWW Document]. FAO. URL <https://www.fao.org/faostat/fr/#home> (accessed 5.21.23).
- [4] Asif M, Rooney LW, Ali R, Riaz MN. Application and opportunities of pulses in food system: A review. *Crit. Rev. Food Sci. Nutr.* 2013, 53(11): 1168-1179.
- [5] Nadjat G-T, Noureddine B, El Hadj O, Didi M, Ezzouar B, Alger A. Mycotoxins: a public health danger. *J. Arid Approx.* 2016; 32:32–49.
- [6] Taruvinga C, Mejia D, Alvarez JS. Appropriate seed and grain storage systems for small-scale

farmers: key practices for DRR practitioners, FAO. 2014


- [7] Torres AM, Barros GG, Palacios SA, Chulze SN, Battilani P. Review on pre- and post-harvest management of peanuts to minimize aflatoxin contamination. *Food Res. Int.* 2014; 62:11–19. <https://doi.org/10.1016/j.foodres.2014.02.023>
- [8] Agriopoulou S, Stamatelopoulou E, Varzakas T. Advances in occurrence, importance, and mycotoxin control strategies: Prevention and detoxification in foods. *Foods.* 2020; 9:137. <https://doi.org/10.3390/foods9020137>
- [9] Yiannikouris A, Jouany JP. Mycotoxins in feeds and their fate in animals: a review. *Anim. Res.* 2002; 51:81–99. <https://doi.org/10.1051/ANIMRES:2002012>
- [10] Aoudou Y, Leopold TN, Michel JDP, Xavier EF, Moses MC. Antifungal properties of essential oils and some constituents to reduce foodborne pathogens. *J. Yeast Fungal Res.* 2010; 1:001–008.
- [11] Adjou ES, Kouton S, Dahouenon-Ahoussi E, Soumanou MM, Sohounhloue DCK. Effect of essential oil from fresh leaves of *Ocimum gratissimum* L. on mycoflora during storage of peanuts in Benin. *Mycotoxin Res.* 2013; 29:29–38. <https://doi.org/10.1007/S12550-012-0150-Y/METRICS>
- [12] Rosenbaum RK, Anton A, Bengoa , Lewis F, Maxime D, Nemecek T, Payet J, Räsänen K, Roux P, Schau EM, Sourisseau S, van Zelm R, von Streit B, Wallman M. The Glasgow consensus on the delineation between pesticide emission inventory and impact assessment for LCA. *Int. J. Life Cycle Assess.* 2015; 20:765–776. <https://doi.org/10.1007/S11367-015-0871-1/TABLES/3>
- [13] Rapp HS, Bellinder RR, Wien HC, Vermeylen FM. Reduced Tillage, Rye Residues, and Herbicides Influence Weed Suppression and Yield of Pumpkins. *Weed Technol.* 2004 ; 18:953–961. <https://doi.org/10.1614/WT-03-136R>
- [14] Mostafalou S, Abdollahi M. Pesticides: an update of human exposure and toxicity. *Arch. Toxicol.* 2016; 912 (91): 549–599. <https://doi.org/10.1007/S00204-016-1849-X>
- [15] Abd-Alla MA, Haggag WM. Use of Some Plant Essential Oils as Post-harvest Botanical Fungicides in the Management of Anthracnose Disease of Mango Fruits (*Mangifera indica* L.) Caused by *Colletotrichum gloeosporioides* (Penz). *Int. J. Agriculture For.* 2013; 1–6. <https://doi.org/10.5923/j.ijaf.20130301.01>
- [16] Zhu X, Lin H, Si Z, Xia Y, Chen W, Li X. Benzothiadiazole-Mediated Induced Resistance to *Colletotrichum musae* and Delayed Ripening of Harvested Banana Fruit. *J. Agric. Food Chem.* 2016; 64: 1494–1502. <https://doi.org/10.1021/acs.jafc.5b05655>
- [17] Kritzingner Q, Aveling TAS, Marasas WFO. Effect of essential plant oils on storage fungi, germination and emergence of cowpea seeds. *Seed Sci. Technol.* 2002; 30:609–619.
- [18] De Vincenzi M, Stammati A, De Vincenzi A, Silano M. Constituents of aromatic plants: carvacrol. *Fitoterapia.* 2004 ; 75:801–804. <https://doi.org/10.1016/J.FITOTE.2004.05.002>
- [19] Vitoratos A, Bilalis D, Karkanis A, Efthimiadou A. Antifungal Activity of Plant Essential Oils Against *Botrytis cinerea*, *Penicillium italicum* and *Penicillium digitatum*. *Not Bot Horti Agrobo.* 2013; 41(1):86-92.
- [20] Yousef SAA. Antifungal Activity of Volatiles from Lemongrass (*Cymbopogon citratus*) and Peppermint (*Mentha piperita*) Oils Against Some Respiratory Pathogenic Species of *Aspergillus*.

Int. J. Curr. Microbiol. Appl. Sci. 2013; 2:261–271.


- [21] Aoudou Y, Etoa E, Dady L. Effect of essential oil of *Cymbopogon citratus* on seed-born fungi and soybean seeds performance. *Int. J. Curr. Res.* 2017; 9.
- [22] Premathilake UGAT, Wathugala DL, Dharmadasa RM. Evaluation of chemical composition and assessment of antimicrobial activities of essential oil of lemongrass (*Cymbopogon citratus* (DC.) Stapf). *Int. J. Minor Fruits, Med. Aroma. Plants.* 2018; 4:13–19.
- [23] Mazari K, Bendimerad N, Bekhechi C, Fernandez X. Chemical composition and antimicrobial activity of essential oils isolated from Algerian *Juniperus phoenicea* L. and *Cupressus sempervirens* L. *J. Med. Plants Res.* 2010; 4: 959–964.
- [24] Ismail A, Lamia H, Mohsen H, Samia G, Bassem J. Chemical composition, bio-herbicidal and antifungal activities of essential oils isolated from Tunisian common cypress (*Cupressus sempervirens* L.) *J. Med. Plants Res.* 2013; 7:1070–1080. <https://doi.org/10.5897/JMPR12.1088>
- [25] Hsieh WC, Chang CP, Gao YL. Controlled release properties of Chitosan encapsulated volatile Citronella Oil microcapsules by thermal treatments. *Colloids Surfaces B Biointerfaces.* 2006; 53: 209–214. <https://doi.org/10.1016/J.COLSURFB.2006.09.008>
- [26] Baptiste Houndou Fokou J, Michel Jazet Dongmo P, Fekam Boyom F., 2020. Essential Oils' Chemical Composition and Pharmacological Properties. *Essent. Oils - Oils Nat.* <https://doi.org/10.5772/intechopen.86573>
- [27] Sessou P, Farougou S, Kaneho S, Djenontin S, Alitonou GA, Azokpota P, Youssao I, Sohounhloúé D. Bioefficacy of *Cymbopogon citratus* essential oil against foodborne pathogens in culture medium and in traditional cheese wagashi produced in Benin. *Int. Res. J. Microbiol.* 2012 ; 3: 406–415.
- [28] Rguez S, Djébalí N, Ben Slimene I, Abid G, Hammemi M, Chenenaoui S, Bachkouel S, Daami-Remadi M, Ksouri R, Hamrouni-Sellami I. *Cupressus sempervirens* essential oils and their major compounds successfully control postharvest grey mould disease of tomato. *Ind. Crops Prod.* 2018; 123: 135–141. <https://doi.org/10.1016/J.INDCROP.2018.06.060>
- [29] Maptue FB, Yaouba A, Ntsomboh-Ntsefong G. Antifungal Effect of Starch and Biochar Based Essential Oil Formulations in the Preservation of Cowpea (*Vigna unguiculata* (L.) Walp) Seeds. *Jacs.* 2021; 1: 1–10. <https://doi.org/10.17303/jacs.2021.1.101>
- [30] Ioannidou O, Zabaniotou A, Antonakou EV, Papazisi KM, Lappas AA, Athanassiou C. Investigating the potential for energy, fuel, materials and chemicals production from corn residues (cobs and stalks) by non-catalytic and catalytic pyrolysis in two reactor configurations. *Renew. Sustain. Energy Rev.* 2009; 13: 750–762. <https://doi.org/10.1016/J.RSER.2008.01.004>
- [31] Liu X, Zhang Yang Li Z, Feng R, Zhang, Yaozhong. Characterization of corncob-derived biochar and pyrolysis kinetics in comparison with corn stalk and sawdust. *Bioresour. Technol.* 2014; 170: 76–82. <https://doi.org/10.1016/J.BIORTECH.2014.07.077>
- [32] Djousse Kanouo BM., Allaire SE, Munson AD. Quality of Biochars Made from Eucalyptus Tree Bark and Corncob Using a Pilot-Scale Retort Kiln. *Waste and Biomass Valorization.* 2018 9: 899–909. <https://doi.org/10.1007/s12649-017-9884-2>
- [33] Camara A. Lutte Contre *Sitophilus Oryzae* L. (Coleoptera: Curculionidae) Et *Tribolium*

Castaneumherbst (Coleoptera: Tenebrionidae) Dans Les Stocks De Riz Par La Technique D'Étuvage Traditionnelle Pratiquée En Basse-Guinée Et L'Utilisation Des Huiles Essentielles Végé. Université du Québec, Montréal. 2009

- [34] ISTA. International rules for seed testing 2023, International rules for seed testing. Zurich, Switzerland. Zurich, Switzerland. 2023
- [35] Champion R. Identifier les champignons transmis par les semences - RÉMI CHAMPION - Achat Livre ou ebook | fnac, INRA. ed, INRA. INRA, Paris, France. 1997
- [36] Warham E, Butler L, Sutton B. Contrôle des semences de maïs et de blé: Guide de laboratoire, International Maize and Wheat Improvement Center. CIMMYT, Lisboa, Mexico. 1997. <https://doi.org/10.3/JQUERY-UI.JS>
- [37] Mathur SB, Kongsdal O. Common laboratory seed health testing methods for detecting fungi. Int. Seed Test. Assoc. Bassend. Switz. 2003. 425. <https://doi.org/10.3/JQUERY-UI.JS>
- [38] Liu WT, Chu CL, Zhou T. Thymol and acetic acid vapors reduce postharvest brown rot of apricots and plums. HortScience. 2002-; 37:151–156. <https://doi.org/10.21273/hortsci.37.1.151>
- [39] Smigielski K, Prusinowska R, Stobiecka A, Kunicka-Styczyńska A, Gruska R. Biological Properties and Chemical Composition of Essential Oils from Flowers and Aerial Parts of Lavender (*Lavandula angustifolia*). 2018—; 21—: 1303–1314. <https://doi.org/10.1080/0972060X.2018.1503068>
- [40] Tchoumboungang F, Michel Jazet Dongmo P, Lambert Sameza M, Gaby Nkouaya Mbanjo E, Bertrand Tiako Fotso G, Henri Amvam Zollo P, Menut C. B A S E Activité larvicide sur *Anopheles gambiae* Giles et composition chimique des huiles essentielles extraites de quatre plantes cultivées au Cameroun. Biotechnol. Agron. Soc. Env. 2009 ; 13 : 77–84.
- [41] Regnier T, Combrinck S, Veldman W, Du Plooy W. Application of essential oils as multi-target fungicides for the control of *Geotrichum citri-aurantii* and other postharvest pathogens of citrus. Ind. Crops Prod. 2014-; 61-: 151–159. <https://doi.org/10.1016/J.INDCROP.2014.05.052>
- [42] Thompson JD, Chalchat JC, Michet A, Linhart YB, Ehlers B. Qualitative and quantitative variation in monoterpene co-occurrence and composition in the essential oil of *Thymus vulgaris* chemotypes. J. Chem. Ecol. 2003 ; 29 : 859–880. <https://doi.org/10.1023>



**Figure 2:** Fungal infection rate of treated peanut seeds based on starch-based essential oil formulations and dose after 70 days of storage. A: Starch formulation at a dose of 3 ug/l. B: Starch formulation at a dose of 7 ug/l.



**Figure 3:** Fungal infection rate of treated peanut seeds based on biochar-based essential oil formulations and dose after 70 days of storage. A: Biochar formulation at a dose of 3 ug/l. B: Biochar formulation at a dose of 7 ug/l.