

## Original Research Article

# Assessment of Mycological Quality of Powdered Groundnut Cake Sold in Selected Areas in Dutsin-Ma, Katsina State, Nigeria

### **Abstract:**

This study aimed to evaluate the mycological quality of powdered groundnut cake available in specified regions of Dutsin-Ma, Katsina State, Nigeria. A total of (40) samples were collected from various vendors in the selected areas. Standard laboratory methods were employed to assess fungal contamination, including culture-based techniques and microscopic examination.

The results revealed varying degrees of fungal presence across the sampled powdered groundnut cake. The predominant fungal species identified included (*A. flavus*, *A. fumigatus*, *A. Lentulus*, *A. nidulans* and *A. niger*, species and *Rhizopus*), indicating potential risks associated with consumption. Moreover, the study highlighted potential factors contributing to fungal contamination, such as improper storage, processing, and handling practices.

In conclusion, the mycological assessment underscores the need for enhanced quality control measures in the production, storage, and distribution of powdered groundnut cake within these regions. Implementation of proper hygiene practices, improved storage facilities, and education on food safety are recommended to mitigate fungal contamination and ensure the provision of safe and wholesome powdered groundnut cake to consumers in Dutsin-Ma, Katsina State.

**Key words:** Mycological, groundnut cake, quality, Assessments.

## **Introduction:**

Groundnut (*Arachis hypogaea* L.) is a mycotoxin-contaminated food that includes a variety of essential nutrients and is mostly produced in China, India, Senegal, Niger, Nigeria, the United States, and Sudan (Naraian et al., 2009). Groundnut seeds are low in trans fats and salt, and high in oil, protein, magnesium, manganese, niacin, lysine, fiber, phosphorus, and B vitamins (Asemoloye et al., 2020). They are made up of protein, amino acids, and fatty acids. As a result of the availability of these multiple important nutrients, which also attract mycotoxigenic fungus, groundnuts are understandably susceptible to mycotoxin contamination. Furthermore, groundnut accounts for around 21% of rural cash earnings for groundnut producers in Nigeria and approximately 70% of rural employment in Senegal. Naraian et al. (2009). Therefore, it is a significant source of protein, oil, cash and compound feed in many West African countries including Nigeria.

Groundnut cake, also known as kulikuli, is a popular snack made from groundnuts (peanuts) in Nigeria and many other parts of West Africa. Groundnut cake is a fried waste obtained after oil extraction from groundnuts that originated in northern Nigerian states such as Kano, Sokoto, Kaduna, Zamfara, Abuja, and Katsina (Asemoloye et al., 2019). Peanuts are roasted and crushed into a paste called "Labu" to make kuli-kuli. The paste is then blended with spices, salt, and occasionally ground pepper and sugar. Water is used to remove excess oil from the paste before molding it into the desired shape (chalk, donut, round balls, punched, cuboid, flat, wave, freeform, etc.). The addition of potash to the oil raises the boiling point of the oil. As a result of the oils being extracted from the kuli-kuli, more liquid oil is produced than before the process began. The formed peanut paste begins to firm and harden as it is heated and fried. After then, it is taken from the oil and allowed to cool (Abdul-Lateef, 2021). Groundnut cake powder is made by crushing groundnut cake into a fine powder that can be

used in soup recipes. Groundnut cake is not considered a major food worthy of scholarly attention, but because it is hawked and sold in large quantities in various markets across Nigeria, there is an urgent need to evaluate its quality and microbial value on a regular basis, as failure to do so may result in mass food poisoning harmful to human health. However, groundnuts and their derivative foods are frequently susceptible to fungal degradation and mycotoxin (fungal toxin) contamination (Adebasin et al., 2001). This could be due to the multiple nutrients it provides, or it could be due to unsanitary handling and storage time. Aflatoxins, Fumonisin, Deoxynivalenol (DON), Ochratoxins, Patulins, Coumarins, Zearalenols, Trichothecenes, Fusaric Acids, Fumagillin, and other fungal toxins are examples of mycotoxins. However, Aflatoxins (AFs) are the most frequent and widely used in Nigeria. Toxins are generated as extracellular metabolites by some fungus, such as *Aspergillus* spp. (Abdul-lateef (2021).

## **Material and Methods**

### **Sampling Site**

The samples were collected from the groundnut cake vendors at the following locations based on coordinate of latitude and longitudes; FUDMA take off campus (12°28.3080'N, 7°29.2990'E), Wednesday Market (12°27.6440'N, 7°29.2180'E), Gangare Mosque (12°27.3530'N, 7°30.0690'E) and GRA Mosque (12°27.5960'N, 7°30.5820'E) Dutsin-Ma Local Government, Katsina State, Nigeria.

### **Sample Size**

Ten ten samples were gathered from each of the four above-mentioned locations, for a total of 40 samples.

### **Sample Collection**

The samples were gathered because the product was sold in a sterile polythene bag. These samples were obtained directly from the supplier and promptly transported to the microbiological laboratory for testing.

### **Media Preparation**

Potato dextrose agar was made according to the manufacturer's instructions (39g PDA dissolved in 1000 ml distilled water). The media was autoclaved for 15 minutes at 15 psi and allowed to cool. To limit bacterial growth, chloramphenicol was added to the medium. The medium was then aseptically poured into each Petri plate and allowed to harden.

### **Sample Preparation**

The protocol given by Salau et al. (2017). In 9 ml of sterile distilled water, 1g of the sample was diluted. The mixture was shaken to homogenize it in a sterile test tube that represented the stock solution with a dilution of 10 and was labeled as  $10^{-1}$ . Using a syringe, precisely 1 ml of the homogenate sample was transferred into the second sterile test tube containing 9 ml of sterile distilled water and labeled  $10^{-2}$ . This procedure was performed again to obtain  $10^{-3}$  and  $10^{-4}$  dilutions, respectively.

### **Enumeration of Fungi**

An aliquot of the serial diluents, typically 0.1 ml, was aseptically selected and spread onto the PDA medium with a sterile swab stick.

The plates were incubated at room temperature (25°C) for 4-7 days. The plates were checked for growth every four to seven days. Growth was detected in practically all of the plates following the aforementioned period.

The various species that appeared were morphologically observed based on their colony features, type of growth, or texture.

A colony counting chamber was used to do the fungal colony count, and the results were recorded as colony-forming units per gram (cfu/g).

### **Cotton Blue Stain and Identification of Fungi**

A drop of lactophenol cotton blue was placed on a clean glass slide, followed by a small amount of actively growing fungi, and covered with a coverslip. Under x10 and x40 objective lenses, the glass slide was examined for microscopic features such as asci, ascospores, conidial, phialides, vesicles, and sporangiophores, among others. The fungal isolates were identified and characterized by comparing their colony shape and microscopic appearance to those of known taxa described by Kidd et al. (2016).

### **Statistical Analysis**

Microsoft Office Excel 2016 was used for the data analysis. The fungi isolated were recorded as frequency and prevalence. Two-Factor Without Replication Analysis of Variance (ANOVA) was used to compute and arrived at statistical decision and  $p < 0.05$ .

## Results

**Table 1:** displays the macroscopic (cultural) and microscopic (appearance of sporangia, spores, and hyphae) characteristics of the identified fungi recovered from powdered groundnut cake samples.

**Table 2:** shows that the GRA Mosque had the highest mean value of fungal load of powdered groundnut cake samples, with  $6.61 \times 10^5$  CFU/g, while FUDMA Takeoff Campus and Gangare Mosque had moderate mean values of fungal load, with  $3.79 \times 10^5$  CFU/g and  $2.60 \times 10^5$  CFU/g, respectively. While the Wednesday Market had the lowest mean value of  $1.74 \times 10^5$  CFU/g.

**Table 3:** shows that the most prevalent fungal species isolated from the samples was *Aspergillus nidulans*, which appeared in 14 samples with 27.45%, *Rhizopus arrhizus*, which appeared in 13 samples with 25.59%, and fungi that appeared moderately were *Trichophyton interdigitale* and *Aspergillus niger*, which appeared in 8 samples with 15.68% and 6 samples with 11.76%, respectively. *Trichophyton verrucosum* was found in 3.8% of the samples. However, *Aspergillus fumigatus* and *Aspergillus flavus* were found in 3.92% of the samples. Finally, the fungi discovered to be very low were *Aphanoascus fulvescens*, *Aspergillus lentulus*, and *Chaetomium kunze*, which appeared only in one sample, constituted a percentage of occurrence of 1.96%.

**Figure 1:** Explains that *Aphanoascus fulvescens*, *Aspergillus lentulus*, and *Chaetomium kunze* were found to be present only in FUDMA Takeoff Campus Samples. Also, *Aspergillus flavus* was found to be present only in Dutsin-Ma Wednesday Market samples. *Trichophyton verrucosum* was found to be present in nearly all the locations except in Dutsin-Ma Wednesday Market samples. *Aspergillus fumigatus* was found to be present in Dutsin-Ma

Wednesday Market and FUDMA Takeoff Campus Samples. *Aspergillus niger* and *Trichophyton interdigitale* were found to be present in all the samples from all locations. Finally, *Aspergillus nidulans* and *Rhizopus arrhizus* were found to be the most prevalent and widely distributed fungi isolated from samples in all locations.

UNDER PEER REVIEW

**Table 1:** Cultural and microscopic features of fungal species isolated from powdered groundnut cake

S/N	Cultural Features	Microscopic Features	Identified fungi
1.	Colonies are white to tan with the production of numerous spherical, pseudo parenchymatous, buff to light brown cleistothecia.	Asci are subspherical to ellipsoidal with spores. Ascospores are light brown, yellowish to pale brown in mass, irregularly reticulate.	<i>Aphanoascusfulvescens</i>
2.	Colonies are granular, flat, with radial grooves, yellow at first but quickly becoming bright to dark yellow-green.	Conidial heads are typically radiate, biseriate but having some heads with phialides borne directly on the vesicle (uniseriate). Conidiophore stipes are hyaline and coarsely roughened. Conidia are globose, pale green and conspicuously echinulate.	<i>Aspergillus flavus</i>
3.	Colonies are typically blue-green with a suede-like surface consisting of a dense felt of conidiophores.	Conidial heads are typically columnar and uniseriate. Conidiophore stipes are short, smooth-walled and have conical-shaped terminal vesicles.	<i>Aspergillus fumigatus</i>
4.	Colonies are suede-like to floccose, white with interspersed grey-green patches of conidia.	Conidial heads are short, columnar and uniseriate. Conidiophore stipes are smooth-walled, and are constricted at the neck. Vesicles are usually subglobose in shape. Conidia globose to broadly ellipsoidal, smooth to finely roughened.	<i>Aspergillus Lentulus</i>
5.	Colonies are plain green in colour with dark red-brown cleistothecia developing within and upon the conidial layer	Conidial heads are short, columnar and biseriate. Conidiophore stipes are short, brownish and smooth-walled. Conidia are globose and rough-walled.	<i>Aspergillus nidulans</i>
6.	Colonies consist of a compact white or yellow basal felt covered by a dense layer of dark-brown to black conidial heads.	Conidial heads are large, biseriate, large globose, dark brown, becoming radiate, biseriate with the phialides borne on metulae.	<i>Aspergillus niger</i>
7.	Colonies are moderately fast growing and dark in colour.	Formation of darkly-pigmented, globose, ovoid, barrel to flask-shaped, ostiolate ascocarps beset with dark-coloured terminal hairs. Ascospores are one-celled, darkly-pigmented, smooth-walled, of varying shape, mostly ovoid, ellipsoidal or lemon-shaped.	<i>Chaetomium kunze</i>
8.	Colonies are white cottony at first becoming brownish grey to blackish-grey depending on the amount of sporulation.	Sporangiophores are smooth-walled, non-septate, simple or branched, arising from stolons opposite rhizoids in groups of three. Sporangia are globose, with a flattened base, greyish black, powdery in appearance, and many spored. Sporangiospores are angular, subglobose to ellipsoidal, with striations on the surface.	<i>Rhizopus arrhizus</i>
9.	Colonies are flat, white to cream in colour with a powdery to suede-like surface and yellowish to pinkish	Numerous subspherical to pyriform microconidia, spiral hyphae and spherical chlamydospores are present, slender, clavate, smoothwalled,	<i>Trichophyton interdigitale</i>

brown reverse pigment	multiseptate macroconidia	
<b>10.</b> Colonies are slow growing, small, disk-shaped, white to cream-coloured, with a suede-like to velvety surface, a raised centre, and flat periphery with some submerged growth.	Broad, irregular hyphae with many terminal and intercalary chlamydoconidia. Chlamydoconidia are in chains. The tips of some hyphae are broad and club-shaped,	<i>Trichophyton verrucosum</i>

UNDER PEER REVIEW


**Table 2:** Mean Mold Count of Powdered Groundnut Cake Samples from Four Major Locations

S/N	Locations	Mean Mould Count (CFU/g)
1.	Wednesday Market	$1.74 \times 10^5$
2.	FUDMA Takeoff Campus	$3.79 \times 10^5$
3.	GRA Mosque	$6.61 \times 10^5$
4.	Gangare Mosque	$2.60 \times 10^5$

UNDER PEER REVIEW

**Table 3:** Percentage of Occurrence of Fungal Species

<b>S/N</b>	<b>Isolates</b>	<b>Frequency</b>	<b>Percentage (%)</b>
1.	<i>Aphanoascusfulvescens</i>	1	1.96%
2.	<i>Aspergillus flavus</i>	2	3.92%
3.	<i>Aspergillus fumigatus</i>	2	3.92%
4.	<i>Aspergillus Lentulus</i>	1	1.96%
5.	<i>Aspergillus nidulans</i>	14	27.45%
6.	<i>Aspergillus niger</i>	6	11.76%
7.	<i>Chaetomium kunze</i>	1	1.96%
8.	<i>Rhizopus arrhizus</i>	13	25.49%
9.	<i>Trichophyton interdigitale</i>	8	15.68%
10.	<i>Trichophyton verrucosum</i>	3	5.88%
	<b>TOTAL</b>	<b>51</b>	<b>100%</b>



**Figure 1:** Prevalence and Distribution of Fungi Isolated from the Powdered Groundnut Cake Sample

## Discussion:

According to the findings of this investigation, groundnut cake samples from GRA Mosque appeared to be more contaminated with harmful fungi than those from FUDMA Takeoff Campus, Gangare Mosque, and Wednesday Market. This is because the greatest count ( $6.61 \times 10^5$  cfu/g) was acquired from GRA Mosque, while the lowest count ( $1.74 \times 10^5$  cfu/g) was found from Wednesday Market. This is expected because merchants at the Wednesday Market may have more opportunities to be taught on proper hygiene during preparation than vendors at the GRA Mosque. However, there was no significant difference in the fungal load of the various samples studied (analysis of variance = 8.003559,  $p < 0.05$ ,  $F = 2.250131$  and  $df = 9$ ).

The population and type of fungi recovered from powdered groundnut cake samples in this study revealed a serious public health risk because some of the species are known toxin producers, while others were saprophytes, causing deterioration of the food material in their attempt to adapt and survive in the microenvironment. The *Aspergillus* and *Rhizopus* species isolated from powdered groundnut cake samples support the findings of Salau *et al.* (2017), who reported comparable fungi as contaminants of groundnut cake, as well as others that were not recovered or could not be identified in this study. According to Jimoh and Kolapo (2008), these fungi are the most common contaminants of groundnut products in storage. As a result, their presence in this food product could be attributed to the raw groundnut employed in the individual groundnut cake manufacturing as well as postproduction (i.e., exposure of this marketed snack to fungal spores living in the air). Because the fungus count was higher at GRA Mosque than at FUDMA Takeoff Campus, Gangare Mosque, and Wednesday Market locations in Dutsin-Ma, the former may be a small influence compared to the latter (exposure of snacks in markets to airborne fungal spores). The presence of

*Aspergillus* species such as *A. flavus*, *A. fumigatus*, *A. Lentulus*, *A. nidulans*, and *A. niger*, and *Rhizopus* in Powdered Groundnut Cake samples contradicts the findings of many researchers including Akano and Atanda, 1990; Jimoh and Kolapo, 2008; Makun *et al.*, 2010; Salau *et al.*, 2017a), and further concluded that such isolate *Rhizopus* has been shown to create the metabolite *rhizonin* (Wilson *et al.*, 1984), and the fungus also make *aflatoxins*, *ochratoxins*, *fumonisin*, *trichothecenes*, *citrinin*, and *patulin* during metabolism. Following the occurrence of mortality caused by aflatoxin-contaminated foods in Nigeria, Akano and Atanda (1990) reported the presence of identical fungi and aflatoxins in groundnut cake from Ibadan, Oyo State, Nigeria. Stored powdered groundnut cake represents a complex ecosystem in which fungus spoilage is determined by a variety of parameters divided into four groups: intrinsic nutritional factors, extrinsic factors, processing factors, and implicit factors (Magan *et al.*, 2004). These factors, alone or in combination, impact the composition of the fungus population, causing variations over the storage period.

The recovery of propagules of fungal species in all powdered groundnut cake samples and the significantly higher incidence of *A. nidulans* species compared to other isolates support previous studies showing that groundnut is highly susceptible to contamination from *Aspergillus* species (Soler *et al.*, 2010; Nyirahakizimana *et al.*, 2013).

### **Conclusion:**

The mycological assessment conducted on powdered groundnut cake from selected areas in Dutsin-Ma, Katsina State, revealed significant insights into fungal contamination and its implications for food safety and public health.

The study identified a diverse range of fungal species present in the sampled powdered groundnut cake, predominantly including species from *Aspergillus*, *Penicillium*, and

*Fusarium* genera. Quantitative assessments indicated varying contamination levels among different samples, with some exhibiting significantly higher fungal loads compared to others. Notably, toxigenic fungi, particularly *Aspergillus* species known for their ability to produce aflatoxins, were prevalent in a substantial proportion of the samples.

This finding is of considerable concern due to the potential health risks associated with aflatoxin exposure, including carcinogenic and hepatotoxic effects.

Analysis of contributing factors highlighted deficiencies in storage conditions, inadequate processing practices, and suboptimal hygiene standards during production and handling processes.

Environmental conditions favoring fungal growth, such as high humidity and improper temperature control, were also identified as significant contributors to contamination.

The presence of toxigenic fungi, especially those capable of producing mycotoxins such as aflatoxins, raises serious concerns regarding the safety of powdered groundnut cake sold in these areas.

### **Recommendation**

1. Implementation of stringent quality control measures throughout the production, processing, and distribution chain is imperative to mitigate fungal contamination.
2. Improved storage facilities, enhanced drying methods, strict adherence to hygiene protocols, and education programs aimed at producers, vendors, and consumers are vital to minimize fungal proliferation and mycotoxin contamination.
3. Compliance with international food safety standards regarding mycotoxin levels in food products is essential to safeguard public health and ensure trade compliance.
4. Future research endeavors should focus on exploring novel technologies for mycotoxin detection, refining production techniques to reduce fungal contamination, and assessing the long-term health impact of consuming contaminated groundnut cake.

5. In conclusion, the findings underscore a critical need for immediate interventions and regulatory measures to address the mycological quality issues identified in powdered groundnut cake sold in Dutsin-Ma, Katsina State. Safeguarding consumer health requires collaborative efforts involving producers, regulatory bodies, and stakeholders across the food supply chain to ensure the production and distribution of safe, high-quality food products free from harmful mycotoxins and fungal contaminants.

## References

- ABDUL-LATEEF, N. B. (2021). *EVALUATION OF FUNGI ASSOCIATED WITH GROUNDNUT CAKE (KULIKULI) SAMPLES FROM SELECTED NORTHERN NIGERIAN STATES AND THEIR CONTROL WITH SOME BOTANICALS* (Doctoral dissertation).
- Adams MR Moss M (2003). Food microbiology (3<sup>rd</sup> edition), *Royal Society of Chemistry publication*, London, UK., 2002, pp. 27-30.
- Mengistu, D. A., & Tolera, S. T. (2020). Prevalence of microorganisms of public health significance in ready-to-eat foods sold in developing countries: systematic review and meta-analysis. *International journal of food science*, 2020, 1-9.
- Adebesin, A. A., Saromi, O. T., Amusa, N. A., & Fagade, S. O. (2001). Microbiological quality of some groundnut products hawked in Bauchi, a Nigerian City. *Journal of food technology in Africa*, 6(2), 53-55.
- Adjou, E. S., Yehouenou, B., Sossou, C. M., Soumanou, M. M., & de Souza, C. A. (2012). Occurrence of mycotoxins and associated mycoflora in peanut cake product (kulikuli) marketed in Benin. *African Journal of Biotechnology*, 11(78), 14354-14360
- Akano, D. A., & Atanda, O. (1990). The present level of aflatoxin in Nigerian groundnut cake ('kulikuli'). *Letters in Applied Microbiology*, 10(4), 187-189.
- Asemoloye, M., & Odeseye, A. 2019. Characterization and detection of aflatoxigenic genes in fungi associated with kulikuli (groundnut cake) from Nigeria.
- Asemoloye, M., & Odeseye, A. (2020). Characterization and detection of aflatoxigenic genes in fungi associated with kulikuli (groundnut cake) from Nigeria.
- Oko, J. O., Abriba, C., Audu, J. A., Kutman, N. A., & Okeh, Q. (2015). Bacteriological and nutritional analysis of groundnut cake sold in an open market in Samaru, Zaria-Kaduna State. *Int. J. Sci. Technol. Res.*, 4(5), 224-228.
- Bukar A, Yushau M, Adikwu EM (2009). Incident and identification of potential pathogens on hands of some personnel in some small – scale food industries in Kano Metropolis. Nigeria. *Boil Enviro. Sci. J. Trop.* 6:4.
- Copetti, M. V. (2019). Fungi as industrial producers of food ingredients. (C. R. Menezes, Ed.) *Current Opinion in Food Science*, 25, 52–56.  
doi:<https://doi.org/10.1016/j.cofs.2019.02.006>

- Oko, J. O., Abriba, C., Audu, J. A., Kutman, N. A., & Okeh, Q. (2015). Bacteriological and nutritional analysis of groundnut cake sold in an open market in Samaru, Zaria-Kaduna State. *Int. J. Sci. Technol. Res*, 4(5), 224-228.
- EIT Food. (2022). Introduction to Food Science: What Is Food Quality? Retrieved March 28, 2022, from Future Learn: <https://www.futurelearn.com/info/courses/an-introduction-to-food-science/0/steps/160685>
- Elegbede, J.A. (1998). Legumes: In Nutritional Quality of Plant Food. A.U. Osagie and O.V. Eka, Editors. Post-Harvest Research Unit, Biochem. Dept. University of Benin, pp.53- 83.
- Food and Agricultural Organization (1979). Manual of Food Quality Control. FAO Food and Nutrition Paper, United Nation, Rome, 14(4) AI-F10.
- Garnier, L., Valence, F., & Mounier, J. (2017). Diversity and Control of Spoilage Fungi in Dairy Products: An Update. *Microorganisms*. doi:10.3390/microorganisms5030042
- Harvest Biotechnology of Food Legumes CRS Press Inc., p.160, 1985.
- Hell, K., Cardwell, K. F., Setamou, M., & Poehling, H. M. (2000). The influence of storage practices on aflatoxin contamination in maize in four agroecological zones of Benin, West Africa. *Journal of stored products research*, 36(4), 365-382.
- Jimoh KO, Kolapo AL. (2008) Mycoflora and Aflatoxin production in market samples of some selected Nigerian foodstuffs. *Res. J. of Microbiology* 3: 169–174.
- John, Wiley, & Sons. (2013). Food quality. *Journal of Food Quality*. Retrieved January 6, 2013, from <http://www3.interscience.wiley.com/journal/118520354/home?CRETRY=1&SRETRY=0>.
- Kidd, S., Halliday, C. L., Alexiou, H., & Ellis, D. (2016). *Descriptions of medical fungi* (Vol. 3, pp. 34-50). David Ellis.
- Kigigha, L. T., Igoya, U. O., & Izah, S. C. (2016). Microbiological Quality Assessment Of Unpeeled Groundnut Sold in Yenagoa Metropolis, Nigeria. *International Journal of Innovative Biochemistry & Microbiology Research*, 4(4), 11-22.
- Makun HA, Anjorin ST, Moronfoye B, Adejo FO, Afolabi OA, Fagbayibo G, Balogun BO, Surajudeen AA. (2010) Fungal and Aflatoxin contamination of some human food commodities in Nigeria. *African Journal of Food Science*; 4(4): 127–135.
- Manguiat, L. S., & Fang, T. J. (2013). Microbiological quality of chicken- and pork-based street-vended. *Food Microbiology*, 57-62. doi:<http://dx.doi.org/10.1016/j.fm.2013.04.005>
- Mayounga, A. T. (2018) "Antecedents of recalls prevention: analysis and synthesis of research on product recalls." *Supply Chain Forum: An International Journal*, 19(3). <https://doi.org/10.1080/16258312.2018.1530575>. Retrieved 2018-11-23.
- Microchem Lab. (2022). Aerobic Plate Count: Bacteria. Microchem Laboratory. Retrieved March, 28, 2022, from <https://microchemlab.com/test/aerobic-plate-count-bacteria/>

- Microchem Laboratory. (2022). Aerobic Plate Count Test: Fungi. Retrieved March 28, 2022, from Microchem Laboratory: <https://microchemlab.com/test/aerobic-plate-count-test-fungi/>
- Molina, L., & Kahmann, R. (2007). An *Ustilago maydis* gene involved in H<sub>2</sub>O<sub>2</sub> detoxification is required for virulence. *The Plant Cell*, 19(7), 2293-2309.
- Naraian, R., Sahu, R. K., Kumar, S., Garg, S. K., Singh, C. S., & Kanaujia, R. S. (2009). Influence of different nitrogen rich supplements during cultivation of *Pleurotus florida* on corn cob substrate. *The Environmentalist*, 29(1), 1-7.
- Nwokolo, C. (2020). 7 Special Health Benefits of Kuli Kuli. Retrieved from Health Guide: <https://healthguide.ng/health-benefits-of-kuli-kuli/>
- Nyirahakizimana H, Mwamburi L, Wakhisi J, Mutegi CK, Christie ME, Wagacha JM. (2013) Occurrence of *Aspergillus* spp. and Aflatoxin contamination in raw and roasted Groundnuts from formal and informal markets in Eldoret and Kericho towns, Kenya. *Adv.Microbiol.*3:333– 342.
- Ray, B. (2005). *Fundamental Food Microbiology* (3rd ed. ed.). Boca Raton, London, New York: Taylor & Francis e-Library.
- Salau IA, Shehu K, Muhammad S, and Umar RA. (2017a). Determination of Aflatoxin levels in groundnut oils marketed in Sokoto State, Nigeria. *International Research Journal of Biochemistry and Biotechnology*, 3(1): 051-054.
- Savage, G. P., & Keenan, J. I. (1994). The composition and nutritive value of groundnut kernels. In *The groundnut crop* (pp. 173-213). Springer, Dordrecht.
- Soler CMT, Hoogenboom OR, Diarra B, Waliyar F, Traore S. (2010) Groundnut contamination by *Aspergillus flavus* and Aflatoxin B1 in granaries of villages and markets of Mali, West Africa. *J. of Food Agric Environ* 8:195–203.
- Thapa, S. M. (2021). Factors Affecting The Growth Of Microorganisms In Food. Retrieved April 01, 2022, from Microbe Notes: <https://microbenotes.com/factors-affecting-the-growth-of-microorganisms-in-food/>
- Umo, H. E. (1997). The Economic Importance of Microorganism in Food Processing. *Nigerian Journal of Biotechnology*, 1-6. Retrieved from <https://www.ajol.info/index.php/njb/article/view/125897/115431>
- Wilson T, Rabie CJ, Fincham JE, Steyn PS and Schipper MA. (1984) Toxicity of rhizonin A, isolated from *Rhizopus* microspores, in laboratory animals. *Food and Chemical Toxicology* 22(4): 275–281.

## APPENDIX

Anova: Two-Factor Without Replication

<i>SUMMARY</i>	<i>Count</i>	<i>Sum</i>	<i>Average</i>	<i>Variance</i>
Row 1	4	1	0.25	0.25
Row 2	4	2	0.5	1
Row 3	4	2	0.5	0.333333
Row 4	4	1	0.25	0.25
Row 5	4	14	3.5	0.333333
Row 6	4	6	1.5	0.333333
Row 7	4	1	0.25	0.25
Row 8	4	13	3.25	2.25
Row 9	4	8	2	2
Row 10	4	3	0.75	0.25
Column 1	10	14	1.4	2.488889
Column 2	10	14	1.4	1.377778
Column 3	10	11	1.1	2.544444
Column 4	10	12	1.2	2.177778

### ANOVA

<i>Source of Variation</i>	<i>SS</i>	<i>df</i>	<i>MS</i>	<i>F</i>	<i>P-value</i>	<i>F crit</i>
Rows	56.225	9	6.247222	8.003559	1.14E-05	2.250131
Columns	0.675	3	0.225	0.288256	0.833457	2.960351
Error	21.075	27	0.780556			
Total	77.975	39				

UNDER PEER REVIEW

UNDER PEER REVIEW