

Orthonormal bases on $L^2(\mathbb{R}^+)$

Abstract

In addition to orthogonal polynomials, orthogonal functions also play an important role. Their applications are, among others, in the fields of signal and data analysis, dynamic modeling. They are related to the solution of differential equations. In this paper we derive the explicit form of one parameter family of orthonormal bases on space $L^2(\mathbb{R}^+)$. The bases are formed by eigenvectors of the self-adjoint extension H_ξ , parametrized by $\xi \in \langle 0, \pi \rangle$, of differential expression $H = -\frac{d^2}{dx^2} + \frac{x^2}{4}$ together with the spectrum $\sigma(H_\xi)$ on the space $L^2(\mathbb{R}^+)$. For each ξ the set of eigenvectors form an orthonormal basis of $L^2(\mathbb{R}^+)$. From the physical point of view, it is a solution of the Schrödinger equation of a harmonic oscillator on a semi-straight line.

1 Introduction

Base of orthogonal functions play essential role in functional data analysis [1]. It is observed that the choice of basis for a given data has an influence on the efficiency of initial data processing.

Another application may be in solving a rational approximation problem using nonlinear parameters see [2].

Orthogonal functions as a eigenfunctions of a generalized second-order differential equation is given in [3]. Well-known orthogonal polynomial systems and orthogonal functional systems are eigenfunctions of differential equation of form

$$f(x)y'' + g(x)y' + \lambda y = 0.$$

A family of orthogonal systems gives some possibilities of using. Moreover, the connection of parabolic cylinder functions with gamma functions gives a decomposition of gamma functions, and it can be applied to Fourier transforms as in [4]. This family of orthogonal functions can be used in a similar way as in [5], but in the space $L^2(\mathbb{R}^+)$. An overview of the use of orthogonal functions can be found in [6].

The Schrödinger operator of perturbed harmonic oscillator acting in the space $L^2(\mathbb{R}^3)$ can be transformed into a direct sum of the Schrödinger operators acting on $L^2(\mathbb{R}^+)$ [7]. To calculate the eigenvalues and eigenfunctions of the operator corresponding to the harmonic oscillator on the half-line, we do not use the Friedrichs extension as in [8]. Therefore we get the full spectrum of the operator.

The basic examples of quantum mechanics is a quantization the harmonic oscillator. A selfadjoint Hamiltonian H_D of the one-dimensional linear harmonic oscillator is generated by the differential expression

$$H = -\frac{d^2}{dx^2} + \frac{x^2}{4} \quad (1)$$

with appropriate definition domain D . It is known that the operator H_D has a pure point spectrum and its eigenfunctions form the orthonormal basis in $L^2(\mathbb{R})$, and H_D is a unique selfadjoint operator generated by H on $L^2(\mathbb{R})$.

The situation is quite different on $L^2(\mathbb{R}^+)$, there is one-parametric set of selfadjoint operators H_ξ , $\xi \in \langle 0, \pi \rangle$ with corresponding definition domains D_ξ and with the same differential expression (1) [9]. All these selfadjoint operators are selfadjoint extensions of the closed symmetric operator \hat{H} with the domain $\hat{D} = \bigcap_{\xi \in \langle 0, \pi \rangle} D_\xi$. Following the theorem [10] all these extension have the same essential spectrum. As in the case of the operator $H_{\xi=0}$, where it applies $\sigma_{ess}(H_{\xi=0}) = \emptyset$, it applies for all operators H_ξ , $\xi \in \langle 0, \pi \rangle$. In other words, for any $\xi \in \langle 0, \pi \rangle$ there exist an orthonormal basis formed by eigenvectors of H_ξ . The objective of this paper is to derive explicit form of the orthonormal basis and express $\sigma(H_\xi)$.

2 Parabolic cylinder functions

Since the core of this paper is parabolic cylinder functions, we first review their properties and the relationships we will used [11, 12, 13].

The parabolic cylinder functions

$$D_\nu(x) = e^{-\frac{x^2}{4}} \left[\frac{\sqrt{\pi} 2^{\frac{\nu}{2}}}{\Gamma(\frac{1-\nu}{2})} {}_1\Phi_1\left(-\frac{\nu}{2}, \frac{1}{2}, \frac{x^2}{2}\right) - \frac{\sqrt{\pi} 2^{\frac{\nu+1}{2}}}{\Gamma(\frac{-\nu}{2})} x {}_1\Phi_1\left(\frac{1-\nu}{2}, \frac{3}{2}, \frac{x^2}{2}\right) \right] \quad (2)$$

are the solutions of the Weber differential equation [11, 12]

$$\left(\frac{d^2}{dx^2} - \frac{x^2}{4} + \nu + \frac{1}{2}\right)D_\nu(x) = 0.$$

that converge to 0 at $+\infty$. These functions are expressed using gamma functions Γ and hypergeometric series Φ . Values $\nu \in \{0, 1, 2, \dots\} \equiv \mathbb{N}_0$

need special attention, because of

$$\frac{1}{\Gamma(\frac{-\nu}{2})} = 0, \quad \Gamma(\frac{1-\nu}{2}) = \infty \quad \text{for } \nu = 1, 3, 5, \dots$$

and

$$\frac{1}{\Gamma(\frac{-\nu}{2})} = \infty, \quad \Gamma(\frac{1-\nu}{2}) = 0 \quad \text{for } \nu = 0, 2, 4, \dots$$

Definition (2) then gives $D_\nu(x) = h_\nu(x)$ known Hermitian functions [11]. For these cases, the parabolic functions are Hermitian functions, and the orthogonal bases on $L^2(\mathbb{R}^+)$ are odd or even Hermitian functions.

The following relations holds for parabolic cylinder functions [11, 13]:

$$\int_0^\infty |D_\nu(x)|^2 dx = \frac{1}{c(\nu)^2}, \quad c(\nu) = \sqrt{\sqrt{\frac{2}{\pi}} \frac{\Gamma(-\nu)}{\beta(-\nu)}}, \quad \beta(-\nu) = \sum_{k=0}^\infty \frac{(-1)^k}{-\nu + k} \quad (3)$$

(note that $c(\nu) D_\nu$ is normalized), and

$$\int_0^\infty D_\nu(x) D_\mu(x) dx = \frac{\pi 2^{\frac{1}{2}(\nu+\mu+1)}}{\mu - \nu} \left[\frac{1}{\Gamma(\frac{1-\mu}{2})\Gamma(-\frac{\nu}{2})} - \frac{1}{\Gamma(\frac{1-\nu}{2})\Gamma(-\frac{\mu}{2})} \right]. \quad (4)$$

3 Family of orthogonal functions

The derivation of the family will be formulated in the form of two Theorems. The derivation of this will be formulated in the form of two sentences. First, we will say two preparatory lemmas. It is known [9] that the differential expression (1)

$$H = -\frac{d^2}{dx^2} + \frac{x^2}{4}$$

with definition domain

$$D_\xi(H) := \{f \in \tilde{\mathcal{D}}, f(0) \cos \xi - f'(0) \sin \xi = 0\}, \quad (5)$$

is a selfadjoint operator on $L^2(\mathbb{R}^+)$ for all $\xi \in \langle 0, \pi \rangle$, and $\tilde{\mathcal{D}} = \{f, f' \in a.c.(0, \infty) : f, Hf \in L^2(\mathbb{R}^+)\}$

So, if D_ν will belong to $\mathcal{D}_\xi(H)$ for some ν then D_ν will be an eigenvector of the considered selfadjoint operator with eigenvalues $\nu + 1/2$. Eq. (3) guarantees that D_ν lies in $L^2(\mathbb{R}^+)$.

The last condition generates the relationship

$$D_\nu(0) \cos \xi - D'_\nu(0) \sin \xi = 0. \quad (6)$$

Although, values $D_\nu(0)$ and $D'_\nu(0)$ can be calculated using definition (2), we have to distinguish two cases:

1. when $\nu \notin \mathbb{N}_0$ we obtain

$$\eta \Gamma\left(-\frac{\nu}{2}\right) - \Gamma\left(\frac{1-\nu}{2}\right) = 0, \quad \eta = \frac{1}{\sqrt{2}} \cot \xi. \quad (7)$$

2. when $\nu \in \mathbb{N}_0$ we obtain

$$h_\nu(0) \cos \xi - h'_\nu(0) \sin \xi = 0. \quad (8)$$

If ν is odd, then $h_\nu(0) = 0$, $h'_\nu(0) = 1$, and Eq. (8) is fulfilled only if $\xi = 0$. If ν is even, then $h_\nu(0) = 1$, $h'_\nu(0) = 0$, and Eq. (8) is fulfilled only if $\xi = \frac{\pi}{2}$. In both cases, condition (8) is fulfilled by the set of Hermitian functions $\{h_0, h_2, h_4, \dots\}$ and $\{h_1, h_3, h_5, \dots\}$, respectively. It is known that both sets form orthonormal bases in $L^2(\mathbb{R}^+)$.

Eq.(7) has to be solved for ν .

First we prove two lemmas.

Lemma 1:

1. If $\nu \in (2M - 1, 2M)$, $M = 1, 2, \dots$ or $\nu < 0$, then $\beta(-\nu) \geq 0$,
2. If $\nu \in (2M - 2, 2M - 1)$, $M = 1, 2, \dots$, then $\beta(-\nu) < 0$.

Proof:

1. Using the relationship

$$\beta(-\nu) = \sum_{k=0}^{\infty} \frac{1}{(-\nu + 2k)(-\nu + 2k + 1)},$$

[11], it is possible to show by elementary calculation that $(-\nu + 2k)(-\nu + 2k + 1) > 0$ for all $k = 0, 1, \dots$ if $\nu \in (2M - 1, 2M)$, $M = 0, 1, \dots$, or $\nu < 0$.

2. In this case we rewrite the sum $\beta(-\nu)$ in the following form:

$$\begin{aligned} \beta(-\nu) &= -\frac{1}{\nu} + \sum_{k=0}^{\infty} \frac{1}{-\nu + k + 1} + \frac{1}{-\nu + k + 2} = \\ &= -\frac{1}{\nu} - \sum_{k=0}^{\infty} \frac{1}{(-\nu + k + 1)(-\nu + k + 2)}. \end{aligned}$$

For considered values of $\nu \in (2M - 2, 2M - 1)$, $M = 1, 2, \dots$ the products $(-\nu + k + 1)(-\nu + k + 2)$ are positive, and therefore all denominators

of the members in the previous sum are positive and so $\beta \leq 0$ (note that $-\frac{1}{\nu} < 0$). \square

Remark: Comparing functions $\Gamma(-\nu)$ and $\beta(-\nu)$ we have the relationship

$$\operatorname{sgn}(\Gamma(-\nu)) = \operatorname{sgn}(\beta(-\nu)), \nu \in \mathbb{R}.$$

It shows that normalization factor $c(\nu)$ (Eq.3) is correctly defined.

Lemma 2:

Function $y(\nu) := \frac{\Gamma(\frac{1-\nu}{2})}{\Gamma(-\frac{\nu}{2})}$ has the following properties:

1. There are asymptotes for $\nu_{\text{as}} \in \{2n+1 | n \in \mathbb{N}_0\}$ and $\lim_{\nu \rightarrow \nu_{\text{as}}^+} y(\nu) = \infty$, $\lim_{\nu \rightarrow \nu_{\text{as}}^-} y(\nu) = -\infty$. Further $\lim_{\nu \rightarrow -\infty} y(\nu) = +\infty$.
2. The set $\{2n | n \in \mathbb{N}_0\}$ consists of all zero points of y .
3. In the intervals $(-\infty, 1)$, $(M, M+1)$, $M = 0, 1, \dots$, y is continuous decreasing function.

Proof:

1. The first assertion is a direct consequence of explicit form [11] of function Γ .

For the remaining assertions it is sufficient to prove that the sequence $\{y(-2n) | n \in \mathbb{N}_0\}$ is growing and $\lim_{n \rightarrow \infty} y(-2n) = +\infty$. As $y(-2n)$ can be expressed as

$$y(-2n) = \frac{\Gamma(n + \frac{1}{2})}{\Gamma(n)} = \frac{\sqrt{\pi}(2n-1)!!}{2^n(n-1)!}$$

[11], the assertion can be easily verified.

2. Γ - function has no zero points. Therefore $y(\nu) = 0$ only if $|\Gamma(-\frac{\nu}{2})| = \infty$, i. e. $\nu = 2n$.

3. For $y'(\nu)$ we obtain

$$\frac{dy(\nu)}{d\nu} = \frac{1}{2} \frac{\Gamma(\frac{1-\nu}{2})}{\Gamma(-\frac{\nu}{2})} [\psi(-\frac{\nu}{2}) - \psi(\frac{1-\nu}{2})], \quad \psi(\mu) = \frac{d}{d\mu} \lg \Gamma(\mu).$$

Using the relationships

$$\psi(-\frac{\nu}{2}) - \psi(\frac{1-\nu}{2}) = -2\beta(-\nu), \quad \text{and} \quad \Gamma(-\nu) = \frac{2^{-\nu-1}}{\sqrt{\pi}} \Gamma(\frac{1-\nu}{2}) \Gamma(-\frac{\nu}{2}),$$

[11] we obtain

$$\frac{dy(\nu)}{d\nu} = -2^\nu \frac{\sqrt{\pi}}{\Gamma(-\frac{\nu}{2})^2} \Gamma(-\nu) \beta(-\nu).$$

As $\Gamma(-\nu) \beta(-\nu) > 0$ (see Remark) the proof is completed. \square

The consequence of this Lemma is a Theorem

Theorem 1:

For any $\eta \in \mathbb{R}$ and any $M \in \mathbb{N} = \{1, 2, \dots\}$ there is just one solution $\nu_\eta^{(M)}$ of Eq. (7) in the interval I_M , where

$$I_1 = (-\infty, 1), \quad I_M = (2M - 1, 2M + 1), \quad M = 2, 3, \dots$$

No further solution of Eq. (7) exists.

Table 1: Example of first 11 values of ν_η . The columns show the eleven lowest values of ν for the corresponding parameter η . The PCFs for a given D_{ν_η} are the first 11 functions of the orthogonal basis.

$\nu_{-2.18}$	$\nu_{-0.51}$	ν_0	$\nu_{0.23}$	$\nu_{0.51}$	$\nu_{0.97}$	$\nu_{2.18}$
0.77051	0.399912	0	-0.311391	-0.875066	-2.33401	-9.95
2.66471	2.26065	2.	1.86885	1.71369	1.5141	1.26337
4.59639	4.20523	4.	3.90249	3.78578	3.62177	3.36297
6.54652	6.1743	6.	5.91892	5.82117	5.67849	5.42659
8.50776	8.15402	8.	7.92911	7.84326	7.715227	7.47292
10.4764	10.1394	10.	9.93622	9.85874	9.74156	9.50897
12.4503	12.1283	12.	11.9415	11.8704	11.7617	11.5382
14.4281	14.1195	14.	13.9457	13.8795	13.7777	13.5626
16.409	16.1123	16.	15.9491	15.887	15.7908	15.5834
18.3922	18.1062	18.	17.9519	17.8932	17.8019	17.6014
20.3773	20.101	20.	19.9543	19.8985	19.8113	19.6172

Let Ω_ξ denote the set

$$\Omega_\xi = \{\nu_{\cot \xi}^{(M)}, M = 1, 2, \dots\}, \quad \xi \in \langle 0, \pi \rangle, \quad \nu_{\cot \xi}^{(M)} = \nu_\eta^{(M)},$$

(we understand $\Omega_0 = \{0, 2, 4, \dots\}$),

and let denote further by \mathcal{E}_ξ the set

$$\mathcal{E}_\xi = \{c(\nu)D_\nu | \nu \in \Omega_\xi\}.$$

The set $\mathcal{E}_\xi \subset \mathcal{D}_\xi$ contains all eigenvectors of the selfadjoint operator H_ξ , and the set $\{\nu + \frac{1}{2}, \nu \in \Omega_\xi\}$ contains all eigenvalues of H_ξ .

Note that orthogonality of two different eigenvectors can be seen also from the Eq.(4). For different μ, ν fulfilling the condition

$$\Gamma\left(\frac{1-\mu}{2}\right)/\Gamma\left(-\frac{\mu}{2}\right) = \Gamma\left(\frac{1-\nu}{2}\right)/\Gamma\left(-\frac{\nu}{2}\right) = \eta,$$

which is our case, Eq. (4) is the scalar product in $L^2(\mathbb{R}^+)$ equal to zero. Moreover, the Eq. (3) guarantees that the eigenvectors are normalized.




Figure 1: To illustrate: for a given value of the parameter η , the values of ν_η are determined by the intersections of the line $y(\nu) = \eta$ with the graph of the function $y(\nu) = \frac{\Gamma(\frac{1-\nu}{2})}{\Gamma(-\frac{\nu}{2})}$.

We denote further by \hat{H} the restriction of H_ξ to domain

$$\hat{D} = \{f \in \tilde{D}, f(0) = f'(0) = 0\} \subset D_\xi(H) \subset L^2(\mathbb{R}^+).$$

Operator \hat{H} is closed, symmetric with deficiency indices (1,1) [9], and H_ξ is a selfadjoint extension of \hat{H} for any $\xi \in \langle 0, \pi \rangle$. Selfadjoint extensions $H_{\xi=0}$ and $H_{\xi=\frac{\pi}{2}}$ have pure point spectra, which is equivalent to the existence of orthonormal bases in $L^2(\mathbb{R}^+)$. The basis is in the case $D_{\xi=0}(H) = \{h_{2n+1} | n = 0, 1, 2, \dots\}$, and it is in the case $D_{\xi=\frac{\pi}{2}}(H) = \{h_{2n+1} | n = 0, 1, 2, \dots\}$. As we mentioned in the introduction, the same is true for all operators H_ξ with any parameter ξ .

Consequently, one can write theorem

Theorem 2

The set \mathcal{E}_ξ consisting of eigenvectors of H_ξ is an orthonormal basis in $L^2(\mathbb{R}^+)$ for any $\xi \in \langle 0, \pi \rangle$, and $\sigma(H_\xi) = \{\nu + \frac{1}{2}, \nu \in \Omega_\xi\}$.

4 Concluding remarks

The results we present can be translated to the case $L^2(\mathbb{R}^-)$. In this case an orthonormal basis in $L^2(\mathbb{R}^-)$ is

$$\tilde{\mathcal{E}}_\xi := \{\tilde{D}_\nu | \nu \in \Omega_\xi\}, \tilde{D}_\nu(x) := D_\nu(-x).$$

These two bases can be combined to the base in $L^2(\mathbb{R})$. As $L^2(\mathbb{R}) = L^2(\mathbb{R}^+) \oplus L^2(\mathbb{R}^-)$. Then for any pair $(\xi, \sigma) \in \langle 0, \pi \rangle \times \langle 0, \pi \rangle$ the set $\mathcal{E}_\xi \oplus \tilde{\mathcal{E}}_\sigma$ is a basis in $L^2(\mathbb{R})$. Explicitly

$$\mathcal{E}_\xi \oplus \tilde{\mathcal{E}}_\sigma = \{(D_\nu, 0), \nu \in \Omega_\xi\} \cup \{(0, \tilde{D}_\nu), \nu \in \Omega_\sigma\}.$$

Of note, the known orthonormal basis $\{h_n, n = 0, 1, \dots\}$ of $L^2(\mathbb{R})$ consisting of Hermitian functions is not contained in this set. Functions h_n are eigenvectors of selfadjoint operator H_D with definition domain

$$D = \{f, f' \text{ absolutely continuous, } f, Hf \in L^2(\mathbb{R})\},$$

and operator H_D is physically interpreted as Hamiltonian of quantum linear harmonic oscillator. It would be interesting to use these bases for numerical solutions of time-dependent partial differential equations, as in publications [5]. Gluing the solutions on the positive and negative axes is suitable for solving an asymmetrical harmonic oscillator and gives the possibility to construct a family of continuous orthogonal functions on $L^2(\mathbb{R})$ [14].

Acknowledgement

All authors acknowledge the financial support from RVO14000 and "Centre for Advanced Applied Sciences", Registry No. CZ.02.1.01/0.0/0.0/16_019/0000778, supported by the Operational Program Research, Development and Education, co-financed by the European Structural and Investment Funds and the state budget of the Czech Republic.

The authors (G.C., M.H.) thank P. Exner for helpfull discussion.

References

- [1] R. Basna, H. Nassar and K. Podgórski, J Multivar Anal, Vol. 189, (2022), 104868
- [2] A. Bultheel, M. Van Barel and P. Van gucht, Orthogonal basis functions in discrete least-squares rational approximation, J Comp Appl Math, Vol. 164–165, (2004), 175-194
- [3] K. W. Kwon and D. H. Lee, Orthogonal functions satisfying a second-order differential equation, J Comp and Appl Math, Vol. 153, 1–2, (2003), 283-293
- [4] M. A. Chaudhry and S.M. Zubair, On the decomposition of generalized incomplete gamma functions with applications to Fourier transforms, J Comp Appl Math, Vol. 59, (1995) 253-284

- [5] A. Iserles and M. Webb, A Family of Orthogonal Rational Functions and Other Orthogonal Systems with a skew-Hermitian Differentiation Matrix. *J Fourier Anal Appl* 26, 19 (2020), <https://doi.org/10.1007/s00041-019-09718-5>
- [6] J.B. Seaborn, Orthogonal Functions. In: *Mathematics for the Physical Sciences*. Springer, New York, (2002), 137-177
- [7] D. Chelkak and E. Korotyaev, The Inverse Problem for Perturbed Harmonic Oscillator on the Half-Line with a Dirichlet Boundary Condition. *Ann. Henri Poincaré* 8, 1115–1150 (2007), <https://doi.org/10.1007/s00023-007-0330-z>
- [8] V. Bonnaillie-Noël, Harmonic oscillators with Neumann condition on the half-line. *Communications on Pure and Applied Mathematics*, 11 (6), (2012), 2221-2237
- [9] J. Blank, P. Exner and M. Havlíček, *Hilbert Space Operators in Quantum Physics*, 2nd edition, Springer, (2008), 127-137
- [10] J. Weidmann, *Linear Operators in Hilbert Space*, Springer Verlag, New York (1980), 246
- [11] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, 4th edition, Fizmatfiz, Moskva (1963), 7.711, 8.314-372, 9.210-253
- [12] I. S. Gradshteyn and I. M. Ryzhik, "Parabolic Cylinder Functions" and "Parabolic Cylinder Functions $D_p(z)$ " § 7.7 and 9.24-9.25 in *Tables of Integrals, Series, and Products*, 6th ed. San Diego, CA: Academic Press, (2000), 835-842, 1018-1021
- [13] N. M. Temme, Parabolic cylinder functions, [DLMF] NIST Digital Library of Mathematical Functions. <https://dlmf.nist.gov/12>
- [14] G. Chadzitaskos and J. Patera, Asymmetric harmonic oscillator, arXiv:2204.06415v1 (2022)