

Atherogenic lipid profile of umbilical cord blood of neonates of gestational diabetic mother: a case-control study

Abstract

Background: Gestational diabetes mellitus (GDM) is a metabolic disorder that occurs during pregnancy and affects both the mother and the fetus. GDM can lead to adverse perinatal outcomes such as macrosomia, shoulder dystocia, neonatal hypoglycemia, polycythemia, hyperbilirubinemia, and congenital malformations. Moreover, GDM increases the risk of long-term complications in the offspring, such as obesity, diabetes and cardiovascular diseases. Lipids are essential biomolecules that have various functions in the body, including energy storage, membrane structure and hormone synthesis. The concentration of lipids in the blood can be altered by maternal hyperglycemia and insulin resistance. The use of insulin or oral antidiabetic drugs can improve glycemic control and reduce the incidence of GDM-related complications, but the neonatal mortality rate remains higher than that of normal pregnancies. The aim of this study was to investigate the impact of maternal diabetes on the lipid profile of umbilical cord blood of their neonates.

Methods: We performed a prospective case-control study on 60 neonates delivered in our hospital. The cases were 30 infants born to mothers with GDM, and the controls were 30 infants born to mothers without GDM. We selected the participants randomly from the labor room.

Results: We observed significant differences between the cases and the controls regarding lipid profile and birth weight. The cases had higher levels of total cholesterol, triglycerides, and LDL-cholesterol in their cord blood than the controls.

Conclusion: Our study demonstrates that maternal diabetes influences the lipid profile of cord blood in neonates. We also find a significant correlation between lipid profile and birth

weight in infants of GDM mothers.

Keywords: Gestational Diabetic mothers, neonates, Cord Serum Lipid

Introduction:

“Gestational diabetes mellitus (GDM) is a condition of impaired glucose metabolism that occurs during pregnancy. GDM can affect the health and development of the fetus, as well as the long-term health of the mother. One of the main complications of GDM is fetal macrosomia, which is defined as a birth weight above the 90th percentile for gestational age. Macrosomia can result from increased fetal insulin production in response to maternal hyperglycemia, which stimulates fetal growth and adiposity” (1). “Fetal insulin also influences the growth hormone/insulin-like growth factor-1 axis, which regulates fetal growth and development and has implications for cardiovascular health later in life” (2). “Moreover, GDM can alter the lipid profile of the mother and the fetus, leading to increased levels of lipoproteins that are associated with atherosclerosis and diabetes in adulthood” (3).

Atherosclerosis is a chronic inflammatory disease of the arteries that causes plaque formation and narrowing of the vessels. Atherosclerosis can manifest clinically as coronary heart disease, stroke, peripheral arterial disease, or renal artery stenosis. Atherosclerosis has its origins in early life, as evidenced by the presence of fatty streaks and fibrous plaques in the arteries of children and adolescents (4). The development and progression of atherosclerosis can be influenced by genetic and environmental factors, such as dyslipidemia, hypertension, obesity, smoking, and diabetes.(5) Non-invasive methods, such as ultrasound, can be used to assess vascular changes related to atherosclerosis in peripheral arteries (6).

“GDM is a risk factor for adverse maternal and neonatal outcomes, such as preterm delivery,

cesarean section, congenital anomalies, preeclampsia, and neonatal hypoglycemia” (7-9). “The incidence of congenital anomalies in infants of diabetic mothers (IDM) ranges from 6% to 10%, depending on the type and severity of maternal diabetes” (10). “GDM also increases the risk of future cardiovascular disease and diabetes in both the mother and the offspring” (11-14). Therefore, early diagnosis and management of GDM are essential to prevent or reduce these complications.

Lipid metabolism is a key process that affects both maternal and fetal health during pregnancy.(15)“In normal pregnancies, lipid metabolism undergoes significant changes to meet the increased energy demands of the mother and the fetus, as well as to support the development of the placenta and the fetal tissues. These changes include increased levels of circulating non-esterified fatty acids (NEFAs), triacylglycerols (TAGs), cholesterol, phospholipids and apolipoproteins (apos)” (16).

“However, in gestational diabetes mellitus (GDM), lipid metabolism is further disrupted by impaired glucose tolerance and insulin resistance, which can have adverse effects on both maternal and fetal outcomes. GDM is associated with higher levels of serum TAGs and apolipoprotein B (apoB), which are the main components of very low density lipoprotein (VLDL) particles” (4, 5). “VLDL particles are responsible for transporting lipids from the liver to the peripheral tissues, and their increased production in GDM may lead to excessive lipid accumulation in the fetus, resulting in macrosomia (large birth weight)” (17).

The mechanisms underlying the altered lipid metabolism in GDM and macrosomia are not fully understood, but they may involve increased glucose and NEFA availability in the maternal circulation, which stimulate hepatic VLDL synthesis and secretion (18). Moreover, hyperinsulinaemia in GDM may enhance lipid and protein synthesis in the liver and other tissues, as well as increase placental transfer of NEFA to the fetus (19). “These factors may contribute to the elevated TAG and apoB levels observed in GDM mothers and macrosomic-

neonates, as well as the positive correlation between maternal glycosylated haemoglobin (HbA1c) and TAG levels in late pregnancy” (14).

Macrosomic newborns are babies who are born with a higher than average birth weight. They may have some health problems related to their size, such as high cholesterol levels. Cholesterol is a type of fat that circulates in the blood. It comes in different forms, such as very-low-density lipoprotein (VLDL), low-density lipoprotein (LDL), and high-density lipoprotein (HDL). VLDL and LDL are considered "bad" cholesterol because they can clog the arteries and increase the risk of heart disease and stroke. HDL is considered "good" cholesterol because it helps remove excess cholesterol from the blood and protects the arteries.

Macrosomic newborns tend to have high levels of VLDL and LDL, which are mainly produced from VLDL by an enzyme called lipoprotein lipase (LPL). This enzyme is also involved in breaking down triglycerides, another type of fat that can be harmful in high amounts. Some studies have shown that macrosomic newborns have higher levels of LPL and another enzyme called hepatic TAG lipase (HTAGL), which may contribute to their high cholesterol levels. These babies may have inherited these traits from their mothers, especially if they have gestational diabetes or type I diabetes.

Macrosomic newborns also tend to have high levels of HDL, which are accompanied by high levels of two proteins called apoA-I and apoA-II. These proteins are important for the structure and function of HDL particles, which carry cholesterol and other fats from the tissues to the liver for disposal. High levels of HDL may reflect the increased need for cholesterol and phospholipids in macrosomic newborns, as these substances are essential for the growth and development of cells, hormones, and surfactants. Surfactants are substances that help the lungs expand and contract during breathing. Macrosomic newborns do not have abnormal levels of an enzyme called lecithin cholesterol acyltransferase (LCAT), which is

responsible for converting free cholesterol into a form that can be carried by HDL.

Cholesterol is a vital component of life, but too much of it can be harmful. Macrosomic newborns may face a higher risk of cardiovascular diseases later in life due to their high cholesterol levels. Therefore, it is important to monitor their cholesterol levels and provide them with appropriate dietary and lifestyle interventions to prevent or reduce the complications associated with high cholesterol.

Methods:

We conducted a case-control study in Topiwala National Medical College and B.Y.L.Nair Charitable Hospital Mumbai from Jan 2021 to June 2021 to compare the lipid profile of umbilical cord blood between infants of gestational diabetes mellitus (GDM) mothers and normal mothers. We collected blood samples from 60 infants (30 neonates of GDM mothers and 30 neonates of normal mothers) who were born via normal vaginal delivery (NVD) or cesarean section (CS). We clamped or milked the umbilical cord before taking the samples and put them in plain bottle.

The inclusion criteria of the study subjects are: (1) Establishing a maternity check-up health card before the 13th week of pregnancy; (2) 18–40 years old.

Exclusion criteria: (1) twin or multiple pregnancy; (2) smoking ; (3) history of hypertension and family history of diabetes; (4) abnormal blood glucose in the first trimester; (5) delivery within 37 weeks of pregnancy; (6) received clinical treatment of GDM. We included only full-term infants (38 weeks or more) with birth weight between 2.5 and 3.5 kg, without any congenital anomalies, and whose mothers were aged 25 to 37 years and had no other systemic diseases besides GDM. We measured the serum levels of total cholesterol (TC), triglyceride (TG), high-density lipoprotein-cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), and very low-density lipoprotein cholesterol (VLDL-C).

We obtained 5 ml of umbilical venous blood from the placental end of the umbilical cord

under sterile conditions after the delivery of the placenta and the clamping of the cord. We let the blood stand for a few minutes, then separated the serum from the clot by centrifuging at 3000 rpm for 30 minutes and analyzed it immediately. We used a computerized automated biochemical XL-300 autoanalyzer and an enzymatic method to measure the serum levels of TC, TG, HDL-C, LDL-C, and VLDL-C with high accuracy.

We used mean values and standard error of mean to describe the data in each group. We compared these values between 30 normal term newborns of GDM mothers and 30 newborns of normal healthy mothers.

Results

Table1	<u>Comparison between the infants of Gestational diabetic and normal mothers</u>		
Variable	Mean±SD	Mean±SD	P-value
Mothers' age	26.86±4.03	30.86±4.24	0.088
Gestational age	38.93±0.827	37.96±1.017	0.764
Neonatal weight(g)	2454.0.0±191.63	4012.6±481.28	<0.001
Random blood sugar (RBS) of mother mg/dl	100.66±15.21	188.23±22.31	<0.001
RBS of neonates mg/dl	72.50±14.81	47.23±8.05	<0.001

Table2. Lipid profile in the two groups

	Neonates of normal mothers	Neonates of Gestational diabetic mothers	
Variable	Mean±SD	Mean±SD	P-value
Serum cholesterol(mg/dl)	70.10±22.56	136.23±26.15	<0.001
Serum triglyceride(mg/dl)	35.4±18.92	72.0±45.02	<0.001
High-density lipoprotein(mg/dl)	39.1 ±11.44	40.50±17.19	0.892

Low-density lipoprotein (mg/dl)	24.8±3.921	81.5±36.15	<0.001
---------------------------------	------------	------------	--------




Figure 1. Correlation between serum cholesterol and neonatal weight in both groups (Newborns of Healthy mother and newborns of gestational diabetic mother) ($r=0.8647, P<0.001$)




Figure 2. Correlation between triglyceride level and neonatal weight in both groups (Newborns of Healthy mother and newborns of gestational diabetic mother) ($r=0.8039, P<0.001$)




Figure 3. Correlation between low-density lipoprotein level and neonatal weight in both groups (Newborns of Healthy mother and newborns of gestational diabetic mother) ($r=0.8431$, $P<0.001$)




Figure 4. Correlation between high-density lipoprotein level and neonatal weight in both groups (Newborns of Healthy mother and newborns of gestational diabetic mother) ($r=-0.2085$, $P=0.1099$)




Figure 5. Correlation between gestational age and triglyceride ($r=-0.0432$, $P=0.7597$)




Figure6.Correlationbetweengestationalageandhigh-densitylipoprotein in both groups ($r=0.2476, P=0.357$)




Figure7.Correlation between gestational age and low-density lipoprotein in both groups ($r=-0.619, P=0.6384$)




Figure8.Correlation between cholesterol level newborns and random blood sugar of mothers in both groups ($r=0.722, P<0.001$) Results of the pearson correlation indicated that there is a significant large positive relationship between Cholesterol of newborns and RBG of

Mothers, ($r(58) = .722, p < .001$)




Figure 9. Correlation between triglyceride and random blood sugar of mothers in both groups ($r = -0.201, P = 0.286$). Results of the Pearson correlation indicated that there is a significant large positive relationship between TG of newborns and RBG of Mothers, ($r(58) = .723, p < .001$).




Figure 10. Correlation between high-density lipoprotein and random blood sugar of mothers in both groups ($r = -0.24, P = 0.854$)




Figure 11. Correlation between low-density lipoprotein and random blood sugar of mothers in both groups ($r = 0.709, P < 0.001$)

Results of the Pearson correlation indicated that there is a significant large positive

relationship between LDL in mg/dl of newborns and RBG of Mothers in both groups ($r(58) = .709, p < .001$).




Figure 12. Correlation between cholesterol and random bloodsugar of neonates in both groups ($r = -0.888, P < 0.001$)

Results of the Pearson correlation indicated that there is a significant large negative relationship between Cholesterol in mg/dl of newborns and RBG of newborns, ($r(58) = .888, p < .001$).




Figure 13. Correlation between triglyceride and random bloodsugar of neonates in both groups ($r = -0.132, P = 0.488$) Results of the Pearson correlation indicated that there is a significant large negative relationship between TG in mg/dl of newborns and RBG of newborns, ($r(58) = .896, p < .001$).




Figure 14. Correlation between low-density lipoprotein and random blood sugar of neonates in both groups ($r=-0.881, P< .001$)

Results of the pearson correlation indicated that there is a significant large negative relationship between LDL in mg/dl of newborns and RBG of newborns, ($r(58) = .881, p < .001$).

Discussion

“Serum lipid profile in childhood is considered as a predictive factor for serum lipid level later in life. This means that not only diet but also other risk factors affect serum lipid level from birth” (22).

“This study showed no significant differences between the two groups of infants regarding gestational age ($P=0.764$) (Table 2). However, as reported by other studies, GDM are at a higher of preterm birth (low gestational age)” (3). The limited number of cases in our study may be the underlying cause of this discrepancy. Moreover, Table 2 demonstrated no significant differences between the two groups regarding mothers’ age ($P=0.088$).

“The results of this study reflect a significant difference between the neonates of GDM and of healthy mothers regarding birth weight ($P<0001$) (the mean birth weight of the IDM was 4012.6 g, while the mean birth weight of the infants of healthy mothers was 2454.0g)” (Table 1). [22]

Other studies have reported a decline in the incidence of macrosomia in IDM from 60% to about 20%-35%, which might be secondary to aggressive diagnosis and treatment of GDM (1). In addition, a study carried out in Japan by Cannon JR *et al* and de Deungria M *et al* respectively, showed that “the incidence of preterm labour and low-birth-weight are associated with GDM” (23, 24). “Whereas, several other studies demonstrate that a small (<5%) number of fetuses, usually carried by mothers with advanced diabetic vascular disease, are at risk for fetal growth restriction (birth weight of less than fifth percentile for gestational age), this inconsistency might be explained by different ways of diabetes management in many studies”.(15, 16)

The results of this study showed a significant difference in the mean RBS of diabetic and healthy mothers. The mean RBS of diabetic mothers was 188.2 mg/dl, while the mean RBS of healthy mothers was 100.6 mg/dl ($P<0.001$) (Table 2). Table 2 also showed a significant difference in the RBS of the neonates of diabetic mothers (IDM) and infants of healthy mothers. “This finding highlights the importance of HbA1c test for diabetes diagnosis. The IDM had a mean RBS of 47.2 mg/dl, while the neonates/babies of healthy mothers had a mean RBS of 72.5 mg/dl ($P<0.001$)”. Rollins *et al.* and Rooney *et al* (25, 26).

Table 2 showed a significant difference in the serum cholesterol levels of IDM and babies of healthy mothers. “The IDM had a mean serum cholesterol level of 136.2 mg/dl, while the babies of healthy mothers had a mean serum cholesterol level of 73.7 mg/dl ($P<0.001$). Table 2 also showed that the levels of TG and LDL were significantly higher in the IDM with means of 72.0 mg/dl and 81.5 mg/dl, respectively, while the means in healthy mothers were 31.4 mg/dl and 12.8 mg/dl, respectively ($P<0.001$). These results may be due to decreased LPL activity in adipose and liver tissues” (27).

“There was no significant difference in the HDL levels between the two groups ($P=0.892$). These findings are partly consistent with previous studies that reported lower levels of total

serum cholesterol, TG, LDL and HDL”.(27)

The figures (1 and 2) show the relationship between maternal lipid levels and neonatal weight in the two groups of healthy and diabetic mothers. The results indicate that there is a strong positive correlation between both serum cholesterol and triglyceride levels and neonatal weight in both groups, with a higher correlation coefficient for cholesterol ($r=0.8647$) than for triglyceride ($r=0.8039$). The correlation is statistically significant at $P<0.001$ level, suggesting that maternal lipid levels are an important factor influencing neonatal weight.

The figures show the correlations between different lipid parameters and neonatal outcomes in the two groups of mothers (healthy and gestational diabetic). Figure 3 shows a strong positive correlation between low-density lipoprotein (LDL) level and neonatal weight, indicating that higher LDL levels in mothers are associated with higher birth weight in newborns. This is in line with the studies by Kalra *et al*(28) and Mathuret *et al*(29), who explained these associations by the placental transport of nutrients and stress factors.

On the other hand, the correlation between neonatal weight and HDL level was weak and negative, but not significant (Figure 4). No significant correlations were found between GA and lipid parameters (TG, HGL and LDL) ($P<0.911$, $=0.357$ and $=0.07$, respectively), as shown in Figures 5-7 for both groups.

We examined the associations between maternal and neonatal blood glucose and lipid levels using pearson correlation analysis. Figures 8 and 9 show that there were strong positive correlations between maternal and neonatal cholesterol ($r(58) = .722$, $p < .001$) and triglycerides ($r(58) = .723$, $p < .001$), respectively. This suggests that higher maternal blood glucose levels may lead to higher neonatal cholesterol and triglycerides levels. Figure 11 shows that there was also a strong positive correlation between maternal blood glucose and neonatal low-density lipoprotein (LDL) levels ($r(58) = .709$, $p < .001$), indicating that maternal hyperglycemia may increase the risk of neonatal dyslipidemia. On the other hand,

Figures 12, 13 and 14 show that there were strong negative correlations between neonatal blood glucose and neonatal cholesterol ($r(58) = .888, p < .001$), triglycerides ($r(58) = .896, p < .001$) and LDL ($r(58) = .881, p < .001$), respectively. This implies that lower neonatal blood glucose levels may be associated with lower neonatal lipid levels.

Conclusion

This study demonstrated that IDM have higher levels of serum cholesterol, triglycerides and LDL than infants of healthy mothers, which may increase their risk of developing cardiovascular diseases later in life. These differences may be attributed to the impaired LPL activity and the placental transport of nutrients and stress factors in IDM. Moreover, this study showed that maternal blood glucose levels are positively correlated with neonatal cholesterol, triglycerides and LDL levels, indicating that maternal glycemic control is important for preventing neonatal dyslipidemia. Further studies are needed to explore the long-term effects of neonatal dyslipidemia and the possible interventions to prevent or treat it. Future research should also assess the serum lipid profile of babies born to diabetic mothers to detect any elevation in their lipid levels. Moreover, babies with high lipid level may require dietary interventions and regular monitoring of their lipid status.

Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

Consent

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

References:

1. KicklighterSD, PotterCF, RosenkrantzT. Infant of Diabetic Mother [Internet]. [Updated 2013 May 10]. Available from: www.emedicine.com/ped/topic485.htm.
2. Widness JA. Fetal Risks and Neonatal Complications of Diabetes Mellitus and Metabolic, and Endocrine Disorders. In: Brody SA, Ueland K, editors. *Endocrine Disorders in Pregnancy*. Norwalk, CT: Appleton-Lang; 1989. P. 273–97.
3. Kliegman RM, Stanton BM, St. Geme J, Schor NF, Behrman RE. *Nelson Textbook of Pediatrics*. 19th ed. Philadelphia: Saunders; 2011.
4. Wagner RK, Nielsen PE, Gonik B. Shoulder dystocia. *Obstet Gynecol Clin North Am*. 1999;26(2):371–83.
5. Lucas MJ. Diabetes complicating pregnancy. *Obstet Gynecol Clin North Am*. 2001;28(3):513–36.
6. Milley JR, Papacostas JS, Tabata BK. Effect of insulin uptake of metabolic substrates by the sheep fetus. *Am J Physiol*. 1986;251(3 Pt 1):E349–59.
7. Philipps AF, Porte PJ, Stabinsky S, Rosenkrantz TS, Ray JR. Effects of chronic fetal hyperglycemia upon oxygen consumption in the ovine uterus and conceptus. *J Clin Invest*. 1984;74(1):279–86.
8. Widness JA, Susa JB, Garcia JF, Singer DB, Sehgal P, Oh W, et al. Increased erythropoiesis and elevated erythropoietin in infants born to diabetic mothers and in hyperinsulinemic rhesus fetuses. *J Clin Invest*. 1981;67(3):637–42.
9. Stonestreet BS, Goldstein M, Oh W, Widness JA. Effect of prolonged hyperinsulinemia on erythropoiesis in fetal sheep. *Am J Physiol*. 1989;257(5 Pt 2):R1199–204.
10. Georgieff MK, Widness JA, Mills MM, Stonestreet BS. The effect of prolonged intrauterine hyperinsulinemia on iron utilization in fetal

- sheep. *Pediatr Res.* 1989;26(5):467–9.
11. Petry CD, Wobken JD, McKay H, Eaton MA, Seybold VS, Johnson DE, et al. Placental transferrin receptor in diabetic pregnancies with increased fetal iron demand. *Am J Physiol.* 1994;267(4Pt1):E507–14.
 12. Georgieff MK, Petry CD, Mills MM, McKay H, Wobken JD. Increased N-glycosylation and reduced transferrin-binding capacity of transferrin receptor isolated from placentas of diabetic women. *Placenta.* 1997;18(7):563–8.
 13. Petry CD, Eaton MA, Wobken JD, Mills MM, Johnson DE, Georgieff MK. Liver, heart and brain iron deficiency in newborn infants of diabetic mothers. *J Pediatr.* 1992;121(1):109–14.
 14. Cianflone K, Roncari DK, Maslowska M, Baldo A, Forden J, Sniderman AD. Adipsin/acylation stimulating protein system in human adipocytes: regulation of triacylglycerol synthesis. *Biochemistry.* 1994;33(32):9489–9495.
 15. Yasruel Z, Cianflone K, Sniderman AD, Rosenbloom M, Walsh M, Rodriguez MA. Effect of acylation stimulating protein on the triacylglycerol synthetic pathway of human adipose tissue. *Lipids.* 1991;26(7):495–9.
 16. Haagsman HP, de Haas CG, Geelen MJ, van Golde LM. Regulation of triacylglycerol synthesis in the liver. Modulation of diacylglycerol acyltransferase activity in-vitro. *J Biol Chem.* 1982;257(18):10593–8.
 17. Germinario R, Sniderman AD, Manuel S, Lefebvre SP, Baldo A, Cianflone K. Coordinate regulation of triacylglycerol synthesis and glucose transport by acylation stimulating protein. *Metabolism.* 1993;40(5):574–80.
 18. Cianflone K, Lu H, Smith J, Yu W, Wang H. Adiponectin, acylation stimulating protein and complement C3 are altered in obesity in very young children. *Clin Endocrinol (Oxf).* 2005;62(5):567–72.

19. Avery ME, Oppenheimer EH, Gordon HH. Renal-vein thrombosis in newborn infants of diabetic mothers—report of two cases. *New Engl J Med*. 1957;265:1134–8
20. Sivan E, Mazaki-Tovi S, Pariente C, Efraty Y, Schiff E, Hemi R, et al. Adiponectin in human cord blood: relation of fetal birth weight and gender. *J Clin Endocrinol Metab*. 2003;88(12):5656–60.
21. Kitamura S, Yokota I, Hosoda H, Kotani Y, Matsuda J, Naito E, et al. Ghrelin concentration in cord and neonatal blood: relation of fetal growth and energy balance. *J Clin Endocrinol Metab*. 2003;88(11):5473–7.
22. Ong K, Kratzsch J, Kiess W, Costello M, Scott C, Dunger D. Size at birth and cord blood levels of insulin, insulin-like growth factor-I (IGF-I), IGF-II, IGF-binding protein-1 (IGFBP-1), IGFBP-3, and the soluble IGF-II/mannose-6-phosphate receptor in term human infants. The ALSPAC study team. Avon longitudinal study of pregnancy and childhood. *J Clin Endocrinol Metab*. 2000;85(11):4266–9.
23. Connor JR, Menzies SL. Relationship of iron to oligodendrocytes and myelination. *Glia*. 1996;17(2):89–93.
24. de Deungria M, Rao R, Wobken JD, Luciana M, Nelson CA, Georgieff MK. Perinatal iron deficiency decreases cytochrome c oxidase (CytOx) activity in selected regions of neonatal rat brain. *Pediatr Res*. 2000;48(2):169–76.
25. Rollins MD, Maxwell AP, Afrasiabi M, Halliday HL, Lappin TR. Cord blood erythropoietin, pH, PaO₂ and haematocrit following caesarean section before labour. *Biol Neonate*. 1993;63(3):147–52.
26. Rooney SA. Regulation of Surfactant-Associated Phospholipid Synthesis and Secretion. In: Polin RA, Fox WW, editors. *Fetal and Neonatal Physiology*. 2nd ed. Philadelphia: Saunders; 1998. P. 1283–99.

27. Deregnier RA, Nelson CA, Thomas KM, Wewerka S, Georgieff MK. Neurophysiologic evaluation of auditory recognition memory in healthy newborn infants and infants of diabetic mothers. *JPediatr.* 2000;137(6):777–84.
28. Kalra B, Gupta Y, Singla R, Kalra S. 2015. Use of oral anti-diabetic agents in pregnancy: a pragmatic approach. *N. Am. J. Med. Sci.* 7, 6–12.
29. Mathur, M.B.; Epel, E.; Kind, S.; Desai, M.; Parks, C.G.; Sandler, D.P.; Khazeni, N. Perceived stress and telomere length: A systematic review, meta-analysis, and methodologic considerations for advancing the field. *Brain Behav. Immun.* **2016**, 54, 158–169.

UNDER PEER REVIEW