

Alteration in Flowering and Fruiting of Commercial Vegetable Crops under Protected Condition

Abstract

India ranks as the second-largest global vegetable producer, yet its current productivity falls short of meeting the nutritional requirements of its expanding population. The country boasts an impressive array of vegetable varieties, spanning from tropical to subtropical and temperate regions. The adoption of protected cultivation for high-value crops has emerged as a pivotal technology to enhance productivity, elevate quality, and secure profitable returns. Optimal production quality can be ensured by implementing precise crop management techniques under controlled conditions, enabling the cultivation of two to three high-yield crops annually. Crop regulation planning entails the identification, selection, implementation, and monitoring of strategies to govern the yield and quality of vegetable crops. The timing of flowering can be strategically manipulated, either advanced or delayed, in response to market demand, leveraging a range of tools and techniques for flowering and fruit regulation. In pursuit of sustainability, it becomes imperative to maximize food production within limited areas. Employing flowering and fruiting regulation methods holds the promise of augmenting yields and, consequently, overall productivity, ultimately addressing the nutritional needs of a larger segment of the population.

Key Words: *Crop Regulation, High Value Crops, Protected Cultivation and Sustainability*

INTRODUCTION:

In the context of market globalization, land scarcity, and climate change, protected cultivation of high-value crops has become a pivotal technology for achieving increased productivity, improved quality, and profitability (Kaushal and Singh, 2019). Flowering is a critical stage in the fruit development process since the presence of flowers directly impacts the presence of fruit. Events during flower formation and fruit set have significant implications for fruitlet development, final fruit size, quality, and overall returns (Singh et al., 2019).

Protected cultivation involves manipulating the microclimate surrounding plants to shield them from adverse weather conditions (Ummyiah et al., 2017). This approach optimizes energy use, particularly solar energy (Gielen et al., 2019). Growing vegetables in protected conditions not only enhances water and nutrient efficiency but also boosts productivity by 3-5 times compared to open-field cultivation in various agro-climatic conditions across the country (Heurn and Post, 2004).

Uncontrolled growth tends to favor vegetative growth over reproductive growth, necessitating the regulation of crops to avoid market oversupply and ensure a consistent vegetable supply (Nir et al., 1972). Increased flowering intensity resulting from stress indicates that flower differentiation occurs during moisture stress, and generative buds remain undeveloped until water is supplied. Moisture stress, followed by relief, effectively triggers vegetative growth (Goell et al., 1981). Extended periods of stress, caused by factors like extreme temperature and humidity, can delay flowering (Lal et al., 2017). Singh and Chadha (1988) argued that inducing stress led to uniform flushing, and the intensity of flushing correlated with the level of stress, as indicated by the plant's relative water content before relief.

In the case of commercial vegetable crops like tomatoes, bell peppers, and cucumbers grown under protected conditions, flowering time can be adjusted as needed, either through early intervention or by employing various tools and techniques for flowering and fruit regulation, including mechanical, environmental, and chemical methods (Anonymous, 2013a).

PRINCIPLE AND OBJECTIVES OF CROP REGULATION:

Crop regulation entails the essential concept of altering a plant's natural flowering and fruiting patterns to optimize fruit yield, quality, and profitability (Boora et al., 2016). This principle revolves around the understanding that the majority of crop flowers appear exclusively on young, tender, and rapidly growing vegetative shoots. The primary objective of crop regulation is to induce the plant to enter a rest phase, prompting it to yield an abundance of blossoms and fruits within a specific season (Bhakti, 2016). The key reasons behind crop regulation are to ensure a consistent supply of high-quality crops and maximize both production and profits for the grower. Without this regulation, continuous and uninterrupted flowering would result in light crop yields throughout the year, significantly increasing monitoring and marketing costs (Lal et al., 2017).

IMPORTANCE OF PROTECTED CULTIVATION:

Protected horticulture, being the most effective solution to counteract the challenges posed by climatic variations, holds the potential to cater to the requirements of small-scale growers. It not only multiplies crop yields but also significantly enhances food quality to meet market demands (Kumar et al., 2014).

Protected cultivation represents a key approach in this endeavor. It constitutes an agrotechnology that involves the use of coverings to control both macro and microenvironments, ultimately fostering optimal plant growth, prolonging the growing period, encouraging early maturation, and bolstering both yield and quality (Gruda and Tanny, 2014, 2015). Employing protected cultivation for crops ensures superior produce quality, increased productivity, year-round cultivation possibilities, controlled pollination, improved management of insect pests and diseases, reduced pesticide residues, simplified plant protection, efficient resource utilization, higher profitability, guaranteed production, and increased nutrient extraction. Additionally, it creates opportunities for educated rural youth to engage in self-employment within the agricultural sector (Pattnaik and Mohanty, 2021). It's important to note that the microclimate within greenhouses significantly influences crop growth, development, and overall productivity (Katsoulas and Kittas, 2008).

During the winter season, the primary systems for controlling the greenhouse climate include heating, ventilation, and CO₂ enrichment (Anonymous, 2003a). In contrast, during the summer months, the main systems utilized consist of ventilation (either natural or forced), shading through screens or whitening, CO₂ enrichment, and cooling through fan and pad systems (Anonymous, 2003a).

FACTORS AFFECTING FLOWERING AND FRUITING UNDER PROTECTED CONDITIONS:

1. Temperature

Because the steel structure of the poly home is wrapped with polythene, the interior temperature can reach 40°C (Bhakti 2016). The ventilation system, as well as cooling pads and fans, are utilised to regulate the temperature within the poly home Pickens and Sibley (2019). However, fruit set is determined by the 24-hour mean temperature and the temperature variation between day and night Hatfield and Prueger (2015).

Table 1. Temperature required for flowering and fruiting

For healthy and maximum growth	
Day temperature	26-30°C

Night temperature	15-18°C
Optimum photosynthesis	21-24°C
Optimum vegetative growth	21-29°C
For yield	24°C
Flowering and fruit setting	16-18°C

Chandani and Dubey (2020)

2. Relative Humidity

Relative humidity (RH) is a measure of the water vapor content in the atmosphere, and it's important to note that the air's capacity to hold water vapor is influenced by temperature (Kong and Singh, 2011). To ensure the health and optimal growth of vegetables, maintaining an appropriate humidity level is crucial. Vegetables typically thrive in an environment with a humidity level ranging from 60 to 65 percent (Anonymous, 2015).

Transpiration, a vital plant process, serves multiple functions, including cooling the plant, facilitating nutrient uptake from the roots, and distributing resources throughout the plant. It plays a pivotal role in optimizing photosynthesis efficiency, the effective transport of nutrients within the plant, and the distribution of these nutrients for overall plant growth (Perez, 2019). Relative humidity plays a significant role in minimizing water loss through evaporation from plants, leading to efficient nutrient utilization. Moreover, it helps maintain cell turgidity, which is advantageous for enzymatic activity, ultimately resulting in improved crop yields (Rajasekar et al., 2013).

Light

Photosynthesis exclusively takes place within the spectrum of visible light. In poly-house technology, the management of light is optimized to ensure that plants receive the maximum amount of visible light while any surplus light is reflected back (Anonymous, 2013b). In general, higher light intensity within the greenhouse leads to increased rates of photosynthesis and transpiration, which in turn elevates humidity levels and solar heat gain (Anonymous, 2008).

Studies have revealed that reducing light levels from 10,000 to 2,500 lux can delay the onset of flowering. Greenhouses are designed to capture a portion of the incident light, allowing up

to 80 percent of the light to reach the crop around midday and an average of 68 percent throughout the day (Brown, 2006).

Carbon dioxide

The ambient environment typically contains a CO₂ concentration of 0.03 percent, which is equivalent to 300 parts per million (Lindsey, 2020). Plants use this CO₂ for photosynthesis. During the night within a poly-house, photosynthesis comes to a halt, but CO₂ is released as a result of respiration (Blom et al., 2002). As a consequence, poly-houses consistently maintain a higher CO₂ concentration compared to the outdoor environment, particularly during the nighttime (Anonymous, 2021a).

This surplus CO₂ becomes a valuable resource for plants in poly-houses, enabling them to undergo more rapid photosynthesis. Research has shown that when the CO₂ concentration in a poly-house reaches 1000 parts per million (ppm), vegetable yields can increase four to fivefold compared to regular conditions. The ideal CO₂ concentration typically falls within the range of 350 to 1000 ppm (Bao et al., 2018).

Wind movement

If a poly-house has a higher humidity level, the risk of illness and pests increases. Under these conditions, the poly-side house's vents are opened to improve air flow within the structure. As a result of the increased air movement, the humidity level lowers and the risk of disease also decreases (Anonymous 2013b).

3. Nutrients required by plants

When conducting a chemical analysis of plants, it has been observed that they contain over 90 different components. However, only 17 of these components have been identified as essential for the successful growth and development of plants, based on the essentiality criterion (Blake et al., 2016). Recent research has revealed the necessity of additional elements like vanadium, silicon, cobalt, sodium, and nickel for the growth of specific plants.

The carbon-to-nitrogen ratio, often denoted as the C/N ratio, in soil refers to the ratio of organic carbon to nitrogen content (Miller, 2000). This ratio plays a significant role in the rate of decomposition of organic matter in residues. Microorganisms responsible for breaking down

residues require nitrogen, along with other crucial components, in addition to carbon. When the residue lacks sufficient nitrogen, the decomposition process is slowed down (Morris, 2004). Furthermore, in cases where the residue has a low nitrogen content, microbes may extract inorganic nitrogen from the soil to meet their nitrogen requirements. This can lead to competition with plants for nitrogen, ultimately reducing the available soil nitrogen for plant growth (Aczel, 2019).

The optimal supply of nutrients has been found to promote early flowering in cucumber due to vigorous vine growth (Bishop et al., 1969). Additionally, the maturity of fruit is significantly influenced by fertilizer application, as demonstrated by prior research (Janapriya et al., 2010), which showed that 100% fertigation can induce early harvest in polyhouse conditions.

Table 2. Environmental control for crop productivity

Crop	Temperature	Relative Humidity	Carbon dioxide concentration
Tomato	22-29°C	55-75%	900-1200ppm
Capsicum	18-27°C	60-80%	900-1200ppm
Cucumber	19-31°C	60-75%	900-1200ppm

REGULATION OF FLOWERING AND FRUITING IN COMMERCIAL CROPS:

1. Flowers

In the typical growth of tomato plants, clusters of flowers are usually left unclipped until three to four well-formed fruits have developed on the cluster. However, in the case of anomalous flowers, like the oversized captivated flower, immediate removal is necessary. These anomalous flowers can give rise to fruits with peculiar characteristics, such as a cat's face (Bhakti, 2016).

Young capsicum plants, when they have 7 to 13 leaves, typically commence flowering within a range of 2 to 6 weeks after being planted (Anonymous, 2021c). It's essential for the temperature conditions during flowering to fall within the ideal range, averaging between 20 and 21 degrees Celsius, both during the day and at night (Paradiso and Pascale, 2014). Low night temperatures can negatively impact pollen viability in capsicum flowers, alter their floral structure, and diminish the effectiveness of self-pollination. Flowers that develop at night temperatures below 18 degrees Celsius tend to produce fruits with a 'tail,' characterized by elongation and a

pointed blossom end. Insufficiently warm temperatures during flowering may lead to a reduction in the number of locules in the fruit, potentially resulting in fruit with only two locules, which is undesirable. Conversely, excessively high temperatures, surpassing 28 degrees Celsius, can contribute to the development of blossom end rot (Elitzur et al., 2009).

Tomato Pollination: Tomato flowers have male and female organs (Lambers 2020). To facilitate pollination in greenhouses, growers use methods like vibrators, air blowers, and bumble bees since natural wind is absent (Picken 1914). Effective pollination is essential to maximize tomato fruit size and weight (Kim et al. 2019). Bees, such as bumble bees or honey bees, are commonly employed as pollinators in greenhouses (Ahmad 2009). Using pollinators ensures better fruit formation, improved harvest, and earlier initial harvest (Bhakti 2016). Stingless bees have also proven to be efficient pollinators for greenhouse crops like tomatoes (Macias et al. 2001) and sweet peppers (Cruz et al. 2003).

Cucumber Pollination: Cucumber pollen is large and sticky, requiring external assistance for transfer between flowers (Sedgley and Scholefield 1980). Adequate pollination leads to well-formed, uniformly ripening fruits, while inadequate pollination results in undersized, malformed fruits and reduced yields (Hodges and Baxendale 1991). In greenhouse settings, gynoecious slicing cucumber varieties do not require pollination. However, monoecious cucumber varieties need pollination, primarily performed by honeybees (Thakur and Rana 2008). Bee pollination in greenhouses can be challenging, as bees need to adapt to the indoor environment. Nonetheless, native and Africanized bees also contribute to greenhouse crop pollination (Velthuis 2002; Malagoli-Braga and Kleinert 2007; Santos, Roselino, and Bego 2008; Roselino, Santos, and Bego 2010). An insect's foraging behavior depends on several factors, including innate behavior and floral characteristics (Free 1970). Successful greenhouse cucumber production relies on the use of pollinators (Godoy and Cardoso 2004).

2. Use of growth regulators

The tomato flower's female organs are encased within the male organs (five anthers form a cone around the female organ) (Lambers, 2020). When anthers mature, they open to the interior, releasing pollen (Melissa, 2021). When mature, the anthers will be a bright yellow colour and the bloom will remain pollination-receptive for around 48 hours. Because natural wind is absent in the greenhouse, tomato growers must pollinate their crop using a variety of methods, including battery-operated vibrators, air blowers, and bumble bees (Picken 1914). Additionally, growers

should make every effort to transport the maximum amount of pollen to the flower's stigma (Bhakti 2016). The tomato fruit's size and weight are positively connected with the amount of pollen transported to the female portion of the flower (Kim et al., 2019). Within the tomato, utilize pollinators such as bumble bees or honey bees inside the greenhouse to assure a set of high-quality crops (Ahmad 2009). Bees ensure that fewer fruits are malformed, that fruit set and first harvest are improved, and that the first harvest occurs slightly earlier (Bhakti, 2016). Recent research demonstrates that stingless bees are also an effective alternative to honeybees for pollinating a variety of greenhouse crops of considerable economic and social importance, including strawberries (Maeta et al. 1992; Kukutani et al. 1993; Malagodi-Braga & Kleinert 2004), tomatoes (Macias et al. 2001), and sweet peppers (Macias et al. 2001; Cruz et al. 2003).

Parthenocarpy:

Auxin induced seedless fruits in cucumbers and watermelon, PCPA 50-100 ppm induced parthenocarpy in tomato and brinjal, and 2,4-D 0.25 percent in lanolin paste applied to the cut end of styles or foliar sprays applied to newly opened flower clusters has been reported to promote parthenocarpy (Prajapati et al. 2015). Plant growth regulators aid in the production of fruit in the absence of fertilisation (Parthenocarpy). Staminate flowers were generated in a parthenocarpic cucumber line using the plant growth regulator GA3 at a concentration of 1500 ppm and silver nitrate at a concentration of 200-300 ppm via four sprays at a 4-day interval Singh and Ram (2004).

3. Training and Pruning

The training approach places a premium on the plant's capacity to obtain sufficient sunlight for growth. Additionally, it is critical to maintain adequate air circulation around the plant to minimise the danger of pest infestation (Kumar et al. 2014).

The interaction between the source and sink influences the growth habit, fruit bearing pattern, and seed yield of tomato (Kang et al. 2010). Tomatoes have a determinate, semi-determinate, or determinate growth behaviour (Grant 2021). Indeterminate varieties/hybrids are preferred for greenhouse hybrid seed production (Pavani et al. 2020). These plants can be grown for an extended period of time and produce numerous fruit trusses (Kugblenu et al. 2013). Seed production of determinate or semi-determinate cultivars is less prevalent and is not favoured in greenhouse settings (Cormank 2004). Training and pruning are an integral part of the greenhouse tomato production (Currey 2017).

In tomato, manipulation of plant architecture by training and optimal spatial arrangement has been identified as a critical management strategy for maximising yield from greenhouse crops

(Cebula 1995; Guo et al. 1991; Lorenzo and Castilla 1995). Tomatoes grown in greenhouses are clipped to a single stem. Plastic string is used to support the plants. With a short, non-slip loop, secure one end of twine loosely to the bottom of plants. The other end is connected to an overhead supporting wire located between 1.8 and 2.5 metres above the plant row. Twist the string around the plant as it grows in one or two gentle rotations for each fruit cluster. When plants develop larger and heavier, “twister” or plastic snap-on clips may be used to secure the plant to the string. The string is untied and lowered frequently when the plant reaches the wire, enabling the lower section of the plant to lie on the ground. Approximately 6-7 weeks before harvesting is to cease, the plant's growth tip is typically clipped off or topped (Ozer 2017). Numerous training methods are used like vertical, arch, V-shape, S-hook, lateral (Anonymous 2003b).

De-shooting: By pinching off any side shoots, prune tomato plants to a single stem. Avoid using a knife. This should be done at least once a week; side shoots should be removed while they are very little. Remove no side shoots above the most recent blossom cluster Novak and Maskova (1979).

De-leafing: Remove the two to three leaves beneath ripe fruit clusters that begin to yellow and wilt. Before removing leaves, ensure that the fruit has reached the mature green stage (Gardner 2015).

Thinning of tomato flowers and fruits is discretionary (Mbonihankuye, 2013). Prune flower clusters and fruits to stimulate tomato growth. Trim excess blooms within clusters once three to four well-formed fruits have developed on each cluster. Remove abnormal flowers and misshapen fruits to reduce competition for fruit production (Link, 2000).

Tipping, which involves the removal of the apical bud in tomato plants, leads to a six-day delay in maturity, similar to the delay resulting from leaf pruning (Knott, 1928). Studies by Tanaka and Fujita (1974) show that photosynthates flow from sources to meet sink demands, and partial leaf removal compensates for the decreased net assimilation rate of remaining leaves, leaving fruit growth unaffected. These findings are in line with similar observations made by Starck (1983). Pruning plays a significant role in carbon partitioning, affecting the ratio and, consequently, the fruitfulness of tomato plants (Resh, 1997). According to Hernandez and Sanches (1992), pruning tomato plants to a single stem result in the longest fruit length and the greatest number of fruits.

Pepper plants usually initiate with a single stem and produce a terminal blossom after 9-13 leaves, at which point the main stem divides into two (Manisha P.). Occasionally, three to four shoots may spontaneously emerge instead of two, and two flowers may develop instead of one.

The blossom at the base of the first branch is known as a crown bud (Peter, 2008). The terminal blossom is promptly removed before maturation. After cutting or pinching the remaining branches, each plant should have two main stems, leaving two leaves and one blossom at each internode. These two stems are trained along strings to the main wire that runs the length of the row at a height of 8–9 feet (Manisha P.). The stems can be trellised either loosely or tightly around the strings, secured with rings or plastic clips. During the period of rapid growth, training and pruning should be performed every two to three weeks (Pramnik et al., 2020). Given the poor photosynthetic efficiency of pepper leaves, a substantial active leaf area is required to produce an adequate amount of dry matter (Anonymous, 1990). Pruning is used selectively when there is profuse growth. In sheltered cultivation, the stem structure of pepper plants is often inadequate to support the plant, necessitating training (Manisha P.b). According to Resh (1996), pruning peppers grown in greenhouses enhances light interception, fruit set, and fruit quality by reducing the number of branches. Jovicich et al. (1999) observed that sweet pepper plants pruned to four stems produced higher marketable yields compared to those pruned to two or one stem. Generally, minimal or no pruning results in excessive vegetative growth on smaller fruiting plants. Bhatt and Rao (1997) demonstrated that removing fruit from bell pepper plants' initial flowering node ten days after fruit set had no effect on the partitioning of dry mass to fruit on the upper nodes of the plant. The first flowering node's fruit serves as a significant sink for photosynthates, accounting for 10.2 percent during the initial 20 days after flowering, before becoming a weaker sink.

Various training methods are employed for greenhouse cucumber plants, primarily aimed at optimizing uniform sunlight interception. Pruning, canopy training, and proper spatial arrangement are essential practices for maximizing yields in greenhouse crop production (Rajalingam et al., 2017).

The choice of system depends on the greenhouse facility, production method, and grower preference. The umbrella system is commonly used for vertical cordon or V-cordon trained plants. Gobeil and Gosselin (1989) found that pruning during the summer resulted in higher fruit production. Duong (1999) observed that pruning had no impact on fruit length or mean fruit weight, but pruned cucumbers yielded more fruits per pound compared to unpruned cucumbers. Hong (2000) noted that timing depends on the training system and spacing.

In the Single Head Training System, vines are trained using a single stem on the overhead wire. All flower buds and lateral branches are removed up to a height of 60cm, and fruits are

allowed to grow, one per axil, on the main stem. When the main vine reaches the overhead wire, it is wound around the wire and left to grow downward (Robert et al., 1996).

In the Low Middle Training System, all blooms and lateral branches are removed up to a height of 70cm from the ground level, and 6–8 fruits are allowed. Subsequently, the vines remain fruitless until they reach the overhead wire. When the main vine reaches the overhead wire, it is wound up to 30cm on the cable and then clipped at the growing point. Three healthy laterals are selected: one grows in the direction of the main stem for 20cm along the wire, while the other two grow in the opposite direction of the main stem for 20cm and 30cm along the wire. These three branches develop downward, one per axil (Shivaraj et al., 2020).

The Umbrella System is straightforward and labor-intensive. Cucumber plants are secured to a 7-foot-tall vertical wire, with the top growth point removed. Support is provided for each fruit on the lower part of the main stem, while all laterals are removed from the main stem's leaf axis. The top two laterals are trained to hang down on either side of the main stem over the wire. These laterals grow to two-thirds of the main stem's length. After harvesting the first laterals, they are pruned back, allowing the second laterals to take over, ensuring continued productivity. Fruit size is typically commercially acceptable by the 11th day after the bloom opens during the spring season (Shivaraj et al., 2020).

All lateral buds up to the sixth node should be removed. Pruning involves the removal of branches or twigs to promote fruiting or flowering. Palada and Chang (2003) reported that removing lateral sprouts increased cucumber yield. After 45 days of emergence, basal leaves should be removed to prevent them from becoming a harbor for pests and diseases. Along with lateral buds, all fruits up to this time should be eliminated. Large-fruited varieties should have only one fruit per node, while small-fruited varieties may leave all fruits intact.

To harvest a large number of cucumbers quickly, cultivars with numerous female flowers, such as multi-pistillate cucumbers or gynoecious parthenocarpic cucumbers, should be grown. However, it's essential to manage the number of fruits per plant and their size to ensure optimal quality and quantity of fruit production. Factors such as differences in plant hormones, sink size, and assimilate distribution influence fruit abortion in cucumbers (Ganeshaiyah and Shaanker, 1994).

SPECIAL PRACTICES:

1. Tomato

Tomato Plant Growth Control: To limit plant growth, remove the terminal growth point above the top flower cluster approximately 45 days before the desired crop termination date, as suggested by Guan (2018). It's essential to maintain 2-3 leaves above the top cluster to offer shade and nourishment to the upper fruits, as recommended by Bhakti (2016).

Shading and Cooling: During hot summer months, retaining as many leaves as possible provides effective shade for growing fruit and aids in cooling the greenhouse. Tomato plants transpire around 1-1.5 liters of water daily per plant, which collectively can lower the greenhouse temperature by at least 10°F (Anonymous 2013b).

Battery-Powered Vibrators for Pollination: Utilize battery-powered vibrators to pollinate tomato flowers by gently vibrating the flower clusters' stems for a few seconds. This vibration effectively disperses pollen and fertilizes most ovules in the ovary (Marin 2019). Pollinate the flowers every other day in well-lit conditions with humidity levels between 60% and 80% (Anonymous 2019b). Be cautious not to touch the flower directly to prevent damage to developing fruit. This method is efficient and suitable for small-scale operations (Rutledge 2015) and ensures the production of larger fruits (Bhakti 2016).

Leaf Blower Pollination: Greenhouse tomato pollination can be achieved using a standard household leaf blower directed at the flower clusters. This method is quicker than using an electric vibrator but may lead to slightly reduced fruit size and weight (Delaplane 2013).

Bumblebee Pollination: Bumblebees are efficient pollinators for greenhouse tomatoes and can be used in one or two greenhouses to save time (Anonymous 2021b). Bumblebees visit blossoms and efficiently collect pollen through brief vibrations, leading to effective pollination and fertilization (Potts 2014). This method is known to result in larger and heavier fruits (Junqueira and Augusto 2016). Bumblebees work diligently throughout the day, and each bee can potentially pollinate up to 350 flowers (Stein 2017). Bumblebee pollination has been observed to produce the highest fruit yield among various pollination methods (Closas et al. 2009).

Capsicum

Fruit Set: Optimal conditions for fruit set and initial growth include increased light and cooler day temperatures (Adams et al. 2001). The entire duration from flowering to fully mature, colored fruit typically ranges from 7 to 12 weeks. For successful fruit development, it's recommended to maintain night temperatures around 18°C, day temperatures between 20-22°C, and a CO₂ concentration of 700-1000 ppm (Rangaswamy et al. 2021). However, for the second flush of capsicum fruit, lowering the temperature is not advisable as it may slow down the ripening of the first flush (Heuvelink et al. 2004). Generally, the subsequent fruits naturally set after harvesting the earlier ones (Kitinoja and Kader 1995).

Fruit Load Management: A young plant can typically support around 4-8 fruits at a time, while a mature crop can handle up to ten fruits per plant. To ensure desirable fruit weight (minimum 150 grams per fruit) and quality, farmers should carefully manage the fruit load. The most effective method to reduce the number of fruits is to remove any excess fruits (Anonymous 2021c).

Cucumber

Fruit Formation and Cucumber Production: In cucumber plants, each node has the potential to yield one or more fruits (Bhakti 2016). Typically, it's advisable to thin multiple-fruit clusters to a single fruit, but in robust plants, multiple cucumbers per node may occasionally occur, while any misshapen fruit should be promptly removed (Hochmuth 2012). Cucumber growth is most rapid during the period between 6 to 14 days after the flower opens (anthesis). By day 14, the fruit reaches its maximum length, followed by an increase in diameter (Hochmuth 2012). Greenhouses play a crucial role in cucumber production, making it one of the most widely cultivated crops in greenhouse settings globally (Robinson; Decker-Walters 1999). Protected environments offer ideal conditions for cucumber growth, surpassing open field circumstances (Smitha and Sunil 2016).

Cultivar Selection in Greenhouses: Special cucumber cultivars in the greenhouse are gynoecious and undergo parthenocarpic fruit development, meaning fruits form without the need for pollination (Boonkorkaew 2008). These cultivars are unsuitable for open field cultivation due to the potential for bitter fruits taking on a club shape at the flower end through natural pollination. Consequently, it is crucial to prevent bees and other pollinators from entering the greenhouse since they may introduce pollen from outdoor gardens or field cucumber plantings (Hochmuth 2012). Cucumber varieties are categorized into three groups based on their flowering patterns:

gynoecious, which exclusively produces female flowers, predominantly gynoecious, which also bears some male flowers, and monoecious, which produces both male and female flowers. The first two types generate fruits parthenocarpically, while monoecious varieties necessitate pollination and can produce seeds.

CONCLUSION

India is the second largest producer of vegetables, which occupy 10.2 million ha producing about 185.88 million tonnes with an average productivity of 18.40t/ha (Anonymous 2018-19) of fresh vegetables, yet the productivity is not sufficient to provide diet to our growing population. Since, the population growth rate continued to be more than 2% per annum but rate of food grain production is less than population growth rate. It was noted food grain production was more than population growth rate till 1991. But, declined to 1.16% per annum in 1991-2001, these factors enhance the need of protected cultivation where more food can be produced in limited area and with the regulation of flowering and fruiting techniques there are chances of increasing the yield and thereby productivity which in turn can fulfil the needs of many people and to an extent resolve the problem of limited vegetable production and eventually increasing the GDP (Agriculture Statistics at glance 2015).

REFERENCES:

1. Aczel MR (2019) What Is the Nitrogen Cycle and Why Is It Key to Life? *Frontiers for Young Minds*.7: 41
2. Adams S, Cockshull AE and Cave CRJ (2001) Effect of Temperature on the Growth and Development of Tomato Fruits. *Annals of Botany* 88(5): 869–877
3. Ahmad Al-abbadi (2009) Efficiency of Different Pollination Treatments on Solanaceae Yields Grown in Plastic House. *Journal of Biological Sciences* 9: 464-469.
4. Al-Harbi AR, Alsadon AA, Khalil SO (1996) Influence of training system and growing media on growth and yield of cucumber cultivars. *Alexandra Journal of Agricultural Research*. 41:355-365.
5. Anonymous (1990) Protected Cultivation in the Mediterranean Region. Manual by: Horticultural Crops Group of the FAO Plant Production and Protection Division, United Nations.
6. Anonymous (2003a) Heating, Ventilating and Cooling Greenhouses, *American Society of Agricultural Engineers* 699-707.

7. Anonymous (2003b) Mississippi Cooperative Extension Service and Experiment Station (MAFES), Vegetable gardening: Staking and training tomatoes.
8. Anonymous (2008) Environmental Factors Affecting Plant Growth. Oregon State University.
9. Anonymous (2011) Indole-3-Butyric Acid (IBA) Crop Production. Technical Evaluation Report Compiled by the Technical Services Branch for the USDA National Organic Program. 1-17pp
10. Anonymous (2013a) Good Agricultural Practices for greenhouse vegetable crops. Food and Agriculture Organization of the United Nations Plant Production and Protection Division, Rome
11. Anonymous (2013b). Horticulture: Greenhouse Cultivation. TNAU Agritech Portal Horticulture.
12. Anonymous (2015) Storage of Fresh Produce. The National Institute of Food and Agriculture, United States Department of Agriculture. Publication #FDNS-E-168-27.
13. Anonymous (2018) Hanna Tomato Book. 29p. Accessed on [September 2, 2021]
14. Anonymous (2019a) National Horticulture Board. [Http://nhb.gov.in](http://nhb.gov.in).
15. Anonymous (2019b) Pollinating Tomatoes in High Tunnels. South Dakota State University, South Dakota Counties, and USDA cooperating.
16. Anonymous (2021A) Hi Tech Polyhouse. Sybex-infotech. Accessed on [September 2, 2021]
17. Anonymous. 2021b. Powerhouse Pollinators: Bumble Bees and Greenhouse Pollination. Nature Fresh Farms.
18. Anonymous. 2021c. Protected Cultivation of Capsicum. Vikaspedia, Ministry of Electronics and Information Technology. Accessed on [September 4, 2021]
19. Anonymous. Agricultural Statistics at Glance. [Http://Agricultural_Statistics_At_Glance-2015.pdf](http://Agricultural_Statistics_At_Glance-2015.pdf). Accessed on [September 2, 2021]
20. Bangerth F (1989) Dominance among fruits/sinks and the search for a correlative signal. *Physiologia Plantarum* 76: 608–14.
21. Bangerth F (2000) Abscission and thinning of young fruit and their regulation by plant hormones and bioregulators. *Plant Growth Regulation* 31: 43–59.
22. Bangerth F, LI CJ and Gruber J (2000) Mutual interaction of auxin and cytokinins in regulating correlative dominance. *Plant Growth Regulation* 32: 205–17.
23. Bao J, Lu WH, Zhao J, Bi XT (2018) Carbon Resource Conservation. *Chinese Roots Global Impact: Science Direct* 183-190pp

24. Bardan K (2016) Crop Regulation: Concepts and its Commercial Uses. Commercial Horticulture, pp 257-275 © 2016, Editors, N.L. Patel, S.L. Chawla and T.R. Ahlawat New India Publishing Agency, New Delhi, India.
25. Bertin N (1995) Competition for assimilates and fruit position affect fruit set in indeterminate greenhouse tomato. *Annals of Botany* 75: 55–65.
26. Bhakti P (2016) Growth Regulation Practices in Important fruits crops, Navsari Agriculture University, Navasri, Gujarat. Accessed on [September 8, 2021]
27. Bhatt R and Rao N (1997) Growth and photosynthesis in bell pepper as affected by sink manipulation. *Biologia Plantarum* 39: 437-439.
28. Bishop RF, Chipmon EW, Mae eachern CR (1969) Effect of nitrogen, phosphorous and potassium on yield and nutrient levels in laminate and petioles in pickling cucumber. *Canadian J. Soil Sci.* 49: 297-404.
29. Blake GR, Steinhardt GC, Pombal XP, Munoz JCN, Cortizas AM, Arnold RW, Schaetzl RJ, Stagnitti F, Parlange JY, Steenhuis TS, Chesworth W, Mualem Y, Seytoux HJM, Spaargaren O, Chesworth W, Soon YK, Orlov DS, Oertli JJ (2016) Plant Nutrients. *Encyclopedia of Soil Science*: 23-33
30. Blom TJ, Straver WA, Ingratta FJ, Khosla S, Omafra WB (2002) Ministry of agriculture, food and rural affairs, Ontario. Accessed on [September 4, 2021]
31. Boora RS, Dhaliwal HS and Arora NK (2016) Crop regulation in guava-A review. *Agricultural Reviews* 37:1-9
32. Brown JW (2006) Light in the Greenhouse: How Much is Enough? Cropping Incorporated. Accessed on [September 9, 2021]
33. Cassin J, Bourdeaut J, Fourgerl A, Furon V, Gillard JP, Montagut G and Morevilli C (1969) The influence of climate upon the blooming of citrus in tropical area. *Proc. Int. Citrus Symp* 1: 315-23.
34. Cebula S (1995) Optimization of plant and shoot spacing in greenhouse production of sweet pepper. *Acta Hort.* 412: 321-328.
35. Chandani and Dubey Abhinav (2020) Management of Environmental Factors in Protected Cultivation. Chapter in: *Research Trends in Agriculture Sciences* 17: 136p
36. Closas ML, Puigdomènech P, sanfeliujll and Pelacho AM (2009) Crop Cycle Influences the Effectiveness of Pollination Techniques in Greenhouse Tomato. *European Journal of Horticultural Science.* 74(6): 241–246.

37. Connor LJ, Martin EC (1969) Honey bee pollination of cucumbers. *American Bee Journal* 109: 389
38. Cormank JHM (2004) Tomato seed production an organic seed production manual for seed growers in the Mid-Atlantic and Southern U.S 15p
39. Couto RHN, Couto LA (2006) *Apicultura: manejo e produtos*. Jaboticabal: FUNEP, 193 p.
40. Cruz DO, Campos LAO (2009) Bee pollination in protected crops. *Brazilian Journal of Agrosience*, Pelotas 15: 1-4 p.
41. Cruz DO, Freitas BM, Silva LA, Silva EMS & Bonfi IGA (2003) Adaptation and grazing behavior of the Jandaíra bee (*Melipona subnitida* Ducke) in a protected environment. *Acta Scientiarum. Animal Sciences* 26: 293-298.
42. Currey CJ (2017) The basics of training and pruning plants. *Produce Growers*.
43. Shivaraj D, Prasanth P, Lakshminarayana D and Ramesh T (2020) Studies on the Effect of Training Systems on Cucumber (*Cucumis sativus* L.) Cv. Malini Grown under Protected Conditions. *Current Journal of Applied Science and Technology* 39(48): 539-544
44. Delaplane KS, Arnon Dag, Robert G Danka, Breno M Freitas, Lucas A Garibaldi, R Mark Goodwin & Jose I Hormaza (2013) Standard methods for pollination research with *Apis mellifera*, *Journal of Apicultural Research* 52(4): 1-28
45. Denna DW (1973) Effects of genetic parthenocarpy and gynoeious flowering habit on fruit production and growth of cucumber *Cucumis sativus* L. *Journal of the American Society for Horticultural Science*, 98: 602–604.
46. Dhillon NS, Sharma P, Kumar P and Singh H (2017) Influence of Training on Vegetative Growth Characteristics and Yield of Polyhouse Grown Cucumber (*Cucumis sativus* L.). 18(1): 1-5.
47. Duong HX (1999) Effect of pruning on yield and quality of cucumber. AVRDC Training Report, Kasetsart University, Bangkok, Thailand. 51p.
48. Singh I and Kaur A (2018) Effect of pruning systems on growth and yield traits of greenhouse grown bell pepper (*Capsicum annuum* L. Var. Grossum). *Indian Journal of Agricultural Research* (52):414-418
49. Egli DB and Bruening WP (2002) Flowering and fruit set dynamics at phloem-isolated nodes in soybean. *Field Crop Research* 79: 9–19.
50. Elitzur T, Hadas Nahum, Yelena Borovsky, Irena Pekker, Yuval Eshed, Ilan Paran (2009) Co-ordinated regulation of flowering time, plant architecture and growth by *Fasciculate*: the pepper orthologue of self pruning, *Journal of Experimental Botany* 60(3): 869–880,

51. Free JB (1970) *Insect Pollination of Crop Plants*, Academic Press, London, 544 pp
52. Fujieda K, Fujita Y, Gunji Y and Takahasi K (1982) The inheritance of plural-pistillate flowering in cucumber. *Journal of the Japanese Society for Horticultural Science* 51: 172–6.
53. Ganeshaiyah KN and Shaanker RU (1994) Seed and fruit abortion as a process of self-organization among developing sinks. *Physiologia Plantarum* 91: 81–9.
54. Garderner L (2015) *Gardening Myth: De-leafing Tomato Plants*. Laidback gardener. Accessed on [September 5, 2021]
55. Gielen D, Boshell F, Saygin D, Brazilian MD, Wagner N, Gorini R (2019) The role of renewable energy in the global energy transformation. *Energy Strategy Reviews* 24: 38- 50
56. Gobeil G and Goselin A (1989) The influence of pruning and season on fruit development in a crop of European cucumber. *Can J. Plant Sci.* 69(3): 1037-1048.
57. Godoy AR & Cardoso AII (2004) Pegamento de frutos em pepino caipira não partenocarpico sob cultivo protegido com aplicação de ácido naftaleno acético. *Bragantia* 63: 25-29.
58. Goell A, Golgmb, Kalmar D, Montell A. and Sharon S (1981) Moisture stresses a potent factor for affecting vegetative growth and tree size on citrus. *Proc. Int. Soc. Citri Cult.*, 2: 503-06.
59. Grant A (2021) Should I Prune My Cucumber Vine – Tips on Pruning Cucumbers in The Garden. *Gardening Know How*.
60. Grant BL (2021) Determinate vs Indeterminate Tomatoes: How to Distinguish a Determinate from An Indeterminate Tomato. *Gardening Know How*. [September 5, 2021]
61. Gruda N and Tanny J (2015). Protected crops e recent advances, innovative technologies, and future challenges. *Acta Horticulturae*. (ISHS.) 1107:271-278. <https://doi.org/10.17660/actahortic.2015.1107.37>
62. Gruda N and Tanny J (2014). Protected crops. In: G.R, Aldous, D.E. (Eds.), *Horticulture: Plants for People and Places*, Dixon. Springer, Netherlands. 1: 327-405pp https://doi.org/10.1007/978-94-017-8578-5_10.
63. Guan W (2018) *Prune Determinate Tomatoes*. Purdue university. [September 12, 2021]
64. Guo FC, Fujime Y, Hirose T and Kato T (1991) Effects of the number of training shoots, raising period of seedlings and planting density in growth, fruiting and yields of sweet pepper. *J. Japan. Soc. Hort. Sci.* 59: 763-770.
65. Rajalingam GV, Rajasree V, Arumugam T and Saraswathi T (2017) Influence of different training systems in cucumber under naturally ventilated poly house. *International Journal of Chemical Studies*; 5(6): 1453-1455

66. Hatfield JL and Prueger JH (2015) Temperature extremes: Effect on plant growth and development. *Weather and climate extremes* 10: 4-10
67. Hernandez CVM and Sanches FCD (1992) Response to planting distance and pruning system in tomatoes growing in hydroponics culture in a basic green house. Deptt. Of de Fitecna, Univerdad Aufonoma Chapingo. 15 (73-74):23-25.
68. Heurn Ernst van and Post Kees Van Der (2004) Construction requirements and use of greenhouses in various climates. Digigrafi, Wageningen, the Netherlands. 90-96pp.
69. Heuvelink E, Marcelis LFM, Körner O (2002). How to reduce yield fluctuations in sweet pepper? *Acta Horticulturae* 633: 349-355
70. Hikosaka S and Sugiyama N (2003) Fruit growth patterns and abortion in multi-pistillate type cucumbers. *Journal of Horticultural Science & Biotechnology* 78: 775-779.
71. Hikosaka S and Sugiyama N (2004) Characteristics of flower and fruit development of multi-pistillate type cucumbers. *Journal of Horticultural Science & Biotechnology* 79: 219-22.
72. Hochmuth R. (2012) "Greenhouse Cucumber Production—Florida Greenhouse Vegetable Production Handbook," EDIS. University of Florida: IFAS Extension Vol 3.
73. Hochmuth RC, Leon LC and Hochmuth GJ (1996) Evaluation of Twelve Greenhouse Cucumber Cultivars and Two Training Systems over Two Seasons in Florida. *Proc. Fla. State. Hort. Soc.* 109:174-177.
74. Hodges L, Baxendale F (1991) Bee pollination of cucurbit crops. Report of University of Nebraska, Lincoln Cooperative Extension, NF91-50
75. Hong H (2000) Effect of pruning and spacing on yield and quality of cucumber. [Http://www.arc-avrdc.org](http://www.arc-avrdc.org).
76. Janapriya S, Palanisamy D and Ramaswamy MV (2010) Soilless media and fertigation for naturally ventilated polyhouse production of cucumber (*Cucumis sativus* L.) Cv. Green Long. *Internal. J. Agri., Environ. & Biotech* 3(2): 199-205.
77. Jovicich E, Cantliffe DJ and Hochmuth GJ (1999) Plant density and shoot pruning on yield and quality of a summer greenhouse sweet pepper crop in North central Florida. 184- 190p. In KD Batal (ed.) 28th National Agricultural Plastics Congress. Proc. Amer. Soc. Plasticulture, Tallahassee FL May 19- 22. ASP, State College, PA.
78. Jullien A, Munier-jolain NG, Malézieux E, Chillet M and Ney B (2001) Effect of pulp cell number and assimilate availability on dry matter accumulation rate in banana fruit [*Musa* sp. AAA group 'Grande Naine' (Cavendish subgroup)]. *Annals of Botany* 88: 321-30.

79. Junqueira CN and Augusto SC (2016) Bigger and sweeter passion fruits: effect of pollinator enhancement on fruit production and quality. *Apidologie* 48:131–140
80. Pramanik K, Mohapatra PP, Pradhan J, Acharya LK and Jena C (2020) Factors Influencing Performance of Capsicum under Protected Cultivation: A Review. *International Journal of Environment and Climate Change* 10(12): 572-588
81. Katsoulas N and Kittas C (2008) Impact of Greenhouse microclimate on Plant Growth and Development with special reference to Solanaceae. *European Journal of Plant Science and Biotechnology* 2(1): 31-44
82. Kaushal Shilpa and Singh Vijay (2019) Potentials and prospects of protected cultivation under Hilly Conditions. *Journal of Pharmacognosy and Phytochemistry* 8(1): 1433-1438
83. Kim JS, Ezura K, Lee J, Ariizumi T and Ezura H (2019) Genetic engineering of parthenocarpic tomato plants using transient *sliaa9* knockdown by novel tissue-specific promoters. *Scientific Reports* 9:18871
84. Kitinoja L and Kader AA (1995) “Small- scale post-harvest handling practices: a manual for horticultural crops. 3rd.Edition, Davis, California Available also at: <http://www.fao.org/inpho/EN/resources/library/index.asp>
85. Knott, JF (1928) The effect of apical pruning of tomato seedlings on growth and early yield. *Amer. Soc. Hort.Sci. Proc.*, 24: 21- 23.
86. Kong F and Singh RP (2011) Chemical Deterioration and physical instability of food and beverages. *Woodhead Publishing Series in Food Science, Technology and Nutrition* 29-62
87. Kugblenu YO, Danso EO, Ofori K, Andersen MN, Mickson SA, Sabi EB, Plauborg F, Abekoe MK, Anim JO, Ortiz R and Søren T. Jørgensen. Screening tomato genotypes for adaptation to high temperature in West Africa. *Acta Agriculturae Scandinavica, Section B - Soil &Plant Science* 37-41
88. Kukutani T, Inoue T & Maeta Y (1993) Pollination of strawberry by the stingless bee, *Trigona minangkabao*, and the honeybee, *Apis mellifera*: An experimental study of fertilization efficiency. *Res. Pop. Ecol.* 35: 95-111.
89. Kumar Patel NB, Saravaiya SN (2016) Response of parthenocarpic cucumber to fertilizers and training systems under naturally ventilated polyhouse in sub-tropical conditions. *International Journal of Current Research* (8):8051 – 8057.
90. Kumar S, Patel NB, Saravaiya SN (2014) Response of parthenocarpic cucumber to fertilizers and training systems under NVPH. *International Journal of Current Research*; 6(8): 8051-8057pp

91. Lal N, Sahu N, Marboh ES, Gupta AK and Patel RK (2017) A Review on Crop Regulation in Fruit Crops. *International Journal of Current Microbiology and Applied Sciences*. 6(7): 4032-4043
92. Lallensack R (2018) Busy Bees Take a Break During Total Solar Eclipses. *Smithsonian Magazine*. Accessed on [September 7, 2021]
93. Lambers H (2020) Plant reproductive system. *Encyclopedia Britannica*.
94. Lindsey R (2020) Climate Change: Atmospheric Carbon Dioxide. *Climate.gov science and Information for a climate-smart nation*.
95. Link H (2000) Significance of flower and fruit thinning on fruit quality. *Plant Growth Regulation* 31:17–26
96. Lorenzo, P. And Castilla, N (1995) Bell pepper yield response to plant density and radiation in unheated plastic greenhouse. *Acta Hort.* 32: 149-158
97. Macias MJO, Juezada-Euan JJG & Parra-Tabla V (2001) Behavior and efficiency of pollination of bees without aguijón (*Nannotrigona perilampoides*) in the cultivation of tomato (*Lycopersicon esculentum* M) under wintering conditions in Yucatán, Mexico, II Mexican Seminar on Bees without Aguijón – a vision on its biological cultivation. Autonomous University of Yucatán – Faculty of Veterinary Medicine and Animal Science Mérida, 119-124p.
98. Maeta Y, Tezuka T, Nadano H & Suzuki K (1992) Utilization of the Brazilian stingless bee *Nannotrigona testaceicornis* as a pollinator of strawberry. *Honey Bee Sci.* 13: 71-78.
99. Malagodi-Braga KS & Kleinert AMP (2004) Could *Tetragonisca angustula* Latreille (Apinae, Meliponini) be used as strawberry pollinator in greenhouses? *Aus. J. Agric. Res.* 55: 771-773.
100. Malagoli-braga KS, Kleinert AMP (2007) How does the behavior of bees on the strawberry flower (*Fragaria x ananassa* Dushesne) influence fruit formation? *Bioscience Journal, Uberlândia*, 23(1):76-81p,
101. Manisha P. How to Cultivate, Produce and Grow Sweet Pepper in Greenhouse? *Agriculture in India*
102. Manisha P.b. How to Produce Quality Vegetable Seeds? | *Agriculture. Agriculture in India.*
103. Marcelis LFM (1992) The dynamics of growth and dry matter distribution in cucumber. *Annals of Botany* 69:487–92.
104. Marcelis LFM (1996) Sink strength as a determinant of dry matter partitioning in the whole plant. *Journal of Experimental Botany*, 47:1281–1291.

105. Marin MV (2019) Buzz pollination: studying bee vibrations on flowers. *National Library of Medicine* 224(3):1068-1074
106. Mbonihankuye C, Kusolwa P and Msogoya TJ (2013) Assessment of the Effect of Pruning Systems on Plant Developmental Cycle – Yield and Quality of Selected Indeterminate Tomato Lines. *Acta Horticulturae* 1007(1007):535-542
107. Mcgregor SE (1976) *Insect Pollination of Cultivated Crops*, U.S. Department of Agriculture, Agriculture Handbook, 496 pp
108. Meena OP (2015) A review: role of plant growth regulators in vegetable production. *International Journal of Agricultural Science and Research* 5:71-84
109. Melissa P (2021) Stamen. *Encyclopedia Britannica*. Accessed on [September10, 2021]
110. Mengzhen Kang, Lili Yang, Baogui Zhang, Philippe de Reffye (2011) Correlation between dynamic tomato fruit-set and source–sink ratio: a common relationship for different plant densities and seasons? *Annals of Botany* 107(5):805–815
111. Miller C (2000) Carbon-nitrogen ratio: understanding chemical elements in organic matter. *Eco- Framing Daily*.
112. Morris LA (2004) Soil Biology and tree Growth: Soil Organic Matter Forms and Functions. *Encyclopedia of Forest Sciences*. 1201-1207.
113. Nagar S, Maurya IB, Hatwal PK (2015) Maintenance of gynoecious line of cucumber, *Cucumissativas L* through induction of staminate flower using silver nitrate and silver thiosulphate. *International Journal of Farm Sciences* 5(3): 67-73.
114. Nandgaonkar AK and Baker LR (1981) Inheritance of multi-pistillate flowering habit in gynoecious pickling cucumber. *Journal of the American Society for Horticultural Science*, 106:755–757.
115. Nir I, Goress R and Leshem B (1972) Effect of water stress, gibberellic acid and 2chloroethyl trimethyl ammonium chloride (CCC) on flower differentiation in Eureka lemon trees. *Amer. Soc. Hort. Sci.*, 97: 774-78.
116. Novak FJ and Maskova I (1979) Apical shoot tip culture of tomato. *Scientia horticulturae* 4: 337-344
117. Ozer AK, Saka and Saribaş HS (2017) The effect of different training systems on greenhouse organic tomato culture. *Acta Horticulturae* 101-106p
118. Boonkorkaew P, Hikosaka SN and Sugiyama (2008) Effect of pollination on cell division, cell enlargement, and endogenous hormones in fruit development in a gynoecious cucumber *Scientia Horticulturae* 116:1-7

119. Palada MC and Chang LC (2003) Suggested cultural practices for bitter gourd. AVRDC. Pub. No. 03-547: 1-5pp.
120. Paradiso R and Pascale SD (2014) Effects of Plant Size, Temperature, and Light Intensity on Flowering of Phalaenopsis Hybrids in Mediterranean Greenhouses. *The Scientific World Journal*. 420807ID
121. Pattnaik RK and Mohanty S (2021) Protected cultivation: importance, scope, and status. *Food and Scientific Reports*. 2(3):19
122. Pavani K, Jena C, Divya VV and Mallikarjunarao K (2020) Cultivation Technology of Tomato in Greenhouse. In book: *Protected Cultivation and Smart Agriculture*. 121-129pp
123. Perez JCD (2019) *Postharvest Physiology and Biochemistry of fruits and vegetables*. Woodhead Publishing: Science Direct. 157-173pp
124. Peter KV (2008) *Basics of Horticulture*. New India Publishing Agency. 9789389907544ISBN
125. Picken AJFL(1914) A Review of Pollination and Fruit Set in the Tomato (*Lycopersicon esculentum* Mill), *Journal of Horticultural Science*. 59(1): 113.
126. Pickens J and Sibley J (2019) *Greenhouse Cooling: An Overview of Fan and Pad Systems*, Crop Production Alabama A and M and Auburn Universities.
127. Potts SG, Breeze T, Herren BG (2014) Crop Pollination. *Encyclopedia of Agriculture and Food Systems* 408-418.
128. Pramanik K and Mohapatra PP (2017) Role of Auxin on Growth, Yield and Quality of Tomato - A Review. *Int.J. Curr.Microbiol.App.Sci* 6(11): 1624-1636
129. Primalatha MGS, Wahundeniya KB, Weerakkody WAP and Wicramathunga CK (2006) Plant Training and Spatial Arrangement for Yield Improvement in Greenhouse Cucumber (*cucumissativusl.*) Varieties. *Tropical Agricultural Research*. 18:346-357.
130. Rao GM, Suryanarayana MC (1988) Studies on the pollination of watermelon *Citrullus lanatus* (Thumb.). *Indian Bee Journal* 50, 5-8
131. Resh HM (1997) *Hydroponic Tomatoes for the Home Gardener*. Published by Woodbridge Press Publishing Company. California.
132. Resh HM (1996) *Hydroponic food production*, 5th edn, Woodridge Press Publ. Co., Santa Barbara, California.
133. Robert C, Hochmuth L and Leon LLC (1996) Evaluation of twelve greenhouse Cucumber cultivars and two training systems over two seasons in Florida. In: *Proceedings of the Florida State Horticultural Society*. Florida, USA. 174-177.

134. Robinson RW, Decker-Walters DS (1999) Cucurbits. Cambridge: CAB International 226 p
135. Roselino AC, Santos SAB, Bego LR (2010) Quality of sweet pepper (*Capsicum annuum* L.) fruits from flowers pollinated by stingless bees (*Melipona quadrifasciata anthidioides* Lepeletier 1836 and *Melipona scutellaris* Latreille 1811) under protected cultivation. *Brazilian Journal of Biosciences*, Porto Alegre 8(2):154-158
136. Rutledge AV (2015) Commercial Greenhouse Tomato Production. Agricultural Extension Service the University of Tennessee.
137. Sahu G, Aslam T, Das SP, Maity TK, Gupta NK (2017) A Study on Pre-flowering Foliar Spray of Plant Growth Regulator on Growth and Yield Parameters in Sweet Pepper (*Capsicum annuum* L.) Under Protected Condition. *International Journal of Current Microbiology and Applied Sciences* 6:3998-4007.
138. Santos SAB, Roselino AC, Bego LC (2008) Pollination of Cucumber, *Cucumis sativus* L. (Cucurbitales: Cucurbitaceae), by the Stingless Bees *Scaptotrigona* aff. *Depilis* Moure and *Nannotrigona testaceicornis* Lepeletier (Hymenoptera: Meliponini) in Greenhouses. *Neotropical Entomology*, Londrina 37(5):506-512
139. Schapendonk AHCM and Brouwer P (1984) Fruit growth of cucumber in relation to assimilate supply and sink activity. *Scientia Horticulturae* 23:21–33.
140. Sedgley M, Scholefield PB (1980) Stigma secretion in the watermelon before and after pollination. *Botanical Gazette* 141(4): 428-434
141. Singh Balraj (2015) Protected Cultivation of Vegetable Crops. Kalyani publishers, New Delhi.
142. Singh HP and Chadha KL (1988) Regulation of flushing and flowering in acid lime (*Citrus aurantifoliaswingle*) through stress management. *Prog. Hort* 20: 1-6.
143. Singh J, Vishwakarma G, Singh RK, Pandey K (2019) Crop Regulation in Fruit Crops. In book: Hi-Tech Horticulture Improved Production Techniques, New India Publishing Agency (NIPA)
144. Smitha K, Sunil KM (2016). Influence of growing environment on growth characters of cucumber (*Cucumis sativus* L.). *Journal of Tropical Agriculture* 54(2):201-203
145. Spehia RS and Sharma IP (2011) Protected cultivation for sustainable Horticulture. Bishen Singh Mahendra Pal Singh.
146. Starck Z (1983) Photosynthesis and endogenous regulation of the source-sink relation in tomato plants. *Photosynthetica* 17: 1–11.

147. Stein K, Coulibaly D, Stenchly K (2017) Bee pollination increases yield quantity and quality of cash crops in Burkina Faso, West Africa. *Sci Rep* 7, 17691
148. Tanaka, A. and Fujita K (1974) Nutriophysiological studies on the tomato plant. IV. Source-sink relationship and structure of the source-sink unit. *Soil Sci. Plant Nutr.* 20:305–315.
149. Tejaswini C. Rangaswamy, Shankarappa Sridhara, Nandini Ramesh, Pradeep Gopakkali, Diaa O. El-Ansary, Eman A Mahmoud, Shaimaa AM Abdelmohsen, Ashraf MM Abdelbacki, Hosam O Elansary and Amal ME Abdel-Hamid (2021) Assessing the Impact of Higher Levels of CO₂ and Temperature and Their Interactions on Tomato (*Solanum lycopersicum* L.). *Plants* 10: 14p
150. Thakur M and Rana RS (2008) Studies on the role of insect pollination on cucumber yield. *Pest Technology*. *Pest Technology*
151. Tiwari AK and Singh DK (2014) Use of Plant Growth Regulators in Tomato (*Solanum lycopersicum* L.) Under Tarai Conditions of Uttarkhand. *Indian Journal of Hill Farming* 27(2):38-40
152. Ummyiah HM, Wani KP, Khan SH and Magray MM (2017) *Journal of Pharmacognosy and Phytochemistry* 6(5): 1629-1634
153. Uzcategui NA and Baker LR (1979). Effects of multiplepistillate flowering on yields of gynoeocious pickling cucumbers. *Journal of the American Society for Horticultural Science* 104:148–51.
154. Velthuis HHW (2002) The historical background of the domestication of the Bumblebee, *Bombus terrestris* and its introduction in agriculture. In: Kevan, PG.; Imperatriz-Fonseca, VL (Ed.). *Pollinating bees: the conservation link between agriculture and nature*. Brasília: Ministério do Meio Ambiente 177- 184.
155. Wayne V (1990) Greenhouse cucumber production. University of Alaska Fairbanks Cooperative Extension Service. HGA – 00434.
156. Yadav S, Ameta KD, Sharma SK, Dubey RB, Rathore RS, Kumar H and Kapuriya VK (2017) Effect of Spacing and Training on Vegetative Growth Characteristics and Yield of Tomato (*Solanumlycopersicum* L.) Grown in Polyhouse. *International Journal of Current Microbiology and Applied Sciences*.6:1969-1976.
157. Prajapati S, Tarun Jamkar, Singh OP, Neelesh Raypuriya, Rishikesh Mandloi and Jain PK (2015) Plant growth regulators in vegetable production: an overview 15(2): 619-626.