

Screening for salt tolerance in *Chenopodium quinoa* genotypes seedlings through germination in a hydroponic system

Abstract

Quinoa (*Chenopodium quinoa* Willd.) is poised to be a global life changer with its ability to adapt to a wide range of abiotic stresses and as a highly nutritious and sustainable food source. Screening for salt tolerance at the germination and seedling stages of quinoa genotypes in control (CK), 200, 300 and 400 mM NaCl after 21 days in MS/2 mixture in 69 genotypes. This results in 16 genotypes with >50% germination at 400 mM NaCl. These were reassessed in germination indices and relative growth. Results indicated that *Chadmo* had the highest germinability of 97% and 32.76% relative height among the 16 genotypes. Considering the germination indices, *Chadmo* had significantly different values (3.05 day^{-1}) in mean germination time, coefficient of variation of the germination time (38.76%), the velocity of germination (0.23 day^{-1}), the uncertainty of germination (0.54 bit), synchrony of germination (0.42) and Timson's index (48.89) with significant differences ($P < 0.05$) among the genotypes. Moreover, *Chadmo* had the highest membrane stability index (MSI) (60.03) at 400 mM NaCl and the least relative change between the CK and 400 mM NaCl with 30.87%. Assessing the stress inhibitory effect of the 16 genotypes, *Chadmo* had the least relative difference between the CK and 400 mM NaCl with shoot length of 34.34%, root length of 25.57%, fresh weight of 22.05%, dry weight of 3.62% and moisture content of 1.99% with Tukey analyses identifying significant differences ($p < 0.05$). To select the salt-sensitive genotype, an assessment was done on five genotypes that exhibited the least germination at 200 mM NaCl. *Kankolla* had the least germinability with 11 and 4% at 100 and 200 mM NaCl, respectively. Juxtaposing all these parameters, *Chadmo* and *Kankolla* were selected as the salt-tolerant and salt-sensitive for further analyses.

Keywords: germination; propagation; quinoa; salt-sensitive; salt-tolerant; *Kankolla*; *Chadmo*

1. Introduction

Germination is a critical stage in a plant's life cycle for propagation, continuation and inevitable for the survival of humans, as it forms the source of indispensable food and other necessities ([Joosen et al., 2010](#); [Larson et al., 2015](#)). [Bewley \(1997\)](#) defined germination as the outgrowth of a radicle through the endosperm and epiderm of a seed. Germination is a dynamic process in which water plays an important role, hence any condition that limits the availability of water will result in delay or hindrance. Salinity and osmotic stress, temperature, light, and pH all influence the germinability of seeds ([Bai et al., 2018](#); [Dekker & Gilbert, 2014](#); [Hanif et al., 2017](#); [Javaid et al., 2018](#); [Mahmood et al., 2016](#); [Tanveer et al., 2013](#)). While some seeds may be susceptible to slight variation, others have evolved to adapt to higher tolerance levels of various conditions. Salinity influences germination based on varieties, species and salt content of the soil and what mechanism is adopted by the plant for its protection and defence ([Batlla & Benech-Arnold, 2014](#); [Florentine et al., 2016](#); [Hanif et al., 2017](#); [Javaid et al., 2018](#)).

While some studies have indicated that salt tolerance during germination is independent of other growth phases in *Triticum aestivum* L. varieties ([Mano & Takeda, 2001](#)) and *Solanum lycopersicum* L. ([Foolad & Lin, 1997](#)). [Prado et al. \(2000\)](#) outlined that once sprouting and rooting have occurred; then the seedlings have a high probability of proliferating successfully in their life cycle. Germination techniques and indices were also employed to identify salt-tolerant and salt-sensitive varieties of rice ([Mondal & Pramanik, 1995](#); [Shi et al., 2017](#); [Wang et al., 2010](#); [Wang et al., 2011](#)). The screening was successful with 182 varieties of quinoa to identify the salt-sensitive and salt-tolerant during germination ([Gómez-Pando et al., 2010](#)). This method of screening was also supported by [Nasir et al. \(2000\)](#) and [Cha-Um et al. \(2012\)](#), they concluded that the seedling stage of sugarcane provides an effective strategy for screening salt-tolerant and susceptible varieties.

Salinity prevents or delays the germination of seeds or seedling growth and development. Germination indices can be used to select tolerant variety at an early stage. Even though many plants have a differential response at various stages to salinity, germination indices can be used as a precursor of selection, while some have posted that germination is independent of further growth and development. Also, some seeds that showed tolerance during germination do not necessarily transcend to seedling growth and further development ([Aflaki et al., 2017](#); [Bybordi, 2010](#); [Foolad & Lin, 1997](#); [Houle et al., 2001](#); [Mano & Takeda, 2001](#); [Sanchez et al., 2014](#)). However,

others have posited that germination and sprouting at the seedling stage are reliable and effective methods to categorize plants as salt-tolerant and sensitive varieties ([Gómez-Pando et al., 2010](#); [Nasir et al., 2000](#); [Prado et al., 2000](#)). For selecting the tolerant variety for this study, germinability (G), mean germination time (GMT), coefficient of velocity of germination (CV_g), uncertainty of germination (U) and synchrony of germination (Z) indices were employed.

2. Material and Methods

2.2 (i) Plant materials and growth conditions

Salt-tolerant and salt-sensitive varieties were selected through rigorous screening from a collection of 97 seeds obtained from the United States Department of Agriculture (USDA). The collection of seeds was sown and propagated for seed proliferation and enhanced quality control. These plants were grown at the Fujian Agriculture and Forestry University glasshouse in ambient light in the temperature-controlled environment at about 24 - 26 °C and average relative humidity of ~65-70 % with 16/8 h light/dark photoperiod ([Goyal et al., 2016](#); [Morales et al., 2011](#); [Panuccio et al., 2014](#)).

After harvest, seeds were stored at 4 °C until the experiment commenced ([Panuccio et al., 2014](#)). Seeds were vapour sterilized with 3 % sodium hypochlorate and HCl in a desiccator placed in a fume hood for 4 ½ h after which they were air blown in a horizontal laminar flow hood for 3 h ([Burrieza et al., 2012](#); [Lindsey et al., 2017](#); [Panuccio et al., 2014](#); [Ruiz-Carrasco et al., 2011](#)).

2.2 (ii) Treatment and Selection

Seeds were tested at 0, 100, 200, 300, and 500 mM NaCl-induced MS/2 media over 21 days ([Lindsey et al., 2017](#); [Murashige & Skoog, 1962](#); [Postnikova et al., 2013](#); [Wang et al., 2009](#)). Fifteen seeds were sown in tissue culture bottles containing MS/2 media with the respective salt concentration and control in three technical replicates placed in a culture room at 22 °C and 60 – 65 % RH ([Joosen et al., 2010](#); [Murashige & Skoog, 1962](#); [Postnikova et al., 2013](#); [Wang et al., 2009](#)). Seed germination was recorded daily for seven days ([Panuccio et al., 2014](#)). After seven days, the bottles were moved to another room with 26 °C and 65 % RH, better elongation and growth-related conditions to facilitate better seedling development ([Joosen et al., 2010](#); [Panuccio et al., 2014](#)).

Seedling height for the technical and biological replicates was measured and was recorded after 21 days.

Salt-tolerant varieties were selected based on the mean germination time, mean germination rate, coefficient of variation of germination time, uncertainty of germination frequency, and synchrony of germination at 400 mM NaCl and the relative growth (height) between the control and maximum treatment ([Aflaki et al., 2017](#); [Kader, 2005](#); [Kader & Jutzi, 2004](#); [Maguire, 1962](#); [Panuccio et al., 2014](#); [Ranal et al., 2009](#); [Scott et al., 1984](#)). The salt-sensitive varieties were selected based on the least germination percentage at the minimum treatment (100 mM NaCl) but with over maximum per cent germination at the control, credence to seed viability.

The first-tier screening resulted in 20 salt-tolerant and 5 salt-sensitive varieties. These were reassessed following the above procedure. The results did not indicate a significant difference from the initial screening. The most salt-tolerant varieties were then subjected to salt conditions at 400 and 500 mM NaCl-induced MS/2 ([Lindsey et al., 2017](#); [Murashige & Skoog, 1962](#); [Postnikova et al., 2013](#); [Wang et al., 2009](#)). This yielded similar results, as with the previous trials, at 400 mM NaCl but germination was poor (6 %) at 500 mM NaCl and with the few (3 seeds) that germinated, no elongation occurred, sprouting followed by death.

2.3 Germination Analysis

In determining the seed for the tolerant variety, the following germination parameters were used with the respective formulas:

2.3.1 Germinability:

Equation 1

$$\sum_{i=1}^n S_i / D_i$$

Where S_i : germinated seeds per time (day), D_i represents seed numbers from the start of the experiment to the i^{th} , n_i : number of seeds germinated in the i^{th} day ([Bai et al., 2018](#); [Maguire, 1962](#); [Panuccio et al., 2014](#); [Ranal et al., 2009](#)).

2.32 Mean germination time:

Equation 2

$$\bar{t} = \frac{\sum_{i=1}^k n_i t_i}{\sum_{i=1}^k n_i}$$

Where t_i : time (day) from the start of the experiment to the i^{th} , n_i : number of seeds germinated in the i^{th} day, and k : last time of germination ([Aflaki et al., 2017](#); [Kader, 2005](#); [Panuccio et al., 2014](#); [Scott et al., 1984](#)).

2.33 Coefficient of variation of the germination time:

Equation 3 (i)

$$s_t^2 = \frac{\sum_{i=1}^k n_i (t_i - \bar{t})^2}{\sum_{i=1}^k n_i - 1}$$

Where t : mean time; t_i : the time between the start of the experiment and the i^{th} day; n_i : number of seeds germinated in the i^{th} day, and k : last day of germination. The variance value will be used to calculate the coefficient of variation of the germination time in the subsequent formula:

Equation 3 (ii)

$$CV_t = \frac{s_t}{\bar{t}} \cdot 100$$

Where s_t : the standard deviation of the germination time and t is: mean germination time ([Kader & Jutzi, 2004](#); [Maguire, 1962](#); [Panuccio et al., 2014](#); [Ranal et al., 2009](#)).

2.34 Uncertainty of germination:

Equation 4

$$U = -\sum_{i=1}^k f_i \log_2 f_i, \text{ being } f_i = \frac{n_i}{\sum_{i=1}^k n_i}$$

Where $n_{i is}$: number of seeds germinated on the i^{th} time, and $k is$: last day of observation ([Ranal et al., 2009](#)).

2.35 Synchrony of germination (Z)

Equation 6

$$Z = \frac{\sum_{i=1}^k C_{n_i,2}}{C_{\sum n_i,2}}, \text{ being } C_{n_i,2} = n_i(n_i-1)/2$$

Where: $C_{ni,2}$ combinations of the seeds germinated in the i^{th} time, two by two, and ni : number of seeds germinated in the i^{th} time. Z is the quotient between the sums of the partial combinations of the number of seeds germinated in each ti , two by two combinations of the total number of seeds germinated at the end of the experiment ([Ranal et al., 2009](#)).

2.4 Results

2.41 (a) Selecting salt-tolerant variety

Seed tolerance screening occurred systematically in a three-tier method. Criteria used to identify and select the tolerant varieties were the germination percentage, mean germination time, coefficient of the velocity of germination, the uncertainty of germination, synchrony of germination and relative growth (height) - the ratio of plantlet height (cm) between the CK and the 400 mM NaCl to select the tolerant variety. Additionally, for validation and quality control, the varieties with $\geq 50\%$ germinability (G_{50}) at 400 mM NaCl were subjected to further testing at 450 and 500 mM NaCl, but they all displayed < 30 and 15% germinability, respectively and those that germinated died thereafter from apparent desiccation. For the highest germinability, varieties PI 614880 and PI 614884 had 97 and 93%, respectively. For the relative growth, they also exhibited the least between the control and 400 mM NaCl with PI 614880 at -32.76 and 69 at -45.89 (Figure 1)




Figure 1 Germinability and relative growth between control and 400 mM NaCl of the salt-tolerant varieties (G50). Data represents means \pm SD of fifteen biological and three technical replicates.

Evaluation of the other germination parameters indicated that the coefficient of variation in germination time (CV_t), germination speed (v) and uncertainty of germination frequency (U) were all the highest for PI 614880 while the same cannot be said for mean germination time (MT) and synchrony (Z) of germination, but they, however, do represent high and low values, respectively (Table1). Hence, observation from the data guided the selection of PI 614880 (Chadmo) as the most salt-tolerant among the varieties at 400 mM NaCl.

Table 1. 400 NaCl treatment with 3 technical replicates constituting 15 biological replicates each. (MT = t :

mean germination time; ν : mean germination rate; CV_t : coefficient of variation of the germination time; U : uncertainty of the germination frequency; Z : synchrony of the germination process).

No.	Accession #	MT (day)	CV_t (%)	ν (day ⁻¹)	U (bit)	Z
1	PI 677096	3.97±0.25	26.52±3.62	0.25±0.02	1.70±0.06	0.31±0.04
2	PI 433232	3.80±0.11	27.76±0.00	0.26±0.01	1.91±0.01	0.22±0.00
3	PI 674266	3.78±0.21	36.98±1.92	0.26±0.01	1.22±0.06	0.09±0.01
4	PI 677100	3.17±0.15	21.61±4.54	0.31±0.01	1.31±0.28	0.33±0.06
5	Ames 13722	4.61±0.42	7.74 ±5.16	0.21±0.02	0.54±0.35	0.75±0.17
6	Ames 13723	5.04±0.15	15.41±1.14	0.19±0.01	1.41±0.08	0.32±0.06
7	Ames 13724	3.47±0.21	21.06±3.83	0.28±0.01	0.98±0.06	0.50±0.02
8	Ames 13725	4.60±0.08	23.48±2.45	0.21±0.00	1.62±0.14	0.28±0.06
9	Ames 13726	4.68±0.19	23.95±4.61	0.21±0.01	1.60±0.28	0.28±0.09
10	Ames 13735	5.04±0.08	14.65±0.83	0.19±0.00	1.44±0.06	0.32±0.04
11	Ames 13736	5.60±0.07	9.13±0.51	0.17±0.00	0.95±0.06	0.47±0.05
12	Ames 13740	3.46±0.17	26.74±4.54	0.29±0.01	1.42±0.29	0.36±0.12
13	Ames 13746	4.52±0.23	19.21±2.25	0.22±0.01	1.65±0.12	0.28±0.05
14	Ames 13747	4.01±0.06	26.58±2.78	0.24±0.00	1.61±0.23	0.30±0.08
15	PI 587173	4.59±0.19	35.90±4.89	0.21±0.01	1.91±0.08	0.24±0.02
16	PI 614880	4.25±0.16	38.76±1.97	0.23±0.01	2.04±0.08	0.12±0.05
17	PI 614884	3.64±0.19	32.22±1.60	0.27±0.01	1.89±0.34	0.23±0.06
18	PI 614920	3.26±0.13	28±0.02	0.30±0.01	1.70±0.12	0.25±0.01
19	PI 634917	3.28±0.09	31.58±2.15	0.30±0.01	1.86±0.18	0.21±0.04
20	PI 634918	3.13±0.03	25.64±1.35	0.31±0.00	1.53±0.02	0.29±0.01

2.41 (b) Selection of salt-sensitive variety

The criteria applied for the selection of the salt-sensitive varieties were the lowest germination percentage at the 100 and 200 mM NaCl and the largest height difference between the control and 100 mM NaCl (68.21%) and between control and 200 mM NaCl (72.24%). However, germination took precedence for varieties 62 and 80 even though the height difference was greater in varieties 40, 41 and 51 because while they indicated a low germination rate at 200 mM NaCl, they also showed evidence of germination at 300 mM NaCl (Table 2). It is important to note that even though germination has occurred; the seedlings did not sprout into well-developed plantlets but died, presumable for the intense stress that led to desiccation. Therefore, upon these observations, Kankolla (PI 510551) was selected as the most salt-sensitive among the tested varieties.

Table 2: Selected salt-sensitive varieties through germination and growth

No	Genotype	Germinability (%)					Height (cm)					Relative growth (height)		
		CK	100	200	300	400	CK	100	200	300	400	100	200	300
1	Ames 13755(40)	97	59	11	2	0	3.99	1.34	1	1.1	0	-66.41	-74.94	-72.43
2	Ames 13756(41)	97	55	15	1	0	3.98	2.35	2.43	1.5	0	-23.70	-38.94	-62.31
3	PI 478418 (51)	95	65	11	2	0	3.64	2	2	1.4	0	-45.05	-45.05	-61.54
4	PI 510551(6) - Kankolla	100	12	4	0	0	5.02	1.6	1	0	0	-68.21	-80.07	0
5	PI 614932(80)	100	46	6	0	0	3.89	2.4	1.9	0	0	-38.30	-51.16	0

2.4 Discussion

Germination is a key process that catapults plants into a continuous cycle of multiple biological and physiological processes that will determine their ability to survive and reproduce, and in many instances, in conditions that may not be conducive. Seed germination is the most critical stage in the growth of the plant and is the most sensitive when exposed to abiotic stress ([Mickky & Aldesuquy, 2017](#); [Yadav et al., 2011](#)). Salinity is an environmental factor that significantly influences germination and hence, the plant's prospect of continuity. Salinity has been shown to delay or prohibit germination in many genotypes of plants and those that survived, in most cases, proceeded to grow, produce and reproduced and hence, are deemed to be tolerant ([Dekker & Gilbert, 2014](#); [Javaid et al., 2018](#); [Mahmood et al., 2016](#); [Tanveer et al., 2013](#)). Despite its halophytic nature, quinoa is rather sensitive to stressful conditions in its vegetative stage, hence, if survived seedling establishment is highly possible ([Ruiz et al., 2017](#); [Ruiz et al., 2019](#)). Among the 16 most tolerant genotypes assessed, *Chadmo* had the highest percentage germinability, among the highest in germination meantime, and the lowest relative growth is calculated as the difference between the CK and 400 mM NaCl. Having the highest germinability (97%) among the 16 genotypes, and with no fatality, is indicative of the ability to germinate in highly saline conditions and progress to the seedling establishment (400 mM NaCl) (Figure 1). Germination is regarded as the most sensitive stage to abiotic stress in the development of a plant and therefore, once they have survived, the seedling establishment will ensue ([Mickky & Aldesuquy, 2017](#); [Yadav et al., 2011](#)). While germination is important in determining tolerance to salinity, if seedling elongation and growth do not proceed then it is irrelevant to the continuity of its life cycle. However, in the assessment of the seedling elongation and growth of the 16 genotypes, *Chadmo*

had the lowest relative growth (32.76%), which is interpreted as the least difference between the CK and 400 mM NaCl among the genotypes (Figure 1).

With these 16 genotypes, root length decreased significantly between the CK and 400 mM NaCl, *Chadmo* exhibited the minimum difference between the CK and 400 mM NaCl, which indicated it is least affected and hence the most tolerant regarding root growth while PI674266 was most affected with 81.19%. Survival of a plant being exposed to salinity mainly depends on how the root system manipulates the intake and distribution of salt as it is the first interface between the plant and that stressful abiotic condition. However, the robust root structure allows them to survive and more so it is sometimes the least affected as compared to shoots, but root elongation is affected by salinity at higher concentrations ([Gómez-Pando et al., 2010](#); [Katembe et al., 1998](#); [Munns & Tester, 2008](#)). Some plants are very well adapted to exclude salt at the level of the root by developing salt filtration mechanism through enhancing hydrophobic barrier deposition, which prevents the absorption of non-selective apoplastic ions ([Kolattukudy, 1984](#); [Krishnamurthy et al., 2014](#); [Lawton et al., 1981](#)). At all levels of plant growth, height attenuation is symptomatic of salinity stress and if a plant can germinate and proceed to growth then it is undoubtedly tolerant of such stressful conditions ([Gómez-Pando et al., 2010](#); [Prado et al., 2000](#)).

To infer from a comparative perspective, the MSI index was assessed at CK, 200, 300 and 400 mM NaCl. While the response was differential among the genotypes and the treatments, *Chadmo* recorded the lowest difference (26.81%) between the CK and 400 mM NaCl while the highest relative difference (42.90%) was observed in Ames 13736. Salinity results in significant to plant cells and more particularly on the membrane and these damages are measured through membrane stability index or electrolyte leakage ([Munns & Tester, 2008](#); [Panta et al., 2014](#); [Sairam & Srivastava, 2002](#); [Shabala & Mackay, 2011](#)). In this study, therefore, *Chadmo* (30.87%) had the least relative difference between the CK and 400 mM NaCl among the 16 salt-tolerant genotypes and hence is designated as the most salt-tolerant. Likewise, the salinity tolerance of two wheat genotypes (*Kharchia 65* and *KRL 19*) was analysed and the conclusion was that MSI decreased under salinity stress in both genotypes at three different stages of sampling but higher in *Kharchia 65* and compared with *KRL 19*, thereby prescribing *Kharchia 65* as the more salt-tolerant genotype ([Sairam et al., 2002](#)). Additionally, nine genotypes of pea plants indicated a decrease in MSI under salinity for all the genotypes at different NaCl treatments (25, 50 and 75

mM NaCl) as compared to the CK ([Shahid et al., 2012](#)). In support, it was also concluded that salt-treated strawberries had a 10% reduction in MSI when treated with 50 mM NaCl ([Avestan et al., 2019](#)).

Germination indices can be used to select tolerant genotypes at an early stage. Even though many plants have a differential response at various stages to salinity, germination indices can be used as a precursor for selection, while some have posited that germination is independent of further growth and development ([Aflaki et al., 2017](#); [Bybordi, 2010](#); [Houle et al., 2001](#); [Sanchez et al., 2014](#)). Quinoa tolerance to salinity during germination results from the changes in the primary metabolites and enzyme activity in response to salinity ([Adolf et al., 2012](#); [González et al., 2015](#)). Many also supported that germination and sprouting at the seedling stage are reliable and effective methods to categorize plants as salt-tolerant and sensitive genotypes ([Gómez-Pando et al., 2010](#); [Prado et al., 2000](#)).

Based on these results on the germination indices, *Chadmois* deemed as the most tolerant genotype among the 16 salt-tolerant genotypes (Table 1)). Hence, the germination process under saline conditions is independent of other biological and physiological processes ([Gómez-Pando et al., 2010](#); [Prado et al., 2000](#)). These germination indices have been used singly or collectively to screen for salt tolerance at the seedling stage in many plants. Germinability and seedling growth were used to assess the responses of three cultivars of bean (*Phaseolus vulgaris* L.) to NaCl and Na₂SO₄ and results showed that both have an inhibitory effect on germination and seedling development ([Kaymakanova, 2009](#)). Furthermore, the responses of *Atriplex prostrata* and *A. patula* after being exposed to NaCl and PEG were determined with the application of germination rate and percentages were used to assess their susceptibility ([Katembe et al., 1998](#)). In support, also worked with quinoa (cv *Titicaca*) to identify germination and seedling tolerance levels to saline water using the germination traits of the coefficient of velocity of germination, germination rate index and mean germination time ([Panuccio et al., 2014](#)). They posited that salinity at a lower concentration does not affect germination percentages but rather increases the germination rate.

The germination indices were irrelevant to select salt-sensitive genotypes because to decide on their sensitivity, it was based on them not germinating and developing into seedlings under saline conditions ([Gómez-Pando et al., 2010](#)). If this were the case, then they would have been sensitive but rather tolerant. Among the five sensitive genotypes assessed, *Kankolla* had the

lowest germinability at 100 mM NaCl and 200 mM NaCl with 12 and 4%, respectively and no germination occurred at 300 mM NaCl. Analysis of variance identified a significant difference between the treatments of each genotype at $P < 0.05$. Therefore, upon these observations, *Kankolla* was selected as the most salt-sensitive among the five tested genotypes. Plants that proliferate in about 200 mM NaCl concentration are referred to as halophyte which makes up about only 1% of all other plants (Flowers & Colmer, 2008; Flowers & Yeo, 1995; Sanchez et al., 2003). *Kankolla* based on these criteria was regarded as the most salt-sensitive genotype (Anjum et al., 2014; Bosque-Sanchez et al., 2003; Glenn et al., 2013; Razzaghi et al., 2015). Therefore, the parameters used in assessing for salt tolerance were relevant and applicable to assist in arriving at an informed decision on the level of tolerance in the quinoa for this study.

Conclusion

It can be concluded that the two contrasting genotypes of salinity tolerance were *Chadmo* (salt-tolerant) and *Kankolla* (salt-sensitive). Further evidence to support this, was entrenched in the origin and locale of both genotypes and the prevailing environmental conditions; *Chadmo* originated from the coastline of Chile (<10 m from the Southern Pacific Ocean), while *Kankolla* originated deeper, and in the upland area of Arapa District in Peru (387 km from the Southern Pacific Ocean). The coastline is normally inundated with saltwater and marshy areas, therefore, causing the soil to be saline. Hence, if *Chadmo* thrives in this area then would have to be halophytic. Conversely, *Kankolla* predominantly grows in the upland areas thereby becoming more adapted and thrives in non-saline soils and is, therefore, more sensitive to salinity. Additionally, the results of germination in response to the different salinity to select the most salt-tolerant genotype, in the above experiment, have been further corroborated in morpho-physicochemical considerations on the salt-treated and control seedlings in the subsequent experiment.

Reference

- Adolf, V. I., Shabala, S., Andersen, M. N., Razzaghi, F., & Jacobsen, S.-E. (2012). Varietal differences in quinoa's tolerance to saline conditions. *Plant and Soil*, 357(1-2), 117-129. <https://doi.org/https://doi.org/10.1007/s11104-012-1133-7>
- Aflaki, F., Sedghi, M., Pazuki, A., & Pessarakli, M. (2017). Investigation of seed germination indices for early selection of salinity tolerant genotypes: A case study in wheat. *Emirates Journal of Food and Agriculture*, 222-226. <https://doi.org/https://doi:10.9755/ejfa.2016-12-1940>
- Anjum, N. A., Ahmad, I., Válega, M., Mohmood, I., Gill, S. S., Tuteja, N., Duarte, A. C., & Pereira, E. (2014). Salt marsh halophyte services to metal-metalloid remediation: Assessment of the processes and underlying mechanisms. *Critical Reviews in Environmental Science and Technology*, 44(18), 2038-2106.

- <https://doi.org/https://doi/abs/10.1080/10643389.2013.828271>
- Avestan, S., Ghasemnezhad, M., Esfahani, M., & Byrt, C. S. (2019). Application of Nano-Silicon Dioxide Improves Salt Stress Tolerance in Strawberry Plants. *Agronomy*, 9(5), 246. <https://doi.org/https://doi:10.3390/agronomy9050246>
- Bai, J., Yan, W., Wang, Y., Yin, Q., Liu, J., Wight, C., & Ma, B. (2018). Screening oat genotypes for tolerance to salinity and alkalinity. *Frontiers in plant science*, 9. <https://doi.org/https://doi:10.3389/fpls.2018.01302>
- Batlla, D., & Benech-Arnold, R. L. (2014). Weed seed germination and the light environment: Implications for weed management. *Weed Biology and Management*, 14(2), 77-87. <https://doi.org/https://doi10.1111/wbm.12039>
- Bewley, J. D. (1997). Seed Germination and Dormancy. *Plant Cell*, 9(7), 1055-1066. <https://doi.org/https://doi10.1105/tpc.9.7.1055>
- Bosque-Sanchez, H., Lemeur, R., Damme, P. v., & Jacobsen, S.-E. (2003). Ecophysiological analysis of drought and salinity stress of quinoa (*Chenopodium quinoa* Willd.). *Food Reviews International*, 19(1-2), 111-119. <https://doi.org/https://doi10.1081/FRI-120018874>
- Burrieza, H. P., Koyro, H.-W., Tosar, L. M., Kobayashi, K., & Maldonado, S. (2012). High salinity induces dehydrin accumulation in *Chenopodium quinoa* Willd. cv. Hualhuas embryos. *Plant and Soil*, 354(1-2), 69-79. <https://doi.org/https://doi10.1007/s11104-011-1045-y>
- Bybordi, A. (2010). The Influence of Salt Stress on Seed Germination, Growth and Yield of Canola Cultivars. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 38(1), 128-133. <https://doi.org/https://doi.org/10.15835/nbha3813572>
- Cha-Um, S., Chuencharoen, S., Mongkolsiriwatana, C., Ashraf, M., & Kirdmanee, C. (2012). *Screening sugarcane (Saccharum sp.) genotypes for salt tolerance using multivariate cluster analysis* (Vol. 110). <https://doi.org/https://doi10.1007/s11240-012-0126-9>
- Dekker, J., & Gilbert, J. (2014). Weedy adaptation in *Setaria* spp.: IX. Effects of salinity, temperature, light and seed dormancy on *Setaria faberi* seed germination. *arXiv preprint arXiv:1408.6187*. [https://doi.org/https://doiarXiv:1408.6187\[q-bio.PE](https://doi.org/https://doiarXiv:1408.6187[q-bio.PE)
- Florentine, S. K., Weller, S., Graz, P. F., Westbrooke, M., Florentine, A., Javaid, M., Fernando, N., Chauhan, B. S., & Dowling, K. (2016). Influence of selected environmental factors on seed germination and seedling survival of the arid zone invasive species tobacco bush (*Nicotiana glauca* R. Graham). *Rangeland Journal*, 38(4), 417-425. <https://doi.org/https://doi10.1071/Rj16022>
- Flowers, T. J., & Colmer, T. D. (2008). Salinity tolerance in halophytes. *New Phytol*, 179(4), 945-963. <https://doi.org/https://10.1111/j.1469-8137.2008.02531.x>
- Flowers, T. J., & Yeo, A. R. (1995). Breeding for Salinity Resistance in Crop Plants: Where Next? *Functional Plant Biology*, 22(6), 875. <https://doi.org/https://doi10.1071/pp9950875>
- Foolad, M., & Lin, G. (1997). Absence of a genetic relationship between salt tolerance during seed germination and vegetative growth in tomato. *Plant Breeding*, 116(4), 363-367. <https://doi.org/https://doi/abs/10.1111/j.1439-0523.1997.tb01013.x>
- Glenn, E. P., Anday, T., Chaturvedi, R., Martinez-Garcia, R., Pearlstein, S., Soliz, D., Nelson, S. G., & Felger, R. S. (2013). Three halophytes for saline-water agriculture: An oilseed, a forage and a grain crop. *Environmental and Experimental Botany*, 92, 110-121. <https://doi.org/https://10.1016/j.envexpbot.2012.05.002>
- Gómez-Pando, L. R., Álvarez-Castro, R., & Eguiluz-de la Barra, A. (2010). Short Communication: Effect of Salt Stress on Peruvian Germplasm of *Chenopodium quinoa* Willd.: A Promising Crop. *Journal of agronomy and crop science*, 196(5), 391-396. <https://doi.org/https://doi10.1111/j.1439-037X.2010.00429.x>
- González, J. A., Eisa, S. S., Hussin, S., & Prado, F. E. (2015). Quinoa: an Incan crop to face global changes in agriculture. *Quinoa: Improvement and sustainable production*, 1-18. Retrieved 26/03/2019, from <https://onlinelibrary.wiley.com/doi/abs/10.1002/9781118628041.ch1>
- Goyal, E., Amit, S. K., Singh, R. S., Mahato, A. K., Chand, S., & Kanika, K. (2016). Transcriptome profiling of the salt-stress response in *Triticum aestivum* cv. Kharchia Local. *Scientific reports*, 6, 27752. <https://doi.org/https://doi10.1038/srep27752>
- Hanif, Z., Naeem, M., Ali, H. H., Tanveer, A., Javaid, M. M., Peerzada, A. M., & Chauhan, B. S. (2017). Effect of Environmental Factors on Germination of *Salsola foetida* : Potential Species for Rehabilitation of Degraded Rangelands. *Rangeland Ecology & Management*, 70(5), 638-643. <https://doi.org/https://doi10.1016/j.rama.2017.02.003>
- Houle, G., Morel, L., Reynolds, C., & Siegel, J. (2001). The effect of salinity on different developmental stages of an endemic annual plant, *Aster laurentianus* (Asteraceae). *American Journal of Botany*, 88(1), 62-67. <https://doi.org/https://doi10.2307/2657127>
- Javaid, M. M., Florentine, S., Ali, H. H., & Weller, S. (2018). Effect of environmental factors on the germination and

- emergence of *Salvia verbenaca* L. cultivars (verbenaca and vernalis): An invasive species in semi-arid and arid rangeland regions. *PLoS One*, 13(3). <https://doi.org/https://doi10.1371/journal.pone.0194319>
- Joosen, R. V., Kodde, J., Willems, L. A., Ligterink, W., van der Plas, L. H., & Hilhorst, H. W. (2010). GERMINATOR: a software package for high-throughput scoring and curve fitting of Arabidopsis seed germination. *Plant Journal*, 62(1), 148-159. <https://doi.org/http://doi10.1111/j.1365-313X.2009.04116.x>
- Kader, M. (2005). A comparison of seed germination calculation formulae and the associated interpretation of resulting data. *Journal and Proceeding of the Royal Society of New South Wales*, 138, 65-75. https://doi.org/https://royalsoc.org.au/images/pdf/journal/138_Kader.pdf
- Kader, M., & Jutzi, S. (2004). Effects of thermal and salt treatments during imbibition on germination and seedling growth of sorghum at 42/19 C. *Journal of Agronomy and Crop Science*, 190(1), 35-38. <https://doi.org/https://doi10.1046/j.0931-2250.2003.00071.x>
- Katembe, W. J., Ungar, I. A., & Mitchell, J. P. (1998). Effect of Salinity on Germination and Seedling Growth of two Atriplex species (Chenopodiaceae). *Annals of Botany*, 82, 167-175.
- Kaymakanova, M. (2009). Effect of salinity on germination and seed physiology in bean (*phaseolus vulgaris* L.). *Biotechnology & Biotechnological Equipment*, 23(1), 326-329. <https://doi.org/https://doi.org/10.1080/13102818.2009.10818430>
- Kolattukudy, P. E. (1984). Biochemistry and function of cutin and suberin. *Canadian Journal of Botany*, 62(12), 2918-2933. <https://doi.org/https://doi10.1139/b84-391>
- Krishnamurthy, P., Jyothi-Prakash, P., A, Qin, L., He, J., Lin, Q., Loh, C.-S., & Kumar, P. P. (2014). Role of root hydrophobic barriers in salt exclusion of a mangrove plant *Avicennia officinalis*. *Plant, Cell & Environment*, 37(7), 1656-1671. <https://doi.org/https://doi.org/10.1111/pce.12272>
- Larson, J. E., Sheley, R. L., Hardegree, S. P., Doescher, P. S., & James, J. J. (2015). Seed and seedling traits affecting critical life stage transitions and recruitment outcomes in dryland grasses. *Journal of Applied Ecology*, 52(1), 199-209. <https://doi.org/https://doi10.1111/1365-2664.12350>
- Lawton, J. R., Todd, A., & Naidoo, D. (1981). Preliminary investigations into the structure of the roots of the mangroves, *Avicennia marina* and *Bruguiera gymnorrhiza*, in relation to ion uptake. *New Phytologist*, 88(4), 713-722. <https://doi.org/https://doi/full/10.1111/j.1469-8137.1981.tb01748.x>
- Lindsey, B. E., 3rd, Rivero, L., Calhoun, C. S., Grotewold, E., & Brkljacic, J. (2017). Standardized Method for High-throughput Sterilization of Arabidopsis Seeds. *Journal of visualized experiments : JoVE*(128), 56587. <https://doi.org/https://doi10.3791/56587>
- Maguire, J. D. (1962). Speed of Germination—Aid In Selection And Evaluation for Seedling Emergence And Vigor 1. *Crop Science*, 2(2), 176-177. <https://doi.org/http://dx.doi.org/10.2135/cropsci1962.0011183X000200020033>
- Mahmood, A. H., Florentine, S. K., Chauhan, B. S., McLaren, D. A., Palmer, G. C., & Wright, W. (2016). Influence of Various Environmental Factors on Seed Germination and Seedling Emergence of a Noxious Environmental Weed: Green Galenia (*Galenia pubescens*). *Weed Science*, 64(3), 486-494. <https://doi.org/https://doi10.1614/Ws-D-15-00184.1>
- Mano, Y., & Takeda, K. (2001). Genetic resources of salt tolerance at germination and the seedling stage in wheat [*Triticum aestivum*]. *Japanese Journal of Crop Science*, 70(2), 215-220. <https://doi.org/https://doi10.1626/jcs.70.215>
- Mickky, B. M., & Aldesuquy, H. S. (2017). Impact of osmotic stress on seedling growth observations, membrane characteristics and antioxidant defense system of different wheat genotypes. *Egyptian Journal of Basic and Applied Sciences*, 4(1), 47-54. <https://doi.org/https://doi.org/10.1016/j.ejbas.2016.10.001>
- Mondal, A., & Pramanik, S. (1995). In vitro studies of salt tolerance of three rice races in Bay Islands. *J Indian Soc Coastal Agric Res*, 13(2), 127-131. <https://doi.org/http://www.iscar.org.in/urls/publication>
- Morales, A. J., Bajgain, P., Garver, Z., Maughan, P. J., & Udall, J. A. (2011). Physiological responses of *Chenopodium quinoa* to salt stress. *International Journal of Plant Physiology and Biochemistry*, 3(13), 219-232. <https://doi.org/https://doi:10.5897/IJPPB11.026>
- Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. *Annu Rev Plant Biol*, 59, 651-681. <https://doi.org/https://doi.10.1146/annurev.arplant.59.032607.092911>
- Murashige, T., & Skoog, F. (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia plantarum*, 15(3), 473-497.
- Nasir, N., Qureshi, R., & Aslam, M. (2000). A quick screening technique for salt tolerance in sugarcane (*Saccharum officinarum* L.). *Pakistan Sugar Journal*, 15(1), 2-7.
- Panta, S., Flowers, T., Lane, P., Doyle, R., Haros, G., & Shabala, S. (2014). Halophyte agriculture: Success stories. *Environmental and Experimental Botany*, 107, 71-83.

- <https://doi.org/https://doi10.1016/j.envexpbot.2014.05.006>
- Panuccio, M. R., Jacobsen, S. E., Akhtar, S. S., & Muscolo, A. (2014). Effect of saline water on seed germination and early seedling growth of the halophyte quinoa. *AoB plants*, 6, plu047. <https://doi.org/https://doi10.1093/aobpla/plu047>
- Postnikova, O. A., Shao, J., & Nemchinov, L. G. (2013). Analysis of the Alfalfa Root Transcriptome in Response to Salinity Stress. *Plant and Cell Physiology*, 54(7), 1041-1055. <https://doi.org/https://doi10.1093/pcp/pct056>
- Prado, F. E., Boero, C., Gallardo, M., & Gonzalez, J. A. (2000). Effect of NaCl on germination, growth, and soluble sugar content in *Chenopodium quinoa* Willd. seeds. *Botanical Bulletin of Academia Sinica*, 41. <https://doi.org/https://ejournal.sinica.edu.tw/bbas/content/2000/1/bot11-05.pdf>
- Ranal, M. A., Santana, D. G. d., Ferreira, W. R., & Mendes-Rodrigues, C. (2009). Calculating germination measurements and organizing spreadsheets. *Brazilian Journal of Botany*, 32(4), 849-855. <https://doi.org/http://www.scielo.br/pdf/rbb/v32n4/a22v32n4.pdf>
- Razzaghi, F., Jacobsen, S.-E., Jensen, C. R., & Andersen, M. N. (2015). Ionic and photosynthetic homeostasis in quinoa challenged by salinity and drought—mechanisms of tolerance. *Functional Plant Biology*, 42(2), 136-148. <https://doi.org/https://doi.org/10.1071/FP14132>
- Ruiz-Carrasco, K., Antognoni, F., Coulibaly, A. K., Lizardi, S., Covarrubias, A., Martínez, E. A., Molina-Montenegro, M. A., Biondi, S., & Zurita-Silva, A. (2011). Variation in salinity tolerance of four lowland genotypes of quinoa (*Chenopodium quinoa* Willd.) as assessed by growth, physiological traits, and sodium transporter gene expression. *Plant Physiology and Biochemistry*, 49(11), 1333-1341. <https://doi.org/https://doi.org/10.1016/j.plaphy.2011.08.005>
- Ruiz, K., Rapparini, F., Bertazza, G., Silva, H., Torrigiani, P., & Biondi, S. (2017). Comparing salt-induced responses at the transcript level in a salares and coastal-lowlands landrace of quinoa (*Chenopodium quinoa* Willd.). 139, 127/142. <https://doi.org/http://dx.doi.org/10.1016/j.envexpbot.2017.05.003>
- Ruiz, K. B., Maldonado, J., Biondi, S., & Silva, H. (2019). RNA-seq Analysis of Salt-Stressed Versus Non Salt-Stressed Transcriptomes of *Chenopodium quinoa* Landrace R49. *Genes*, 10(12), 1042. <https://doi.org/https://doi:10.3390/genes10121042>
- Sairam, R. K., Rao, K. V., & Srivastava, G. C. (2002). Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration. *Plant Science*, 163(5), 1037-1046. [https://doi.org/https://doi10.1016/s0168-9452\(02\)00278-9](https://doi.org/https://doi10.1016/s0168-9452(02)00278-9)
- Sairam, R. K., & Srivastava, G. C. (2002). Changes in antioxidant activity in sub-cellular fractions of tolerant and susceptible wheat genotypes in response to long term salt stress. *Plant Science*, 162(6), 897-904. [https://doi.org/https://doi10.1016/s0168-9452\(02\)00037-7](https://doi.org/https://doi10.1016/s0168-9452(02)00037-7)
- Sanchez, H. B., Lemeur, R., Damme, P. V., & Jacobsen, S. E. (2003). Ecophysiological Analysis Of Drought And Salinity Stress Of Quinoa (*Chenopodium quinoa* Willd.). *Food Reviews International*, 19(1-2), 111-119. <https://doi.org/https://doi10.1081/FRI-120018874>
- Sanchez, P. L., Chen, M.-k., Pessarakli, M., Hill, H. J., Gore, M. A., & Jenks, M. A. (2014). Effects of temperature and salinity on germination of non-pelleted and pelleted guayule (*Parthenium argentatum* A. Gray) seeds. *Industrial crops and products*, 55, 90-96. <https://doi.org/https://doi10.1016/j.indcrop.2014.01.050>
- Scott, S., Jones, R., & Williams, W. (1984). Review of Data Analysis Methods for Seed Germination 1. *Crop Science*, 24(6), 1192-1199. <https://doi.org/https://doi.org/10.2135/cropsci1984.0011183X002400060043x>
- Shabala, S., & Mackay, A. (2011). Ion transport in halophytes. *Advances in botanical research*, 57, 151-199. <https://doi.org/https://doi10.1016/B978-0-12-387692-8.00005-9>
- Shahid, M., Balal, R., Pervez, M., Abbas, T., Ashfaq, M., Ghazanfar, U., Afzal, M., Rashid, A., Garcia-Sanchez, F., & Mattson, N. (2012). Differential response of pea (*Pisum sativum* L.) Genotypes to salt stress in relation to the growth, physiological attributes antioxidant activity and organic solutes. *Australian Journal of Crop Science*, 6(5), 828. https://doi.org/www.cropj.com/shahid_6_5_2012_828_838.pdf
- Shi, Y. Y., Gao, L. L., Wu, Z. C., Zhang, X. J., Wang, M. M., Zhang, C. S., Zhang, F., Zhou, Y. L., & Li, Z. K. (2017). Genome-wide association study of salt tolerance at the seed germination stage in rice. *BMC plant biology*, 17(1), 92. <https://doi.org/https://doi10.1186/s12870-017-1044-0>
- Tanveer, A., Mumtaz, K., Javaid, M. M., Chaudhry, M. N., Balal, R. M., & Khaliq, A. (2013). Effect of ecological factors on germination of horse purslane (*Trianthema portulacastrum*). *Planta Daninha*, 31(3), 587-597. <https://doi.org/https://doi10.1590/s0100-83582013000300011>
- Wang, W.-B., Kim, Y.-H., Lee, H.-S., Kim, K.-Y., Deng, X.-P., & Kwak, S.-S. (2009). Analysis of antioxidant enzyme activity during germination of alfalfa under salt and drought stresses. *Plant Physiology and Biochemistry*, 47(7), 570-577. <https://doi.org/https://doi.org/10.1016/j.plaphy.2009.02.009>
- Wang, Z. F., Wang, J. F., Bao, Y. M., Wu, Y. Y., Xuan, S. U., & Zhang, H. S. (2010). Inheritance of rice seed

- germination ability under salt stress. *Rice Science*, 17(2), 105-110. [https://doi.org/https://doi.org/10.1016/S1672-6308\(08\)60112-2](https://doi.org/https://doi.org/10.1016/S1672-6308(08)60112-2)
- Wang, Z. F., Wang, J. F., Bao, Y. M., Wu, Y. Y., & Zhang, H. S. (2011). Quantitative trait loci controlling rice seed germination under salt stress. *Euphytica*, 178(3), 297-307. <https://doi.org/https://doi10.1007/s10681-010-0287-8>
- Yadav, P. V., Kumari, M., & Ahmed, Z. (2011). Seed priming mediated germination improvement and tolerance to subsequent exposure to cold and salt stress in capsicum. *Research Journal of Seed Science*, 4(3), 125-136. <https://doi.org/https://DOI:10.3923/rjss.2011.125.136>

UNDER PEER REVIEW