

# A COMPARATIVE STUDY OF PUBERTAL DEVELOPMENT IN HIV-INFECTED AND NON-HIV-INFECTED BOYS IN PORT HARCOURT NIGERIA

## Abstract

**Background:** Pubertal abnormalities can occur in HIV-infected boys and have negative effects on reproductive health, final adult height as well as psychosocial health and quality of life. This study was aimed at comparing the pubertal development in HIV-infected and non-HIV-infected boys attending two tertiary hospitals in Port Harcourt.

**Materials and methods:** A Comparative cross sectional study was carried out involving 80 HIV-infected boys aged 10-18 years and 80 non-HIV-infected boy who were matched for age, sex and socioeconomic class.

**Results:** The mean age of the HIV-infected boys was  $13.22 \pm 2.31$  while that of the non-HIV-infected boys was  $13.24 \pm 2.30$  and this difference was not statistically significant. On physical evaluation using the Tanner staging method, only 64 (80%) HIV-infected boys had attained puberty as against 76 (95%) in the non-HIV-infected group ( $\chi^2 = 13.091$ ,  $p\text{-value} = 0.011$ ). Pubertal onset occurred at a significantly later age in HIV-infected boys when compared to the non-HIV-infected ones ( $11.76 \pm 1.33$  years vs  $10.81 \pm 0.77$  years). Pubertal delay was seen in 2 (2.5%) and this was significantly more common among those who were underweight as well as those in WHO clinical stages 3 and 4.

**Conclusion:** Pubertal onset occurred at a significantly later age in HIV-infected boys when compared to the non-HIV-infected boys. There is need to monitor pubertal development in HIV-infected boys so that deviations from normal can be promptly identified and necessary interventions given.

**Keywords:** Puberty, HIV-infected-boys, Port Harcourt.

## 1. INTRODUCTION

Puberty is a complex developmental process which begins in late childhood and is triggered by maturation of the hypothalamic-pituitary-gonadal axis [1]. Pulsatile activity of the neuronal network of the gonadotropin releasing hormone leads to a surge in gonadotropin secretion leading to the appearance of secondary sexual characteristics, growth spurt and ultimately the capacity for fertility and reproduction [2-4].

The commonest clinical system for describing normal pubertal development and its variation is the five-stage system developed by Tanner in 1969 [5]. In boys, the onset of puberty is marked by testicular enlargement followed by the appearance of pubic hair, enlargement of the penis and first seminal emission (spermarche) [5].

Disorders of pubertal development may occur at any stage of the maturational process leading to either precocious or delayed puberty [4]. Pubertal development is influenced by genetic factors, nutrition, environmental factors and chronic diseases [1, 6].

Human Immunodeficiency Virus (HIV) / Acquired Immunodeficiency Syndrome (AIDS) and the attendant chronic immunosuppression have been shown to affect the pubertal development of children [7-10]. In HIV-infected children, pubertal delay which is defined as the absence of secondary sexual characteristics at an age  $\geq 2$  standard deviation above the mean for age and sex is the most common pubertal abnormality seen [11, 12]. In boys, puberty is considered delayed if testicular enlargement does not occur by 14 years of age, there is lack of pubic hair by 15 years or more than 5 years are required to complete genital development [1, 13].

The mechanisms of pubertal delay in HIV-infected children are largely unknown. However, mechanisms postulated include nutritional deficiencies, recurrent infections and chronic immune activation which may all distort the endocrine axes [8].

Similarly, precocious puberty has also been reported in HIV-infected boys [11]. This is defined as the appearance of secondary sexual characteristics before the age of nine years in boys [11].

With a large number of HIV-infected children now surviving into adolescence as a result of advancements in the management of HIV and AIDS, pubertal developmental disorders may pose an increasing challenge [14].

Unidentified pubertal disorder may be associated with risk for psychosocial problems including low self-esteem and depression and may also have negative implications for reproductive health and final adult height [10,15-16]. This thus underscores the need to study pubertal development in HIV-infected children.

When identified early, adolescents with pubertal delay can benefit from interventions such as psychological support, nutritional support and possibly low dose of gonadal steroid hormones or growth hormone in addition to Highly Active Antiretroviral therapy (HAART) [8,9,17,18]. Similarly, HIV-infected children with precocious puberty may benefit from psychological support in addition to HAART to help limit the impact of chronic immunosuppression on the endocrine axis. In some cases, gonadotropin releasing hormone agonists such as leuprolide may help restore final adult height [8,19]

This study was aimed at evaluating the pattern of pubertal development among HIV-infected boys in comparison to non-HIV-infected boys in Port Harcourt, Nigeria.

## **2. MATERIALS AND METHODS**

### **2.1 Study Design**

A comparative cross sectional study design was used. The study was carried out over a four month period (July 1<sup>st</sup> –October 31<sup>st</sup> 2017).

### **2.2 Study Area**

The study location was the Paediatric departments of the University of Port Harcourt Teaching Hospital (UPTH) and the Rivers State University Teaching Hospital (RSUTH). The two hospitals are both tertiary care facilities located in the city of Port Harcourt and providing Antiretroviral (ARV) care for HIV-infected.

### **2.3 Study Population**

This study involved 80 boys age 10-18 years with confirmed HIV infection and on follow up at the Paediatric ARV clinic. Similarly, 80 non-HIV-infected boys were recruited from the children's outpatient clinic and matched for age, sex and socioeconomic class to serve as a comparison group. HIV-infected boys who had other chronic disease conditions like sickle cell anaemia which can also affect pubertal development were excluded from the study.

### **2.4 Study Procedure**

HIV-infected boys receiving follow up care at the paediatric ARV clinic were consecutively recruited over a four month period. For each HIV-infected subject, a comparison matched for age, sex and socioeconomic class was recruited. HIV infection was excluded through rapid diagnostic testing before enrolment of the comparison group. History and physical examination were used to exclude the presence of other chronic diseases that can affect pubertal development such as chronic kidney diseases and bronchial asthma.

Relevant sociodemographic data and information on age of attainment of pubertal indices were obtained using a semi-structured interviewer administered questionnaire. This was followed by physical examination of the boys to determine the pubertal stage status using the Tanner staging system for testicular and pubic hair development. Testicular volume was determined using an Orchidometer.

### **2.5 Statistical Analysis**

Data analysis was done using SPSS version 20.0. Results were presented as tables and charts in simple proportions. Student t-test was used for the comparison of means while Chi square and Fisher exact test was used to test for associations between categorical variables. Statistical significance was at 95% confidence interval with p-value < 0.05.

### 3 Results

The study participants consisted of 80 HIV-infected boys and 80 non-HIV-infected boys. The mean age of HIV-infected boys was  $13.33 \pm 2.42$  years while that of non-HIV-infected boys was  $13.34 \pm 2.43$  years ( $t=0.0261$ ,  $p=0.9792$ ).

Using both recall of pubertal events and physical examination method (Tanner staging), pubertal onset occurred at significantly later ages in HIV-infected boys compared to the non-HIV-infected ones (Tables 1 and 2). Other pubertal events such as pubic hair development, axillary hair development and spermarche also occurred at significantly later ages in the HIV-infected group (Table 1).

Table 3 shows that a lower proportion of HIV-infected boys had attained puberty (Tanner stage 2 and above for genital development) in comparison to the non-HIV-infected boys ( $\chi^2=13.091$ ,  $p\text{-value}=0.011$ ). There was however no significant difference between the two groups in terms of the final pubertal outcome (Table 4).

**Table 1: Comparison of Mean Age at onset of Pubertal Events (by recall) in Male Study**

**Subjects.**

---

**Males**

---

<b>Pubertal event recall</b>	<b>HIV-infected</b>	<b>Non HIV-infected</b>
------------------------------	---------------------	-------------------------

---

	Mean age ± S.D (years)	Mean age ± S.D (years)	t	p-value
Genitals	12.04 ± 1.05	11.46 ± 0.88	2.943	0.004*
Pubic hair	12.04 ± 1.04	11.40 ± 1.25	2.767	0.007*
Axillary hair	12.42 ± 1.08	11.89 ± 0.90	2.403	0.019*
Spermarche	14.80 ± 0.41	13.86 ± 0.71	5.162	0.0001*

\*Statistically significant  
S.D-Standard Deviation

**Table 2: Comparison of Mean Ages of Male Study Subjects at various Tanner Stages**

Tanner stages	Males		t	p-value
	HIV-infected Mean age ± S.D (years)	Non HIV-infected Mean age ± S.D (years)		
<b>Genitals</b>				
G1	11.46 ± 1.61	10.80 ± 1.02	0.769	0.452
G2	11.76 ± 1.33	10.81 ± 0.77	2.558	0.015*
G3	13.14 ± 2.04	12.73 ± 1.72	0.699	0.489
G4	15.14 ± 1.72	14.13 ± 1.71	1.679	0.103

G5	16.94 ± 0.68	16.30 ± 1.45	1.318	0.199
<b>Pubic hair</b>				
PH1	11.66 ± 1.50	10.59 ± 0.63	1.799	0.086
PH2	11.87 ± 1.63	11.40 ± 1.28	1.057	0.296
PH3	13.50 ± 2.00	12.87 ± 2.03	0.782	0.442
PH4	14.04 ± 1.79	13.38 ± 0.66	1.155	0.262
PH5	16.52 ± 1.07	16.21 ± 1.32	0.806	0.425
*Statistically significant ;      S.D-Standard deviation;      G-Genitals;      PH-pubic hair				

**Table 3: Comparison of Tanner Stages among Male Study Subjects**

Tanner stages	Males		Total N=160(%)
	HIV-infected n=80(%)	Non HIV-infected n=80(%)	
<b>Genitals</b>			
G1	16 (20.0)	4 (5.0)	20 (12.5)
G2	20 (25.0)	16 (20.0)	36 (22.5)
G3	16 (20.00)	26 (32.5)	42 (26.3)
G4	18 (22.5)	15 (18.7)	33 (20.6)
G5	10 (12.5)	19 (23.8)	29 (18.1)

*Chi Square=13.091,p-value=0.011\**

**Pubic hair**

PH1	16 (20.0)	7 (8.8)	23 (14.4)
PH2	23 (28.8)	21 (26.2)	44 (27.5)
PH3	10 (12.5)	17 (21.2)	27 (16.9)
PH4	13 (16.2)	11 (13.8)	24 (15.0)
PH5	18 (22.5)	24 (30.0)	42 (26.2)

*Chi Square=6.451,p-value=0.168*

---

\*Statistically significant      G-Genitals;      PH-pubic hair

**Table 4: Comparison of Pubertal Stage Status in Male Study Subjects**

Pubertal Stage Status	Males		Total N=160(%)
	HIV-infected n=80(%)	Non-HIV-infected n=80(%)	
<b>Recall</b>			
Normal	73 (91.2)	75 (93.8)	148 (92.5)
Delayed	4 (5.0)	0 (0.0)	4 (2.5)
Precocious	1 (1.3)	2 (2.5)	3 (1.9)
Can't remember	2 (2.5)	3 (3.7)	5(3.1)
<i>Fishers exact= 3.955; p-value = 0.358</i>			
<b>Examination</b>			
Normal	78 (97.5)	80 (100.0)	158 (98.8)
Delayed	2 (2.5)	0 (0.0)	2 (1.2)
<i>Fisher's exact p-value=0.497</i>			

---

#### 4. Discussion

Our study found the mean age of pubertal development in HIV-infected boys using both the recall and physical examination of Tanner stages to be significantly higher in comparison to non-HIV-infected boys. This is understandable as HIV infection and the attendant immunosuppression is known to exert a negative effect on the pubertal development process. Similar findings have also been reported in some previous studies [9,20,21].

The mean age of pubertal onset in HIV-infected boys in the index study was 11.76 years. This compares favorably with that reported by Buchacz *et al*, [9] and Williams *et al*, [20] both in the USA. It is however lower than the 13.5 years reported by Mbwire, [7] in Tanzania and 12.8 years reported by Szubert *et al*, [8] in Uganda and Zimbabwe. Other factors influencing pubertal development such as nutrition and environmental exposures may have contributed to the variations in the mean age of pubertal onset among the various studies in comparison.

The mean age of completion of pubertal development as evidenced by attainment of Tanner stage 5 for genital and pubic hair development was higher in HIV-infected boys compared to non-HIV-infected ones. This difference was however not statistically significant. A significantly lower proportion of HIV-infected boys than non-HIV-infected ones were found in the more advanced pubertal stages (Tanner stages 3-5). Similar findings were reported in previous studies [9,22]. This may be explained by the deleterious effects of HIV infection on the hypothalamic-pituitary-gonadal axis leading to delays in progression through the pubertal stages.

With regard to pubertal stage status, majority of the children studied had normal pubertal stage status. Pubertal delay was however found in 5% of the HIV-infected boys and none in

the non-HIV-infected. The higher prevalence of pubertal delay seen in HIV-infected boys is similar to that reported by Shadabet *al*, [23] in India. The proportion of HIV-infected boys with pubertal delay in our study also compares favorably with that reported in the Indian study. In contrast, Mbono et al, [24] reported normal pubertal development among all the HIV-infected boys evaluated in a study done in Cameroun.

One HIV-infected child reported precocious puberty in the index study. This is an uncommon finding and the explanation for this is unknown. However, Horner and Bhumbra, [11] have also reported a case of precocious puberty diagnosed in a perinatally HIV-infected boy in the USA. It would be important to exclude other underlying causes of precocious puberty in such children.

## **Conclusion**

Pubertal onset occurred at a significantly later age in HIV-infected boys when compared to the non-HIV-infected boys. There is need to monitor pubertal development in HIV-infected boys so as to identify deviations from normal early and evaluate need for intervention.

## **Consent**

Written informed consent was obtained from all the parents and caregivers of the study participants before enrolment into the study.

## **Ethical Approval**

Ethical approval was obtained from the Research Ethics Committee of the University of Port Harcourt Teaching Hospital.

## **5. References**

1. Zacharin M, Banerjee I, Patel L. Puberty: normal and abnormal. In: Zacharin M (ed.) Practical Paediatric Endocrinology in a Limited Resource Setting. 1<sup>st</sup> ed. Melbourne: Merck Serono;2011.26-55.

2. Herbison AE. Control of puberty onset and fertility by gonadotropin-releasing hormone neurons. *Nat Rev Endocrinol* 2016; 12(8):452-66.
3. Plant T. The mystery of puberty. Presented at the society for endocrinology BES 2017. Harrowgate, UK. *Endocrine Abstracts* 50 SE1.3.
4. Kakarla N, Bradshaw K. Disorders of Pubertal Development: precocious puberty. *Semin Reprod Med* 2003;21:339-51.
5. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. *Arch Dis Child* 1970;45:13-23.
6. Loomba-Albrecht LA, Styne DM. The physiology of puberty and its disorders. *Pediatr Ann* 2012;41:1-9.
7. Mbwile GR. Growth and Pubertal Development among HIV-infected Children aged 8-18 years in Dar es Salaam. A Dissertation submitted in Partial Fulfillment of the Requirement for the award of Degree of Master of Medicine (Paediatrics and Child Health) of the Muhimbili University of Health and Allied Sciences. Muhimbili University of Health and Allied Sciences Dar es Salaam; 2012.
8. Szubert AJ, Musiime V, Bwakura-Dangarembizi A, Nahirya-Ntege P, Kekitinwa A, Gibb DM, *et al.* Pubertal Development in HIV-infected African Children on First-line Antiretroviral therapy. *AIDS* 2015;29:609-18.
9. Buchacz K, Cervia JS, Lindsey JC, Hughes MD, Seage GR, Dankner WM, *et al.* Impact of Protease Inhibitor-containing combination antiretroviral therapies in HIV-infected children. *Pediatrics* 2001;108:72-9.
10. Majaliwa ES, Mohn A, Chiarelli F. Growth and Puberty in Children with HIV Infection. *J Endocrinol Invest* 2009;32:85-90.
11. Horner JM, Bhumbra N. Congenital HIV infection and precocious puberty. *J Pediatr Endocrinol Metab* 2003;16:791-793.
12. Pozo J, Argente J. Delayed Puberty in Chronic illness. *Best Pract Res Clin Endocrinol Metab* 2002;16:73-90.

13. Rochelle N. Puberty: Normal and Delayed Puberty. Paediatrics clinical curriculum. Available at [www.uchicago.com](http://www.uchicago.com) accessed on 2<sup>nd</sup> November 2023.
14. Gortmaker SL, Hughes M, Cervia J, Brady M, Johnson GM, Seage GR, *et al.* Effects of Combination Therapy including Protease inhibitors on Mortality among Children and Adolescents with HIV-1. *N Engl J Med* 2001;345:1522-8.
15. Hirschfield S. Dysregulation of growth and development in HIV-infected children. *J Nutr* 1996;126:2641-50.
16. Mckillop KA. An exploration of the psychosocial consequences of delayed puberty in children who attend the Royal Hospital for sick children Endocrine clinic : A qualitative study and clinical research portfolio. D ClinPsy thesis. University of Glasgow; 2013.
17. Kaufman FR, Gertner JM, Sleeper LA, Donfield SM. Growth hormone secretion in HIV- positive versus HIV-negative hemophiliac males with abnormal growth and pubertal development: the Hemophilia Growth and Development study. *J Acquir Immune Defic Syndr Hum Retrovirol* 1997;15:137-44.
18. Neinstein LS, Kaufman FR. Abnormal physical growth and development. In: Neinstein LS (ed.) *Adolescent healthcare, a practical guide*. 3<sup>rd</sup> ed. Baltimore: Williams and Wilkins; 1996.165-93.
19. Carel JC, Lahlou N, Roger M, Chaussain JL. Precocious puberty and statural growth. *Hum Reprod* 2004;10:135-47.
20. Williams PL, Abzug MJ, Jacobson DL, Wang J, Van Dyke RB, Hazra R, *et al.* Pubertal onset in HIV-infected children in the era of combination antiretroviral treatment. *AIDS* 2013;27:1959-70.
21. Ndiokwelu CO, Uwaezuoke SN, Iloh KK. Physical growth and sexual maturation of perinatally HIV-infected adolescent males in a southeast Nigerian tertiary hospital: a comparative cross-sectional study. *BMC Pediatr* 2022;22:573.
22. Gertner JM, Kaufman FR, Donfield SM, Sleeper LA, Shapino AD, Howard C, *et al.* Delayed somatic growth and pubertal development in human immunodeficiency virus-

infected hemophiliac boys: Hemophilia growth and development study. J Pediatr 1994;124:896-902.

23. Shadab R, Seth A, Gupta RA. Growth and Pubertal development in adolescents infected with HIV. Presented at the 20<sup>th</sup> international AIDS Conference. Melbourne, 2014.

24. Mbono RC, Ngo Um SS, Edongue M, Ndombo P. Does HIV infection affect growth and puberty of Cameroonian children? *Acchive de Pediatrie* 2021;28(3):238-41.

UNDER PEER REVIEW