

## Review Article

### **An Overview of Leaf Litter Decomposition and Nutrient Dynamics of Multipurpose Tree Species**

#### **Abstract:**

The multipurpose tree species is a plant species that is purposefully grown so as to provide two or more than two products and their leaf litter decomposition in our ecosystems has a play major role in nutrient dynamics in the atmosphere. Abiological factors such as temperature, rainfall, humidity, moisture, evaporation rate and other seasonal variations affect the rate of leaf litter decomposition. From the this review paper attempt to understand the influence of various soil physical factors like-soil pH, texture, organic matter, chemical factor of leaf litter content like cellulose, hemicelluloses, tannin, lignin and biological factor like major soil fauna and microbes such as algae, actinomycetes, bacteria and fungi communities etc. to stimulating microbial action such as physical breakdown of leaf litter, transfer of organic matter to nutrients, release of CO<sub>2</sub> to the environment etc.

**Keyword:** Nutrient dynamics, leaf litter, multipurpose tree species, macro-invertebrates.

#### **Introduction:**

The dead plant materials such as leaves, bark, needles, flower and twigs that have fallen on the ground are called litter. As the majority of organic matter (OM) produced by plants is returned to the soil as litter, the transfer of nutrients and energy from living biological components to the soil is the important mechanism of nutrient recycling. In the various components of the plant-soil-microorganism mechanism, decomposition processes have a crucial vital role by releasing a complex organic compound into the simple usable form for plants growth and soil fertility. Tree leaf tissue accounted for more than 70% of above ground litter fall in the ecosystems and rest is composed stem, twigs, flower, fruits and other components. The decomposition is a biological disintegration process of dead organic materials whereby mineralization of complex organic matter into simple inorganic form (1). Tree leaf litter decomposition has physical, chemical, and

biological impact and is an important biological process that regulates the complex nutrient content to simple form and net productivity of ecosystems (2). Litter decomposition also affects the soil moisture status, light, and temperature conditions required for the growth of surrounding plants (3), and it impacts the microbial community structure, soil physiochemical properties, and organic matter escapism and content of soil sediments. Generally correlation of main three factors ~~are viz.~~ Climatic condition of site, leaf litter quality and soil biota. Out of ~~it~~ ~~these~~, climate ~~plays~~ dominant crucial role followed by leaf litter quality and soil biota as well as release of nutrient ~~are~~ directly correlated to decomposition process (4). Therefore, the main aim of this review paper is to summarize the role of microbes for degradation of leaf litter as well as correlation of leaf litter quality, climatic conditions, soil biota and importance of nutrient dynamics process.

### **Studies of leaf litter decomposition process:**

Leaf litter decomposition plays a very vital and important role for cycling of nutrient in ecosystems, where different types of flora on the basis of quality ~~influence~~ most significantly. The interaction among soil microbes, climatic conditions and quality of leaf litter will result in the decomposition of litter, in which soil microbes fungi and bacteria ~~is~~ ~~are~~ the primary decomposers of leaf litter and broken down into smaller ~~pieces~~ ~~particles~~ and modified mineralized form into inorganic ~~compounds~~, because microbes feed ~~on~~ the organic remains of dead plants materials (5,6). After that the following major processes take place.

1. During the decomposition process soluble materials are moved to lower soil layer for further process.
2. Soil ~~microbes~~ breakdown the ~~larger~~ ~~of~~ ~~pieces~~ of leaf litter into smaller ones.
3. Decomposer recognizes the molecules of litter to change the chemical alteration in soil.

During the decomposition process time, ~~S~~soil fauna, microorganism play a very important role to break the complex compound of leaf litter and change the chemical composition of leaf litter structure and soluble ~~compounds~~ (7,5). ~~Atmospheric~~ features, like temperature, rainfall, relative humidity and other seasonal variations may impact the existence of soil microbes and other soil features such as pH, organic carbon, nutrient availability, soil texture

and structure that significantly affect the rate of litter decomposition as well as leaf litter quality like litter structure, roughness, smoothness, waxes **structure**, etc. directly influence the activity of soil communities and process during the decomposition (8). A graphical representation of litter degradation is shown in Fig. 1.




Fig.1 Graphical representation of factors affecting litter degradations.

#### Factors affecting leaf litter decomposition:

Leaf litter decomposition depends on various biotic such as chemical composition of litter and soil **microorganisms** and abiotic factors like temperature, relative humidity, and rainfall. Both **factors** directly influenced on mineralization and humification process of structural complex of litter such as lignin, cellulose, hemicelluloses, pectine, **starch**, etc. through microorganisms and gradually mineralization of soluble **nutrients** compounds into soil by leaching process. However, from the above discussion we **can concluded** that the main three leading features are regulating the litter decomposition like leaf litter quality traits, composition of soil microorganism activities and physio-chemical environments. **ete.** (9). In abiotic **process**, factor such as temperature is a prime factor in determining the rate of leaf litter decomposition and more sensitive **compared to others** because abiotic factor like temperature **influences** soil temperature and soil temperature

directly ~~influences~~ soil microbial activity rises exponentially. The fresh and old leaf litter position is a readily available substrate for soil microorganisms because leaf litter quality directly ~~influences~~ the decomposition process, as it reduces the throughout litter decomposition due to the loss of readily accessible carbon and accumulation of recalcitrant compounds ~~according to~~ (10), ~~also~~ shown in Fig.1.

### **Role of trees and leaf litter quality:**

Tree leaf litter contains have different organic compounds, mainly four important soluble organic material present in litter such as phenolics, hydrocarbons, sugars and glycerides, ~~in~~ which soluble sugar, oligosaccharides and primarily monosaccharide are difficult to metabolize. The above proportions of all compounds ~~varyare-varies~~ with the plant part such as leaves, stems, roots and bark as well as different tree species. The tree leaf litter ~~quality~~ chemical composition like nitrogen, phosphorus, potassium and cell wall components like lignin, cellulose and hemicelluloses etc, directly influence the litter decomposition and nutrient release (5,11). The components account ~~of-for~~ chief cell wall like lignin content 15 to 40% of total litter quantity. ~~It~~ may be in extreme cases low as 4% or high as 50%. ~~As-well-as-in~~ In addition to lignin, other compound like cellulose, hemicelluloses, ~~and~~ pectin are common constituents in tree leaf litter in term of quantity. The ~~account~~ amount of cellulose (10 to 50%) of litter quantity ~~are is~~ linked with  $\beta$ -1-4 bonds and hemicelluloses are polymers of sugar, ~~which~~ create long chains of molecules organized into fibres (12). ~~The-a~~ All the components such as litter traits-leaf toughness, N, lignin, polyphenol, C/N ratio, lignin: nitrogen ratio etc. directly and indirectly influence the decomposition process and various traits, nitrogen and lignin content of plant litter significantly, regulating the decomposition rate (13,14).

### **Role of soil properties and soil fauna:**

Different soil physical and chemical characteristics like soil texture, soil pH, cation exchange capacity, organic matter, nutrients contents, bulk density, etc. have a significantly role in leaf litter decomposition process. In decomposition process, soil texture stimulates nutrients and water dynamics, porosity, permeability and soil surface area. ~~According to~~ (15) Soil organic matter quality and quantity also play important role to increasing the population density of soil macro-organism, which plays important role in jumble the leaf litter and decomposition rate (15,

16,17). Table 1 shows the significant role of major soil fauna and microbes such as algae, actinomycetes, bacteria and fungi communities, etc. to stimulating microbial action such as physical breakdown of leaf litter, transfer of organic matter to nutrients, release of CO<sub>2</sub> to the environment, etc. (18,19,20,21). Some soil micro-arthropods includes small invertebrate animals with an exoskeleton and segmented body found on or near the surface of soil and in plant litter. Their population range is between 10<sup>3</sup> m<sup>-2</sup> in agriculture soils and 10<sup>6</sup> m<sup>-2</sup> in forest soils. It is a very essential part of the ecosystems due to their significant role in organic compound decomposition process and mineralization activity, nutrient cycling and pedogenesis (22). The labile structural compound like cellulose, sugar, amino acid quickly cleaved and absorbed by exo-enzymes and rapid decay (23).

### Role of atmosphere factors:

From the studies of different literature we concluded that the atmosphere factors such as temperature, moisture, humidity, rainfall directly influence the chemical component of soil structure and affect the leaf litter decomposition rate in the soil (24). It is also reported that the rate of decomposition of leaf litter is slow in winter season and faster in rainy season, due to the presence of sufficient rainfall, soil moisture and microbial load (25).

**Table 1: Role of soil fauna and microbes in leaf litter decomposition.**

S.no.	Decomposer communities	Roles	References
1.	Soil fauna	Microalgae fix nitrogen and produce organic matter via photosynthesis	(17)
		Increase the surface area of substrate for microbial use	(20)
		Releases soil enzymes, which can help to process root-driven carbon, small organic matter, and fresh aboveground litter, as energy source for bacteria (for example, fungi)	(16,21)
		Augment the nutrient in soil by adding nitrogenous compounds present in their excreta and dead tissue	(21)
2.	Soil microbes	Decompose the fragmented litter and release nutrients	(19)
		Release soil enzymes for the purpose of breaking the larger compounds	(18)

### **Leaf litter decomposition assessing techniques:**

For determination of leaf litter decomposition and decay rate, various techniques are used by the authors like mass balance technique, litter bag technique, Tethered leaves technique and Cohort layered screen technique, etc. Mass balance technique is used to evaluate the leaf litter decay on the assumption of constant fraction,  $k$  of the detrital leaf litter quantity decays because mass balance technique depends on natural litter fall and it does not efficiently explain the role of atmosphere factors like temperature and moisture. In litter bag technique, mesh size of more than 2 mm and litter bag of 20x20 cm are common and efficient to permit of entry of macrofauna for contact of leaf litter and decay process (26). Another method of leaf litter decay evaluation is tethered leaves techniques. This technique is similar to the litter bag method. In this method, specific leaf litter is tightly packed with nylon thread or monofilament fishing line and permit the leaf intake by macro-invertebrates such as crabs and snails (27). A last Cohort layered screen technique is most appropriate and accurate evaluation of leaf litter decomposition and decay rate. It takes normally three to more years for leaf litter decay studies because there are separate techniques to determination of actual litter fall and decay condition through 1x1 m fiberglass or aluminum window screen with a mesh size of 2-3 mm are used (28).

### **Conclusion:**

Leaf litter decomposition of multipurpose tree species is highly significantly plays a role in the ecosystem, as it is a most important way of nutrient dynamics, specially C, N and other nutrients elements in the ecosystems. Different multipurpose tree leaf litter decomposition is a very high complex process that involves a number of soil physical factors like soil pH, soil texture, humification etc. chemical factors such as composition of leaf litter-cellulose, hemicelluloses, pectin, lignin, chitin, etc. and last and most important biological factors like atmosphere factor and soil fauna and microbes. However, from the study it is very difficult to understand that the which one is the most important factor for litter decomposition. All the physical, chemical and biological factors are inter-correlated to each other for appropriate leaf

litter decomposition and decay. From ~~the~~ this review paper ~~we~~ focuses on different factors that affect the leaf litter decomposition and decay as well as different techniques emphasized for assessing litter degradation.

## References:

1. Saha S, Rajwar G S, Kumar M, Upadhaya K. Litter production, decomposition and nutrient release of woody tree species in Dhanaulti region of temperate forest in Garhwal Himalaya. *Eurasian Journal of Forest Science*. 2016; 4 (1): 17–30.
2. Cornelissen JHC, Cornwell WK, Freschet G T, Weedon J T, Berg M P, Zanne, A E. Coevolutionary legacies for plant decomposition. *Trends Ecol. Evol.* 2023; 38, 44–54.
3. Zhang W P, Fornara D, Yang H, Yu R P, Callaway R M, Li L. Plant litter strengthens positive biodiversity–ecosystem functioning relationships over time. *Trends Ecol. Evol.* 2023 ;38, 473–484.
4. Bhalawe S, Kukadia M U, Nayak D. Nutrient release pattern of decomposed leaf litter in different multipurpose trees. *Indian Forester*, 2013; 139 (3): 212-213.
5. Bhalawe S, Nayak D, Kukadia M U, Gayakvad, P. Leaf litter decomposition pattern of trees. *The Bioscan*. 2013; 8(4):1135-140.
6. Geethanjali P A, Jayashankar M. A review on litter decomposition by soil fungal community. *Journal of Pharmacy and Biological Sciences*. 2016; 11(4) 01-03.
7. Krishna M P, Mohan M. Litter decomposition in forest ecosystems: a review. *EnergEcol Environ*. 2007; 2(4):236–49.
8. Chapman S K, Koch G W. What type of diversity yields synergy during mixed litter decomposition in a natural forest ecosystem. *Plant Soil*. 2007; 299 : 153– 162.
9. Dechaine J, Ruan H., Sanchez deLeon, Y, Zou X. . Correlation between earthworms and plant-litter decomposition in a tropical wet forest of Puerto Rico. *Pedobiologia*2005; 49 (6) : 601–607.
10. Dilly O, Munch J C. Shifts in physiological capabilities of the microbiota during the decomposition of leaf litter in black alder (*Alnus glutinosa* (Gaertn.) L.) forest. *Soil BiolBiochem*. 2001;33:921–930.

11. Swift M J, Heal O W, Anderson J M. Decomposition in terrestrial ecosystems. In: Anderson DJ, Greig-smith P, Pitelka FA (eds) *Studies in ecology*, vol 5. University of California Press, Berkeley. **1979**; pp 167–219.
12. Akpor B O, Okoh A I, Babalola G.O. Culturable microbial population during decomposition of *Cola nitida* leaf litters in a tropical soil setting. *J Biol Sci.* 2005; 18(3):313–319.
13. Perez-Harguindeguy N, Diaz S, Cornelissen J H C, Venramini F, Cabido M, Castellanos A. Chemistry and toughness predict leaf litter decomposition rates over a wide spectrum of functional types and taxa in central Argentina. *Plant Soil.* 2000; 218:21–30.
14. Cadish G, Giller K E. *Driven by nature: plant litter quality and decomposition.* CAB International, Wallingford. 1997; p 432.
15. Akpor B O, Okoh A I, Babalola G O. Culturable microbial population dynamics during decomposition of *Theobroma cacao* leaf litters in a tropical soil setting. *J Biol Sci.* 2006; 6(4):768–774.
16. Schinner F. Introduction. In, Schinner F, Ohlinger R, Kandeler E, Margesin R, editors. *Methods in soil biology.* Berlin: Springer-Verlag.1996; p. 3–6.
17. Crawford D L. Biodegradation of agricultural and rural wastes. In: Goodfellow M, Williams ST, Mordaski M, editors. *Actinomycetes in biotechnology.* London: Academic;. 1988; p. 433–9.
18. Brady N C, Wei RR. *The nature and properties of soils.* Pearson prentice hall, upper saddle river. 2010; NY.
19. Laganier J, Par D, Bradley R L. How does a tree species influence litter decomposition. Separating the relative contribution of litter quality, litter mixing, and forest floor conditions. *Can J For Res.* 2010; 40:465.
20. Gonzalez G, Ley R E, Schmidt S K., Zou X, Seastedt TR. Soil ecological interactions: comparison between tropical and subalpine forests. *Oecologia.* 2001; 128:549–56.
21. Gonzalez G, Zou X. Earthworm influence on N availability and the growth of *Cecropia schrebiana* in tropical pasture and forest soils. *Pedobiologia.* 1999; 43:824–9.
22. Imler U. Litter fall and nitrogen turnover in an Amazonian black water inundation forest. *Plant Soil.* 1982; 67:355–358.

23. Hobbie S E. Effects of plant species on nutrient cycling. *Trends EcolEvol.* 1996; 7:336–339.
24. Taylor B R, Parkinson D, Parsons W F J. Nitrogen and lignin content as predictors of litter decay rates: a microcosm test. *Ecology.* 1989; 70:97.
25. Kumar M, Joshi M, Todaria N P. Regeneration status of a sub-tropical *Anogeissus latifolia* forest in Garhwal Himalaya, India. *J For Res.* 2010; 21(4):439–44.
26. Robertson G P, Paul E A. Decomposition and soil organic matter dynamics. In: Sala OE, Jackson RB, Mooney HA, Howarth RW (eds) *Methods of ecosystem science.* Springer, New York. 1999; pp 104–116.
27. McKee K L, Faulkner P L. Restoration of biogeochemical function in mangrove forests. *Restor Ecol.* 2000;8 (3):247–259.
28. Binkley D. Ten year decomposition in a loblolly pine forest. *Can J For Res* 2002; 32(12):2231–2235.