

## Original Research Article

### **Effect of Foliar Application of Zinc on Zn-fortification and morph-physiological responses of rainfed wheat Yield**

#### **Abstract**

Wheat is a staple food in Pakistan and plays a vital role in the country's agriculture. It is one of the country's most widely cultivated and consumed crops, providing millions of people's food and livelihoods. The study reviewed the literature on the effect of foliar application of zinc on zinc-fortification and the yield of wheat grown under rainfed conditions. The review analyzed a range of studies that investigated the impact of the foliar application of zinc on wheat growth and grain quality. The results showed that foliar application of zinc significantly improves the zinc content of the wheat grain, which has important implications for addressing zinc deficiency in areas where it is prevalent. Additionally, the review revealed that zinc treatment leads to a noticeable increase in wheat yield, demonstrating its potential to improve agricultural productivity in rainfed farming systems. The review concludes by highlighting the need for further research to optimize the application rate and timing of foliar zinc application to maximize its benefits for wheat production under rainfed conditions.



**Key Words:** Zinc, Foliar application, rainfed, wheat, grain quality, yield.

## **INTRODUCTION**

Zinc is an important micronutrient for animals, human beings, as well as plants. Zn is significant element of different enzyme catalyzing many metabolic reactions in plants (Rudani et al., 2018). Furthermore, zinc is important for photosynthesis, protein synthesis, cell membrane integrity, pollen development, disease resistance, and boosting antioxidant enzymes and chlorophyll levels in plant tissues (Hussain et al., 2015). Zn produces heat resistance in water-stress plants (Hera et al., 2018) because the cell membrane is first affected by water stress and is a main part of the plant, therefore, its maintenance is necessary during water stress (Bajji et al., 2002 ; ). Stomata play major role in regulation of temperature during water stress condition (Karam et al., 2007) and Zinc play important role in the regulation of stomata in water stress condition (Bajji et al., 2002).

Zinc is a vital nutrient that plays a key role in human health, and is the 5th leading cause of illness and death in developing-country populations (Khalid et al., 2014). At the cellular level, it plays a major role in apoptosis and differentiation (Maret & Sandstead, 2006). Zinc is required for different functions including taste, respiration, DNA metabolism, vision, and behavior in the human body (Fraker & King, 2004). It is important for growth and development, particularly in children and pregnant women (Araujo et al., 2021). It is involved in numerous metabolic processes, including DNA synthesis, cell division, and protein metabolism (Chasapis et al., 2020). Zinc also helps to maintain a strong immune system and promotes skin, eye, and reproductive health (Dhok et al., 2020). Good food sources of zinc include meat, seafood, dairy products, and whole grains (Dussiot et al., 2022) and its deficiency is estimated to affect approximately 2 billion people worldwide (Singh et al., 2017) and in Pakistan, 37% of the population affected by zinc deficiency (Younas et al., 2022). Zinc deficiency is caused by a lack of zinc-rich foods in the diet, poor zinc absorption, and increased requirements during growth and illness (Gupta et al., 2020). Zinc deficiency adversely impacts immune function, slows wound healing, and inhibits taste and smell (Suman et al., 2020). Supplements are available for people who do not get enough zinc from their diet (Ceylan et al., 2021).


Cereals are an important part of a balanced diet and provide numerous health benefits (Kaur et al., 2014). According to the Food and Agriculture Organization (FAO) of the United Nations, cereals account for over 50% of the total calorie intake for almost half of the world's population (FAO, 2022).

Cereals are also a good source of carbohydrates, which provide the body with energy and are also rich in fiber, vitamins, and minerals such as iron and zinc (Sarita & Singh. 2016).

Wheat is one of the most important cereal crops in the world, providing a significant source of food for both humans and livestock (Poutanen et al., 2022). It is a staple food in many countries, and is a critical component of the diets of billions of people around the world, providing essential nutrients such as carbohydrates, proteins, vitamins, and minerals (Awika. 2011). According to the United Nations Food and Agriculture Organization (FAO), Wheat is one of the most widely cultivated crops in the world, grown on over 220 million hectares with a global production of over 740 million metric tons in 2021, and Pakistan is one of the world's largest wheat producers, with a production of around 25 million metric tonnes in 2021 (FAOSTATE, 2021). In Pakistan, approximately 75% of the cultivated area is irrigated, with the remaining 3.99 M ha being rainfed. Wheat is grown on 8.797 million hectares, with an annual production of 25.07 million tonnes. (Anjum et al., 2022).

Pakistan's rainfed areas are essential to the country's agriculture and economy because they provide a major source of food production, including wheat (Baig et al., 2013). Rainfed area contributes only about 12% of the total wheat production of the country (Qureshi et al., 2010). Drought is a major issue that has a significant impact on Pakistan's rainfed wheat yield (Ullah et al., 2021). Over the past three decades, heat waves on agricultural lands have increased, which has an impact on crop productivity worldwide (FAO. 2018). Mostly the grain-filling stage of wheat results in grain shrink that automatically affect the yield of wheat (Bodner et al., 2015). A lack of rainfall can also cause soil moisture stress, which can limit crop nutrient absorption and reduce overall yield (Ullah et al., 2019). Furthermore, drought can increase the risk of plant diseases and pest outbreaks, which can reduce crop yield (Tudi et al., 2021). Water stress crop losses can have serious consequences for the country's food security and economy (Newton et al., 2011). Foliar zinc application can be an effective way to mitigate the effects of drought stress in crops by providing essential micronutrients and improving plant tolerance to stress conditions (Toor et al., 2020).

**Picture 1 :Soil Factors affecting the availability of zinc to roots.**



This picture is taken from a study by Rudani et al., (2018). This picture shows the factors like high CaCO<sub>3</sub>, high pH, Clay particles, low organic matter, low soil moisture, and high Al oxide that affect the zinc level in soil and retard the availability of zinc to the plant roots.

Wheat crops grown in the field at a global scale have grain Zn ranges from 20.4 to 30.5 mg/kg and the human requirement for good health is 40 mg/kg which shows a solid gap. Foliar application of zinc can provide an effective and efficient method of delivering this essential micronutrient directly to the leaves of the plant. This method allows the plant to absorb the zinc quickly and directly, without having to rely on the slower process of soil uptake. As a result, foliar application of zinc can help to overcome soil-related zinc deficiencies and alleviate the effects of drought stress on crops.

## ISSUES

### Zinc in soil

Zinc levels are low in a variety of soils. Soils with high pH, sandy soils, calcareous soils, and soils derived from highly weathered parent materials all have low total Zn levels (Natasha et al., 2022). Similarly, due to saline soils, vertisol, and highly weathered soils have low plant-available zinc (Mossa et al., 2020). Zinc soil deficiency is a very common issue at the global level (Akther et al., 2020). The Zn soil deficiency has been reported in 57% of the soil samples of Iraq, 35% of Turkey, and 20% of Pakistan (Alloway, 2009). Almost 30% of agricultural soils show zinc deficiency in the World, and in Asia, 30% of soils are zinc deficient (Alloway, 2008; Cakmak, 2008). Low zinc in soils mostly affects main crops like wheat, oat, sorghum, and maize (Alloway, 2009).

**Table 1 list of sensitive crop species to zinc deficiency**

High	Medium	Low
Bean	Barley	Alfalfa
Citrus	Cotton	Carrot
Flax	Lettuce	Clover
Fruit trees	Potato	Grass
Grape	Soybean	Oat
Corn	Tomato	Pea
Rice	Sugar Beet	Rye
Sorghum		Wheat

Martens and Westermann (1991)

Globally, Zn-deficient soils identified in mostly arid and semi-arid areas of Pakistan and India, and in China, turkey and Australia due to low organic soils (Alloway,2008).

### Effect of Low Zinc on Plant Growth

Low zinc availability in soil has a negative impact on plant growth and development its deficiency stress becomes more sensitive to Zn deficiency stress became more distinct when plants in drought-stressed (Bagci et al., 2007). For example, a study by Suganya et al. (2020) found that zinc deficiency in soil reduced the growth and biomass production of corn plants. Similarly, a study of Rehman et al. (2012) reported that a low zinc supply decreased the growth and grain yield of rice plants. The study of Mousavi et al. (2012) reported that Zinc deficiency reduces plant growth by impairing the plant's ability to absorb and use essential nutrients like nitrogen and phosphorus, as well as by altering its hormone balance. The study by Rahman et al. (2019) reported that low zinc availability in soil reduces wheat growth and if wheat grows in supra optimal temperature. These findings highlight the importance of adequate zinc supply for optimal plant growth and productivity.

### Low Zinc Effect on Plant Height

A deficiency of zinc can significantly impact plant growth and development, including plant height (Hong et al., 2007). Several studies have demonstrated the negative effect of zinc deficiency on plant height. For instance, a study by Rag and Nadarajah (2022) reported that zinc-deficient maize plants exhibited a significant reduction in plant height compared to plants grown under an adequate zinc supply. Similarly, a study by Gonzalez-Caballo et al. (2022) showed that zinc-deficient wheat plants had shorter stems and reduced shoot length compared to plants grown under an adequate zinc supply. The negative impact of zinc deficiency on plant height is

attributed to its role in the production of auxins, which are plant hormones that regulate cell division and elongation. Zinc is required for the activity of enzymes involved in auxin biosynthesis and transport (Gondal et al., 2021). Zinc deficiency leads to a decrease in the production of auxins, which subsequently reduces stem elongation and plant height (Sharma et al., 2013).

### Effect of Low Zinc on Leaf Area Index

The study showed that rice crops grown under zinc-deficient soil significantly reduced growth in terms of Leaf Area Index, Leaf Area Duration, Crop Growth Rate, Total Dry Matter accumulation, and Net Assimilation Rate (Sawver et al 2013). The study of Hafeez et al. (2013) resulted that zinc deficiency shows a negative effect on plant growth by decreasing the number of tillers, small leaf area, and by increasing the crop maturity period. Moreover, a recent study by Ahmad et al. (2021) investigated the effect of different zinc concentrations on the growth and development of rainfed wheat. The study found that low zinc concentrations resulted in reduced leaf area and stunted growth.

### Effect of Low Zinc on Relative Water Contents

Rainfed agriculture relies on rainfall for water supply, but it is often erratic and unreliable, leading to water stress in plants (Zahoor et al., 2019). Low levels of zinc in the soil reduce the plant's ability to produce ABA, leading to inefficient stomatal closure and increased water loss. This can lead to wilting, stunted growth, and reduced yield (Dubey. 2018). The study of Kumar et al. (2008) showed that low zinc in soil decreases water potential and relative water contents also have fewer chloroplast pigments and high tissue Mn concentration. Vazin. (2012) found that zinc spray significantly increased the thousand-grain weight of maize under water stress, improving the effects of drought stress. This suggests that zinc spray can be used to reduce the effects of drought stress.

### Effect of Zinc Deficiency on Crop Yield

Zn is mainly a plant micronutrient that is involved in several physiological functions and insufficient supply reduces crop yields in calcareous soil, sandy soils, and soils with high phosphorus conditions (Hafeez et al., 2013). The study by Hassan et al. (2021) demonstrated the number of tillers, chlorophyll contents, plant growth, and crop yield affected by zinc deficiency. Similarly, the growth and yield of crops are reduced by different abiotic stresses as well as water stress in rainfed areas during

the booting, heading and flowering stages (Hera et al., 2018). It is also resulted the skipping of irrigation at the grain-filling stage causing a reduction in yield. It is concluded that zinc-deficient plants showed leaf chlorosis, necrosis bronzing, resetting dwarf, and stunting of plants. However, 40% yield was reduced by zinc-deficient soil without the existence of diverse symptoms in plant leaves (Vadlamudi et al., 2020).

### Effect of Zinc Deficiency on Human Health

The study of Plum et al. (2010) Proved that the Zinc deficiency disturb the normal health of the human like neural development, growth and immunity and in severe cases its consequences are more lethal. The study of Gibson. (2012) resulted that the competence of the immune system, risk of stunting, childhood mortality, diarrhea, respiratory diseases, and abortions caused by zinc deficiency.

### Zinc essentiality for human

Zinc was discovered to be essential in humans in 1963 (Prasad et al., 2014). Zinc is required to properly operate thousands of transcriptions (Maret. 2017). Zinc is a second messenger of immune cells, and intracellular free zinc participates in signaling events in these cells (Haase & Rink. 2014). Zinc deficiency in pregnant women during growth periods shows no proper growth. The gastrointestinal, central nervous, skeletal, and reproductive systems are the organs affected by zinc deficiency (Roohani et al. 2013). Nutritious necessities during lactation are greater than during pregnancy (Roba et al., 2018).

## POSSIBLE SOLUTIONS

### Plant Height Increased by Foliar Zinc

The study of Abdoli et al. (2014) Revealed that the height of wheat crops significantly increased by foliar application of zinc sulfate at development and grain filling stages however non-significant on the harvest index. A study by Gull et al., (2011) revealed that the foliar application of nitrogen, potassium, and Zn at the rate of 0.5% applied two times significantly increases the number of tillers, and gets maximum plant height (100.50cm). The study of Ali et al. (2015) concluded that the combined foliar application of Boron and Zinc on the summer tomato significantly increases the growth rate, plant height, leaf area number of branches, and overall growth and yield of the summer tomato.

### Crop Yield Increase by Foliar Zinc

The study of Sattar et al. (2022) on the wheat crop grown under different water stress


conditions resulted that foliar zinc of 15 mM reduce the water stress impact also increasing the grain yield by increasing the number of grains per spike, grain weight, and biological yield. The study by Noreen & Kamran (2019) resulted that foliar application of 4 and 6 mM of zinc significantly increases the leaf area, plant height, number of grains per spike, spike length, yield, and grain zinc content. It also concluded that plant fresh weight increased by zinc spray. According to literature studied by Gite et al. (2021), it is observed that  $ZnSO_4 \cdot H_2O$  and  $ZnSO_4 \cdot 7H_2O$  significantly increase the wheat biomass, yield, and zinc contents in grain. The study by Ma et al. (2017) revealed that the soil application of zinc in wheat grown under rainfed conditions during severe drought increased grain yield and zinc concentration by 28.2 and 32.8% respectively. Furthermore, the scientist showed that zinc fertilization plays an important role in alleviating drought stress in wheat plants by increasing zinc fertilization in photosynthetic pigment and active oxygen-scavenging substances.

### Plant Resistance to Water Stress

### Zinc Fertilization

To overcome the zinc deficiency in the soil, zinc-containing fertilizers are spread or sprayed on the surface of the soil, side dressed in standing crop, applied as foliar sprays, used as a seed coating, and in case of transplanted rice plants by dipping the roots before transplanting (Allowy. 2009). Zinc sulphate is the most commonly used fertilizer compound ( $ZnSO_4 \cdot 7H_2O$  with 27% Zn, and  $ZnSO_4 \cdot H_2O$  containing 33% Zn).

Fig 1 :Effect of Zinc Fertilization



### Effect of foliar zinc on grain weight

Grain weight play major role in health of grain and crop yield. Different scientists work on

This parameter. The study of Amanullah et al. (2021) on Integrated foliar nutrients application improve wheat (*Triticum Aestivum* L.) productivity under calcareous soils in drylands revealed that foliar application of Zn + boron by the rate of (0.2%) at booting stage of wheat grown under calcareous soil of semiarid region increase the grain weight and more spikelet's spike<sup>-1</sup> higher grain yield.

#### **Effect of Foliar zinc on grains spike<sup>-1</sup>:**

The study of Esfandiari et al. (2016) on Impact of foliar zinc application on agronomic traits and grain quality parameters of wheat grown in zinc deficient soil revealed that the foliar application of zinc sulfate at booting and milking stage significantly increase the number of grains spike<sup>-1</sup>. Furthermore, it also found that by the application of zinc sulphate agronomic traits also increased.

#### **Conclusion**

#### **References**

Rudani, L., Vishal, P., & Kalavati, P. (2018). The importance of zinc in plant growth-A review. *International Research Journal of Natural and Applied Sciences*, 5(2), 38-48.

Hussain, A., Arshad, M., Zahir, Z. A., & Asghar, M. (2015). Prospects of zinc solubilizing bacteria for enhancing growth of maize. *Pakistan journal of agricultural sciences*, 52(4).

Hera, M. H. R., Hossain, M., & Paul, A. K. (2018). Effect of foliar zinc spray on growth and yield of heat tolerant wheat under water stress. *Int J Biol Environ Eng*, 1(1), 10-16.

Bajji, M., Kinet, J. M., & Lutts, S. (2002). The use of the electrolyte leakage method for assessing cell membrane stability as a water stress tolerance test in durum wheat. *Plant growth regulation*, 36(1), 61-70.

Karam, F., Lahoud, R., Masaad, R., Kabalan, R., Breidi, J., Chalita, C., & Rouphael, Y. (2007). Evapotranspiration, seed yield and water use efficiency of drip irrigated sunflower under full and deficit irrigation conditions. *Agricultural water management*, 90(3), 213-223.

Maret, W., & Sandstead, H. H. (2006). Zinc requirements and the risks and benefits of zinc supplementation. *Journal of trace elements in medicine and biology*, 20(1), 3-18.

Fraker, P. J., & King, L. E. (2004). Reprogramming of the immune system during zinc deficiency. *Annual review of nutrition*, 24, 277.

Chasapis, C. T., Ntoupa, P. S. A., Spiliopoulou, C. A., & Stefanidou, M. E. (2020). Recent aspects of the effects of zinc on human health. *Archives of toxicology*, 94, 1443-1460.

Dhok, A., Butola, L. K., Anjankar, A., Shinde, A. D. R., Kute, P. K., & Jha, R. K. (2020). Role of vitamins and minerals in improving immunity during Covid-19 pandemic-A review. *Journal of Evolution of Medical and Dental Sciences*, 9(32), 2296-301.

Araújo, L. A. D., Veloso, C. F., Souza, M. D. C., Azevedo, J. M. C. D., & Tarro, G. (2021). The potential impact of the COVID-19 pandemic on child growth and development: a systematic review. *Jornal de pediatria*, 97, 369-377.

Dussiot, A., Fouillet, H., Wang, J., Salomé, M., Huneau, J. F., Kesse-Guyot, E., & Mariotti, F. (2022). Modeled healthy eating patterns are largely constrained by currently estimated requirements for bioavailable iron and zinc—a diet optimization study in French adults. *The American journal of clinical nutrition*, 115(3), 958-969.

Singh, S. P., Keller, B., Gruissem, W., & Bhullar, N. K. (2017). Rice NICOTIANAMINE SYNTHASE 2 expression improves dietary iron and zinc levels in wheat. *Theoretical and Applied Genetics*, 130, 283-292.

Younas, N., Fatima, I., Ahmad, I. A., & Ayyaz, M. K. (2022). Alleviation of zinc deficiency in plants and humans through an effective technique; biofortification: A detailed review. *Acta Ecologica Sinica*.

Gupta, S., Brazier, A. K. M., & Lowe, N. M. (2020). Zinc deficiency in low- and middle- income

countries: prevalence and approaches for mitigation. *Journal of Human Nutrition and Dietetics*, 33(5), 624-643.

Suman, P. B. M. (2020). A review on role of zinc deficiency in humans and its rectification through Horti-agro food supplements.

Ceylan, M. N., Akdas, S., & Yazihan, N. (2021). Is zinc an important trace element on bone-related diseases and complications A meta-analysis and systematic review from serum level, dietary intake, and supplementation aspects. *Biological Trace Element Research*, 199, 535-549.

Khalid, N., Ahmed, A., Bhatti, M. S., Randhawa, M. A., Ahmad, A., & Razaqat, R. (2014). A question mark on zinc deficiency in 185 million people in Pakistan—possible way out. *Critical reviews in food science and nutrition*, 54(9), 1222-1240.

FAOSTAT Data. Available online: <https://www.fao.org/faostat/en/#home> (accessed on 13, December 2021).

Anjum, M. M., & Arif, M. (2022). Perspectives of wheat hybrid yield and quality under limited irrigation supply and sowing windows. *Gesunde Pflanzen*, 74(4), 761-770.

Baig, M. B., Shahid, S. A., & Straquadine, G. S. (2013). Making rainfed agriculture sustainable through environmental friendly technologies in Pakistan: A review. *International Soil and Water Conservation Research*, 1(2), 36-52

Qureshi, A. S., McCornick, P. G., Sarwar, A., & Sharma, B. R. (2010). Challenges and prospects of sustainable groundwater management in the Indus Basin, Pakistan. *Water resources management*, 24(8), 1551-1569.

Ullah, N., Ditta, A., Imtiaz, M., Li, X., Jan, A. U., Mehmood, S., & Rizwan, M. (2021). Appraisal for organic amendments and plant growth-promoting rhizobacteria to enhance crop productivity under drought stress: A review. *Journal of Agronomy and Crop Science*, 207(5), 783-802.

Bodner, G., Nakhforoosh, A., & Kaul, H. P. (2015). Management of crop water under drought: a review. *Agronomy for Sustainable Development*, 35, 401-442.

Ullah, H., Santiago-Arenas, R., Ferdous, Z., Attia, A., & Datta, A. (2019). Improving water use efficiency, nitrogen use efficiency, and radiation use efficiency in field crops under drought stress: A review. *Advances in agronomy*, 156, 109-157.

Tudi, M., Daniel Ruan, H., Wang, L., Lyu, J., Sadler, R., Connell, D., & Phung, D. T. (2021). Agriculture development, pesticide application and its impact on the environment. *International journal of environmental research and public health*, 18(3), 1112.

Newton, A. C., Johnson, S. N., & Gregory, P. J. (2011). Implications of climate change for diseases, crop yields and food security. *Euphytica*, 179, 3-18.

Toor, M. D., Adnan, M., Javed, M. S., Habibah, U., Arshad, A., Din, M. M., & Ahmad, R. (2020). Foliar application of Zn: Best way to mitigate drought stress in plants; A review. *International Journal of Applied Research*, 6(8), 16-20.

Suganya, A., Saravanan, A., & Manivannan, N. (2020). Role of zinc nutrition for increasing zinc availability, uptake, yield, and quality of maize (*Zea mays* L.) grains: An overview. *Commun. Soil Sci. Plant Anal*, 51(15), 2001-2021.

Rehman, H. U., Aziz, T., Farooq, M., Wakeel, A., & Rengel, Z. (2012). Zinc nutrition in rice production systems: a review. *Plant and soil*, 361, 203-226.

Mousavi, S. R., Galavi, M., & Rezaei, M. (2012). The interaction of zinc with other elements in plants: a review. *International Journal of Agriculture and Crop Sciences*, 4(24), 1881-1884.

World Health Organization. (2018). *The state of food security and nutrition in the world 2018: building climate resilience for food security and nutrition*. Food & Agriculture Org.

Poutanen, K. S., Kårlund, A. O., Gómez-Gallego, C., Johansson, D. P., Scheers, N. M., Marklinder, I. M & Landberg, R. (2022). Grains—a major source of sustainable protein for health. *Nutrition reviews*, 80(6), 1648-1663.

- Rehman, A., Farooq, M., Asif, M., & Ozturk, L. (2019). Supra-optimal growth temperature exacerbates adverse effects of low Zn supply in wheat. *Journal of Plant Nutrition and Soil Science*, 182(4), 656-666.
- <https://www.fao.org/3/cc0639en/cc0639en.pdf>
- Hong, W. A. N. G., & JIN, J. Y. (2007). Effects of zinc deficiency and drought on plant growth and metabolism of reactive oxygen species in maize (*Zea mays* L). *Agricultural Sciences in China*, 6(8), 988-995.
- Raj, S. R. G., & Nadarajah, K. (2022). QTL and Candidate Genes: Techniques and Advancement in Abiotic Stress Resistance Breeding of Major Cereals. *International Journal of Molecular Sciences*, 24(1), 6.
- González-Caballo, P., Barrón, V., Torrent, J., del Campillo, M. C., & Sánchez-Rodríguez, A. R. (2022). Wheat and maize grown on two contrasting zinc-deficient calcareous soils respond differently to soil and foliar application of zinc. *Journal of Soil Science and Plant Nutrition*, 22(2), 1718-1731.
- Gondal, A. H., Zafar, A., Zainab, D., Toor, M. D., Sohail, S., Ameen, S & Younas, N. (2021). A detailed review study of zinc involvement in animal, plant and human nutrition. *Indian Journal of Pure & Applied Biosciences*, 9(2), 262-271.
- Sharma, A., Patni, B., Shankhdhar, D., & Shankhdhar, S. C. (2013). Zinc—an indispensable micronutrient. *Physiology and Molecular Biology of Plants*, 19, 11-20.
- Sarwar, N., Ali, H., Maqsood, M., Ullah, E., Shahzad, M., Mubeen, K., & Ahmad, S. (2013). Phenological response of rice plants to different micronutrients application under water saving paddy fields on calcareous soil. *Turkish Journal of Field Crops*, 18(1), 52-57.
- Kumar Tewari, R., Kumar, P., & Nand Sharma, P. (2008). Morphology and physiology of zinc-stressed mulberry plants. *Journal of Plant Nutrition and Soil Science*, 171(2), 286-294.
- Plum, L. M., Rink, L., & Haase, H. (2010). The essential toxin: impact of zinc on human health. *International journal of environmental research and public health*, 7(4), 1342- 1365.
- Chen, X. P., Zhang, Y. Q., Tong, Y. P., Xue, Y. F., Liu, D. Y., Zhang, W., & Zou, C. Q. (2017). Harvesting more grain zinc of wheat for human health. *Scientific Reports*, 7(1), 7016.
- Gibson, R. S. (2012). Zinc deficiency and human health: etiology, health consequences, and future solutions. *Plant and soil*, 361, 291-299.
- Hafeez, B. M. K. Y., Khanif, Y. M., & Saleem, M. (2013). Role of zinc in plant nutrition-a review. *American journal of experimental Agriculture*, 3(2), 374.
- Hassan, M. U., Aamer, M., Nawaz, M., Rehman, A., Aslam, T., Afzal, U., & Guoqin, H. (2021). Agronomic bio-fortification of wheat to combat zinc deficiency in developing countries. *Pakistan Journal of Agricultural Research*, 34(1), 201.
- Kaur, K. D., Jha, A., Sabikhi, L., & Singh, A. K. (2014). Significance of coarse cereals in health and nutrition: a review. *Journal of food science and technology*, 51, 1429-1441.
- Sarita, E. S., & Singh, E. (2016). Potential of millets: nutrients composition and health benefits. *Journal of Scientific and Innovative Research*, 5(2), 46-50.
- Ahmad, G., Khan, A. A., & Mohamed, H. I. (2021). Impact of the low and high concentrations of fly ash amended soil on growth, physiological response, and yield of pumpkin (*Cucurbita moschata* Duch. Ex Poiret L.). *Environmental Science and Pollution Research*, 28, 17068- 17083.
- Hafeez, B. M. K. Y., Khanif, Y. M., & Saleem, M. (2013). Role of zinc in plant nutrition-a review. *American journal of experimental Agriculture*, 3(2), 374.
- Bagci, S. A., Ekiz, H., Yilmaz, A., & Cakmak, I. (2007). Effects of zinc deficiency and drought on grain yield of field-grown wheat cultivars in Central Anatolia. *Journal of Agronomy and Crop Science*, 193(3), 198-206.
- Vadlamudi, K., Upadhyay, H., Singh, A., & Reddy, M. (2020). Influence of zinc application in plant growth: an overview. *Eur. J. Mol. Clin. Med*, 7, 2321-2327.

- Rudani, L., Vishal, P., & Kalavati, P. (2018). The importance of zinc in plant growth-A review. *International research journal of natural and applied sciences*, 5(2), 38-48.
- Abdoli, M., Esfandiari, E., Mousavi, S. B., & Sadeghzadeh, B. (2014). Effects of foliar application of zinc sulfate at different phenological stages on yield formation and grain zinc content of bread wheat (cv. Kohdasht). *Azarian Journal of Agriculture*.
- Sattar, A., Wang, X., Ul-Allah, S., Sher, A., Ijaz, M., Irfan, M., & Skalicky, M. (2022). Foliar application of zinc improves morpho-physiological and antioxidant defense mechanisms, and agronomic grain biofortification of wheat (*Triticum aestivum* L.) under water stress. *Saudi Journal of Biological Sciences*, 29(3), 1699-1706.
- Noreen, S., & Kamran, A. (2019). Foliar application of zinc sulphate to improve yield and grain zinc content in wheat (*Triticum aestivum* L.). *African Journal of Agricultural Research*, 14(20), 867-876.
- Gite, D., Gite, H., & Gite, M. D. (2021). Zinc sulphate a potential micronutrient for wheat growth: A review.
- Roohani, N., Hurrell, R., Kelishadi, R., & Schulin, R. (2013). Zinc and its importance for human health: An integrative review. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*, 18(2), 144.
- Prasad, A. S. (2014). Impact of the discovery of human zinc deficiency on health. *Journal of trace elements in medicine and biology*, 28(4), 357-363.
- Maret, W. (2017). Zinc in cellular regulation: the nature and significance of “zinc signals”. *International journal of molecular sciences*, 18(11), 2285.
- Haase, H., & Rink, L. (2014). Zinc signals and immune function. *Biofactors*, 40(1), 27-40.
- Roba, K. T., O'Connor, T. P., Belachew, T., & O'Brien, N. M. (2018). Concurrent iron and zinc deficiencies in lactating mothers and their children 6–23 months of age in two agro-ecological zones of rural Ethiopia. *European journal of nutrition*, 57, 655-667.
- Mossa, A. W., Young, S. D., & Crout, N. M. (2020). Zinc uptake and phyto-toxicity: Comparing intensity-and capacity-based drivers. *Science of the Total Environment*, 699, 134314.
- Roohani, N., Hurrell, R., Kelishadi, R., & Schulin, R. (2013). Zinc and its importance for human health: An integrative review. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*, 18(2), 144.
- Akther, M. S., Das, U., Tahura, S., Prity, S. A., Islam, M., & Kabir, A. H. (2020). Regulation of Zn uptake and redox status confers Zn deficiency tolerance in tomato. *Scientia Horticulturae*, 273, 109624.
- Natasha, N., Shahid, M., Bibi, I., Iqbal, J., Khalid, S., Murtaza, B., ... & Arshad, M. (2022). Zinc in soil-plant-human system: A data-analysis review. *Science of the Total Environment*, 808, 152024.
- Alloway, B. J. (2009). Soil factors associated with zinc deficiency in crops and humans. *Environmental geochemistry and health*, 31(5), 537-548.
- Alloway, B. J. (2008). Zinc in soils and crop nutrition.
- Cakmak, I. (2008). Enrichment of cereal grains with zinc: agronomic or genetic biofortification?. *Plant and soil*, 302, 1-17.
- <http://nmsp.cals.cornell.edu/publications/factsheets/factsheet32.pdf>
- Ma, D., Sun, D., Wang, C., Ding, H., Qin, H., Hou, J., & Guo, T. (2017). Physiological responses and yield of wheat plants in zinc-mediated alleviation of drought stress. *Frontiers in plant science*, 8, 860.
- Alloway, B. J. (2009). Soil factors associated with zinc deficiency in crops and humans. *Environmental geochemistry and health*, 31(5), 537-548.
- Awika, J. M. (2011). Major cereal grains production and use around the world. In *Advances in cereal science: implications to food processing and health promotion* (pp. 1-13). American Chemical Society.
- Zahoor, S. A., Ahmad, S., Ahmad, A., Wajid, A., Khaliq, T., Mubeen, M., & Nasim, W. (2019). Improving water use efficiency in agronomic crop production. *Agronomic Crops: Volume 2: Management Practices*, 13-29

Dubey, R. S. (2018). Photosynthesis in plants under stressful conditions. In *Handbook of photosynthesis* (pp. 629-649). CRC Press.

Gul, H., Said, A., Saeed, B., Mohammad, F., & Ahmad, I. (2011). Effect of foliar application of nitrogen, potassium and zinc on wheat growth. *ARPJ Agric Biol Sci*, 6(4), 56-59.

Ali, M. R., Mehraj, H., & Jamal Uddin, A. F. M. (2015). Effects of foliar application of zinc and boron on growth and yield of summer tomato. *Journal of Bioscience and Agriculture Research*, 6(1), 512-517.

Vazin, F. (2012). Effect of zinc sulfate on quantitative and qualitative characteristics of corn (*Zea mays*) in drought stress.

Amanullah, Ilyas, M., Nabi, H., Khalid, S., Ahmad, M., Muhammad, A., ... & Parmar, B. (2021). Integrated foliar nutrients application improve wheat (*TriticumAestivum* L.) productivity under calcareous soils in drylands. *Communications in Soil Science and Plant Analysis*, 52(21), 2748-2766.

Esfandiari, E., Abdoli, M., Mousavi, S. B., & Sadeghzadeh, B. (2016). Impact of foliar zinc application on agronomic traits and grain quality parameters of wheat grown in zinc deficient soil. *Indian Journal of Plant Physiology*, 21, 263-270.