

Navigating the Complexities of HIV Care Engagement and the Care Cascade: A Perspective

Comment [Dr Uduma 1]: Suggested Title: "Navigating Complexities in HIV Care: Challenges, Solutions, and Strategies."

Abstract:

HIV/AIDS remains a global public health challenge, despite significant progress in prevention and treatment. Achieving the UNAIDS 95-95-95 targets, which aim to have 95% of all people living with HIV know their status, 95% of those diagnosed receiving antiretroviral therapy (ART), and 95% of those on ART achieving viral suppression, requires overcoming numerous challenges and barriers. In this viewpoint, we discuss the multifaceted obstacles that hinder HIV care engagement and the care cascade, encompassing issues related to testing, linkage to care, retention in care, and viral suppression. We also propose potential solutions and strategies to address these challenges and move closer to the Sustainable development goal of ending the HIV epidemic by 2030.

Comment [Dr Uduma 2]: consider adding a dedicated subsection/statement in the abstract to summarize proposed solutions and strategies.

Keywords: Care Cascade, Complexities, Engagement, HIV Care Perspective

Introduction

HIV/AIDS remains a global health crisis, with approximately 38 million people living with the virus worldwide. In recent years, the UNAIDS 95-95-95 targets have been established to accelerate progress towards ending the HIV epidemic. While these targets have the potential to transform the global response to HIV, substantial challenges and barriers persist in achieving them [1]. This viewpoint aims to shed light on the complexities surrounding HIV care engagement and the care cascade and to propose solutions to overcome these obstacles.

Challenges in HIV Testing:

HIV testing is a pivotal entry point into the care cascade, where individuals learn about their HIV status and take the first steps toward managing their health. Despite notable advancements in the field and existence of well curated national health programs for HIV/AIDS, testing rates still fall short in many regions, largely due to the persistence of multifaceted challenges that need attention [2].

Comment [Dr Uduma 3]: write as : challenges that need attention. Remove "s" from needs

a. Stigma and Discrimination: A persistent and damaging social stigma attached to HIV continues to be a formidable barrier to testing. Some have also witnessed stigma when visiting healthcare facilities to access HIV related or general healthcare. The stigma could persist in three forms; internalized, anticipated and enacted. The fear of negative consequences, such as discrimination, ostracism, or a loss of social support, often deters individuals from seeking an HIV test. This stigma can create a climate of silence and secrecy surrounding HIV, making it difficult for those at risk or in need of testing to come forward [3].

b. Lack of Access: Accessibility to HIV testing facilities and healthcare services is another major concern, particularly in rural and underserved areas. In these regions, the limited availability of testing centers and healthcare infrastructure restricts the opportunities for individuals to get tested. Non-availability of testing centers near one's locality, discourages individuals to get themselves tested as daily wagers have to bear economic losses for the same. The resulting low testing rates exacerbate the overall HIV burden in these communities, making it challenging to identify cases and link individuals to appropriate care [4].

c. Testing Modalities: To overcome the barriers associated with traditional testing methods, there is a growing need for more accessible and user-friendly testing modalities. Initiatives such as self-testing and community-based testing have shown great promise in reaching populations that are often marginalized and under-tested [5]. These approaches empower individuals to take control of their own health by offering them the option to test in the privacy of their own homes or in community settings. Widening the implementation of these innovative testing modalities is vital to ensure that a broader cross-section of society has access to testing and can engage with the care cascade [6].

Addressing these challenges in HIV testing requires a comprehensive approach that not only focuses on improving the availability and accessibility of testing services but also aims to combat the stigma and discrimination that hinders people from taking this crucial first step toward managing their HIV status [7]. It is essential to create an environment where individuals feel safe, supported, and empowered to know their HIV status and access the care they need [8].

Barriers to Linkage to Care:

Once individuals test positive for HIV, the next essential step is linking them to appropriate care and treatment. However, numerous barriers can hinder this crucial transition, impacting both individual health outcomes and the broader epidemic [9].

a. Delayed Linkage: One significant barrier faced by individuals who test positive is delayed linkage to care. The delay in the linkage to appropriate care could be due to individual's own health seeking behavior, stigma associated or due to lacunas in the health systems itself. This delay can lead to a myriad of adverse consequences, including increased morbidity, deteriorated quality of life and the heightened risk of viral transmission [10]. The longer the gap between diagnosis and care initiation, the greater the potential harm to the individual's health, as well as

the risk of transmission to others. Reducing these delays is paramount to improving individual health outcomes and preventing further HIV transmission [11].

b. Sociodemographic Factors: The sociodemographic factors surrounding an individual can play a substantial role in their ability to successfully link to HIV care. Factors such as poverty, homelessness, and substance abuse can significantly hinder this linkage process. Individuals facing economic hardships or unstable housing may struggle to access the necessary care facilities and support services. Substance abuse issues can exacerbate these challenges, making it even more challenging to engage with care. Despite treatment being available free of cost in public healthcare facilities, still linkage to treatment presents a challenge as there are financial implications when a person has to come over to treatment center for medicines as he/she has to take a loss on their wages (applicable to those who are daily wage earners). To address this, after initial assessment of person's overall health, efforts should be made to deliver the medicines at the infected person's doorstep if he/she is not willing to come over to health facilities due to any reason possible [12].

c. Mental Health: Co-occurring mental health issues often go unaddressed and can impede the engagement of individuals in HIV care. Mental health disorders, such as depression or anxiety, can create significant barriers to accessing and adhering to treatment regimens [13]. This can further compound the complexity of managing HIV and achieving viral suppression.

To overcome these barriers to linkage to care, a holistic approach is essential. This approach involves not only streamlining the process from diagnosis to care initiation but also addressing the social determinants of health that influence an individual's access to care. Implementing strategies to combat poverty, homelessness, and substance abuse, integrating mental health services into HIV care programs and support within the society can help break down these barriers, enabling more individuals to engage with care and successfully manage their HIV infection [14].

Retention in Care:

Sustaining engagement in HIV care is crucial to achieving positive health outcomes and maintaining viral suppression. However, numerous challenges can undermine retention in care, perpetuating the risk of HIV transmission and disease progression [15].

a. Socioeconomic Factors: Socioeconomic hardships, such as unstable housing, unemployment, and financial insecurity, represent significant barriers to retaining individuals in HIV care. The stress and uncertainty associated with these challenges can divert a person's focus away from managing their health, making it difficult to consistently attend medical appointments and adhere to prescribed treatment regimens [16].

b. Mental Health and Substance Abuse: Co-occurring mental health issues and substance abuse can substantially contribute to disengagement from HIV care. When get unnoticed and

unmanaged, these conditions can disrupt an individual's ability to maintain treatment adherence and regularity in attending medical appointments [17]. The struggle to address mental health challenges or substance abuse issues often takes precedence over maintaining HIV care, posing a substantial threat to both individual well-being and public health [18].

c. Stigma and Discrimination: The enduring stigma attached to HIV can lead to patients discontinuing care due to fear, shame, or a desire to keep their HIV status hidden. This fear of discrimination can result in individuals disengaging from care, jeopardizing their own health and the progress of the epidemic. Many a times, healthcare delivery system is itself responsible for making infected person feel stigmatized. Overcoming stigma and discrimination is pivotal to retaining patients in HIV care and ensuring their overall well-being [19].

Addressing these barriers to retention in HIV care necessitates a comprehensive approach. It requires not only providing medical support but also addressing the broader social determinants of health that influence an individual's ability to sustain care [20]. Efforts should be made to tackle socioeconomic hardships, integrate mental health and substance abuse services into HIV care programs, the healthcare providers should be trained on avoidance of stigma when dealing with HIV infected persons, and to create an environment where individuals feel safe, supported, and encouraged to remain in care, leading to better health outcomes and reduced HIV transmission [21].

Barriers to Achieving Viral Suppression:

The ultimate goal in the HIV care cascade is to achieve viral suppression through antiretroviral therapy (ART). However, numerous challenges can impede this crucial step in managing HIV, potentially leading to poor health outcomes and continued transmission of the virus [22].

a. Medication Adherence: Sustained adherence to medication regimens is often challenging for individuals living with HIV. The need for lifelong treatment, coupled with the side effects of ART and inadequate counseling can lead to non-adherence, treatment interruptions, and suboptimal outcomes. Non adherence increases the risk of viral rebound, progression of the disease, and potential transmission to others. Improving medication adherence is central to achieving viral suppression [23].

b. Drug Resistance: The development of drug resistance is a significant concern if ART regimens are not well-tailored or adhered to. This can occur when the virus mutates and becomes less responsive to the medications prescribed. In cases of drug resistance, treatment options may become limited, and more complex, expensive, and less effective regimens may be required to manage the virus [24].

c. Comorbid Conditions: Patients with comorbid conditions, such as tuberculosis or other infections, may face additional difficulties in achieving viral suppression. Coexisting health issues can complicate the management of HIV and may require specialized care and coordination

to ensure that both the primary HIV infection and the comorbid conditions are effectively treated [25].

To overcome these barriers to achieving viral suppression, a patient-centered approach is crucial. This approach should involve comprehensive healthcare, regular monitoring of medication adherence, and the development of tailored treatment plans. It is essential to provide ongoing support, education, and resources to empower individuals to adhere to their ART regimens and effectively manage comorbid conditions, ultimately leading to viral suppression and improved overall health [26].

Proposed Solutions and Strategies:

To address the multifaceted challenges within the HIV care cascade, a comprehensive and multi-pronged approach is essential. The following strategies can help overcome these obstacles and improve the overall management of HIV:

a. Reducing Stigma: Tailored public health campaigns and education should be utilized to reduce the stigma and discrimination associated with HIV. Healthcare delivery systems should be made more inclusive. By fostering understanding and empathy, individuals are more likely to seek testing and care without the fear of social repercussions. This not only benefits individual health but also helps reduce the overall transmission of the virus [27].

b. Expanding Access: To improve access to testing and care, there should be a wider implementation of community-based testing, mobile clinics, and telemedicine. These approaches can reach underserved and remote populations and to all others who need them, making it easier for individuals to access services in their communities [28].

c. Supportive Services: Integrating mental health and substance abuse treatment within HIV care settings is crucial. Addressing co-occurring conditions can significantly improve engagement in care. Comprehensive care, which includes both physical and mental health support, helps individuals better manage their HIV [29].

d. Social Support: Providing social programs, housing assistance, and employment support can address socioeconomic factors that disrupt retention in care. By assisting individuals in finding stable housing and employment, they are more likely to remain engaged in care, leading to better health outcomes [30].

e. Treatment Tailoring: Personalized treatment plans based on patient needs and preferences can enhance medication adherence and reduce the development of drug resistance. Recognizing that each individual's journey with HIV is unique allows for tailored care that is more effective [31].

f. Integrated Care: Combining HIV care with primary healthcare services is another strategy to enhance overall health outcomes. This approach ensures that individuals receive comprehensive

healthcare, addressing not only HIV but also other healthcare needs. Integrated care helps to manage comorbid conditions more effectively. As improved treatment adherence will lead to HIV infected population living longer, the emergence of non-communicable diseases presents considerable challenges in management of these comorbidities as they usually take longer to get treated. Hence, integrated care becomes essential to overcome such challenges [32].

By implementing these strategies, it is possible to address the challenges and barriers in the HIV care cascade comprehensively. This can lead to improved outcomes for individuals living with HIV and contribute to the global effort to end the HIV epidemic [33].

Conclusion

The challenges and barriers within the HIV care engagement and care cascade are complex and interwoven, demanding a comprehensive, all-encompassing approach. While strides have been taken in the fight against HIV, there is still substantial work to be done to achieve the ambitious UNAIDS 95-95-95 targets and, ultimately, to bring an end to the HIV epidemic. To reach this goal, it is imperative to address the structural, social, and individual impediments that hinder progress while embracing innovative solutions. Only by tackling these multifaceted challenges head-on can we draw closer to a world where HIV is no longer a global health crisis, but rather a triumph of public health and human resilience.

REFERENCES

1. Bouabida, K., Chaves, B. G., & Anane, E. (2023). Challenges and barriers to HIV care engagement and care cascade: viewpoint. *Frontiers in Reproductive Health, Section HIV and STIs, Volume 5*, 1-5.
2. Hammer, S. M., Squires, K. E., Hughes, M. D., Grimes, J. M., Demeter, L. M., Currier, J. S., ... et al. (1997). A controlled trial of two nucleoside analogues plus indinavir in persons with human immunodeficiency virus infection and CD4 cell counts of 200 per cubic millimeter or less. *New England Journal of Medicine*, 337(11), 725–733. doi: 10.1056/NEJM199709113371101.
3. Medeiros, P., Warren, L., Kazemi, M., Massaquoi, N., Smith, S., Tharao, W., ... et al. (2023). HIV care cascade for women living with HIV in the greater Toronto area versus the rest of Ontario and Canada. *International Journal of STD & AIDS*, 34(1), 4–17. doi: 10.1177/09564624221108034.

4. Keene, C. M., Ragnathan, A., Euvrard, J., English, M., McKnight, J., Orrell, C., ... InCARE Stakeholder Group. (2022). Measuring patient engagement with HIV care in sub-Saharan Africa: a scoping study. *Journal of the International AIDS Society*, 25(10), e26025. doi: 10.1002/jia2.26025.
5. Cohen, M. S., Chen, Y. Q., McCauley, M., Gamble, T., Hosseinipour, M. C., Kumarasamy, N., ... et al. (2011). Prevention of HIV-1 infection with early antiretroviral therapy. *New England Journal of Medicine*, 365(6), 493–505. doi: 10.1056/NEJMoa1105243.
6. Yuvaraj, A., Mahendra, V. S., Chakrapani, V., Yunihastuti, E., Santella, A. J., Ranauta, A., ... et al. (2020). HIV and stigma in the healthcare setting. *Oral Diseases*, 26(Suppl 1), 103–111. doi: 10.1111/odi.13585.
7. Corrigan, P. W., & Rao, D. (2012). On the self-stigma of mental illness: stages, disclosure, and strategies for change. *Canadian Journal of Psychiatry*, 57(8), 464–469. doi: 10.1177/070674371205700804.
8. Logie, C. H., Kaida, A., de Pokomandy, A., O'Brien, N., O'Campo, P., MacGillivray, J., ... et al. (2020). Prevalence and correlates of forced sex as a self-reported mode of HIV acquisition among a cohort of women living with HIV in Canada. *Journal of Interpersonal Violence*, 35(21–22), 5028–5063. doi: 10.1177/0886260517718832.
9. Seid, A., Seid, O., Workineh, Y., Dessie, G., & Bitew, Z. W. (2023). Prevalence of undernutrition and associated factors among adults taking antiretroviral therapy in sub-Saharan Africa: a systematic review and meta-analysis. *PLoS One*, 18(3), e0283502. doi: 10.1371/journal.pone.0283502.
10. Mafi, J. N., Reid, R. O., Baseman, L. H., Hickey, S., Totten, M., Agniel, D., ... et al. (2021). Trends in low-value health service use and spending in the US Medicare fee-for-

service program, 2014–2018. *JAMA Network Open*, 4(2), e2037328. doi: 10.1001/jamanetworkopen.2020.37328.

11. Barre-Sinoussi, F., Chermann, J. C., Rey, F., Nugeyre, M. T., Chamaret, S., Gruest, J., ... et al. (1983). Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science*, 220(4599), 868–871. doi: 10.1126/science.6189183.
12. Mellors, J. W., Munoz, A., Giorgi, J. V., Margolick, J. B., Tassoni, C. J., Gupta, P., ... et al. (1997). Plasma viral load and CD4+ lymphocytes as prognostic markers of HIV-1 infection. *Annals of Internal Medicine*, 126(12), 946–954. doi: 10.7326/0003-4819-126-12-199706150-00003.
13. The INSIGHT START Study Group. (2015). Initiation of antiretroviral therapy in early asymptomatic HIV infection. *New England Journal of Medicine*, 373(9), 795–807. doi: 10.1056/NEJMoa1506816.
14. Genberg, B. L., Shangani, S., Sabatino, K., Rachlis, B., Wachira, J., Braitstein, P., & Operario, D. (2016). Improving Engagement in the HIV Care Cascade: A Systematic Review of Interventions Involving People Living with HIV/AIDS as Peers. *AIDS and Behavior*, 20(10), 2452–2463. doi: 10.1007/s10461-016-1307-z.
15. Gardner, E. M., McLees, M. P., Steiner, J. F., Del Rio, C., & Burman, W. J. (2011). The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clinical Infectious Diseases*, 52(6), 793–800. doi: 10.1093/cid/ciq243.
16. Kranzer, K., Govindasamy, D., Ford, N., Johnston, V., & Lawn, S. D. (2012). Quantifying and addressing losses along the continuum of care for people living with HIV infection in sub-Saharan Africa: a systematic review. *Journal of the International AIDS Society*, 15(2), 17383. doi: 10.7448/IAS.15.2.17383.

17. Thompson, M. A., Mugavero, M. J., Amico, K. R., et al. (2012). Guidelines for improving entry into and retention in care and antiretroviral adherence for persons with HIV: evidence-based recommendations from an International Association of Physicians in AIDS Care panel. *Annals of Internal Medicine*, 156(11), 817–833. doi: 10.7326/0003-4819-156-11-201206050-00419.
18. Okeke, N. L., Ostermann, J., & Thielman, N. M. (2014). Enhancing linkage and retention in HIV care: a review of interventions for highly resourced and resource-poor settings. *Current HIV/AIDS Reports*, 11(4), 376–392. doi: 10.1007/s11904-014-0226-7.
19. Simoni, J. M., Franks, J. C., Lehavot, K., & Yard, S. S. (2011). Peer interventions to promote health: conceptual considerations. *American Journal of Orthopsychiatry*, 81(3), 351–359. doi: 10.1111/j.1939-0025.2011.01103.x.
20. Higgins, J. P. T., & Green, S. (2011). *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.1.0. The Cochrane Collaboration.
21. Thomas, B. H., Ciliska, D., Dobbins, M., & Micucci, S. (2004). A process for systematically reviewing the literature: Providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing*, 1(3), 176–184.
22. Hatcher, A. M., Turan, J. M., Leslie, H. H., et al. (2012). Predictors of Linkage to Care Following Community-Based HIV Counseling and Testing in Rural Kenya. *AIDS and Behavior*, 16(5), 1295–1307. doi: 10.1007/s10461-011-0056-4.
23. Richter, L., Rotheram-Borus, M. J., Van Heerden, A., et al. (2014). Pregnant women living with HIV (WLH) supported at clinics by peer WLH: a cluster randomized controlled trial. *AIDS and Behavior*, 18(4), 706–715. doi: 10.1007/s10461-014-0725-3.
24. Enriquez, M., Cheng, A. L., Banderas, J., et al. (2014). A Peer-Led HIV Medication Adherence Intervention Targeting Adults Linked to Medical Care but Without a Suppressed Viral Load. *Journal of the International Association of Providers of AIDS Care*, 14(5), 441–448. doi: 10.1177/2325957414548931.

25. Simoni, J. M., Huh, D., Frick, P. A., et al. (2009). Peer support and pager messaging to promote antiretroviral modifying therapy in Seattle: a randomized controlled trial. *Journal of Acquired Immune Deficiency Syndromes*, 52(4), 465–473. doi: 10.1097/QAI.0b013e3181bdc61e.
26. IOM (Institute of Medicine). (2008). *Methodological Challenges in Biomedical HIV Prevention Trials*. Washington, DC: The National Academies Press.
27. Koester, K. A., Morewitz, M., Pearson, C., et al. (2014). Patient navigation facilitates medical and social services engagement among HIV-infected individuals leaving jail and returning to the community. *AIDS Patient Care and STDs*, 28(2), 82–90. doi: 10.1089/apc.2013.0288.
28. Arem, H., Nakyanjo, N., Kagaayi, J., et al. (2011). Peer health workers and AIDS care in Rakai, Uganda: a mixed methods operations research evaluation of a cluster-randomized trial. *AIDS Patient Care and STDs*, 25(12), 719–724. doi: 10.1089/apc.2011.0142.
29. Buchacz, K., Chen, M. J., Parisi, M. K., Yoshida-Cervantes, M., Antunez, E., Delgado, V., Moss, N. J., & Scheer, S. (2015). Using HIV surveillance registry data to re-link persons to care: the RSVP project in San Francisco. *PLOS ONE*, 10(3), e0118923.
30. Centers for Disease Control and Prevention. (2015). HIV infection: detection, counseling, and referral.
31. Skarbinski, J., Rosenberg, E., Paz-Bailey, G., Hall, H. I., Rose, C. E., Viall, A. H., Fagan, J. L., Lansky, A., & Mermin, J. H. (2015). Human immunodeficiency virus transmission at each step of the care continuum in the United States. *JAMA Internal Medicine*, 175(4), 588–596.
32. International Association of Providers of AIDS Care. (2016). News release: fast-track cities unveil 90–90–90 and HIV care continuum data at AIDS 2016 demonstrating progress towards goal of ending AIDS as a public health threat by 2030.

33. Genberg, B. L., Shangani, S., Sabatino, K., Rachlis, B., Wachira, J., Braitstein, P., & Operario, D. (2016). Improving Engagement in the HIV Care Cascade: A Systematic Review of Interventions Involving People Living with HIV/AIDS as Peers. *AIDS and Behavior*, 20(10), 2452-2463. doi: 10.1007/s10461-016-1307-z. PMID: 26837630; PMCID: PMC4970970.

UNDER PEER REVIEW