

Varietal effect of different nitrogen levels on yield attributing characters and quality parameters of wheat (*Triticumaestivum* L.) crop

Comment [I1]: Please rephrase the title.

Abstract

The field experiment was conducted at Agronomy Research Farm, Narendra Deva University of Agriculture and Technology, Narendra Nagar (Kumarganj) Faizabad (U.P.), during Rabi season of 2014-2015. The yield components like number of spikes m^{-2} , spike length (cm), number of grain spike $^{-1}$, grain weight spike $^{-1}$ (g), grain yield ($q\ ha^{-1}$), straw yield ($kg\ ha^{-1}$) and nitrogen content in grain (%), nitrogen content in straw (%), nitrogen uptake in grain ($kg\ ha^{-1}$), nitrogen uptake in straw, protein content in grain (%) were maximum under 160 $kg\ N\ ha^{-1}$ and among the varieties over PBW-373, NW-1014 being at par with HD-2327.

Comment [I2]: Please rephrase and clarify the sentence:
among the varieties over PBW-373, NW-1014 being at par with HD-2327

Key words- number of spikes m^{-2} , spike length (cm), number of grain spike $^{-1}$, grain weight spike $^{-1}$ (g), Test weight (1000-grain weight), N content in grain (%), N content in straw (%), Protein content in grain (%).

Introduction

Wheat (*Triticumaestivum* L.) belongs to family Poaceae, is a staple food of the world. India is one of the principle wheat producing and consuming country in the world. Its importance in Indian agriculture is second after rice. About 55% of the world population depends on wheat for intake of about 20% of food calories. Globally, according to USDA (2014-15) reported that wheat is grown in the world with an area of 220.88 million hectares, production of 725.47 million tonnes with a productivity of 3.28 tonnes per hectare. In India it is grown in an area of 30.47 million hectares, production of 95.85 million tonnes with a productivity of 3.15 tonnes per hectare. About 91% of the total wheat production is contributed by six states viz., U.P., Punjab, Haryana, M.P., Rajasthan, and Bihar. Among them, Uttar Pradesh ranks first with respect to area 9.2mha and production 24.50mt with a productivity of 2.7 tonnes per hectare respectively. In India numbers of wheat varieties are cultivated but due to fast changing ecosystem these varieties become susceptible to different insects, pests and diseases which

Comment [I3]: Please rephrase and clarify the sentence:
Wheat (*Triticumaestivum* L.) belongs to family Poaceae, is a staple food of the world

Comment [I4]: countries

Comment [I5]: world's

Comment [I6]: number

cause a decline in yield. It was thus decided to generate a steady flow of new varieties, deriving resistant from diverse sources, to replace the old varieties for sustainable higher production.

Comment [I7]: resistance

Wheat revolution was used with the introduction of high yielding dwarf varieties which need high level of management.

Comment [I8]: ahigh

Material and methods

The experiment was laid out in randomized block design with four varieties (PBW-373, HD-2327 and NW-1014) and six nitrogen levels (0, 40, 80, 120, 160 and 200 kg N ha⁻¹) with three replications. Experimental design Randomized Block Design (Factorial), replication 03, varieties 03, nitrogen levels 06, treatment combinations 18.

Comment [I9]: It is preferable to include the paragraph below under the heading of statistical design, along with the test used to compare averages. Please add soil texture and its nutrient content. Please add the type of fertilizer used and the source of nitrogen. Please add the location where the experiment was conducted, along with site coordinates.

Number of spikes per meter square:

Number of spikes per meter square was counted before harvesting of crop at three places and average value was taken and expressed in number of spikes per metre square.

Comment [I10]: Only three varieties are mentioned in the table! Please explain the statement.

Spike length (cm):

Length of five selected ears from each plot as measured carefully from the neck node to the tip of last grain and averaged out to get the length of single ear.

Comment [I11]: Please rephrase the sentence. It was mentioned previously:

Experimental design Randomized Block Design (Factorial), replication 03, varieties 03, nitrogen levels 06, treatment combinations 18

Number of grains per spike:

Three randomly selected spikes were threshed and seed were counted and averaged to get the number of grains spikes.

Comment [I12]: anaverage

Comment [I13]: anumber

Comment [I14]: was

Comment [I15]: seeds

Grain weight spike⁻¹ (g):

The total number of grain weight of five selected spikes was measured and averaged to get the grain weight per spike and expressed in grams.

Comment [I16]: grains per spike

1000-grain weight (g):

One thousand grains from net plot were counted and weighed to get 1000 grains weight (g).

Comment [I17]: thenet

Nitrogen uptake by crop:

First of all, thoroughly powdered grain and straw of wheat from each plot were analysed by modified micro-Kjeldahls methods, as given by Jackson (1967).

Comment [I18]: Jackson et al.

Nitrogen content in grain and straw thus obtained was multiplied separately with respective yields. The values of both were expressed as total nitrogen uptake by the crop in kg ha⁻¹.

$$\text{Nitrogen uptake by grain (kg ha}^{-1}\text{)} = \frac{\text{N content in grain (\%)} \times \text{Grain yield (kg ha}^{-1}\text{)}}{100}$$

$$\text{Nitrogen uptake by straw (kg ha}^{-1}\text{)} = \frac{\text{N content in straw (\%)} \times \text{Straw yield (kg ha}^{-1}\text{)}}{100}$$

Protein content in grain:

Total Protein content in grain as estimated multiplying the nitrogen content with a factor 6.25.

Result and Discussion

The effect of treatments on growth and yield of crop, the influence of soil and climate on crop yield is also concerned, which cannot be ignored. Therefore, the existing edaphic and weather conditions of the experimental area were also taken into consideration prior to discussing the findings of treatments obtained during the investigation. Yield was the resultant of co-ordinated interplay of yield attributes. Vigorously growing plants are able to absorb larger quantity of mineral nutrients through well-developed root system. The variety NW-1014 gave higher number of spikes m⁻², length of spike, number of grain spike⁻¹ and 1000 grain weight than other varieties. It might be due to the genetic character of the variety like more reproductive tillers producing capacity, more spike length etc. Minimum yield contributing characters were credited to PBW-373. It was due to less reproductive tillers, less spike length as well as less number of grains spike⁻¹. The results were in conformity with those of Singh and Singh (1991) and Brijkishor (1998).

Comment [I19]: The

Comment [I20]: a concerned

Comment [I21]: into

Comment [I22]: result

Comment [I23]: a well-developed

Comment [I24]: a higher

The highest grain and straw yield was credited to variety NW-1014 followed by variety HD-232. The reason behind this may be because of good plant stand, more number of spike

bearing tillers, long shoots head and more number of grains spike⁻¹ with more test weight. Minimum grain and straw yield recorded with variety PBW-373 might be due to less number of spike bearing tillers, small shoots head and less number of grains spike⁻¹ and poor grain development. The results obtained in the present investigation in accordance with those obtained by Singh (1998) and Sardana *et al.* (1999).

Comment [I25]: heads

Comment [I26]: are in

number of effective shoots, spike length, number of grains spike⁻¹, grain weight spike⁻¹, test weight (1000-grain weight) increased with increase in nitrogen levels. The significant increase of these characters obtained only up to 160 kg N ha⁻¹ (Table 1). This might be due to enhanced shoots, photosynthetic area and increased sink size in presence of adequate nitrogen. Similar research findings were reported by Singh and Singh (1989), Patel *et al.* (1994) and Patel *et al.* (1995).

Comment [I27]: thepresence

Interaction effect between varieties and nitrogen levels was not found significant. This might be due to the fact that short statured high yielding wheat varieties performed equally good to optimum level of nitrogen. The similar findings were reported by Singh and Uttam (1992) and Singh(1998).

Comment [I28]: short-statured

Comment [I29]: nitrogen. Similar

UNDER PEER REVIEW

Table 1: Different Yield attributing characters of wheat crop

Treatments	Number of spikes (m ⁻²)	spike length (cm)	No. of grains spike ⁻¹	Grain weight spike ⁻¹ (g)	Test weight (1000-grain weight)	N content in grain (%)	N content in straw (%)	Protein content in grain (%)
(A) Varieties								
PBW-373	280.04	10.66	49.17	1.52	35.11	1.80	0.50	11.25
HD-2327	303.88	10.77	49.67	1.64	35.25	1.78	0.49	11.14
NW-1014	309.85	11.20	51.67	1.66	36.00	1.84	0.51	11.44
SEm±	4.19	0.14	0.66	0.03	0.48	0.01	0.00	0.075
CD (P=0.05)	12.05	0.42	1.90	0.1	1.39	0.03	0.01	NS
Nitrogen levels (kg ha ⁻¹)								
0	262.17	9.78	45.50	1.00	34.50	1.70	0.46	10.65
40	281.13	10.52	48.50	1.40	35.07	1.75	0.49	10.95
80	297.92	10.78	49.50	1.62	35.57	1.79	0.50	11.18
120	310.80	11.40	52.50	1.87	35.98	1.83	0.51	11.45
160	321.75	11.60	53.50	1.96	35.99	1.87	0.52	11.68
200	321.80	11.62	53.60	1.98	35.61	1.89	0.53	11.83
SEm±	5.93	0.20	0.93	0.05	0.68	0.01	0.005	0.405
CD (P=0.05)	17.04	0.60	2.68	0.1	1.96	0.05	0.01	NS

Conclusion

Yield attributes viz. number of spikes per square meter, length of spike, number of grain spike⁻¹ and weight of grains spike⁻¹ were significantly higher with variety NW-1014 followed by variety HD2327. 1000-grain weight was not influenced significantly by different varieties. number of spikes per square meter, length of spike, number of grains spike⁻¹ and weight of grains spike⁻¹ were significantly higher with 160 kg N ha⁻¹ being at par with 120 and 200 kg N ha⁻¹ while significantly superior over rest of the nitrogen levels. 1000-grain weights were significantly higher with 160 kg N ha⁻¹ being at par with 120 and 200 kg N ha⁻¹ while significantly superior over rest of the nitrogen levels. Protein content in grain was not influenced significantly with nitrogen levels. Among the varieties NW-1014 was found most appropriate for eastern plain zone

Comment [I30]: therest

Comment [I31]: significantly by nitrogen

Comment [I32]: theeastern

References

Anonymous (1999). The Hindu Survey of Indian Agriculture. Printed & Published at the National Press, Chennai.

Brijkishor (1998). To assess the performance of newly developed strains of wheat under zero tillage condition with varying nitrogen levels. M.Sc. (Ag.) thesis submitted to N.D.U.A & T., Kumarganj, Faizabad.

Jackson, Kjeldahls; Pande, I.B.; Dwivedi, D.K. and Prakash S.C. (1967). Impact of method and level of fertilizer application and weed management on nutrient economy and yield of wheat (*Triticumaestivum*). *Indian J. Agron.* **51**(3):193-198.

Patel, N.R.; Thaker, K.R. and Barevadia, T.N. (1994). Influence of different proportions of organic and inorganic nitrogen on irrigated wheat (*Triticumaestivum*). *Indian J. Agron.* **39**(4):2641-643.

Patel, N.M.; Sadaria, S.G.; Keneria, B.B. and Khanpara, V.D. (1995). Effect of nitrogen, potassium and zinc on growth and yield of wheat. *Indian J. Agron.* **40**(2):290-292.

Sardana, V.; Sharma, S.K. and Randhawa, A.S. (1999). Performance of wheat cultivars under different sowing dates and levels of nitrogen under rainfed conditions. *Ann. Agri. Res.* **20**(1): 60-63.

Singh, B. (1998). To assess the performance of new wheat varieties under late-sown condition with different nitrogen levels. M.Sc.(Ag) thesis submitted to N.D.U.A.&T., Kumarganj, Faizabad.

Singh, B. And Singh, G. (1998). Effect of nitrogen on wheat cultivars under late-sown conditions. *Indian J. Agron.* **34**(3):387-389.

Singh, S.P. and Singh, H.B. (1991). Effect of irrigation time and nitrogen level on wheat (*Triticumaestivum*) under late-sown condition of western UP. *Indian J. Agron.* **36**:41-42.

Comment [133]: It does not appear in the text.

Singh, V.P.N. and Uttam, S.K. (1992). Response of wheat cultivars to different nitrogen levels under early sown conditions. *Crop Res.*5:82-86.

UNDER PEER REVIEW