

# Effect of Designer Seeds and Fertilizer Micro-dosing on Productivity of Cowpea

## ABSTRACT

An experiment was conducted at the Instructional Farm of the College of Agriculture, Kerala Agricultural University(KAU), Vellanikkara, Thrissur from October to December 2021. The objective of the study was to assess the effect of fertilizer **micro-dosing** and designer seeds on the growth and yield of cowpea (*Vigna unguiculata* (L.) Walp). The experiment was laid out in Completely Randomized Design (Factorial), with 3 replications. The treatments consisted of five levels of fertilizer **micro-dosing** (micro-dose 25%, 50%, 75% of NPK as per package of practices (POP) of KAU, NPK as per POP of KAU and absolute control) and three levels of seed treatments (designer seeds, rhizobium treated seeds, and untreated seeds). The results of the study indicated that, micro-dose and designer seeds significantly increased the **biometric and yield parameters** of cowpea. Among fertilizer micro-dosing, micro-dose (50 %) showed minimum days (39.78 days) to 50% flowering which was on par with micro-dose (25 %) (39.89 days). Designer seeds showed a minimum of days to 50% flowering (37.27 days) as compared to other seed treatments. The combination of micro-dose (50%) along with designer seeds emerged as the most potent, yielding the highest pod weight per plant and seed yield (43.80g and 969.87 **kg/ha**, respectively).

*Keywords: Designer seeds, fertilizer micro-dosing, biometric **parameters**, seed yield*

## 1. INTRODUCTION

Cowpea, *Vigna unguiculata* (L.) Walp, is one of the most important grain legumes which is widely cultivated in the semi-arid tropics and **subtropics**, that contributes significantly to the nutritional security of the country. Cowpea yield remains low (less than 1 **t/ha**) in most areas of India (Justin, *et al.*, 2015) [5]. It is highly essential to increase the **yield** of cowpea. Application of fertilizers in micro-doses permits more precise and better timed fertilizer placement and proper seed treatment techniques enhance vigour and viability. These techniques can be adopted to increase the productivity and nutrient use efficiency of cowpea.

Mineral fertilizers have been recognized as an entry point for a sustainable agriculture production system. However, overdosing on mineral fertilizers has several harmful effects. Applying fertilizers in micro-doses permits more precise and better timed fertilizer placement and appropriate management of fertilizers. According to Tsegay *et al.* (2019) [14], fertilizer micro-dosing lowers input costs while increasing production and efficiency of the usage of fertilizer. Fertilizer micro-dosing technology consists of the application of a small quantity of mineral fertilizer together with seeds of the target crop in the planting hole at sowing or 3 to 4 weeks after planting (ICRISAT, 2009) [4].

Seed fortification through designer seed technology increases seed production and quality which can be achieved by treating the seeds with nutrient solution, insecticides, bioagents and polymers. Designer seed is an integrated pre-sowing seed treatment that involves the addition of nutrients, plant protectants, and bioinoculants to enhance seed quality, field emergence and yield parameters (Sujatha and Ambika, 2018) [12].

There is a felt need to increase the productivity of cowpea with limited resources to achieve nutrient security for the growing population. Resource conservation technologies such as designer seeds and fertilizer micro-dosing, to increase productivity and nutrient use efficiency are lacking. Hence, with this background, the present study was undertaken on cowpea with designer seed technology and fertilizer micro-dosing to maximize its productivity.

## 2. MATERIALS AND METHODS

The study was conducted at Instructional Farm, College of Agriculture, Kerala Agricultural University (KAU), Vellanikkara, Thrissur (India) from October to December, 2021. The experiment was laid out in Completely Randomized Design (Factorial), with 3 replications. Cowpea variety Anaswara was used for the study. The treatments consisted of five levels of fertilizer micro-dosing (micro-dose 25%, 50%, 75% of NPK as per POP of KAU, NPK as per POP of KAU and absolute control) and three levels of seed treatments (Designer seeds, rhizobium treated seeds, and untreated seeds). Designer seeds were prepared by soaking seeds in a micronutrient solution (KAU sampoorna micronutrient mix) of 100ppm for 3 hours and drying back to original moisture content and then treating the seeds with polymer @ 3ml/kg of seed, imidacloprid @ 2ml/kg of seed, PGPR (plant growth promoting rhizobacteria) mix I @ 4g/kg of seed and rhizobium @ 20g/kg of seed. Rhizobium treated seeds were prepared by mixing rhizobium culture @ 20g/kg of seed along with rice gruel and shade dried for a few hours. For fertilizer micro-dosing, fertilizers were applied along with seeds at sowing, and top dressing was applied as hill placement. Conventional fertilization was done as basal placement. The growth parameters viz., plant height, number of branches and dry matter content per plant were recorded at 20 days interval. The number of days from sowing to first flower bud appearance of 50% of the plants per treatment was recorded as days to 50% flowering. Yield parameters viz., pod weight per plant (g) and yield (kg/ha) were recorded at harvest. The experimental data on various growth and yield parameters of cowpea were subjected to analysis of variance using statistical package 'GRAPES (General R based Analysis Platform Empowered by Statistics)' developed by Gopinath *et al.* (2020). This platform is based on R software.

### 3. RESULTS AND DISCUSSION

The highest plant height, number of branches, and dry matter content per plant (54.90 cm, 7.00, and 28.15g) were recorded for micro-dose (50%) (Table 1). When compared to broadcasting, fertilizer micro-dosing increased the lateral root density of maize in the topsoil by 72% and 40% at corresponding lateral distances of 25 cm and 50 cm (Okebalama, *et al.*, 2017)[6]. Abdalla *et al.* (2015)[1] reported that, micro-dosing of phosphatic fertilizer improves plant height by 8 % and 3 % in sorghum and sesame respectively.

Among seed treatments, designer seeds showed the highest plant height, number of branches and dry matter content per plant (53.09cm, 6.13 and 23.90g respectively) (Table 1). Pelleting materials in designer seeds supply early micro and macronutrient supplements to plants during their early establishment. This could enhance the root and shoot growth in plants, which enabled the plants to absorb water and nutrients more effectively and grow more luxuriantly (Sujatha and Ambika, 2018)[11]. The KAU sampoorna micronutrient mix, consisting of essential elements such as Mg, S, B, Zn, Cu, Fe, Mn, and Mo was utilized as the micronutrient solution for the designer seed production. These elements are critical for various physiological functions in plants. Zinc, for instance, is an essential component of many enzymes, serving as a functional, structural, and regulatory cofactor. Copper acts as an enzyme activator and plays a crucial role in the absorption of nitrogen compounds, indirectly promoting chlorophyll production and increasing sugar content. Manganese and Zinc regulate protein metabolism and influence protein biosynthesis by adjusting the activity of peptidases. Iron promotes chlorophyll formation and enzyme mechanisms that work with cell respiratory systems, playing a role in cell growth and division. The synergistic effects of these elements were vital in the overall functioning and growth of cowpea (Saquee *et al.*, 2023)[9].

Among fertilizer micro-dosing, micro-dose (50 %) showed minimum days (39.78 days) to 50% flowering which was on par with micro-dose (25 %) (39.89 days) (Table 1). Through fertilizer micro-dosing tiny amounts of critical nutrients were added to deficient soil to help root systems to grow and absorb more water, which increases growth and development. Designer seeds showed a minimum of days to 50% flowering (37.27 days) as compared to other seed treatments (Table 1). The polymer coating and nutrients of designer seeds could account for the beneficial effect observed in the field.

This effect might be due to the activation of cells, which leads to a boost in mitochondrial activity. This process creates more high energy compounds and vital biomolecules that are available during the early stages of germination, thereby improving field emergence and early flowering (Seema *et al.*, 2020)[10].

**Table 1. Effect of treatments on plant height, number of branches, drymatter content per plant at harvest and days to 50% flowering of cowpea**

The highest pod weight per plant (35.16 g) and seed yield (640.24 kg/ha) were found in micro-

Treatments	Plant height (cm)	Number of branches	Dry matter content per plant(g)	Days to 50 % flowering
<b>Fertilizer micro-dosing</b>				
25% micro-dose	43.69 <sup>c</sup>	4.89 <sup>b</sup>	18.86 <sup>b</sup>	39.89 <sup>c</sup>
50% micro-dose	54.90 <sup>a</sup>	7.00 <sup>a</sup>	28.15 <sup>a</sup>	39.78 <sup>c</sup>
75% micro-dose	42.37 <sup>c</sup>	3.89 <sup>c</sup>	17.20 <sup>c</sup>	40.67 <sup>ab</sup>
NPK as per POP	46.29 <sup>b</sup>	5.56 <sup>b</sup>	19.96 <sup>b</sup>	41.11 <sup>a</sup>
Absolute control	33.83 <sup>d</sup>	3.11 <sup>c</sup>	12.33 <sup>d</sup>	40.11 <sup>bc</sup>
CD (0.05)	2.33	0.88	1.28	0.62
SE(m)±	0.81	0.31	0.44	0.22
<b>Seed treatments</b>				
Designer seeds	53.09 <sup>a</sup>	6.13 <sup>a</sup>	23.90 <sup>a</sup>	37.27 <sup>b</sup>
Rhizobium treated seeds	43.59 <sup>b</sup>	4.86 <sup>b</sup>	18.65 <sup>b</sup>	41.73 <sup>a</sup>
.27Untreated seeds	35.97 <sup>c</sup>	3.67 <sup>c</sup>	15.27 <sup>c</sup>	41.93 <sup>a</sup>
CD (0.05)	1.81	0.68	0.99	0.49
SE(m)±	0.63	0.24	0.34	0.17

dose (50 %)(Table 2). Observational data suggests that crop yield enhanced when micro-dosing techniques are employed. A reduction of fertilizer application by 50% resulted in a 19% increase in seed yield. The practice of micro-dosing fertilizer enables plants to receive precise amounts of nutrients, minimizing wastage and optimizing uptake. This targeted approach ensures balanced growth, avoiding nutrient imbalances that are caused from excess or deficiency. As a result, nutrient efficiency is enhanced, leading to healthier plants and overall improved yield. According to Ibrahim, *et al.* (2015)[3], lateral root growth within the topsoil allows for better nutrient utilization, which in turn leads to a good effect of fertilizer micro-dosing in improving millet yield. Among seed treatments, designer seeds showed the highest pod weight per plant (36.56g) and seed yield (683.43 kg/ha). Designer seed treatment resulted in 38% and 114 % increase in yield as compared to rhizobium treated seeds and untreated seeds respectively. The inclusion of inoculants, protectants, nutrients, and polymer in the coating treatment may have contributed to the yield increase. Bioinoculants used for seed treatment were also contributed in plant growth by the solubilization and sequestration of various plant nutrients, and their subsequent delivery to the plants (Dinesh *et al.*, 2018)[2].

**Table 2. Effect of treatments on pod weight per plant and seed yield at harvest**

Treatments	Pod weight per plant (g)	Seed yield (kg/ha)
<b>Fertilizer micro-dosing</b>		
25% micro-dose	33.59 <sup>b</sup>	579.16 <sup>b</sup>
50% micro-dose	35.16 <sup>a</sup>	640.24 <sup>a</sup>
75% micro-dose	28.53 <sup>d</sup>	482.73 <sup>d</sup>
NPK as per POP	31.01 <sup>c</sup>	536.49 <sup>c</sup>
Absolute control	21.53	258.25 <sup>e</sup>
CD (0.05)	1.41	15.28
SE(m)±	0.49	5.29
<b>Seed treatments</b>		
Designer seeds	36.56 <sup>a</sup>	683.43 <sup>a</sup>
Rhizobium treated seeds	28.75 <sup>b</sup>	494.88 <sup>b</sup>
Untreated seeds	24.59 <sup>c</sup>	319.82 <sup>c</sup>
CD (0.05)	1.09	11.83
SE(m)±	0.38	4.09

The results of the experiment showed that the treatment combination of micro-dose (50 %) with designer seeds (T4) yielded the most significant positive effects on plant height (66.70 cm), number of branches (8.00), and dry matter content per plant (36.11g) (Table 3). On the other hand, the lowest values were observed in absolute control with untreated seeds, with plant height, number of branches, and dry matter content per plant measuring 30.40cm, 2.33, and 10.53g, respectively (Table 3). Treatment combinations T1 (micro-dose (25%) + designer seed), T4 (micro-dose (50 %) + designer seeds) and T13 (absolute control + designer seed) were recorded minimum days (35.67 days) for 50% flowering compared to absolute control with untreated seeds (42.33 days) (Table 3). According to Rakotoson *et al.*, (2021)[8], vigorous growth, early flowering and accumulation of nutrients was observed in rice seedlings under fertilizer micro-dosing. Designer seeds of black gram showed increased plant growth, vigour and took minimum days for 50 per cent flowering as compared to untreated control (Sujatha and Ambika, 2016)[12]. Thus, it was found that the micro-dose (50 %) and designer seed combination performed better in terms of biometrics and early flowering.

**Table 3. Interaction effect of treatments on plant height, number of branches, dry matter content per plant at harvest and days to 50% flowering of cowpea**

Treatments		Plant height (cm)	Number of branches	Dry matter content per plant (g)	Days to 50% flowering
T1	25% micro-dose + designer seed	52.43 <sup>b</sup>	6.33 <sup>c</sup>	25.48 <sup>b</sup>	35.67 <sup>c</sup>
T2	25% micro-dose + rhizobium treated	42.23 <sup>cd</sup>	5.00 <sup>de</sup>	18.13 <sup>f</sup>	41.67 <sup>a</sup>
T3	25% micro-dose + untreated seed	35.40 <sup>fg</sup>	3.33 <sup>fg</sup>	15.33 <sup>gh</sup>	42.00 <sup>a</sup>
T4	50% micro-dose + designer seed	66.70 <sup>a</sup>	8.00 <sup>a</sup>	36.11 <sup>a</sup>	35.67 <sup>c</sup>
T5	50% micro-dose + rhizobium treated	53.60 <sup>b</sup>	6.67 <sup>b</sup>	21.40 <sup>de</sup>	41.67 <sup>a</sup>

T6	50% micro-dose + untreated seed	44.40 <sup>cd</sup>	5.33 <sup>cd</sup>	20.67 <sup>e</sup>	42.33 <sup>a</sup>
T7	75% micro-dose + designer seed	51.93 <sup>b</sup>	5.00 <sup>de</sup>	23.10 <sup>d</sup>	39.33 <sup>b</sup>
T8	75% micro-dose + rhizobium treated	41.80 <sup>de</sup>	4.00 <sup>fg</sup>	18.10 <sup>f</sup>	41.33 <sup>a</sup>
T9	75% micro-dose + untreated seed	33.37 <sup>gh</sup>	2.67 <sup>ij</sup>	16.43 <sup>fg</sup>	41.33 <sup>a</sup>
T10	NPK as per POP+ designer seed	55.53 <sup>b</sup>	6.67 <sup>b</sup>	27.67 <sup>b</sup>	40.00 <sup>b</sup>
T11	NPK as per POP + rhizobium treated	47.03 <sup>c</sup>	5.33 <sup>cd</sup>	16.70 <sup>fg</sup>	41.67 <sup>a</sup>
T12	NPK as per POP+ untreated seed	36.30 <sup>fg</sup>	4.67 <sup>ef</sup>	13.50 <sup>hi</sup>	41.67 <sup>a</sup>
T13	Absolute control + designer seed	38.83 <sup>ef</sup>	3.67 <sup>gh</sup>	13.57 <sup>hi</sup>	35.67 <sup>c</sup>
T14	Absolute control + rhizobium treated	32.27 <sup>gh</sup>	3.33 <sup>hi</sup>	12.63 <sup>ij</sup>	42.33 <sup>a</sup>
T15	Absolute control + untreated seed	30.40 <sup>h</sup>	2.33 <sup>jk</sup>	10.53 <sup>j</sup>	42.33 <sup>a</sup>
	CD (0.05)	4.04	1.31	2.21	1.08
	SE(m)±	1.40	0.41	0.76	0.38

The highest yield **parameters** viz., pod weight per plant (44.55g) and seed yield (969.87kg) were found in the treatment combination, 50% micro-dose with designer seed treatment (T4) (Table 4). The seed yield recorded by NPK as per POP with designer seeds (T10) was 736.60 kg/ha. There was a 32 % improvement in yield when fertilizer application was reduced by 50%. Fertilizer micro-dosing led to a significant rise in the leaf area index and leaf chlorophyll content in millets ( Ibrahim, *et al.*, 2015)[3]. These could improve photosynthesis and lead to increased production when fertilizer is micro-dosed. Tovihoudjiet *al.* (2017)[13] also reported increased productivity by 50 % in maize under fertilizer micro-dosing.

The surge in yield in designer seeds may be attributed to the improved photosynthetic efficiency, which stabilizes chlorophyll and increases the production of photosynthates. Consequently, organic material is transported more efficiently from the source to the sink in the treated plants. The application of coating treatment containing inoculants, protectants, nutrients, and polymers also appears to have contributed to the observed yield increase. Plant development and yield were enhanced by biologically active products and these products facilitate biological nitrogen fixation, phosphate mobilization, and nutrient uptake (Ponnuswamy and Vijayalakshmi., 2011)[7]. Similar findings are also reported in paddy, and black gram where, polymer coating accelerated the uptake of water, gasses, minerals, and hormones that promote early panicle emergence, faster development, and swift seedling emergence (Sujatha and Ambika, 2018)[11].

**Table 4. Effect of treatments on pod weight per plant and seed yield of cowpea at**

Treatments		Pod weight per plant (g)	Seed yield (kg/ha)
T1	25% micro-dose + designer seed	41.06 <sup>b</sup>	842.67 <sup>b</sup>
T2	25% micro-dose + rhizobium treated	29.62 <sup>ef</sup>	559.10 <sup>e</sup>
T3	25% micro-dose + untreated seed	25.63 <sup>h</sup>	313.77 <sup>h</sup>
T4	50% micro-dose + designer seed	44.55 <sup>a</sup>	969.87 <sup>a</sup>
T5	50% micro-dose + rhizobium treated	34.23 <sup>d</sup>	596.26 <sup>d</sup>
T6	50% micro-dose + untreated seed	26.70 <sup>gh</sup>	364.10 <sup>g</sup>
T7	75% micro-dose + designer seed	34.15 <sup>d</sup>	562.17 <sup>e</sup>
T8	75% micro-dose + rhizobium treated	26.92 <sup>gh</sup>	513.70 <sup>f</sup>
T9	75% micro-dose + untreated seed	24.53 <sup>h</sup>	372.33 <sup>g</sup>
T10	NPK as per POP+ designer seed	37.78 <sup>c</sup>	736.60 <sup>c</sup>
T11	NPK as per POP + rhizobium treated	31.59 <sup>e</sup>	559.60 <sup>e</sup>
T12	NPK as per POP+ untreated seeds	28.14 <sup>fg</sup>	335.70 <sup>h</sup>
T13	Absolute control + designer seed	25.27 <sup>h</sup>	315.33 <sup>h</sup>
T14	Absolute control + rhizobium treated	21.37 <sup>i</sup>	246.23 <sup>i</sup>
T15	Absolute control + untreated seed	17.97 <sup>j</sup>	213.20 <sup>j</sup>
	CD (0.05)	2.45	26.47
	SE(m)±	0.85	9.16

**harvest**

#### **4. CONCLUSION**

The study results indicate that, the application of fertilizers at micro dose (50%) significantly increased cowpea yield by 148%, compared to absolute control. Furthermore, the implementation of designer seeds led to a 114% increase in cowpea yield compared to untreated seeds. The combination of microdose (50 %) with designer seeds (T4) proved to be the most effective, showing superior performance in both biometric and yield parameters. Notably, a substantial 189% increase in yield was recorded compared to the application of NPK as per POP of KAU with untreated seeds (T12).

#### **ACKNOWLEDGEMENTS**

The authors are highly thankful to the College of Agriculture, Vellanikkara, Thrissur, Kerala for providing all assistance and research material.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **REFERENCES**

1. Abdalla EA, Osman AK, Maki MA, Nur FM, Ali SB, Aune, JB. The response of sorghum, groundnut, sesame, and cowpea to seed priming and fertilizer micro-dosing in South Kordofan State, Sudan. *Agron.* 2015; 5(4): 476-490.
2. Dinesh R, Biraris Eugenia P, Lal. Integration of biofertilizers with in-organic fertilizers and zinc for growth, yeild and biochemical parameters of sweet corn. *Int. J. Chem. Stud.* 2018; 6(5):705-709.
3. Gopinath, P. P, Parsad, R, Joseph, B, Adarsh, V. S. GRAPES: General Rshiny Based Analysis Platform Empowered by Statistics. 2020.<https://www.kaugrapes.com/home>. version 1.0.0. DOI: 10.5281/zenodo.4923220.
4. Ibrahim A, Abaidoo RC, Fatondji D, Opoku A. Hill placement of manure and fertilizer micro-dosing improves yield and water use efficiency in the Sahelian low input millet-based cropping system. *Field Crops Res.* 2015; 180: 29-36.
5. ICRISAT [International Crops Research Institute for the Semi-Arid Tropics]. *Fertilizer Micro-dosing Boosting Production in Unproductive Lands* [online]. Available: <http://www.icrisat.org/impacts-stories/icrisat-is-fertilizer-micro-dosing> 2009; [Access date: 04/November2023]
6. Justin CGL, Anandhi P, Jawahar D. Cataloguing, screening and assessing the effect of sowing time on the incidence of black gram pests under dryland condition. *Cogent Food and Agric.* 2015; 1(1): 1022632.
7. Okebalama CB, Ibrahim A, Safo EY, Yeboah E, Abaidoo RC, Logah V, Monica UF. Fertilizer micro-dosing in West Africa low-input cereals cropping : benefits, challenges and improvements strategies. *Afr. J. Agric. Res.* 2017; 12(14):1169-1176.
8. Ponnuswamy AS, Vijayalakshmi V. Effect of seed fortification with Bio -inoculants, nutrients and growth regulators on seed germination and seedling vigour of tomato (*Lycopersicon esculentum*), Brinjal (*Solanum melongena*) and Chilli (*Capsicum annum*). *Madras Agric. J.* 2011; 98: 251-252
9. Rakotoson T, Rinasoa S, Andriantsiorimanana A, Razafimanantsoa MP, Razafimbelo T, Rabeharisoa L, Tsujimoto Y. and Wissuwa M. Effects of fertilizer micro-dosing in nursery on rice productivity in Madagascar. *Plant Prod. Sci.* 2021; 24(2):170-179.
10. Saquee FS, Diakite S, Kavhiza NJ, Pakina E, Zargar, M. The Efficacy of Micronutrient Fertilizers on the Yield Formulation and Quality of Wheat Grains. *Agron.* 2023; 13(2): 566.
11. Seema AN, Koti VHMR, Basavaraja B. Augmenting the Productivity of Maize (*Zea Mays* L.) Through Designer Seed. *Pharma J.* 2023; 12(7): 1702-1708
12. Sujatha K, Ambika S. Designer seed treatment techniques on enhancement of yield in paddy. *Oryza* 2018; 55(4): 602-607.
13. Sujatha K, Ambika, S. Designer seed for enhancement of yield in black gram (*Vigna mungo* L.). *Indian J. Agric. Res.* 2016; 50(5):479-482.
14. Tovihoudji PG, Akponikpè PI, Agbossou EK, Bertin P, Biolders CL. Fertilizer micro-dosing enhances maize yields but may exacerbate nutrient mining in maize cropping systems in northern Benin. *Field Crops Res.* 2017; 213:130-142.
15. Tsegay A, Kiros G, Tesfay G. Gebregziabher Y. 2019. Comparative Analysis of Maize Yield and Economic Returns of Fertilizer Micro-dosing in Tigray, Northern Ethiopia. *Tigray Agricultural Research Institute (TARI) Agricultural Growth Program-II (AGP-II)* 2019; 7(1):177.