

Antimicrobial activity and GC-MS analysis of bioactive compounds of *Tribulusterrestris*L.

ABSTRACT

Aims: To examine the potential phytoconstituents present in leaves and fruits of *Tribulusterrestris* and its antimicrobial activity against *Enterobacter cloacae*.

Study design: Plant extracts preparation, determination of total phenolic content, total antioxidant activity and antimicrobial activity. GC-MS analysis

Place and Duration of Study: Sample: Fresh plant parts of *Tribulusterrestris* were collected from the campus of Banasthali Vidyapith, Rajasthan, gram-negative bacterial strain, *Enterobacter cloacae* BVDWH38 (Gene bank Acc. No. OQ772170) was used for antimicrobial study.

Methodology: Plant extracts preparation; free radical scavenging of sample extract antimicrobial activity was done by standard methods. All experiments were repeated at least thrice. The data represented are Mean \pm standard deviations (SD) of all the three replicates.

Results: Alcoholic and aqueous leaves and fruits extracts of *T. terrestris* were prepared. Total phenolic compound (TPC) and antioxidant activity was determined and found to be higher in aqueous extract of both plant parts of *T. terrestris*. Antimicrobial activity against water borne pathogenic *E. cloacae* strains was evaluated and found that aqueous extract of both leaves and fruits of *T. terrestris* was more effective. Gas chromatography- Mass spectrophotometer (GC-MS) with alcoholic extract of leaves and fruits of *T. terrestris* identified Digitoxin (6.37%) and Ethyl iso-allocholate (22.90%) respectively among other compounds.

Conclusion: The leaves and fruits of *T. terrestris* can be used as alternative medicine as it has vital bioactive secondary metabolites.

Keywords: Anti-microbial, Anti-oxidant, GC-MS, *Tribulusterrestris*, medicinal plant

1. INTRODUCTION

Fresh water plays an essential role in all living organisms for their survival and growth however an ever increasing population, industrialization, urbanization chemical fertilization associated with water-borne diseases [1]. Water associated infectious disease like acute gastrointestinal, cholera, diarrhea, dysentery, typhoid etc. diseases are principally caused by pathogenic microorganisms such as *Enterobacter* sp., *Enterococcus* sp., *Vibrio cholera*, *Escherichia coli*, *Giardiasis* sp., *Shigellosis* sp., *Salmonella* sp. and many more other bacterial pathogens [1, 2].

Enterobacter cloacae, the gram negative, rod-shaped bacteria, is majorly found in contaminated water, sewage, hospital environment as well as intestinal tracts of human and animals. *Enterobacter* species were rarely found as pathogens but these groups of bacteria progressively encountered and causing urinary tract infection (UTI) and bacteremia[3].

Medicinal plants play an important role in the widely available primary treatment of healthcare remedies. Herbal remedies are in high demand due to their ease of access, low cost, and safer use than modern pharmaceuticals[4]. *Tribulusterrestris* commonly known as 'Gokhru' grows in tropical climates and mostly found in Australia, India, and America. In traditional medicine, Gokhru extract is used to control cholesterol and blood pressure, as a tonic, aphrodisiac, analgesic, astringent, stomachic, anti-hypertensive, tonic, aphrodisiac, analgesic, diuretic lithon-triptic and urinaryinfection[5]. The fruits are known for their cooling diuretics and tonic properties, and they are used to treat painful micturition, UTI, and affections. Gokhru fruits have traditionally been used to treat kidney disease, gout, and as diuretics in Punjab. The leaves are used as an aperient and tonic in southern France and Europe [6].

The present work is conducted to evaluate the antimicrobial and antioxidant activity of ethanolic and aqueous extract of *T. terrestris* and characterize bio-active compounds by GC-MS analysis.

2. MATERIAL AND METHODS

2.1 Plant material

Fresh *Tribulusterrestris* plants were collected from the campus of BanasthaliVidyapith, Rajasthan, India and voucher specimen submitted and the herbarium (BURI 1731/2023). The leaves and fruit were properly washed and shade dried. Dried leaves and fruit powder were ground to fine powder and kept in airtight container at room temperature in the dark. Voucher specimen is submitted to the BanasthaliVidyapith Herbarium (BURI).



Figure 1. A twig and fruits of *Tribulusterrestris*

2.2 Preparation of Plant Extracts

20g of dried powder of *T. terrestris* leaves and fruits were stored for further use. The extraction was conducted by 250 ml of the solvent (95% ethanol, aqueous) for 72 hours, filtered by Whatmann No.41 filter paper and evaporated to Dryness at room temperature then stored the dried crude extract at 4°C[7].

2.3Determination of Total Phenolic Content (TPC)

Extraction of phenolic compounds was done [8,9].To determine the total phenolic concentration, freshly harvested leaves were used. Total phenolics were determined with Folin-ciocalteau reagent and expressed as "gallic acid (GA) equivalents" [10].

2.4Determination of antioxidant activity

Free radical scavenging of sample extract was determined using DPPH(1, 1 dihydroxy 2-picrylhydrazyl), a stable free radical [11].The total antioxidant activity of the sample extract was determined [11] and Hydrogen peroxide scavenging assay was conducted [12]. Ascorbic acid was used as standard.

2.5Determination of antimicrobial activity

A gram-negative bacterial strain, *Enterobacter cloacae* BVDWH38 (Gene bank Acc. No.OQ772170, was used for studying the antimicrobial activity by broth dilution method [13]. An aliquot of individual bacterial stock of *E. cloacae* was grown in 2ml liquid LB medium overnight at 37°C.The plant extracts (50-350µg/mL) were added to the bacterial culture. Bacteria were allowed to grow for 24h under standard conditions and absorbance taken at 540nm. MIC₅₀ value was determined [14].

2.6 GC-MS Analysis

The ethanolic extract was used for GC-MS analysis. The extract was filtered on a Whatmann no.41 filter paper, evaporated and dried using vacuum concentrator unit and stored at 4°C. GC-MS(Thermo Fischer) with HP-5MS capillary column (film thickness of 0.25µm; 30m X 250µm) fitted in a split/split less injection system was used. Samples of 1µl aliquot were injected automatically. The temperature variations were: 280°C (injector temperature); 50°C (initial oven temperature) rising up to 300°C at 25°C/min followed by 10 min hold. Carrier gas (Helium) was used with flow rate of 2.1 ml/min at 17.69 psi pressure.

2.7 Statistical analysis

All experiments were repeated at least thrice. The data represented are Mean ± standard deviations (SD) of all the three replicates. Difference between values are significantly (P=0.05) compared using LSD following ANOVA.

3. RESULTS AND DISCUSSION

3.1 Total phenolic content (TPC)

Maximum total phenolic content was observed in aqueous extract of both leaves and fruits of *T. terrestris*. Leaf aqueous extract of *T. terrestris* showed higher TPC value as compared to fruit aqueous extract of *T. terrestris*(Table 1).



Table 1. Total phenolic content of *T. terrestris* leaves and fruit's extracts

Extract/ plant part	Total phenolic content GA equivalent (mM g ⁻¹ fwt)	
	leaves	fruit
Alcoholic	0.762±0.016 ^b	0.557±0.018 ^a
Aqueous	1.098±0.011 ^b	0.857±0.022 ^a

Values followed by the same alphabet in a column are statistically not significantly different at P=0.05 following ANOVA and LSD

3.2 Antioxidant activity

High free radical scavenging activity and DPPH and phoshomolybdenum activity was observed in the aqueous of both plant parts followed by alcoholic extract. However, both alcoholic and aqueous leaves extract of *T. terrestris* showed higher antioxidant activity as compared to its fruits extract (Figure 2, 3, 4 and 5). Minimum H₂O₂ production was found in leaves and fruits aqueous followed by alcoholic extracts of *T. terrestris* (Figure 4.a and 4.b).






Figure 4. Phoshomolybdenum antioxidant activity of (a) alcoholic extract, (b) aqueous extract of *T. terrestris* leaves




Figure 5. Phoshomolybdenum antioxidant activity of (a) alcoholic extract, (b) aqueous extract *T. terrestris* fruit





Figure 6 Reducing power of (a) alcoholic extract, (b) aqueous extract of *T. terrestris* leaves



3.3 Antimicrobial activity

Antimicrobial activity of plant against *E. cloacae* strain was evaluated and compared with standard tetracycline. Antibacterial activities of *T. terrestris* fruit extracts were better in comparison to its leaves extracts.

Table 2 MIC₅₀ and MIC₉₀ in different extract of *T. terrestris* leaves and fruits against *E. cloacae*

Plant part	Extract	MIC ₅₀ (µg/mL)	MIC ₉₀ (µg/mL)
leaves	Aqueous	137.974	302.588
	Alcoholic	133.666	306.826
fruits	Aqueous	109.828	281.502
	Alcoholic	144.106	300.351
Tetracycline		69.83	

3.4 GC-MS Analysis

GC-MS chromatogram (Figure 8 and 9) revealed various vital compounds present in *T. terrestris* leaves and fruit and is represented in Table 3 with their medicinal properties.





Table 3.GC-MS analysis of Ethanolic extract of *T. terrestris* leaves

Compound Name	Molecular Formula	MW	RT (min)	Peak Area %	Compound Class	Bioactivity	Ref.
Digitoxin	C ₄₁ H ₆₄ O ₁₃	764.9	4.02	6.37	Glycoside	Treating Heart failure	[15,16]
2,6-Dihydroxyacetophenone, 2TMS derivative -	C ₁₄ H ₂₄ O ₃ Si ₂	296.5	4.47	4.38	Alkaloids	Antimicrobial, Anti-oxidant, Anti-inflammatory	[17]
Octadecane, 1,1'-[1,3-propanediylbis(oxy)]bis	C ₃₉ H ₈₀ O ₂	581.1	4.56	7.59	fatty acids	hypertension, musculoskeletal disorders, gastrointestinal distress, cough, amenorrhea	[18,19]
2-Trimethylsiloxy-6-hexadecenoic acid, methyl ester	C ₂₀ H ₄₀ O ₃ Si	356.6	5.37	13.82	fatty acids	urinary passage, lung and spleen diseases	[20]
2,7-Diphenyl-1,6-dioxypyridazino[4,5:2',3']pyrrolo[4',5'-d]pyridazine	C ₂₀ H ₁₃ N ₅ O ₂	355.3	7.43	13.75	Alkaloids	Antimicrobial Antibacteria, Antifungal and Anti-Diabetic	[21,22]
9-Octadecenoic acid, (2-phenyl-1,3-dioxolan-4-yl)methyl ester, cis	C ₂₈ H ₄₄ O ₄	444.6	8.00	5.76	fatty acids	anti-histaminic, anticoronary, insectifuge, anti-eczemic, anti-acne.	[23,24]
Glycine, N-[(3à,5á)-24-oxo-3-[(trimethylsilyloxy)cholan-24-yl]-, methyl ester	C ₃₀ H ₅₃ NO ₄ Si	519.8	13.12	11.77	Glycosides, fatty acid	Antimicrobial	[25]

Table 4.GC-MS analysis of Ethanolic fruit's extracts of *Tribulusterrestris*

Compound Name	Molecular Formula	MW	RT (min)	Peak Area %	Compound Class	Bioactivity	Ref.
Phosphonoacetic Acid, 3TMS derivative	C ₂ H ₅ O ₅ P	140.03	5.36	6.10	Secondary metabolites	Antiviral agent	[26]
Oleic Acid	C ₁₈ H ₃₄ O ₂	282.47	13.09	24.28	Fatty acid	Prevent heart disease, reduce cholesterol.	[25]
9,12-Octadecadienoyl chloride, (Z,Z)-	C ₁₈ H ₃₁ ClO	298.89	14.96	10.60	linoleoyl chloride	Antimicrobial, antioxidant and cytotoxic	[15,20]
Ethyl iso-allocholate	C ₂₆ H ₄₄ O ₅	436.6	27.28	22.90	Steroids derivative	Antimicrobial activity	[15,18]
Trilinolein	C ₅₇ H ₉₈ O ₆	879.4	28.66	13.32	linoleoyl chloride	Antimicrobial, Antioxidant, anti-ischemic	[27]

4. CONCLUSION

Tribulusterrestris plant extract contains polyphenolic compounds, antioxidants and antimicrobials. The presence of bioactive compounds in the ethanolic fractions as detected by GC-MS and their biological roles in *T. terrestris* support the ethnobotanical claims and medicinal value of this plant.

REFERENCES

1. Patil PN, Sawant DV, Deshmukh RN. Physico-chemical parameters for testing of water—A review. *Int J Environ Sci.* 2012;3(3):1194-1207.
2. Marale M, Gavali RS, Rao KR. Evaluation of water quality with waterborne diseases for assessing pilgrimage impact along river Indrayani, Pune (India). *Int J Environ Prot.* 2012;2:1-14
3. Cabral JP. Water microbiology, bacterial pathogens and water. *Int J Environ Res Public Health.* 2010;7(10):3657-3703.
4. Siddique MH, Andleeb R, Ashraf A, Zubair M, Fakhar-e-Alam M, Hayat S, Afzal M. Integration of *in silico* and *in vitro* approaches to evaluate antioxidant and anticancer properties of *Tribulusterrestris* extracts. *Arabian J Chem.* 2022;15(8):103984.
5. Sridhar K, Rajesh B, Sangeetha K. Phytochemical screening and GC-MS analysis of ethanolic extract of *Tribulusterrestris*. *Int J Pharmacol Toxicol.* 2016;6(1):44-50.
6. Abirami P, Rajendran A. GC-MS analysis of *Tribulusterrestris* L. *Asian J Plant Sci. Res.* 2011;1(4):13-16.
7. Widyaningrum I, Wibisono N, Kusumawati AH. Effect of extraction method on antimicrobial activity against *Staphylococcus aureus* of Tapakliman (*Elephantopus scaber* Linn.) leaves. *Int J Health Med Sci.* 2020;3(1):105-110.
8. Harborne JB. *Phytochemical Methods*, Chapman & Hall, London, UK. 1998.
9. Chakraborty D, Mandal, SM, Chakraborty J. Antimicrobial activities of leaf extract of *Basilicum polystachyon* (L) Moench. *Indian J Exp Biol.* 2007;45(8):744-748.
10. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods Enzymol.* 1999;299:152-178.
11. Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J.* 2013;21(2):143-152.
12. Ruch RJ, Cheng SJ, Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcino.* 1989;10(6):1003-1008.
13. Bais HP, Walker TS, Schweizer HP, Vivanco JM. Root specific elicitation and antimicrobial activity of rosmarinic acid in hairy root cultures of *Ocimum basilicum*. *Plant Physiol Biochem.* 2002;40(11):983-995.
14. Kambli P, Ajbani K, Sadani M, Nikam C, Shetty A, Udwadia Z, Rodrigues C. Correlating minimum inhibitory concentrations of ofloxacin and moxifloxacin with gyrA mutations using the genotype MTBDRsl assay. *Tuberculosis.* 2015;95(2):137-141.
15. Al-Gara NI, Abu-Serag NA, Shaheed KAA, Al Bahadly ZK. Analysis of bioactive phytochemical compound of (*Cyperus alternifolius* L.) By using gas chromatography–mass spectrometry. In IOP conference series: Mater. Sci. Eng. 2019;571(1):012047.
16. Melo P, Machado R, Teixeira HM. Analysis of digoxin and metildigoxin in whole blood using solid-phase extraction and liquid chromatography tandem mass spectrometry. *Int J Anal Chem.* 2012:975824.
17. Abdulkader OM, Sharaf AEA, Fouda HM, Elhaw MH. Phyto-biochemical constituents and GC-MS analysis of *Bidens pilosa* leaves and stems. *Egypt. Acad J Biol Sci H Bot.* 2022;13(2):1-9.

18. Karthikeyan K, Dhanapal CK. GC-MS analysis of ethyl acetate extract of *Alysicarpusmonilifer*-whole plant. *Der Pharm Lett.*2016;8(13):106-114.
19. Negm W, Ibrahim AR, Aboelsauod K, Ragab A, Attia GI. GC-MS analysis of petroleum ether extract and volatiles of *Cycas revolute*Thunb. growing in Egypt. *InventiRapid:PlantaActiva.* 2016(1):1-5.
20. Joshi UP, Wagh RD. GC-MS analysis of phytochemical compounds present in the bark extracts of *Ehretialaevis*Roxb. *Int J ResDevPharmLife Sci.*2018;7(6):3150-3154.
21. Brintha S, Rajesh S, Renuka R, Santhanakrishnan VP, Gnanam R. Phytochemical analysis and bioactivity prediction of compounds in methanolic extracts of *Curculigoorchioides*Gaertn. *JPharmacognPhytochem.*2017;6(4):192-197.
22. Altameme HJ, Hameed IH, Kareem MA. Analysis of alkaloid phytochemical compounds in the ethanolic extract of *Daturastramonium* and evaluation of antimicrobial activity. *Afr J Biotechnol.* 2015;14(19):1668-1674.
23. Kadhim MJ, Al-Rubaye AF, Hameed IH. Determination of bioactive compounds of methanolic extract of *Vitisvinifera* using GC-MS. *Int J ToxicolPharmacol Res.* 2017;9(2):113-126.
24. Gazwi HS, Shoeib NA, Mahmoud ME, Soltan OI, Hamed MM, Ragab AE. Phytochemical Profile of the ethanol extract of *Malvaviscusarboreus*red flower and investigation of the antioxidant, antimicrobial, and cytotoxic activities. *Antibiot.* 2022;11(11):1652.
25. Ali B, Rawal YK, Ahmad F, Ali A. Phytochemistry, GC-MS, and FTIR analysis of ethanolic extract of alfalfa (*Medicagosativa*) leaves from cold desert Ladakh, India. *Int J Bot Stud.*2016;6(6):241-246.
26. Rakib A, Paul A, Chy MNU, Sami SA, Baral SK, Majumder M, Simal-Gandara J. Biochemical and computational approach of selected phytocompounds from *Tinosporacrispa* in the management of COVID-19. *Mol.*2020;25(17):3936.
27. Srivastava R, Mukerjee A, Verma A. GC-MS analysis of phytocomponents in pet ether fraction of *Wrightiatinctoria* seed. *Pharmacogn J.*2015;7(4):249-253.