

Biochemical studies on evaluation of Sunflower (*Helianthus annuus* L.) genotypes for heat stress

Abstract

Crops are facing heat stress because of rapid climate change caused by global warming. We examined these issues in sunflower by exposing the plants growing with mean maximum temperature of 32.2°C (sowing to flowering) and 36.1°C (flowering to harvest) at normal temperature (S1) and 35.3°C (sowing to flowering) and 38.3 °C (flowering to harvest) at high temperatures (S2) by staggered sowings. Antioxidative responses of sunflower were also explored by studying the Superoxide dismutase, Catalase, Peroxidase and Ascorbate peroxidase activities. A significant increase was observed in antioxidant enzyme activities under high temperature stress. The final oil composition proved to be sensitive to the timing of heat stress and reduced (13%) by high temperatures. Some innovative steps should be taken on an emergency basis to prepare plants for such stressful conditions.

Keywords: *high temperature, oil composition, staggered sowing, sunflower, stress*

Introduction

Rising global earth surface temperature is one of the most enthralling factor emerging from the changing climate and is the major environmental **parameter** which affects plant growth, development, and yield (Saleem et al. 2020) Globally, annual temperature is expected to rise by 1.8–4.0 °C at the end of the 21st century (Hassan et al. 2021). The heat stress is many fold high in arid to tropical zones of the globe affecting vital physiological processes of crops resulting in the reduction of food quantity and quality (Fischer and Knutti, 2015). The **high** temperature is different for crops like for corn 29°C, soybean is 30°C, cotton 32°C, sunflower 33°C, and maize 36°C (Gornall et al., 2010).

Sunflower (*Helianthus annuus* L.) is an important lipid and protein-rich oilseed crop and the fourth largest oilseed crop in the world cultivated in more than 70 countries (Killi et al. 2020). In *H. annuus*, previous research has demonstrated that brief period of heat stress during grain filling negatively impact on oil yield and fatty acid composition (Catiempo et al. 2021)

High temperatures (HT) damage the activity of membrane proteins and lipids, thus affecting the activity of chloroplast- and mitochondria-based enzymes and membrane integrity (Ul Hassan et al. 2021). Stress conditions can disturb the dynamic equilibrium of reactive oxygen species (ROS) production and elimination under normal growth in plants (Saleem et al. 2020) which promotes ROS accumulation, membrane lipid peroxidation, and disrupt the structure and function of cell membrane system (Saleem et al. 2020).

Different stress condition causes variations in plant metabolism and results in formation of reactive oxygen species (ROS) which is removed by antioxidant enzymes like ascorbate peroxidase (APX), catalase (CAT) and superoxide dismutase (SOD) (Taghvaei et al. 2015). In a cell SOD enzyme is the first line of defence against reactive oxygen species which detoxifies superoxide anion to hydrogen peroxide and molecular oxygen (Yan et al., 2016). The main objective of this study was to investigate the direct effects of constant increase of high temperature on biochemical traits in sunflower.

Materials and methods

This experiment was conducted in field at Narkhoda farm, ICAR- Indian Institute of Oilseeds Research, Hyderabad (17_1501600 N, 78_1803000 E; 542 m above sea level) during *late rabi* (February- June) 2020 in split-plot design with two temperature treatments using staggered sowings (S1- Timely sowing and S2- Delayed sowing). Fourteen cultivars of *H. annuus* named as AKSF 6-3B, CMS 17B, CMS 42B, CMS 70B, CMS 107B, CMS 125B, CMS 127B, CMS 135B, CMS 144B, ARM 243B along with four checks CO 2, CSFH 12205, DRSH 1, KBSH 44 were selected from the TIR study (Aparna et al., 2023). Each genotype was sown in 6 x 3.6 m plots with a spacing of 60 cm (between rows) x 15 cm (between plants); there were three replicates for each treatment. Sowing was done by dibbling, and the recommended fertilizer dose (60 Kg N, 90 Kg P₂O₅, and 30 kg K₂O /ha) (Malligawad et al., 2004) was applied; other standard practices and need-based plant protection measures were followed to ensure a healthy crop. The soil structure is sandy loam with low organic carbon (0.4%) and nitrogen (235 kg/ ha) content, and relatively high phosphorus (22 kg/ha) and potassium (405 kg/ha) content. For enzyme activity Enzyme extracts were prepared by freezing a weighed amount of leaf samples (0.1g) in liquid nitrogen to prevent proteolytic activity, followed by grinding in a 0.1 M phosphate buffer at pH 7.5 containing 0.5 mM EDTA and 1 mM ascorbic acid at a 1:10 (w/v) ratio (Dhindsa et al. 1981). The homogenate separated then passed through four layers of gauze, and the filtrate was centrifuged at 15,000

rpm for 20 min. The resulting supernatant was used as an enzyme source. Biochemical traits like CAT (Sinha 1972), SOD (Beauchamp and Fridovich 1971), POX (Rao *et al.*, 1996), APX (Nakano and Asada 1981) and OC were studied. Cleaned seed sample of 15 gram each of fourteen lines were oven dried at 70°C for 3 hours to determine the oil content using NMR (Nuclear Magnetic Resonance) spectrometer (MARS, UAS, Raichur).

Statistical Analysis: Results were analyzed by analysis of variance and LSD values were calculated for cultivars, treatments and their interactions.

Results and Discussion

Two sowings were taken up with timely (S1) and delayed (S2) sowing. The maximum (Tmax) and minimum temperature (Tmin) from sowing to flowering was 32.2°C, 16.4°C and 35.3°C, 20.2°C and from flowering to harvest was 36.1°C, 20.8°C and 38.3°C, 23.8°C for S1 and S2 respectively. The difference in Tmax recorded in the two sowings was 3.1°C at sowing to flowering and 2.2°C at the flowering to harvest. The results of the biochemical study are presented below (Table 1)

Biochemical parameters: Significant variation among the genotypes along with TxG interaction to HT was observed for CAT activity at the vegetative stage, and with temperature treatments for CAT activity at the flowering stage, SOD activity at the vegetative stage, POX activity at the vegetative and flowering stages and APX activity at the vegetative and flowering stages (Table 1).

Catalase (CAT) (EC 1.11.1.6) ($\mu\text{mol min}^{-1} \text{g}^{-1}$):

During the vegetative stage CAT activity varied from 0.75 to 1.27 (1.09) and from 0.76 to 1.84 (1.31) in S1 and S2. Checks DRS1 1 (1.27) being at par with checks CSFH 12205 (1.26), CO 2 (1.24), KBSH 44 (1.16), inbred CMS lines-70B (1.14), -107B (1.21), AKSF 6-3B (1.11) in S1 while inbred CMS 70B (1.84) being at par with CMS 144B (1.73) in S2 has higher CAT activity. Inbred CMS lines-70B (62%), -144B (49%), -17B (30%), -127B (30%), and AKSF 6-3B (26%) have recorded more increase in CAT activity compared to checks at the vegetative stage. At the flowering stage, CAT activity varied from 0.77 to 1.45 (1.19) and from 1.06 to 1.99 (1.58) in S1 and S2. Inbred AKSF 6-3B (1.45) being at par with checks DRS1 1 (1.45), CO 2 (1.3), CSFH 12205 (1.27), KBSH 44 (1.24) and inbred CMS 107B (1.35) in S1 while, in inbred CMS lines-70B (1.99), -107B (1.81), -144B (1.80), -17B (1.75) and checks CO 2 (1.74), DRS1 1 (1.72) in S2 has higher CAT activity. Inbred CMS lines -

70B (69%), -125B (67%), -17B (51%), -144B (45%), and -42B (41%) has recorded more increase in CAT activity compared to checks at the flowering stage.

The results were in accordance with canola (Zhang *et al.*, 2015), Brassica sps (Soengaset *al.*, 2018).

The activities of CAT, POX, SOD, and APX were increased with exposure to heat stress. Production of ROS often induces the production of abscisic acid and regulates the gene expressions that control the production of enzymatic antioxidants such as SOD and CAT (Thalman M, Santelia D 2017). CAT enzyme plays a key role in maintaining H₂O₂ concentration at normal level, inhibiting the chain reaction of ROS.

Superoxide dismutase (SOD) (EC 1.15.1.1)(nmol g⁻¹):

During the vegetative stage, SOD activity varied from 13.89 to 26.81 (19.3) in S1 and from 17.93 to 31.85 (23.98) in S2. Inbred CMS 70B (26.81) being at par with inbred CMS lines -42B (25.77), -144B (25.21) in S1 and CMS 42B (29.75) in S2 has higher SOD activity. At flowering SOD activity varied from 20.06 to 33.69 (25.21) and from 24.7 to 37.73 (29.67) under S1 and S2. Higher SOD activity was recorded in inbred CMS 70B (33.69 in S1 and 37.73 in S2) being at par with CMS 42B (33.33 in S1 and 36.03 in S2). Inbred AKSF 6-3B at vegetative stage (33%) and flowering (42%) recorded more percent increase in SOD activity compared to checks. The present results were also in accordance with Brassica sps (Soengaset *al.*, 2018).

Aerobic metabolism in plants releases ROS as by-products; however, environmental stresses prompt ROS production. Superoxide dismutase (SOD) is considered to be the first line of defence to safeguard plants against environmental fluctuations (Raja *et al.* 2017) by converting superoxide anion (O₂⁻) into O₂ and H₂O₂. H₂O₂ still being toxic is converted into H₂O and oxygen by CAT (Carvalho *et al.* 2020).

Peroxidase (POX) (EC 1.11.1.7)(nmol min⁻¹ g⁻¹):

During the vegetative stage, POX activity varied from 14.04 to 17.91 (15.49) and from 16.08 to 27.99 (20.69) under S1 and S2. POX activity is higher in check CO 2 (17.91) being at par with inbreds AKSF 6-3B (16.91), CMS 107B (16.37), ARM 243B (16.15) in S1, and in inbred AKSF 6-3B (27.99) at par with check CO 2 (26.99) in S2. At the flowering stage, POX

activity varied from 14.64 to 20.38 (17.47) under S1 and from 16.27 to 29.75 (23.19) under S2. Higher POX activity was recorded in check CO 2 (20.38) being at par with inbreds AKSF 6-3B (19.6), CMS 107B (18.44), ARM 243B (18.55), and checks DRSH 1 (19.08), CSFH 12205 (18.16) under S1 and in check CO 2 (29.75) and inbred AKSF 6-3B (29.75) under S2 respectively. Inbred AKSF 6-3B at vegetative stage (66%) and flowering (52%) recorded a more percent increase for the trait POX activity compared to checks.

Similar results were reported in Sorghum (Gosavi *et al.*, 2014) and wheat (Sarkar *et al.*, 2016) under HT stress (30 °C). Increased levels of activities of peroxidase (POX) and ascorbate peroxidase (APX) were observed in lablab (*Dolichos lablab*) seedlings (D'Souza and Devaraj, 2013). Maintenance of a high antioxidant capacity to scavenge the toxic ROS has been linked to increase in tolerance of plants to these environmental stresses (Hasanuzzaman *et al.*, 2013).

Ascorbate peroxidase (APX) (EC 1.11.1.11) (nmol min⁻¹ g⁻¹):

APX activity varied from 4.12 to 7.16 (5.2) in S1 and from 5.79 to 9.22 (7.11) in S2 at vegetative stage. Inbreds CMS 42B (7.16) in S1 and ARM 243B (9.22) in S2 has higher APX activity. At the flowering, APX activity varied from 4.54 to 7.26 (5.52) under S1 while, and from 6.36 to 9.42 (7.42) under S2. Higher APX activity was recorded in inbreds CMS 42B (7.26) being at par with inbred ARM 243B (6.63) in S1. Similarly, inbred ARM 243B (9.42) being at par with check CSFH 12205 (8.92) and inbred CMS 42B (8.71) in S2 has higher APX activity. Inbred AKSF 6-3B at vegetative stage (54%) and at flowering (57%) recorded a more percent increase for the trait APX activity compared to checks.

APX efficiently scavenges H₂O₂ due to its higher affinity for H₂O₂ compared to CAT (Sharma *et al.* 2012). When the crop is exposed to continuous stress, the equilibrium between H₂O₂ production and the antioxidant activity becomes unbalanced, leading to excessive accumulation of H₂O₂ which further increases the levels of CAT and APX as they are continually involved in rebalancing the equilibrium.

The generation of destructive reactive oxygen species, including superoxide radical (O₂⁻), singlet oxygen (1 O₂), hydroxyl radical (OH⁻) and hydrogen peroxide (H₂O₂) was probably the reason of enhanced enzyme activities (Rivero *et al.*, 2014). An increase in CAT, SOD, POX, APX activity may be a manifestation of the adaptive response of plants to abiotic stress without which the plant growth reduction could be more severe. An increase in

temperature leads to the increased expression of these antioxidative enzymes until a pre-determined temperature after which they decline. Tolerant varieties could maintain increased activities at higher temperatures than susceptible ones (Chakraborty and Pradhan 2011). This suggests that there is a common regulatory (biochemical and molecular) framework of mechanism(s) underlying the induction of enzyme activity or possibly the entire suite of changes, including other antioxidative defenses in response to the different abiotic stress conditions (Leung et al., 2018)

Increased APX activity in tomato plants proposing an effectual H₂O₂ scavenging capacity under HS. However, this enzyme activity under high temperatures could be inadequate to scavenge the surplus of H₂O₂ when the activity of CAT is not initiated, triggering oxidative damage (Raja et al., 2020).

Table1 Enzymes activity of sunflower genotypes during late *Rabi*, 2020

	CAT		SOD		POX		APX		OC (%)
	Vegetative	Flowering	Vegetative	Flowering	Vegetative	Flowering	Vegetative	Flowering	
Temperature	NS	0.23	1.79	4.15	0.61	2.14	0.09	1.08	1.8
CV (%)	28	17.4	8.8	16.08	3.6	11.22	1.6	17.8	5.7
Genotypes	0.2	0.28	2.3	3.01	1.88	1.88	0.5	0.8	3.7
CV (%)	14.27	17.5	9.18	9.46	8.97	7.98	7.06	10.6	9.4
T*G	0.28	NS	NS	NS	2.66	2.66	0.71	NS	NS

CAT-catalase, SOD- super oxide dismutase, POX-peroxidase, APX- ascorbate peroxidase, OC- Oil concentration

Oil Concentration (%):

OC varied from 31.9 to 39.9% (35.7 %) in S1 and from 24.8 to 37.4 % (31.1 %) in S2. Under both situations, the checks CSFH 12205 and DRSH 1 recorded the highest OC whereas inbreds CMS -135B &-17B secured the lowest OC respectively. Inbred AKSF 6-3B (5%) recorded the lowest percent reduction.

Oil concentration was significantly altered by the HT. This decrease in oil concentration with an increase in temperature was in accordance with previous reports (Rondanini et al. 2014). The cause of the reduction in OC could be attributed to a shortening of the grain-filling period and reduced rate of seed maturation and duration of oil deposition in the grain at the HT (Van der Merwe et al., 2015).

Conclusion: Our results demonstrate that sunflower (*Helianthus annuus* L.) primary leaf grown under high temperature increased the activity of key enzymes and the oxidation status

of the plants during leaf ontogeny. The biochemical traits under S1 and S2 differed among the sunflower genotypes studied. The inbreds and hybrids with different genetic backgrounds resulted in trait variation. Variation in the S2 condition of specific traits measured among genotypes aids the selection of these traits. These trait-specific genotypes could be used in sunflower breeding programs to develop location-specific varieties.

References

- Aparna, V., Lakshmi Prayaga., Arti Guhe., Lakshamma, P., 2023. "Identification of temperature tolerant sunflower (*Helianthus annuus* L.) inbreds." *Journal of Oilseeds Research* 37. 10.56739/jor.v37iSpecialissue.141195.
- Beauchamp, C., Fridovich, I., 1971. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical biochemistry* 44(1), 276-287.
- Carvalho, F.E., Silveira, J.A., 2020. H₂O₂-retrograde signaling as a pivotal mechanism to understand priming and cross stress tolerance in plants. In *Priming-mediated stress and cross-stress tolerance in crop plants*. Academic Press 57-78
- Catiempo, R.L., Photchanachai, S., Bayogan, E.R.V., Vichitsoonthonkul, T.J.C.S., 2021. Possible role of nonenzymatic antioxidants in hydro-primed sunflower seeds under heat stress. *Crop Science* 61, 1328-1339.
- Chakraborty U, Pradhan D., 2011. High temperature-induced oxidative stress in *Lens culinaris*, role of antioxidants and amelioration of stress by chemical pre-treatments. *Journal of Plant Interactions*. 6:43–52
- Dhindsa, R.A., Plumb-Dhindsa, P., Thorpe, T.A. 1981. Leaf senescence: Correlated with increased permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *Journal of experimental Botany*. 126: 93-101.
- D'Souza, M., Devaraj, V.R., 2013. Induction of thermotolerance through heat acclimation in lablab bean (*Dolichos lablab*). *African Journal of Biotechnology* 12, 5695–5704.
- Fischer, E.M., Knutti, R., 2015. Anthropogenic contribution to global occurrence of heavy-precipitation and high-temperature extremes. *Nature climate change* 5(6), 560-564.
- Gornall, J., Betts, R., Burke, E., Clark, R., Camp, J., Willett, K., Wiltshire, A., 2010. Implications of climate change for agricultural productivity in the early twenty-first century. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365(1554), 2973-2989.

Hasanuzzaman, M., Nahar, K., Anee, T.I., Fujita, M., 2017. Glutathione in plants: biosynthesis and physiological role in environmental stress tolerance. *Physiology and Molecular Biology of Plants* 23, 249-268.

Hassan, M.U, Chattha, M.U, Khan, I, Chattha, M.B., Barbanti, L., 2021. Heat stress in cultivated plants: Nature, impact, mechanisms, and mitigation strategies—A review. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* 155, 211-234.

Killi, D., Raschi, A., Bussotti FJjoms., 2020. Lipid peroxidation and chlorophyll fluorescence of photosystem II performance during drought and heat stress is associated with the antioxidant capacities of C3 sunflower and C4 maize varieties. *International Journal of Molecular Sciences* 21, 4846.

Leung, D.W.M., 2018. Studies of Catalase in Plants Under Abiotic Stress. In: Gupta, D., Palma, J., Corpas, F. (eds) *Antioxidants and Antioxidant Enzymes in Higher Plants*. Springer, Cham. https://doi.org/10.1007/978-3-319-75088-0_2

Malligawad, L. H., Parameshwarappa, K. G., Giriraj, K. 2004. Studies on the effect of ratios and level of NPK fertilizer nutrients of the productivity of hybrid sunflower under rainfed farming situations. In *Proc. 16th Int. Sunflower Conference, Fargo, ND, USA* (Vol. 1, pp. 377-386).

Nakano, Y., Asada, K., 1981. Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and cell physiology* 22(5), 867-880.

Rao, M.V., Paliyath, G., Ormrod, D.P., 1996. Ultraviolet-B-and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*. *Plant physiology* 110(1), 125-136.

Raja, V., Majeed, U., Kang, H., Andrabi, K.I., John, R., 2017. Abiotic stress: interplay between ROS, hormones and MAPKs. *Environmental and Experimental Botany* 137,142–157.

Raja, V., Qadir, S.U., Alyemeni, M.N., Ahmad, P., 2020. Impact of drought and heat stress individually and in combination on physio-biochemical parameters, antioxidant responses, and gene expression in *Solanum lycopersicum*. *Biotechnology* 10(5), 208.

Rivero, R.M., Mestre, T.C., Mittler, R.O.N., Rubio, F., Garcia- Sanchez, F.R.A.N.C.I.S.C.O. Martinez, V., 2014. The combined effect of salinity and heat reveals a specific physiological, biochemical and molecular response in tomato plants. *Plant, cell and environment* 37(5), 1059-1073.

Rondanini, D.P., Castro, D.N., Searles, P.S., Rousseaux, M.C., 2014. Contrasting patterns of fatty acid composition and oil accumulation during fruit growth in several olive varieties and locations in a non-Mediterranean region. *European journal of agronomy* 52, 237-246.

Saleem, M.H., Rehman, M., Fahad, S., Tung, S.A, Iqbal, N., 2020. Leaf gas exchange, oxidative stress, and physiological attributes of rapeseed (*Brassica napus* L.) grown under different lightemitting diodes. *Photosynthetica* 58 (3), 836

Sharma, P., Jha, A.B., Dubey, R.S., Pessarakli, M., 2012. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of botany*.

Sinha, A.K., 1972. Colorimetric assay of catalase. *Analytical biochemistry*, 47(2), 389-394.

Taghvaei, M., Jafari, S.M. 2015. Application and stability of natural antioxidants in edible oils in order to substitute synthetic additives. *Journal of Food Science and Technology*. 52, 1272–1282.

Thalman, M., Santelia, D., 2017. Starch as a determinant of plant fitness under abiotic stress. *New Phytologist*, 214(3), 943-951.

Ul Hassan, M., Rasool, T., Iqbal, C., Arshad, A., Abrar, M., 2021. Linking Plants Functioning to Adaptive Responses Under Heat Stress Conditions: A Mechanistic Review. *Journal of Plant Growth Regulation* 1-18.

Van der Merwe, R., Labuschagne, M., Herselman, L., Hugo, A., 2015. Effect of heat stress on seed yield components and oil composition in high- and mid-oleic sunflower hybrids. *South African Journal of Plant and Soil*. 32, 1-8.

Yan, H. F., Mao, P. S., Sun, Y., Li, M. L., 2016. Impacts of Ascorbic Acid on Germination, Antioxidant Enzymes and Ultrastructure of Embryo Cells of Aged *Elymus sibiricus* Seeds with Different Moisture Contents. *International Journal of Agriculture & Biology*, 18(1).