

## Effect of integrated nutrient management on yield and yield attributes in Scented Rice (*Oryza sativa* L.)

### ABSTRACT

The present experiment was conducted at Barrister Thakur Chhedilal College of Agriculture and Research Station, Bilaspur, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh during *Kharif* season 2019-20 entitled with integrated nutrient management in scented rice including organic and inorganic nutrients source with recommended dose of fertilizers (RDF) 100: 60: 40 NPK kg ha<sup>-1</sup>. Experiment was layout at randomized block design (RBD) with the ten treatments and three replications, taking variety of scented rice Chhattisgarh SugandhitBhog. The investigation of result found that between the different organic and inorganic treatments T<sub>6</sub> (75% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia), result indicated that the highest grain yield 35.78 q ha<sup>-1</sup> was observed which was significantly superior over other treatments but was at par with T<sub>5</sub> (75% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) with 34.24 q ha<sup>-1</sup> grain yield closely followed by the treatments T<sub>4</sub>, T<sub>3</sub>, T<sub>2</sub> and T<sub>10</sub>. The higheryield and yield attributes characters under the T<sub>6</sub> has been affiliated, highest no. of panicles per m<sup>2</sup> (349.36) also significantly superior.

**Keyword:** Integrated nutrient management, scented rice,

**Comment [WU1]:** Abstract need more elaboration

### INTRODUCTION

Rice (*Oryza sativa* L.) is the world's largest staple food crop. It is the rich energy source and contains reasonable protein (6-10 percent), carbohydrate (70-80 percent), minerals (1.2-20%) and vitamins (Riboflavin, Thiamine, Niacin and Vitamin E). (Anonymous 2014). This data shows that in the 2018 crop year about 167.13 million hectare. India covers more than 30% of total cultivated area and contributes more than 40% of total production. In India rice production for 2015-16 amounts to 104.41 million tonnes, 2400 kg ha<sup>-1</sup> of production covering an area of 434.99 lakh hectares. The field size decreased to 431.94 lakh hectare in 2016-17 with an increase in 110.15 million tonnes of output and 2500 kg ha<sup>-1</sup> productivity (Annual report of 2017-18). India has 42.95 million hectare acreage with 111.01 million tonnes of production. According this data published by Indian stat (Anonymous, 2018). The state leader in rice in India is Uttar Pradesh, the rice is cultivated in an area of 5.95 million hectare (mha) with gross output of 13.53 hectare. In rice responsible for its scent, 2-Acetyl-1 pyrroline (2-AP) is most common; almost every state in the country has its own set of aromatic rice that has a good performance in.

**Comment [WU2]:** Well in introduction ,there is no role of consortia on rice production. What kind of consortia and their mechanism of action

Chhattisgarh is historically known as an Indian bowl of rice. More than 23,250 rice varieties were reported in state. In Chhattisgarh, rice is grown in 3.79 million hectare (mha) area and covers 8.58 percent of India, with average yield of 6.91 million tonne. (Anonymous, 2018). In the state of Chhattisgarh, there are many scented rice varieties that are cultivated and popular for their cultivation. Rice varieties are being grown in Chhattisgarh *i.e.* Jeeraphool, Dubraj, Vishnubhog and Jawaphool. Indira Gandhi Krishi Vishwavidyalaya developed medium length and short height of scented rice "Chhattisgarh SugandhitBhog". Several studies have highlighted the role of integrated nutrient management

activities to increase crop yields and improve soil quality in long term. The use of organic manures plays an significant role in the use of fertilizers performance, lower cost of supply of nutrients, increase production under rainfed conditions and require less capital investment, particularly under unfavourable weather conditions.

**Comment [WU3]:** Reference?

Organics supplies nutrients during peak adsorption and also supplies micro nutrients and alters soil physical behaviour and increase the performance of applied nutrients (Pandey *et al.*, 2018). Use of organic and inorganic fertilizers mixture which increases the nutrient use efficiency and minimum losses of nutrients. The nitrogen loss processes occur by leaching and de-nitrification. Reasonable and fertilizer use can increase markedly the yield and the rice quality improved. PSB (Phosphorus Solubilizing Bacteria) plays an important role amongst soil microorganisms in the solubilization of P for plants and use of phosphatic fertilizers.

**Comment [WU4]:** Introduction need some more clarification that why this study was planned, what were the objectives behind all this

## MATERIALS AND METHODS

Field experiments were conducted at Barrister Thakur Chhedilal College of Agriculture and Research Station, Bilaspur (C.G.) during *Kharif* season 2019-20. The experimental plot soil has been classified as black group, known as clayey loam soil. pH in soil was 6.2, Electrical conductivity was recorded 0.25 dSm<sup>-1</sup>, OC 0.64 and available nitrogen was 150 kg ha<sup>-1</sup>, phosphorus and potassium was 13.88 kg ha<sup>-1</sup> and 204.96 kg ha<sup>-1</sup> respectively. The present experiment was laid out in Randomized Block Design (RBD) with ten treatments and three replications. Ten treatments of the experiment *viz.*, T<sub>1</sub> – Control, T<sub>2</sub> – 100% RDF (100:60:40 kg NPK ha<sup>-1</sup>), T<sub>3</sub> – 75% RDF + 5 tonne FYM ha<sup>-1</sup>, T<sub>4</sub> – 75% RDF + 5 tonne Vermicompost ha<sup>-1</sup>, T<sub>5</sub> – 75% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia, T<sub>6</sub> – 75% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia, T<sub>7</sub> – 50% RDF + 5 tonne FYM ha<sup>-1</sup>, T<sub>8</sub> – 50% RDF + 5 tonne Vermicompost ha<sup>-1</sup>, T<sub>9</sub> – 50% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia, T<sub>10</sub> – 50% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia. Rice variety “Chhattisgarh sugandhitbhog” used as a experimental variety. Recommended dose of fertilizers for scented rice *i.e.*, 100:60:40kg NPK ha<sup>-1</sup> was application as urea, single super phosphate (SSP) and murate of potash (MOP). Available nitrogen, phosphorus and potassium content sample was determined by Alkaline permanganate method (Subbaiah and Asija, 1956), Olsen’s method (Olsen, 1954) and Flame photometer (Jackson, 1967). Data subjected was statistical analysis as prescribed by Gomez and Gomez, 2010.

**Comment [WU5]:** Unit?

**Comment [WU6]:** Is author telling about NPK fertilizer dose or NPK soil fertility status, if its about soil fertility status then unit is wrong

**Comment [WU7]:** Nothing mentioned about consortia origin, its species and density how much it was used. Methodology lacking the data measurement details that at which plant stage physical, physiological and yield parameters taken and with what ways?

## RESULT AND DISCUSSION

### Panicle Length

The data on the length of the panicles (cm) is shown in Table- 1 . Panicle length was at par with different treatment. However, the longest panicles (24.39 cm) were reported in the T<sub>6</sub> (75% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia) nearly followed by

T<sub>5</sub>(75% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) (23.50 cm), T<sub>4</sub> and T<sub>3</sub>. The shortest panicle length reported in T<sub>1</sub> (control) 20.29cm.

The data showed that the integrated nutrient management methods and the application on **of** only chemical fertilizers had a positive impact on the length of panicles but was statistically at par. **however**, the maximum plant height, highest grain yield and maximum panicle length observed in treatment T<sub>6</sub> but treatment T<sub>2</sub> is closely followed without the bio-fertilizer application which is totally inorganic fertilizers.

Comment [WU8]: Confusing statement

Comment [WU9]: Font color should be similar

### Number of Panicle per m<sup>2</sup>

Data on various yield attributes are shown in Table- 1 and shown in Table It was clear from the data that there was a substantial difference in the number of ear-bearing panicles per square meter between different treatments. Results indicated that the highest number of effective panicles per square meter (349.36) was observed in T<sub>6</sub> (75% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia) which was significantly superior over other treatments but was at par with T<sub>5</sub> (75% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) 327.97. Minimum number of effective panicles observed in T<sub>1</sub> (Control) with 232.40.

### Test Weight

Data on the weight of 1000grain as determined by different treatments are shown in Table- 1. The test weight of seed differed significantly between the different treatments. Result indicated that the highest test weight (24.84g) was observed in treatment T<sub>6</sub> (75%RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia) which was significantly superior over other treatments but was at par with treatment T<sub>5</sub> (75%RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) 24.54g recorded the minimum test weight of seed (20.18g) was reported in treatment T<sub>1</sub> (Control) which was significantly different from the other treatments.

In integrated nutrient management treatments, the number of panicles and test weight of seed nearly equal to the other significantly differed between single application of the required chemical or inorganic fertilizers. Its increase or decrease combination with FYM and Bio-fertilizers. Same results also found by Sahu *et al.*,(2017). Pandey *et al.*,(2018). Farooq *et al.*,(2018). Raj *et al.*,(2013). Sravan and singhet *et al.*,(2019).

### Table- 1 Effect of INM of Yield Attributing Characters

	Treatments	Length of panicles (cm)	No. of panicles/m <sup>2</sup>	Test weight
T <sub>1</sub>	Control	20.29	232.40	20.18
T <sub>2</sub>	100% RDF ( 100:60:40 kg N:P:K ha <sup>-1</sup> )	22.34	293.30	21.76
T <sub>3</sub>	75% RDF + 5 tonne FYM ha <sup>-1</sup>	22.79	293.62	21.83
T <sub>4</sub>	75% RDF + 5 tonne Vermicompost ha <sup>-1</sup>	22.59	307.59	23.76
T <sub>5</sub>	75% RDF + 5 tonne FYM ha <sup>-1</sup> enriched with consortia	23.50	327.97	24.54
T <sub>6</sub>	75% RDF + 5 tonne Vermicompost ha <sup>-1</sup> enriched with consortia	24.39	349.36	24.84
T <sub>7</sub>	50% RDF + 5 tonne FYM ha <sup>-1</sup>	21.4	265.19	21.36
T <sub>8</sub>	50% RDF + 5 tonne Vermicompost ha <sup>-1</sup>	22.06	279.25	21.42
T <sub>9</sub>	50% RDF + 5 tonne FYM ha <sup>-1</sup> enriched with Consortia	22.15	282.76	21.51
T <sub>10</sub>	50% RDF + 5 tonne Vermicompost ha <sup>-1</sup> enriched with Consortia	22.29	286.51	21.58
<b>SEm±</b>		<b>0.71</b>	<b>12.54</b>	<b>0.64</b>
<b>CD(0.05%)</b>		<b>NS</b>	<b>37.26</b>	<b>1.89</b>

**Comment [WU10]:** Statistical analysis needs revision

### Grain Yield

Result indicated that the highest grain yield (35.78 qha<sup>-1</sup>) was observed in treatment T<sub>6</sub> (75%RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) which was significantly superior over other treatments but was at par with T<sub>5</sub> (75%RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) with 34.24 qha<sup>-1</sup>). Minimum grain yield observed in treatment T<sub>1</sub> (Control) 13.76 qha<sup>-1</sup>.

In results indicated that the higher grain yield observed in treatment T<sub>6</sub> over other treatments but which was significantly at par with T<sub>5</sub> and followed by T<sub>4</sub>, T<sub>3</sub>. Organic sources such as vermicompost, bio-fertilizers and inorganic fertilizers have contributed to the constant supply of nutrients, hormones and related enzymes eventually reducing nutrient losses and increasing nutrient quality, yield and also increased the use efficiency. Same

results also found by Rao *et al.*,(2013). Kumar *et at.*,(2014). Kandeshwari and Thavaprakash (2016). Sahu *et al.*,(2017). Harikesh *et al.*,(2017). Pandey *et al.*,(2018). Kumar *et al.*,(2018).

#### 4.2.5. Straw Yield

In relation to grain yield, straw yield was also affected by various treatments, as shown in Table- 2. Result indicated that the highest straw yield ( $60.66 \text{ qha}^{-1}$ ) was observed in treatment T<sub>6</sub> (75% RDF + 5 tonne Vermicompost  $\text{ha}^{-1}$  enriched with consortia) which was significantly superior over other treatments but was at par with T<sub>5</sub> (75% RDF + 5 tonne FYM  $\text{ha}^{-1}$  enriched with consortia)  $57.50 \text{ qha}^{-1}$ . Minimum straw yield observed in treatment T<sub>7</sub> and T<sub>1</sub>  $38.57$  and  $24.70 \text{ qha}^{-1}$  in descending order.

Higher yields of grain and straw could be attributed to the combined use of organic and inorganic nutrients sources resulting in increased essential nutrients availability to the plant and improved soil quality which facilitated improved root spread this contributes to improved nutrient absorption and eventually to higher yields.

Comment [WU11]: Same para

#### 4.2.7. Harvest Index

The harvest index for various treatments is shown in Table- 2. Result indicated that the highest harvest index (39.77) was observed in treatment T<sub>1</sub> (Control) which was significantly superior over other treatments but was at par with T<sub>9</sub> (50% RDF + 5 tonne FYM  $\text{ha}^{-1}$  enriched with consortia) 39.07 followed by T<sub>8</sub>, T<sub>10</sub> and T<sub>2</sub>. The minimum harvest index recorded in treatment T<sub>7</sub> (50% RDF + 5 tonne FYM  $\text{ha}^{-1}$ ) 35.55.

The results indicated that the highest grain yield observed in T<sub>6</sub> and harvest index was observed in treatment T<sub>10</sub> (50% RDF + 5 tonne Vermicompost  $\text{ha}^{-1}$ ) which was significantly superior over other treatments. The use of organic nutrients with prescribed dose of fertilizer showed strong response for all characters.

Higher yields of grain and straw could be attributed to the combined use of organic and inorganic nutrients sources resulting in increased essential nutrients availability to the plant and improved soil quality which facilitated improved root spread this contributes to improved nutrient absorption and eventually to higher yields.

Comment [WU12]: Repetition of same para at end of each result.

Each treatment was observed different harvest index but in treatment T<sub>4</sub> (75% RDF + 5 tonne Vermicompost  $\text{ha}^{-1}$ ) was lower straw yield than other rest treatments and harvest index was increased. In results indicated that the higher grain yield observed in treatment T<sub>6</sub> over other treatments but which was significantly at par with T<sub>5</sub> and followed by T<sub>4</sub>, T<sub>3</sub>. Organic sources such as vermicompost, bio-fertilizers and inorganic fertilizers have contributed to the constant supply of nutrients, hormones and related enzymes eventually reducing nutrient losses and increasing nutrient quality, yield and also increased the use efficiency. Same results also found by Rao *et al.*,(2013). Kumar *et at.*,(2014).

Kandeshwari and Thavaprakash (2016). Sahu *et al.*,(2017). Harikesh *et al.*,(2017). Pandey *et al.*,(2018). Kumar *et al.*,(2018).

**Straw: Grain ratio**

Straw: grain ratio data are presented in Table-2. Between the treatments, T<sub>7</sub> (50% RDF + 5 tonne FYM ha<sup>-1</sup>) observed in highest straw: grain ratio (1.81) which was significantly superior over other treatments but was at par with T<sub>1</sub>. The treatment T<sub>9</sub> (50% RDF + 5 tonne FYM ha<sup>-1</sup> enriched with consortia) observed the minimum straw: grain ratio of 1.55.

UNDER PEER REVIEW

**Table- 2 Effect of integrated nutrient management Grain and Straw Yield**

	Treatments	Grain Yields (qha <sup>-1</sup> )	Straw: Grain ratio	Straw Yield (qha <sup>-1</sup> ) <sup>1)</sup>	Harvest Index (%)
T <sub>1</sub>	Control	13.67	1.80	24.70	39.77
T <sub>2</sub>	100% RDF (100:60:40 kg NPK ha <sup>-1</sup> )	28.35	1.66	45.71	38.27
T <sub>3</sub>	75% RDF + 5 tonne FYM ha <sup>-1</sup>	28.88	1.61	46.60	38.26
T <sub>4</sub>	75% RDF + 5 tonne Vermicompost ha <sup>-1</sup>	31.30	1.64	51.48	37.81
T <sub>5</sub>	75% RDF + 5 tonne FYM ha <sup>-1</sup> enriched with consortia	34.24	1.67	57.50	37.32
T <sub>6</sub>	75% RDF + 5 tonne Vermicompost ha <sup>-1</sup> enriched with consortia	35.78	1.69	60.66	37.10
T <sub>7</sub>	50% RDF + 5 tonne FYM ha <sup>-1</sup>	21.28	1.81	38.57	35.55
T <sub>8</sub>	50% RDF + 5 tonne Vermicompost ha <sup>-1</sup>	24.32	1.58	38.58	38.66
T <sub>9</sub>	50% RDF + 5 tonne FYM ha <sup>-1</sup> enriched with Consortia	26.81	1.55	41.81	39.07
T <sub>10</sub>	50% RDF + 5 tonne Vermicompost ha <sup>-1</sup> enriched with Consortia	27.96	1.59	44.57	38.54
<b>SEm±</b>		<b>0.72</b>	<b>0.04</b>	<b>0.98</b>	
<b>CD(0.05%)</b>		<b>2.13</b>	<b>0.11</b>	<b>2.91</b>	

## CONCLUSION

From the experimental result, it could be enlightened that application of 75% RDF + 5 tonne Vermicompost ha<sup>-1</sup> enriched with consortia could be considered as a better option for obtained higher production under integrated nutrient management and poor experimental results occurs in control. Integrated nutrient management was found to significantly superior results in all stages of crop, combination of organic and inorganic fertilizers without influenced the yield of rice in addition to increase in the sustainability of the production

capacity of soil. Hence, integrated nutrient management, it is possible more successful and profitable, for scented rice.

## REFERENCES

- Anonymous 2014. Agricultural Statistics at a glance, Directorate of Economics and Statistics Department of Agriculture and Cooperation, Pp.72-73.
- Anonymous. 2012. Agricultural Statistics at a glance, Directorate of Economics and statistics, Department of Agriculture and Cooperation. Pp.65-66.
- Pandey, A., Parihar, R. and Srivastava, V.K. (2018). Effect of Integrated nutrient management practices on soil chemical and biological properties under rice (*Oryza sativa* L.) cultivation. *International Journal of Chemical studies*; 6(4) : 1871-1874.
- Harikesh, Ali A., Shivam, Yadav, R.P., Kumar, S., Kumar,A., and Yadav,A. (2017). Effect of Integrated Nutrient Management and Plant Geometry on Growth and Quality of Rice (*Oryza sativa* L.) Varieties under SRI Technique.*International Journal of Current Microbiology and Applied Science* ISSN : 2319-7706 Volume 6 No.10 pp. 2503-2515.
- Farooq, K., Nabi, B.B., Rashid, M., Hamid, F. (2018). Studies on integrated nutrient management in rice (*Oryza sativa* L.). *International Journal of Advance Research in Science and Engineering*. Volume No.07, Special Issue No. 04, (744-749). ISSN: 2319-8354.
- Kandeshwari, M. and Thavaprakash, N. (2016). Influence of integrated nutrient management practices on yield and nutrient uptake in rice (*Oryza sativa* L.) under system of rice intensification. *International Journal of Agricultural Sciences and Research*.ISSN(P) : 2250-0057; ISSN(E): 2321-0087. Vol. 6, Issue 2, pg. 123-130.
- Kumar, K., Sridhara, C.J., and Nandini, K.M. (2018). Effect of Integrated use of organic and inorganic fertilizers on soil fertility and uptake of nutrients in aerobic rice (*Oryza sativa* L.). *International Journal of Chemical Studies*; 6(4) : 417-421.

- Kumar, A., Meena, R.N., Yadav, L. and Gilotia, Y.K. (2014). Effect of Organic and Inorganic Sources of Nutrient on Yield, Yield Attributes and nutrient Uptake of Rice CV.PRH-10. *The Bioscan* 9(2) :595-597.
- Pandey, A., Parihar, R. and Srivastava, V.K.(2018). Effect of Integrated nutrient management practices on soil chemical and biological properties under rice (*Oryza sativa* L.) cultivation. *International Journal of Chemical studies*; 6(4) : 1871-1874.
- Rao, T.K., Rao, U.A., Ramu, S.P., Sekhar, D. and Rao, V.N. (2013). Effect of organic manures on performance of scented varieties of rice in high altitude areas of Andhra Pradesh. *International Journal of Current Microbiology and Applied Science*. ISSN: 2319-7706 Volume 2 Number 11 . pp. 339-346.
- Sahu, G., Chatterjee, N., and Ghosh, K.G. (2017). Integrated Nutrient Management in Rice (*Oryza sativa* L.) in Red and Lateritic Soils of West Bengal. *Indian Journal of Ecology* .44 (Special Issue-5): 349-354.
- Sravan, S.U. and Singh, P.S. (2019). Effect of Integrated Nutrient Management on Yield and Quality of Basmati Rice Varieties. *Journal of Agricultural Science*. Vol. 11, No. 5. ISSN 1916-9752 E-ISSN 1916-9760. Pg. 93-103.
- Raj, K.S., Mathew, R., Jose, N. and Leenakumary, S.(2013). Integrated Nutrient Management Practices for Enhancing Yield and Profitability of Rice (*Oryza Sativa* L.). *Madras Agriculture . Journal*. 100 (4-6): pg. 460-464.

UNDER PEER REVIEW