

QUALITATIVE PHYTOCHEMICAL SCREENING AND LARVICIDAL EFFICACY OF PHYSIC NUT (*Jatropha curcas*) LEAVES, STEM-BARK AND ROOT EXTRACTS ON MOSQUITO LARVAE.

ABSTRACT

The advance of battle by mosquito species to artificial chemicals has obliged the unrelenting pursuit for....of nontoxic ones from plants. This study screened the qualitative phytochemicals and evaluated the larvicidal efficacy of extracts of *Jatropha curcas* leaves, stem-bark and root extracts against third and fourth mosquito instar larvae. The *J. curcas* obtained was...were subjected to extraction by maceration. Qualitative phytochemical screening of ethanolic stem-bark, aqueous and ethanolic root extracts plant...plants showed the presence of phenols, saponins, steriods...steroids flavonoids, alkaloids, glycosides, carbohydrate...carbohydrates and terpenoids. The aqueous extracts of the leaves and stem-bark showed the presence alkaloids...presence of alkaloids, cardiac glycosides, phenols, saponins, and terpenoids whereas, ethanolic leaves extract had similar phytochemicals with the leaves and stem-bark aqueous extracts but did not have steriods..steroids. Twenty-four (24) hours exposure of different concentrations (2.0mg/ml, 4.0mg/ml, 6.0mg/ml, 8.0mg/ml, 10 mg/ml) of various extracts showed that larval mortality increased significantly ($P < 0.05$) with increase in extracts concentration. Aqueous stem-bark extract showed highest mortality with 87% against third mosquito instar larvae followed by aqueous, ethanolic root extracts against third and fourth mosquito instar larvae respectively with 85% larval mortality. However, least mortality was observed in ethanolic leaf extract with 5% larval mortality against fourth mosquito instar larvae. Low LC_{50} and LC_{90} values were noticed in ethanolic root and aqueous stem-bark extracts with 2.19mg/ml and 11.51mg/ml respectively. Whereas the highest LC_{50} and LC_{90} values were noticed in both ethanolic leaves extracts with 14.09mg/ml and 26.20mg/ml respectively against fourth mosquito instar larvae. Conclusively, aqueous stem-bark and ethanolic root extracts can be harnessed to control 50% and 90% mosquito instar larval mortality respectively. However, there is a need to ascertain the quantities of bioactive components of *J. curcas* and its toxicity to non-target organisms.

Key Words: *Jatropha curcas* , LARVICIDAL EFFICACY , PHYTOCHEMICAL , MOSQUITO LARVAE.

INTRODUCTION

Mosquitoes are very important vectors of several human diseases including malaria, lymphatic filariasis, dengue virus and Zika virus worldwide (Giovanni, 2016). Mosquito-borne diseases are the major public health problems in developing countries. Malaria parasites are conveyed to humans by female *Anopheles* mosquitoes (CDC, 2012). The parasites are propagated in humans through the bites of infected *Anopheles* mosquitoes. There are more than 400 species of *Anopheles* mosquitoes have been known and of these, 60 species serve as host...hosts to *Plasmodium*. *Plasmodium* is a parasitic protozoan that causes malaria. There are five types of human malaria parasites: *Plasmodium falciparum*, *Plasmodium ovale*, *Plasmodium*

vivax, *Plasmodium malariae* and *Plasmodium knowlesi* naturally infects...infect macaques in Southeast Asia and also infects humans, causing malaria that is transmitted from animal to humans “zoonotic malaria” (CDC, 2012). Yellow fever is endemic in most part of Nigeria which causes lives and economy loss. Lymphatic filariasis which is transmitted by *Aedes* and *Culex* species, affects more than 120 million people, and about 1.4 billion are at risk of these diseases in 73 countries which is rampant in African and Asian countries. Although, these diseases are associated with health and socioeconomic consequences (WHO, 2019).

Mosquito can be controlled by preventing mosquito bite using mosquito repellants, killing mosquitoes and causing larval mortality (CDC, 2018). The larval stage is the most vulnerable stage to attack mosquitoes as they are concentrated in smaller areas. Thus one of the approaches for control of malaria transmission is by interrupting the mosquito life cycle at larval stage...at the larval stage. Vector control for the prevention of malaria includes insecticide treated bed nets, indoor residual spraying and source reduction (larval control) (CDC, 2018).

Resistance to these insecticide classes has emerged in anopheles’ mosquitoes (WHO, 2016) and its rapid spread has become a major obstacle to vector control (Ranson and Lissenden, 2016). Nigeria has a vision...has the vision to become a malaria-free country with a mission to free every Nigerian citizen from the menace of malaria through effective leadership, environmental sanitation, health system strengthening and coordination in developing efficient policies, strategies and guidelines (National Malaria Strategic Plan, 2014).

The unpredictable mosquito larvae and pupae population transition dynamics, especially during transmission seasons remains...remain a critical issue in the tropics. Ability to carefully address and smashed this vague indicator has eventually determined the level of success attaining malaria..success in attaining the malaria elimination agenda. This should be significantly reduced through correct, consistent and effective application of Larval Source Management (LSM), which targets mosquito larvae as they mature in aquatic habitats (Oluwasogo and Godson, 2016). However, extract from plants may serve as an alternative source of bioactive compounds that are biodegradable into non-toxic products and are potentially suitable for use to control mosquitoes. Plant extracts in general have best recognized as an important natural resource of insecticides (Ghosh *et al.*, 2012). The efficacy of phytochemicals against mosquito larvae can vary significantly depending on plant species, plant parts used, age

of plant parts (young, mature or senescent), solvent used during extraction as well as upon the available vector species (Ghosh *et al.*, 2012; Rattanam *et al.*, 2014).

Phytochemicals derived from plant sources act as larvicides, insect growth regulators, repellents, oviposition attractants and can play an important role in the interruption of the transmission of mosquito-borne diseases to the individual as well as at the community level (Velu *et al.*, 2015). Therefore, larviciding is one of the successful way of reducing mosquito densities in their breeding places before they emerge into adults. Larviciding largely depends on the use of synthetic chemical insecticides such as organophosphates. Although effective and its repeated use has disrupted natural biological control systems and sometimes result in the wide spread development of resistance (Tehri and Singh, 2015). These problems have warranted the need for developing alternative strategies using eco-friendly products. Plants often have an alternative source of insect control because they contain a range of bioactive chemicals many of which are selective and have little or no harmful effect on non-target organisms and the environment. Much effort has, therefore, been focused on plant extracts or phytochemicals as potential sources of mosquito control agents or as lead compounds (Ghosh *et al.*, 2012).

MATERIALS AND METHODS

3.1 Collection of Plant Materials

A portion of *Jatropha curcas* containing leaves and fruits were collected within Sangere-FUTY premises and were....and was transported to the laboratory of plant science...of the plant department, school of life science, Modibbo Adama University of Technology Yola, for identification.

Phytochemical analysis

Ethanollic and aqueous extracts of *Jatropha curcas* leaves, stem-bark and root were qualitatively screened for the presence of tannins, saponins, phenol, carbohydrate, alkaloids, flavonoids, sterioids....steroids, terpenoids and cardiac glycosides were identified by characteristic colour....color change as described (Sofowora, 1993; Trease and Evans, 2002).

Collection and rearing of mosquito larvae

Mosquito larvae were collected from the study area (MAUTECH) campus and transported to Biochemistry laboratory (Old School of Pure and Applied Science) of Modibbo Adama University of Technology Yola, for rearing in a plastic bowl containing water covered with nylon mesh. Glucose was used in feeding the mosquito larvae (WHO, 2005).

Plant extraction process

For the crude aqueous extraction, 100 g of the leaves, stem-bark and root of *Jatropha curcas* were pounded with mortar and pestle. The filtrate (juice) was squeezed out using muslin...using a muslin cloth and concentrated to dryness at room temperature with rotary evaporator...with a rotary evaporator. While for the ethanolic extraction of 100 g of shade-dried leaves, stem-bark and root at room temperature ($27\pm 2^{\circ}\text{C}$), *Jatropha curcas* were macerated in 90% ethanol for 72 h. The filtrate was then concentrated to dryness using a rotary evaporator and stored in amber bottles in a refrigerator (-4°C) until it was ready for use.

Preparation of stock solution

0.1 gram of ethanolic and aqueous extracts of *Jatropha curcas* leaf, stem-bark and root each were dissolved in 100 millilitre of distilled water to form a stock solution. Serial dilution to obtain different concentrations of 10mg/ml, 8mg/ml, 6mg/ml, 4mg/ml and 2mg/ml were prepared by pipetting 5ml, 4ml, 3ml, 2ml and 1ml respectively of stock solution and they were added to 45ml, 46ml, 47ml, 48ml and 49ml of distilled water containing mosquito instar larvae respectively.

Larvicidal Bioassay

The third and fourth mosquito larvae, twenty were exposed to different concentrations with slight modifications (WHO, 2005). Three replicates along with control...along with the control were run simultaneously on each trial. The larvae were fed with glucose. The number of dead larvae at 24 hours were...was recorded. Moribund larvae were also counted as dead larvae. The dead larvae were removed soon after mortality in order to...to prevent decomposition, which may cause rapid...cause the rapid death of the remaining larvae. Total...A total of three trials was carried out. The corrected mortality was analysed using Abbot Formulae.

$$\text{Percentage mortality (\%)} = \frac{\text{Number of Dead larvae}}{\text{Number of larvae introduced}} \times 100$$

3.7 Data analysis

Larval mortality data of the third and fourth mosquitoes' larvae was....were subjected to Probit analysis to determine the median lethal concentration (LC₅₀) and lethal concentration ninety (LC₉₀) of each extracts against mosquito larvae. Fiducial limits of upper and lower confidence limits of LC₅₀ was....were determined (Abbot, 1925). All data were tested at P<0.05 level of significant....significance.

RESULTS

Phytochemical Analysis

The results ethanolic...results of ethanolic stem-bark, ethanolic and aqueous root extracts of *Jatropha curcas* showed the presence of alkaloids, cardiac glycosides, saponins, terpenoids, carbohydrate....carbohydrate, phenols, flavonoids and steroid but tannins were absent (Table 1). Leaf and stem-bark aqueous showed nearly same....nearly the same phytochemicals with....as the above extracts but tannins, flavonoids and carbohydrates were absent. Whereas, ethanolic leaf extracts revealed only alkaloid...only alkaloids, cardiac glycosides, phenols, saponins and terpenoids. Tannin, steriods, flavonoids and carbohydrates were absent.

Toxicity of *Jatropha curcas* aqueous extracts on fourth and third mosquito larvae after twenty-four (24) hours exposure....hours of exposure

The toxic effect of aqueous leaf extracts of *Jatropha curcas* against fourth...against the fourth instar was observed at highest concentration (10mg/ml) with 30% mortality, followed by 25%, 20%, 20% and 15% at 8mg/ml, 6mg/ml, 4mg/ml and 2mgml respectively. However, it showed highest larval mortality of 70% against third mosquito instar larvae at 10mg/ml followed by 53%, 33%, 15% and 10% at respective concentrations of 8mg/ml, 6mg/ml, 4mg/ml and 2mg/ml (Table 2).

The toxic effect of aqueous stem-bark extracts of *Jatropha curcas* against fourth instar was observed at highest concentration (10mg/ml) with 85% mortality, followed by 65%, 55%, 45% and 35% at 8mg/ml, 6mg/ml, 4mg/ml and 2mgml respectively. However, it showed highest larval mortality of 87% against third mosquito instar larvae at 10mg/ml followed by 75%, 60%, 40% at respective concentrations of 8mg/ml, 6mg/ml, 4mg/ml and the least mortality was observed at 2mg/ml with 40% mortality (Table 2).

The toxic effect of aqueous root extracts of *Jatropha curcas* against fourth instar was observed high at highest concentration (10mg/ml) with 60% mortality, followed by 40%, 40%,

30% and 25% at 8mg/ml, 6mg/ml, 4mg/ml and 2mg/ml respectively. However, it showed highest larval mortality of 85% against third mosquito instar larvae at 10mg/ml followed by 60%, 45%, 35% at respective concentrations of 8mg/ml, 6mg/ml, 4mg/ml and the least mortality was observed at 2mg/ml with 25% mortality (Table 2).

Table 1: Phytochemical constituents of leaves, stem-bark and root extracts of *J. curcas*

Phytochemicals	ELE	ALE	ESBE	ASBE	ERE	ARE
Alkaloids	+	+	+	+	+	+
Carbohydrates	-	-	+	-	+	+
Cardiac glycosides	+	+	+	+	+	+
Flavonoids	-	-	+	-	+	+
Phenols	+	+	+	+	+	+
Saponins	+	+	+	+	+	+
Steroids	-	+	+	+	+	+
Terpenoids	+	+	+	+	+	+
Tannins	-	-	-	-	-	-

Key; + = Present; - = Absent

ELE = Ethanol leaves extract; ALE = Aqueous leaves extract;

ESBE = Ethanol stem-bark extract; ASBE = Aqueous stem-bark extract;

ERE = Ethanol root extract; ARE = Aqueous root extract

Table 2: Mean Mortality of mosquito larvae exposed to different doses of *Jatropha curcas* aqueous extracts

Larvae	Concentrations (mg/ml)	Aqueous Leaves extract	Aqueous stem-bark extract	Aqueous Root extract
Third	2	2 (10%)	8 (40%)	5(25%)
	4	3 (15%)	8 (40%)	7(35%)
	6	7 (35%)	12(60%)	9(45%)
	8	11(55%)	15(75%)	12(60%)
	10	14(70%)	18(87%)	17(85%)
Fourth	2	3(15%)	7(35%)	5(25%)
	4	4(20%)	9(45%)	6(30%)
	6	4(20%)	11(55%)	8(40%)
	8	5(25%)	13(65%)	8(40%)
	10	6(30%)	17(85%)	12(60%)

Toxicity of *Jatropha curcas* ethanolic extracts on fourth and third mosquito larvae after twenty (24) hours exposure....hours of exposure

The toxic effect of ethanolic leaves extract of *Jatropha curcas* against fourth instar was observed at highest concentration (10mg/ml) with 45% mortality, followed by 40%, 25%, 15% and 5% at 8mg/ml, 6mg/ml, 4mg/ml and 2mg/ml respectively. However, it showed highest larval mortality of 60% against third mosquito instar larvae at 10mg/ml and 8mg/ml followed by 55%, 45% and 35% at respective concentrations 6mg/ml, 4mg/ml and 2mg/ml (Table 3).

The toxic effect of ethanolic stem-bark extracts of *Jatropha curcas* against fourth instar was observed at highest concentration (10mg/ml) with 80% mortality, followed by 70%, 65%, 50% and 40% at 8mg/ml, 6mg/ml, 4mg/ml and 2mg/ml respectively. However, it showed highest larval mortality of 65% against third mosquito instar larvae at 10mg/ml followed by 55%, 50%, 50% at respective concentrations of 8mg/ml, 6mg/ml, 4mg/ml and the least mortality was observed at 2mg/ml with 45% mortality (Table 3).

The toxic effect of ethanolic root extracts of *Jatropha curcas* against both third and fourth mosquito larvae was the same. At highest concentration (10mg/ml) 85% larval mortality was observed, followed by 75%, 65%, 60% at 8mg/ml, 6mg/ml, 4mg/ml respectively and the least mortality was observed at 2mg/ml with 50% mortality (Table 3).

Table 3: Mean Mortality of mosquito larvae exposed to different doses of *Jatropha curcas* ethanolic extracts.

Larvae	Concentrations (mg/ml)	Ethanolic Leaves extract	Ethanolic Stem-bark extract	Ethanolic root extract
Third	2	7 (35%)	9(45%)	10(50%)
	4	9 (45%)	10(50%)	12(60%)
	6	11(55%)	10(50%)	13(65%)
	8	12 (60%)	11(55%)	15(75%)
	10	12 (60%)	13(65%)	17(85%)
Fourth	2	1(5%)	8(40%)	10(50%)
	4	3(15%)	10(50%)	12(60%)
	6	5(25%)	13(65%)	13(65%)
	8	8(40%)	14(70%)	15(75%)
	10	9(45%)	16(80%)	17(85%)

LC₅₀ and LC₉₀ of *Jatropha curcas* plant against third and fourth mosquito instar larvae

The highest and lowest LC₅₀ value of ethanolic leaf extract were 10.054mg/ml and 7.702mg/ml respectively; the highest and lowest LC₅₀ of aqueous leaf extract of *J. curcas* were 14.090mg/ml and 7.744mg/ml respectively. Whereas, the lowest and highest LC₅₀ value of ethanolic stem-bark is 3.794mg/ml both (Table 4). But, the aqueous stem-bark showed 3.794mg/ml and 4.672mg/ml as the lowest and highest values. However, the lowest and highest value of aqueous root extract were 6.357mg/ml and 2.189mg/ml respectively while the highest and lowest LC₅₀ of ethanolic root extract of *J. curcas* were 8.317mg/ml and 5.831mg/ml and. Highest and lowest values of aqueous extract were 6.357mg/ml and 2.289 respectively (Table 4). While highest and lowest LC₉₀ of *Jatropha curcas* ethanolic leaf extract were 17.183mg/ml and 13.011mg/ml; the highest and lowest LC₉₀ of aqueous leaf extract of were 126.196mg/ml and 13.291mg/ml respectively. Whereas, the highest and lowest LC₉₀ value of ethanolic stem-bark 13.218mg/ml both. However, the lowest and highest LC₉₀ value of aqueous root extract were 12.648mg/ml and 15.260mg/ml while highest and lowest LC₉₀ of ethanolic root extract of *Jatropha curcas* were 16.441mg/ml and 12.648mg/ml respectively.

Table 4: Probit Analysis of *Jatropha curcas* extracts against third and fourth mosquito instar larvae

Parts of plants solvent	Instar larvae		LC ₅₀ (mg/ml)	LC ₉₀ (mg/ml)	Probit Regression equation
Leaves	Aqueous	L3	7.744	13.291	$y=-1.789+0.231x$
		L4	14.090	26.196	$y=-1.492+0.106x$
	Ethanol	L3	7.702	13.011	$y=-1.859+0.241x$
		L4	10.054	17.183	$y=-1.807+0.180x$
Stem-bark	Aqueous	L3	3.981	11.507	$y=-0.678+0.170x$
		L4	4.672	12.141	$y=-0.802+0.173x$
	Ethanol	L3	3.794	13.218	$y=-0.266+0.057x$
		L4	3.794	13.218	$y=-0.516+0.136x$
Root	Ethanol	L3	5.831	11.767	$y=-0.516+0.136x$
		L4	8.317	16.441	$y=-1.312+0.158x$
	Aqueous	L3	2.189	12.648	$y=-0.268+0.123x$
		L4	6.357	15.260	$y=-0.915+0.144x$

Key: L4 = Fourth mosquito instar larvae
L3 = Third mosquito instar larvae
LC₅₀ = Lethal Concentration fifty

LC₉₀ = Lethal Concentration ninety

DISCUSSION

The result of this study reveals the efficacy of aqueous and ethanolic extracts of root, stem bark and leaf of *Jatropha curcas* against third and fourth instar larvae. The ethanolic extract of all the plant parts recorded the highest larvicidal activity on both third and fourth mosquito instar larvae than the aqueous extracts. Although ethanol is polar, but it also has the ability to attract non-polar molecules due to ethyl group of ethanol being non-polar. This contributes to its ability to extract highly polar and non-polar components from the plant material. It has very low toxicity, completely miscible in water, volatile and easily removed from plant material at low temperature (Roy, 2014).

The phytochemical screening of ethanolic stem-bark, ethanolic root and aqueous root extracts, recorded the presence of carbohydrates, cardiac glycosides, phenols, terpenoids, flavonoids, sterioids, saponins and alkaloids. This is in agreement with the research findings, who..which suggested that the bioactive compounds are responsible for antiplasmodial properties (Abdullahi *et al.*, 2017; Naqab, 2018) In addition to the absence of tannins in all the extracts, aqueous and ethanolic leaf extracts lack carbohydrates and flavonoids and sterioids were additionally absent in ethanolic leaf extract which is in accordance with....following the findings carried by Nwokocha, *et al.* (2011), also in consonance to research findings of Zainab, (2016), who found those bioactive compounds. These bioactive compounds found in the extracts were responsible for antibacterial activities (Zainab, 2016). These discrepancies in bioactive compounds could be as result...as a result of method....the method of plant material extraction, solvent...the solvent used and the geographical location of the plant. Since the soil where plants are established have different...have a different metallic and non-metallic compound (Bhattacharyya *et al.*, 2008). The aqueous stem-bark extracts revealed the presence of alkaloid, cardiac glycoside, sterioids, saponins, terpenoids and phenols. In addition, carbohydrates, flavonoids and tannins were found absent. This could be as a result of plant sample's proximity to the soil. Ethanolic and aqueous root extracts recorded 85% larval mortality both on third mosquito instar larvae and 87% larval mortality was shown by aqueous stem-bark extract against fourth mosquito instar larvae. The ethanolic leaf extract are...is responsible for antimalarial and anti-insecticidal (Juliet *et al.*, 2012). This is contrary to the report that, says only ethanolic root

extract elicits much.....more larval mortality than leaves and stem-bark extract due to phytochemical variations (Oseni and Alphone, 2011).

After exposing the mosquito instar larvae for twenty-four hours, the mean mortality of aqueous extracts was insignificant against third and fourth mosquito instar larvae. There was an increase in mortality of the larvae across the concentration gradient, that is, an increase in concentration elicits an increase in larval mortality. This observation supports the result of Rahuman *et al.*, 2008, who reported significant larvicidal activities of *Jatropha curcas* against different mosquito species. The larvicidal activities may be due to the presence of bioactive compounds which are toxic to mosquito....to mosquitoes.

The LC₅₀ lowest value of 2.19mg/ml was noticed in ethanolic root extract whose LC₉₀ value is 11.77mg/ml. This suggest....suggests the potency of *Jatropha...of the Jatropha curcas* part used against mosquito larvae. Whereas, the lowest LC₉₀ value of 11.51mg/ml was noticed in aqueous stem-bark extract whose LC₅₀ value is 3.98mg/ml against third instar larvae This is in consonance to the findings that recorded lowest....recorded the lowest LC₅₀ value of 3.715mg/ml in the subspecies of *Jatropha* (Buduwara *et al.*, 2021). Similar....A similar study was carried by Gawande *et al.* (2013), he....who observed LC₅₀ value of *Jatropha curcas* aqueous leaf extract at 20.41mg/ml whereas....whereas the LC₉₀ value as.....was 69.01mg/ml. This could be as result of the strain of the insect pest and part of the plant used. Although direct....Although the direct comparison may be impossible because of the differences in bioassay....in the bioassay method used. The result of this study is in consonance with the findings of Manjara *et al.* (2014), who reported that the acetone leaf extracts of *Clausena dentata* are very toxic to *Culex quinquefasciatus*, *Aedes aegypti* and *Anopheles stephensi* fourth instar larvae of mosquito. He recorded....recorded an LC₅₀ value of 0.15mg/ml and...and an LC₉₀ value of 7.20mg/ml against culex larvae. The LC₅₀ value of 0.17mg/ml and LC₉₀ value of 1.10mg/ml against *Aedes* larvae; LC₅₀ value of 0.05mg/ml and LC₉₀ value of 1.05mg/ml against *Anopheles stephensi* larvae. This disagrees with the findings of Alzeir *et al.* (2019), who noticed LC₅₀ value, 12.11µg/ml and LC₉₀ value of 30.21µg/ml of aqueous leaf extract of *Annona muricata* to be effective on *Aedes aegypti* larvae; and LC₅₀ value of 3.41µg/ml and LC₉₀ value of 6.17µg/ml to be effective on *Aedes albopictus* larvae. However, Johnson *et al.* (2018), who reported that, the chloroform extract of *Dichanthium foveolatum* LC₅₀ value is 277.03mg/ml whereas the LC₅₀ value of *Leptochloa uniflora*, *Molinera trichocarp*, *Pancratium triflorum* was

recorded as 300.56mg/ml, 306.60mg/ml, 318.42mg/ml respectively against culex species larvae. This could be as a result of different parameters used. Thus, the plants extracts that have a promised potency against test organisms.

Conclusion

In conclusion, presence of the following bioactive compounds such as alkaloids, flavonoids, saponins, steroid, phenol, carbohydrates, terpenoids, cardiac glycosides in solvent extracts *Jatropha curcas* leaves, stem-bark and root were responsible for its larvicidal activities against both the third and fourth mosquito instar larvae. Stem-bark extract was more potent against the third and fourth mosquito instar larvae. However, ethanolic leaves extract of *Jatropha curcas* was less potent against both the third and fourth mosquito instar larvae.

Recommendations

- i. Further work is needed to identify the potent active component in the extracts responsible for observed larvicidal effects.
- ii. There is also the need to investigate the effects of the solvent extracts on other life stages (eggs, pupae and adults) and **species mosquito....species of mosquito.**
- iii. **There is need....there is a need** to include plant extracts for mosquito control operations after safety tests.
- iv. There is need to investigate the effects of these extracts on mosquito co-inhabitant non-target aquatic organisms to determine specificity.

REFERENCES

- Abbott, W.S. (1952). A method of computing the effectiveness of an insecticide. *Journal Economic Entomology*, **18**: 265-267.
- Abdullahi, A.I. Amina, S., Ibrahim, A., K. and Hadiza, A. (2017). Photochemistry and in-vitro antiplasmodial properties of aqueous and ethanol stem bark extracts of *Jatropha curcas*. *Bayero Journal of Pure and Applied Sciences*, **10**(2): 2006-6996.
- Alzeir, M.R., Antonio, A.S., Cleonilda, C.P., Dayanne, L.S., Jose, C.C., Victor, E.P. and Selene, M. (2019). Larvicidal and enzymatic inhibition effect of *Annona muricata* seed extract and main constituents of Annonacin against *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Pharmaceutical Journal*, **12**(112): 1-12.
- Bhattacharyya, P., Chakrabarti, K., Chakraborty, A., Tripathy, S. and Powell, M.A. (2008). Fractionation and bioavailability of lead (Pb) in municipal solid waste compost and lead (Pb)

uptake by rice straw and grain under submerged condition in amended soil. *Geosciences Journal*, **12** (1):41 –45.

Buduwara, J. H., Adiel, T., Samy, R. and Tafem, M. L (2021). Phytochemical Screening and Larvicidal Assessment of Bellyache Bush (*Jatropha gossypifolia*) Leaf Extracts Against *Culex quinquefasciatus* Larvae. *Nigerian Journal of Parasitology*, **42**(1): 142-146.

Centers for Disease Control and Prevention (2012). Dengue and the *Aedes albopictus* mosquito. *Emergence of Infectious Diseases*, **18**(2): 345-347.

Centre for Disease Control and Prevention (2018). Malaria 5-8

Gawande, D., Wasu, Y. and Raja, I. (2013). Plant derived toxicants to control *Castor semicooper*, *Achaca Janata* (Noctuidae: Lepidoptera). *Asian Journal of Biology and Biotechnology*, **2**(12): 1-9.

Geovanni, R.M. (2016). Diversity and distribution in the world of Megalopodidae Latreille (Coleoptera: Chrysomeloidea). *International Congress of Entomology*. 25-30

Ghosh, A., Chowdhury, N., and Chandra, G. (2012). Plants extracts a potential mosquito larvicides. *The Indian Journal of Medical Research*, **135**(5): 581-598.

Juliet, S., Ravindran, R., Ramankutty, S.A. Gopalan, A.K. and Nair, S.N. (2012). *Jatropha curcas* (Linn) leaf extract a possible alternative for population control of *Rhipicephalus (Boophilus) annulatus*. *Asian Pacific Journal of Tropical Diseases*, **2**(3): 225–229

Manjara, M.S., Karhi, S., Ramkumar, G., Muthusamy, R., Natarajan, D. and Shivakumar, M.S. (2014). Chemical composition and larvicidal activity of plant extract from *Clausena dentata* (Wild) (Rutaceae) against dengue, malaria and filariasis vectors. *Parasitology Research*; **113** (7): 2475-2481

Naqab, K., Abdul, H.S., Ejaz, A.K., Muhammad, S., Samiullah, K. and Natashs, B. (2018). Phytochemical analysis and biological activities of tem bark extract of *Jatropha curcas*. Linn. *International Journal of Pharmacognosy*, 69-75

National Malaria Strategic Plan, 2014

Nwokocha, A., Blessing, I. and Okoli, O. (2011). Comparative phytochemical screening of *Jatropha L.* species in the Niger Delta. *Research Journal of phytochemistry*, **5**(2): 107-114.

- Oluwasogo, A.O. and Godson, K.C. (2016) Promoting Larval Source Management as a Vital Supplemental Addendum and More Likely Cost-Effective Approach for Malaria Vector Control in Nigeria. *Journal of Prevention & Infection Control*, **2**(2): 67-81
- Oseni, L.A. and Alphonse, P.K. (2011). Comparison of Antimicrobial Properties of Solvent Extracts of Different Parts of *Jatropha curcas*, *International Journal of Pharmacy and Pharmacology*
- Rahuman, A.A., G. Gopalakrishnan, P. Venkatesan and Geetha, K. (2008). Larvicidal activity of some Euphorbiaceae plant extracts against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). *Journal of Parasitology Research*, **102**(5): 867-873.
- Ranson, H. and Lissenden, N. (2016) Insecticide resistance in African Anopheles mosquitoes: A worsening situation that needs urgent action to maintain malaria control. *Trends Parasitology*, **32**(3), 187–196.
- Rattanam, A., Wan-Fatma, Z. and Jeevandran, S. (2014). Larvicidal efficacy of different plant parts of Railway creeper, *Ipomoea cairica* extract against dengue vector mosquitoes, *Aedes albopitus* (Diptera: Culicidae) and *Aedes aegypti* (Diptera: Culicidae). *Journal of Insect Science*, **14**(1):180-183
- Roy, P. (2014). Life cycle Assessment of Ethanol produced from Lignocellulosic Biomass: Techno-economic and Environmental evaluation. University Guelph, Ontario. 16-24
- Sofowora, A. (1993). Medicinal plants and Traditional Medicine in Africa. Second Edition. Spectrum Books Limited, Ibadan Nigeria, pp 231-234.
- Tehri, K. and Singh, N. (2015). The role of botanical as a green pesticide in Integrated Mosquito management-A review. *International Journal of Mosquito Research*, **2**(1): 18-23
- Trease, G.E and Evans, W.C. (2002). Pharmacognosy 15th Edition. Saunders, pp 214-393.
- Velu, K., Elumalai, D., Hemalatha, P., Babu, M., Janaki, A. and Kaleena P.K. (2015). Phytochemical screening and larvicidal activity of peel extracts of *Arachis hypogea* against Chikungunya and malaria vector. *International Journal of Mosquito research*, **2**(1): 1-8
- World Health Organization (2016). World malaria report. Geneva: World Health Organization. 3-8
- World Health Organization (2019). Guidelines for malaria vector control. Geneva.4-9

World Health Organization, (2005). Guideline for laboratory and field testing of mosquito larvicides, Communicable diseases control, Prevention and eradication, pesticide evaluation scheme. Geneva. 7-14

UNDER PEER REVIEW