

## Original Research Article

# Evaluating Weed Management Strategies and the Differential Effects of Herbicides on Weed dynamics and crop performance of foxtail millet [*Setaria italica* (L.) Beauv] grown under the irrigated conditions of Rayalaseema, Andhra Pradesh-India

**Comment [I1]:** There is no need of scientific name in the title please

### ABSTRACT

At the wetland farm of S.V. Agricultural College, Acharya N.G. Ranga Agricultural University, Tirupati, Andhra Pradesh, India, a field experiment was carried out during the (rainy season) of kharif 2020 to evaluate ~~ing~~ weed management strategies and the differential effects of herbicides on weed dynamics and crop performance of foxtail millet [*Setaria italica* (L.) Beauv] grown under the irrigated conditions of Rayalaseema. The current study has shown that the application of PE pretilachlor 500 g ha<sup>-1</sup> fb intercultivation at 20 DAS was followed by ~~HW~~ twice at 20 and 40 DAS and it led to the highest ~~WCE~~, including foxtail millet's nutrient uptake, and the lowest weed density and dry weight. In ~~unweededuntreated~~ check, a heavy weed infestation reduced the amount of nitrogen, phosphorus, and potassium absorbed by crop by 45.67, 18.03, and 35 kg ha<sup>-1</sup>, respectively. Grain production was increased by hand weeding twice, however the ~~benefit-cost~~ ~~cost-benefit~~ ratio lagged behind the most effective weed management strategy, which is applying PE of pretilachlor 500 g ha<sup>-1</sup> fb ~~intercultivation~~ ~~inter cultivation~~ at 20 DAS.

**Formatted:** Font: Italic

**Comment [I2]:** Write full words followed by abbreviation in the abstract please

**Comment [I3]:** Write full words followed by abbreviation in the abstract please

**Comment [I4]:** Write full words followed by abbreviation in the abstract please

**Comment [I5]:** Write concrete conclusion at the end of abstract as its last part

**Keywords:** ~~[Broad-spectrum, Pre-emergence herbicides, Pretilachlor, Inter~~ ~~cultivation, Hand weeding, Nutrient uptake]~~

**Comment [I6]:** Key words should be written alphabetically

### 1. INTRODUCTION

Foxtail millet [*Setaria italica* (L.) Beauv] is cultivated in India as a rain-fed Kharif crop. Appropriate weed control is regarded as a crucial component of agronomic procedures because to significant losses brought on by weed infestation (Munirathnam and Sawadhkar 2007) [4]. The highly diverse weed flora associated with foxtail millet varies according to the season, the state of the agro-ecological system, and the degree of control. During its early growth, foxtail millet is vulnerable to weed competition due to its slow-growing canopy. In general, especially in the early phases of the crop, minor millets are less competitive with weeds for growth resources. Foxtail millet has a severe weed invasion because of its initial delayed development during the rainy season. The first four to six weeks following seedling emergence were thought to be crucial for weed eradication (Nanjappa and Hosmani, 1985) [5]. According to Ning et al. (2015) [6], the presence of weeds during the crop season on calcareous soils caused a 56% reduction in the grain production of foxtail millet. Major millets have larger seeds and are sown at a deeper depth than minor millets, pre-emergence herbicides increase the efficiency of productivity and weed control in major millets (Mishra 2016) [3]. Very little research has been done on the topic of chemical weed control in foxtail millet. Farmers are now more likely to use herbicides in minor millet fields to effectively manage weeds because hand weeding has become more expensive in recent years. Hence,

the present study was undertaken to evaluate weed management strategies and the differential effects of herbicides on weed dynamics and crop performance of foxtail millet [*Setaria italica* (L.) Beauv] grown under the irrigated conditions of Rayalaseema.

## 2. MATERIAL AND METHODS

At the wetland farm of S.V. Agricultural College, Acharya N.G. Ranga Agricultural University, Tirupati, Andhra Pradesh, India, a field experiment was carried out during the kharif 2020 rainy season. The texture of the soil was sandy clay loam, with a neutral reaction, low levels of accessible nitrogen and organic carbon, and medium levels of available potassium and phosphorus. Eleven treatments were included in the randomized block design trial, which was conducted three times. On August 14, 2020, foxtail millet was sowed with a 30 x 10 cm spacing. Pre-emergence application (PE) of 500, 500, and 15  $\text{g/ha}$  of pretilachlor, isoproturon, and pyrazosulfuron-ethyl, respectively, as well as twice-hand weeding and unweeded check were the weed management treatments (Table 1). Twenty days after sowing (DAS), intercultivation or post-emergence application (PoE) of penoxsulam 20  $\text{g/ha}$  was added to all pre-emergence herbicides. Herbicides for pre-emergence were treated at 1 DAS, whereas penoxsulam, an inter-cultivation/post-emergence herbicide, was applied at 20 DAS. Applying all pre- and post-emergence herbicides required the use of a knapsack sprayer with a 500  $\text{L/ha}$  spray volume and a flat fan nozzle. All of the plots received an equal dosage of 20 kg N and 20 kg P in the form of urea and single super phosphate, respectively. The complete dose of phosphorous was administered as basal at the time of sowing, and the nitrogen was treated in two splits: half the dose was applied as basal and the other half as top dressing at 30 DAS. The recommendations of the Acharya N.G. Ranga Agricultural University were followed in adopting the remaining packages of practices. Using a 0.25  $\text{m}^2$  quadrat, category-specific weed density and biomass were randomly recorded. To standardize the distribution of the weed density and biomass data, a square root ( $\sqrt{X + 0.5}$ ) transformation was applied. Based on the methodology proposed by (Mani et al. 1973) [2], the efficiency of weed suppression was calculated. At harvest, every component of yield was documented. The  $\frac{\text{benefit} - \text{cost}}{\text{cost} - \text{benefit}}$  ratio was computed by dividing the cost of cultivation by the gross returns. November 5, 2020, was the harvest date of the crop. The statistical analysis of the weed and crop data was conducted using the analysis of variance for randomized block design, as recommended by Panse and Sukhatme (1985) [9]. At harvest in foxtail millet, the nitrogen uptake ( $\text{kg ha}^{-1}$ ) by weeds and crop was determined using the techniques recommended by Subbiah and Asija (1956) [12] and Olsen et al. (1954) [7], respectively. Following the normal protocol, data on plant height, dry matter production, yield qualities, and grain yield and straw yield were recorded at harvest. The net returns were calculated by taking the gross returns for each treatment and subtracting the cultivation cost. The  $\frac{\text{benefit} - \text{cost}}{\text{cost} - \text{benefit}}$  ratio was computed by dividing the cost of cultivation by the gross returns.

## 3. RESULTS AND DISCUSSION

### 3.1 Dry weight and weed density

Weed management significantly influenced rainfed foxtail millet. Hand weeding twice and pre-emergence pretilachlor at 500  $\text{g ha}^{-1}$  were most effective, except at 20 DAS. Hand weeding twice at 20 and 40 DAS might have controlled all the categories of weeds including perennial sedge *Cyperus rotundus* L. These results are in agreement with the findings of Jawahar et al. (2020) [1] with HW twice on 20 and 40 DAT in kodo millet. Pretilachlor 500  $\text{g ha}^{-1}$  followed by intercultivation ( $W_4$ ), which was at par with PE pyrazosulfuron-ethyl 15  $\text{g ha}^{-1}$  fb intercultivation ( $W_6$ ) and isoproturon 500  $\text{g ha}^{-1}$  fb intercultivation ( $W_5$ ) in order of ascent, at all the stages of observation (Table 1). This might be due to broad spectrum weed control by the PE herbicides

**Comment [17]:** Please add at least 6 latest references inline with your paper contents and title, the references must be from 2018-2023

**Comment [18]:** Divide this section in heading and sub heading  
For example  
Location of the experiment (Provide google map)  
Procedure  
Experimental design  
Procedure  
Statistical analysis  
Etc.  
You can write your own, it was just a suggestions

**Formatted:** Superscript

**Formatted:** Superscript

**Formatted:** Superscript

supplemented with [intercultivation](#) at 20 DAS. These results are in agreement with the findings of Patil et al. (2013) [10]. Isoproturon 500 g ha<sup>-1</sup> followed by penoxulam 20 g ha<sup>-1</sup> (W<sub>8</sub>) showed the higher dry weight and density of total weeds due to its poor performance in controlling all the categories of weeds associated with foxtail millet. These outcomes agree with those of Mishra et al. (2016) [3]. The highest dry weight and weed density were found in the [unweededuntreated](#) check. The most effective method of controlling weeds was hand weeding twice at 20 and 40 DAS (W<sub>10</sub>). The only exception was pre-emergence application of pretilachlor at 500 g ha<sup>-1</sup> with [intercultivation](#) at 20 DAS had a higher WCE than other pre-emergence herbicides.

**Table-1: Weed density m<sup>-2</sup>, dry weight (g m<sup>-2</sup>) and Weed control efficiency (%) as influenced by different weed management practices in foxtail millet**

Treatments	Dose (g ha <sup>-1</sup> )	Time of application (DAS)	Weed density m <sup>-2</sup>			Weed dry weight (g m <sup>-2</sup> )			WCE (%)		
			20 DAS	60 DAS	At harvest	20 DAS	60 DAS	At harvest	20 DAS	60 DAS	At harvest
Pretilachlor	500	1	47.67 (6.67)	100.87 (10.06)	(131.67) 11.49	11.13 (3.32)	79.70 (9.12)	(94.37) 10.06	85.01	29.18	27.34
Isoproturon	500	1	63.53 (7.86)	102.17 (10.13)	(133.33) 11.57	24.25 (4.97)	81.27 (9.21)	(95.93) 10.14	67.35	27.78	26.13
Pyrazosulfuron-ethyl	15	1	76.35 (8.92)	101.33 (10.09)	(132.50) 11.53	36.15 (6.05)	80.93 (9.19)	(95.60) 10.12	51.32	28.08	26.39
Pretilachlor <i>fb</i> IC	500	1+20	48.00 (6.75)	52.33 (7.27)	(69.33) 8.36	12.60 (3.57)	15.47 (3.99)	(32.07) 5.70	83.03	86.26	75.31
Isoproturon <i>fb</i> IC	500	1+20	63.67 (7.89)	55.00 (7.45)	(72.00) 8.51	24.61 (5.01)	19.10 (4.39)	(36.77) 6.07	61.47	83.03	71.69
Pyrazosulfuron-ethyl <i>fb</i> IC	15	1+20	76.00 (8.86)	53.67 (7.36)	(70.00) 8.40	35.21 (5.95)	18.77 (4.32)	(35.73) 6.01	52.59	83.32	72.48
Pretilachlor <i>fb</i> penoxsulam	500 + 20	1+20	48.33 (6.87)	80.63 (9.01)	(104.33) 10.24	12.40 (3.54)	58.40 (6.60)	(68.40) 7.39	83.30	48.10	47.33
Isoproturon <i>fb</i> penoxsulam	500+ 20	1+20	63.50 (7.85)	124.17 (11.15)	(161.00) 12.70	25.21 (5.07)	100.77 (10.29)	(117.27) 11.56	66.05	10.46	9.70
Pyrazosulfuron-ethyl <i>fb</i> penoxsulam	15+ 20	1+20	77.01 (8.89)	81.00 (9.03)	(106.50) 10.34	35.81 (6.01)	58.80 (6.67)	(69.60) 7.45	51.78	47.75	46.41
Hand weeding		20+40	136.67 (11.71)	18.33 (4.34)	(21.33) 4.67	72.13 (8.50)	2.90 (1.84)	(8.50) 2.98	-	97.42	93.45
<a href="#">UnweededUntreated</a> check			137.27 (11.74)	163.83 (12.82)	(192.33) 13.89	74.27 (8.64)	112.53 (11.58)	(129.87) 13.20			
LSD (p=0.05)			0.88	0.92	1.06	0.57	0.73	0.84			

**Comment [I9]:** Provide lettering in all tables with figure as you have used ANOVA

Data given in parenthesis are original values. Original data subjected to square root transformation.  
WCE: weed control efficiency; IC: [intercultivation](#); *fb*: followed by

## 3.2 Crop growth parameters

### 3.2.1 Phytotoxicity and Plant population m<sup>-2</sup>

Isoproturon at 500 g ha<sup>-1</sup> (W<sub>2</sub>, W<sub>5</sub>, W<sub>8</sub>) exhibited mild phytotoxicity, (Table 2) disappearing within 20 days. On average, it reduced initial and final plant population by 18.55% and 20.50%. Pre-emergence pretilachlor at 500 g ha<sup>-1</sup> and pyrazosulfuron-ethyl at 15 g ha<sup>-1</sup> showed no phytotoxicity. Post-emergence penoxulam at 20 g ha<sup>-1</sup> at 20 DAS resulted in severe crop injury and stunted growth exhibited phytotoxicity ratings of '1' and '5', respectively, on a 0-10 scale. Penoxsulam inhibit the acetolactate synthase enzyme activity, a key enzyme responsible for biosynthesis of branched chain amino acids, which intum making crop plants deficit in protein synthesis in meristematic tissues lead to discolouration of foliage and stunted growth. These outcomes agree with Prasuna and Rammohan (2015)

[11] findings with PE application of pretilachlor 0.75 kg ha<sup>-1</sup>fbPoE application of penoxsulam 25 g ha<sup>-1</sup> at 20 DAS in aerobic rice.

### 3.2.2 Plant height and LAI

Pre-emergence pretilachlor at 500 g ha<sup>-1</sup> with intercultivation (W<sub>4</sub>) showed optimal growth parameters, akin to pre-emergence pyrazosulfuron-ethyl at 15 g ha<sup>-1</sup> with intercultivation (W<sub>6</sub>) and isoproturon at 500 g ha<sup>-1</sup> with intercultivation (W<sub>5</sub>). Notably effective among pre-emergence herbicides, pretilachlor at 500 g ha<sup>-1</sup> performed comparably to pyrazosulfuron-ethyl (W<sub>3</sub>). During the key phase of crop-weed competition, which results in enhanced plant height and dry matter production, as well as higher crop uptake of nutrients, these three weed management practices provided broad-spectrum weed control. These outcomes agree with those of Pandey et al. (2018) [8]. However, sequential isoproturon at 500 g ha<sup>-1</sup> followed by penoxsulam at 20 g ha<sup>-1</sup> at 20 DAS (W<sub>8</sub>) exhibited reduced growth due to penoxsulam phytotoxicity, falling behind the unweededuntreated check (W<sub>11</sub>), which displayed significantly lower growth parameters than other weed management practices.

**Comment [I10]:** Please check the figure given in the results with figures given in the all tables. Make sure that they are inline

**Table-2: Crop growth parameters as influenced by different weed management practices in foxtail millet**

Treatments	Dose (g ha <sup>-1</sup> )	Time of application (DAS)	Phytotoxicity scoring		Plant population m <sup>-2</sup>		Plant height (cm)	Leaf area index
			PE herbicides	PoE herbicides	Initial	Final		
Pretilachlor	500	1	0	-	30.67	30.00	105	1.76
Isoproturon	500	1	1	-	26.67	25.67	105	1.70
Pyrazosulfuron-ethyl	15	1	0	-	31.67	31.00	105	1.74
Pretilachlor fb IC	500	1+20	0	-	31.67	31.67	117	1.91
Isoproturon fb IC	500	1+20	1	-	26.67	25.33	116	1.88
Pyrazosulfuron-ethyl fb IC	15	1+20	0	-	31.67	31.00	116	1.90
Pretilachlor fb penoxsulam	500+ 20	1+20	0	5	31.67	30.00	77	1.36
Isoproturon fb penoxsulam	500+ 20	1+20	1	5	25.67	24.33	67	1.31
Pyrazosulfuron-ethyl fb penoxsulam	15+ 20	1+20	0	5	31.33	31.00	76	1.32
Hand weeding		20+40	-	-	31.33	31.00	128	2.16
<u>UnweededUntreated</u> check			-	-	32.23	32.00	93	1.55
LSD (p=0.05)					3.17	3.01	10	0.047

### 3.3.3 Yield and Yield components

The highest yield was produced by hand weeding twice at 20 and 40 DAS (W<sub>10</sub>) followed by pre-emergence application of pretilachlor 500 g/ha<sup>g</sup> ha<sup>-1</sup> with inter-cultivation at 20 DAS (Table 3) because there was less competition for growth resources, which increased photosynthates translocation to developing grains. These outcomes concurred with the direct-seeded finger millet research conducted by Yathisha et al. (2020) [13]. The unweededuntreated check (W<sub>11</sub>) outperformed sequential pre-emergence herbicides with penoxsulam at 20 g ha<sup>-1</sup> (W<sub>7</sub>, W<sub>9</sub>, W<sub>8</sub>) due to penoxsulam phytotoxicity and poor weed control efficiency.

**Formatted:** Superscript

**Table-3: Yield and Yield components as influenced by different weed management practices in foxtail millet**

Treatments	Dose (g ha <sup>-1</sup> )	Time of application (DAS)	Number of panicles (m <sup>-2</sup> )	Dry matter production (kg ha <sup>-1</sup> )	Grain yield (kg ha <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )	Harvest index (%)	Gross returns (₹ha <sup>-1</sup> )	Net returns (₹ ha <sup>-1</sup> )	B:C
Pretilachlor	500	1	47.00	4617	1309	3008	30.37	40775	18815	1.86
Isoproturon	500	1	47.00	4604	1182	2962	28.02	36947	14900	1.68
Pyrazosulfuro n-ethyl	15	1	47.00	4611	1284	2978	30.12	39996	18363	1.85
Pretilachlor fb IC	500	1+20	62.67	5754	1961	3592	35.42	60637	35077	2.37
Isoproturon fb IC	500	1+20	56.00	5298	1660	3348	33.55	51460	25813	2.11
Pyrazosulfuro n-ethyl fb IC	15	1+20	58.33	5380	1745	3435	33.75	54067	28834	2.14
Pretilachlor fb penoxsulam	500+ 20	1+20	37.33	3297	779	2438	24.22	26999	2241	1.09
Isoproturon fb penoxsulam	500+ 20	1+20	32.00	3220	690	2250	23.47	25317	472	1.02
Pyrazosulfuro n-ethyl fb penoxsulam	15+ 20	1+20	33.33	3256	724	2397	22.92	25914	1483	1.06
Hand weeding		20+40	70.00	6496	2353	3944	37.47	72547	34867	1.92
Unweeded Untreated check			2.292	3951	861	2881	22.92	27257	7077	1.35
LSD (p=0.05)			6.72	4617	315	419	4.11	3220	4634	0.17

Hand weeding twice ( $W_{10}$ ) resulted in the highest gross returns, followed by pre-emergence pretilachlor at 500 g ha<sup>-1</sup> with [intercultivation/inter cultivation](#) ( $W_4$ ), showing comparable net returns. However, hand weeding twice had a lower [benefit-cost-cost-benefit](#) ratio due to increased weeding costs. The lowest gross and net returns, along with a lower [benefit-cost-cost-benefit](#) ratio, were observed with pre-emergence isoproturon at 500 g ha<sup>-1</sup> with penoxulam at 20 g ha<sup>-1</sup> ( $W_8$ ), significantly lower than the [unweededuntreated](#) check ( $W_{11}$ ) in terms of net returns and [benefit-cost-cost-benefit](#) ratio but comparable in gross returns.

### 3.3.4 Nutrient uptake by weed and crop

Outperforming other treatments, hand weeding twice ( $W_{10}$ ) reduced weeds' intake of nutrients (Table 4) and maximised the crop uptake of nutrients. [Intercultivation/inter cultivation](#) after pre-emergence pretilachlor at 500 g ha<sup>-1</sup> ( $W_4$ ) successfully decreased weeds' nutrient outflow and increased the crop uptake of nutrients. This could be because these weed management techniques eliminated all weeds, which decreased the dry weight of all weeds and hence reduced nutrient intake. These outcomes agree with those of Pandey et al. (2018) [8]. Nutrient absorption by weeds was enhanced by isoproturon at 500 g ha<sup>-1</sup> and penoxulam at 20 g ha<sup>-1</sup> ( $W_8$ ). In comparison to other techniques, the [unweededuntreated](#) check ( $W_{11}$ ) had the maximum nutrient uptake by weeds, with losses of 45.67, 18.03, and 35 kg ha<sup>-1</sup> of nitrogen, phosphate, and potassium, respectively.

**Table-4: Nutrient uptake (kg ha<sup>-1</sup>) by weeds and crop as influenced by different weed management practices in foxtail millet**

Treatments	Dose (g ha <sup>-1</sup> )	Time of application (DAS)	Nutrient uptake by weeds (kg ha <sup>-1</sup> )			Nutrient uptake by crop (kg ha <sup>-1</sup> )		
			Nitrogen	Phosphorus	Potassium	Nitrogen	Phosphorus	Potassium
Pretilachlor	500	1	30.00	13.33	22.67	41.33	11.17	48.33
Isoproturon	500	1	32.33	14.00	24.17	36.67	10.33	46.00
Pyrazosulfuron-ethyl	15	1	30.50	13.00	23.67	40.00	11.00	45.67
Pretilachlor fb IC	500	1+20	11.67	6.00	12.00	63.33	16.93	57.00
Isoproturon fb IC	500	1+20	14.20	6.67	14.17	59.67	15.83	55.00
Pyrazosulfuron-ethyl fb IC	15	1+20	13.33	7.00	13.33	62.00	16.33	55.00
Pretilachlor fb penoxsulam	500 + 20	1+20	24.67	10.33	19.00	25.67	6.00	29.33
Isoproturon fb penoxsulam	500+ 20	1+20	36.33	16.23	30.67	21.00	5.33	26.67
Pyrazosulfuron-ethyl fb penoxsulam	15+ 20	1+20	26.33	10.63	19.33	23.00	5.00	28.33
Hand weeding		20+40	5.00	2.00	5.67	73.00	19.67	64.67
Unweeded/Untreated check			45.67	18.03	35.00	25.33	6.17	31.00
LSD (p=0.05)			2.53	1.07	2.37	4.51	1.27	4.60

#### 4. CONCLUSION

In order to effectively manage weeds in foxtail millet, the field experiment showed that pre-emergence pretilachlor at 500 g ha followed by ~~intercultivation~~ inter cultivation at 20 DAS produced higher grain production and ~~benefit-cost~~ cost-benefit ratio. The grain yield was higher when hand weeding twice, but the benefit-to-cost ratio was not as high as it would have been with the ideal pre-emergence pretilachlor technique. In areas with abundant and economical labour for hand weeding, it remains a viable option. Notably, pre-emergence isoproturon at 500 g ha<sup>-1</sup> and post-emergence penoxsulam at 20 g ha<sup>-1</sup> at 20 DAS exhibited phytotoxicity ratings of '1' and '5', respectively, on a 0-10 scale.

**Comment [I11]:** Either add discussion to the existing results in the current format or add separate discussion section of atleast 2 pages and support your discussion with latest references

#### REFERENCES

1. Jawahar S, Ramesh S, Vinod Kumar SR, Kalaiyaran C, Arivukkarasu K. and Suseendran K. Effect of weed management practices on weed indices in transplanted kodo millet. Plant Archives. 2020;20(1):3196-3198.
2. Mani VS, Malla ML, Gautam KC and Bhagavandas. Weed killing chemicals in potato cultivation. Indian Farming. 1973;23:17-18.
3. Mishra JS, Singh VP and Rao SS. Bioefficacy and phytotoxicity of pre- and post-emergence herbicides in grain sorghum. Indian Journal of Weed Science. 2016;48(1):70-73.

4. Munirathnam P and Sawadhkar SM. Integrated weed management in foxtail millet (*Setariaitalica* (L.) Beauv). Indian Journal of Weed Science. 2007;39(1&2):144–145.
5. Nanjappa HV and Hosmani MM. Critical stage of crop weed competition in finger millet. Journal of Farming Systems. 1985;1:89-92.
6. Ning, N, Yuan X, Dong S, Wen Y, Gao Z, Guo M and Guo P. Grain yield and quality of foxtail millet (*Setariaitalica* (L.) in response to tribenuronmethyl. Plos One. 2015;10(11):1– 12.
7. Olsen SR, Cole CV, Watanable FS and Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. 1954;United States Department of Agriculture, circular number:939.
8. Pandey S, Sonboir HL and Thawait D. Evaluation of post emergence herbicides on growth parameters of finger millet. International Journal of Current Microbiology and Applied Sciences. 2018;7(3):1126-1134.
9. Panse VG and Sukhatme PV. Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research, New Delhi. 1985;100–174.
10. Patil B, Reddy VC, Ramachandra PTV, Shankaralingappa BC, Devendra, R and Kalyanamurthy, KN. Weed management in irrigated organic finger millet. Indian Journal of Weed Science. 2013;45(2):143-145.
11. Prasuna JG, and Rammohan J. Effect of weed management practices on growth and yield attributes of aerobic rice. Journal Crop and Weed. 2015;11(1): 229-231.
12. Subbiah BV and Asija GL. A rapid procedure for determination of available nitrogen in soil. Current Science. 1956;25:259-260.
13. Yathisha K, Yogananda S, Thimmegowda P, Sanjay M and Prakash S. Growth and yield of direct seeded finger millet (*Eleusine coracana* L.) as influenced by weed management practices. Journal of Crop and Weed. 2020;16(3):67-72.