

ASSOCIATION BETWEEN RISK FACTORS AND TRICHOMONIASIS AMONG WOMEN IN PORT HARCOURT RIVERS STATE ,NIGERIA

Comment [A1]: Title should be in font 18.

Abstract

Comment [A2]: ABSTRACT (Arial, Bold, 11 font, left aligned, caps)

Historically, the presence of *T. vaginalis* has been viewed to be associated with certain risk factors. This study is focused as assessing the association between selected risk factors and trichomoniasis. The cross sectional study composed of 650 women, 450 of them were tested for trichomonal infection using urine sample while the remaining 200 were tested for trichomonal infection using high vagina swab (HVS). These women were randomly selected among pregnant women, women with HIV, Out-patient, and healthy volunteer. After consenting to participate in the study, data concerning their exposure to selected risk factors were gathered through filled questionnaire. The results revealed that there was an association between “engaging in unprotected sex” and trichomoniasis ($p < 0.05$). Similarly, there were associations between prevalence of trichomoniasis and “wearing air-tight underwear” ($p < 0.05$), “repeating underwears” ($p < 0.05$), “knowledge of trichomoniasis” ($p < 0.05$) and “source of information” ($p < 0.05$). This study has shown that engaging in unprotected sex, wearing air-tight underwear, repeating underwear, knowledge of trichomoniasis and source of infection are significant risk factors of trichomoniasis among women in RSUTH.

Comment [A3]: In abstract separate into parts (objectives, methodology, results and conclusion). If you have any doubts, there are archives of recently published articles that show this structure. For example: **Methodology:** **Results:** **Conclusion**

Keywords: high vagina swab, *T. vaginalis* urine, trichomoniasis

Comment [A4]: Put four to eight keywords (font Arial, inclined, 10 font, justified)

Introduction

Comment [A5]: 1. INTRODUCTION (Arial, Bold, 11 font, left aligned, caps)

Historically, the presence of *T. vaginalis* has been viewed as a risk marker for other sexually transmitted agents such as *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, or bacterial vaginosis [1] Infection with *T. vaginalis* is increasingly been recognized to be associated with reproductive tract complications including sepsis that occur after abortion and after Cesarean Section, as well as adverse pregnancy outcome. Trichomoniasis is the major problem in developing countries where parental testing and antibiotics use are not adequately available [2]

Comment [A6]: Could be in this paragraph some reference about % os incidence in alguns countries in development.

In women, the disease encompasses symptoms ranging from a severe inflammation and irritation with frothy malodorous discharge to a relatively asymptomatic carrier state. The outcome of infection with *Trichomonas* may be due to genetic variability of the isolates and the host immune response. The vagina is the most common site of infection in women. The parasite is sexually transmitted through penis-to-vagina intercourse or vulva-to-vulva (the genital area outside the vagina) contact with an infected partner [3] Women can acquire the disease from infected men or

Comment [A7]: Why? It may be related to the pH of a woman's vagina

women, but men usually contract it only from infected women. Pregnant women with trichomoniasis may have babies who are born early or with low birth weight (low birth weight is less than 5.5 pounds). [2,9] Prevention of trichomoniasis has not been a priority due to lack of understanding of its public health implications and lack of resources. For long it has been considered a 'minor' STD. [3]

Comment [A8]: The most recent study by the World Health Organization provides data on more frequent STIs, including TV. It would be interesting to mention these values.

Trichomonas vaginalis is the most common Sexually Transmitted Parasitic Diseases (STPDs) in Nigeria[4;5]but had neither been the focus of intensive study nor of active control program and the negligence is as a result of relatively mild symptoms of the disease on initial exposure especially in African descent. Therefore, the study was aimed at assessing the association of trichomoniasis and selected risk factors in urine HVS of women attending RSUTH.

Materials and Methods

Comment [A9]: 2. material and methods (Arial, Bold, 11 font, left aligned, caps)

Study Design

Comment [A10]: 2.1 Subheading Subheading (ARIAL, BOLD, 11 FONT, LEFT ALIGNED) - second level heading.
For example: **2.1 Study Design** (arial, bold, font 11 ...)

A descriptive cross sectional study was adopted and 650 participants[6]were randomly selected [7]from females within the age of 15 and 55 years attending Rivers State University Teaching Hospital (RSUTH), Port Harcourt, Nigeria. The participants were selected among pregnant women, HIV patients, out patients and healthy volunteers. Out of the 650 participants, 450 were tested for trichomoniasis using urine while 200 participants were tested using high vagina swab (HVS). Data pertaining possible risk factors were obtained via well-structured questionnaire supported by verbal interview[8].

Comment [A11]: Why were 450 women tested with urine and 200 were tested with vaginal swabs? Why wasn't this number the same? For example 300/300 or others.

Ethical Consideration

Ethical approval was sought and obtained from Rivers State University Teaching Hospital (RSUTH) and informed written consent was obtained from each participant before their enrolment in the study.

Comment [A12]: Ethical approval obtained from Rivers State University Teaching Hospital (RSUTH) could be added in the supplementary material.

Eligibility Criteria

Inclusion Criteria

Comment [A13]: 2.1.1 Sub-subheading (ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.
For example: **2.3.1 Inclusion Criteria** (arial, 10 font, bold ...)

Only subjects who gave informed and written consent were included in the study. Females within the reproductive age of 15-55 years of age and those who were sexually active were enrolled.

Comment [A14]: If woman doesn't have relation sexual it's not participate of study? describe

Exclusion Criteria

Females who were not within the reproductive age of 15-55 years and females did not give written consent were not recruited. Females who were not registered with RSUTH were also not selected.

Sample Collection and Analysis

A total of 200 High vaginal swab (HVS) specimen using sterile cotton swab sticks were aseptically collected from consented participants across the study area by qualified female medical laboratory scientists and 450 urine specimens were collected by the patients themselves using clean sterile sample bottles as directed. The HVS and urine samples were moved to the laboratory for microscopic smear examination for the trophozoites of *T. vaginalis*. Prior to smear preparation, the samples were cultured using *T. vaginalis* medium[6,9]

Statistical Analysis

Data obtained was entered into Microsoft Excel and analyzed using SPSS version 25. Descriptive statistics including frequencies and percentages were used to describe the population and also to estimate the prevalence of trichomoniasis in the studied population. Chi-square test was used to investigate associations among categorical variables. Confidence level was set at 95%.

Comment [A15]: The statistics agree. Very good.

Results

Table 1 shows the association between awareness and trichomoniasis among women attending RSUTH. The results presents that there is a significant association ($p < 0.05$) between knowledge of trichomoniasis and trichomonal infection. Similarly, there was a significant association ($p < 0.05$) between source of information and trichomoniasis among women attending RSUTH. There is no relation ($p > 0.05$) between knowledge of STI and trichomoniasis.

Comment [A16]: RESULTS (...)

Comment [A17]: And I know it is not the focus of the work, but it would be interesting to mention the general prevalence of VT found in the study, this is data that makes the study multiplyable and with citation potential.

Comment [A18]: Table or figure numbers should be mentioned in parentheses for better understanding.

Table 2 shows the association between personal hygiene and trichomoniasis among women attending RSUTH. The results presents that there is a significant association ($p < 0.05$) between wearing air-tight underwear and trichomonal infection. Similarly, there was a significant association ($p < 0.05$) between repeating underwear and trichomoniasis among women attending RSUTH. There was no association ($p > 0.05$) between the infection and other studied personal hygiene practices (hand washing, vagina washing, vagina washing method and clean after urination).

Comment [A19]: Reserve the paragraphs by entering the table number inside.

Table 3 shows the association between lifestyle and trichomoniasis among women attending RSUTH. The results presents that there is a significant association ($p < 0.05$) between engaging with unprotected sex and trichomonal infection. There was no association ($p > 0.05$) between the infection and other studied lifestyle (sexual activeness, protection use, number sexual partners, STI screening on partners and STI screening on self).

UNDER PEER REVIEW




Table 1.0: Prevalence of Trichomoniasis based on Awareness among the study population

Awareness	Responses	NE		NP (%)								x ²	P-value			
		Urine	HVS	OP		HIV		PTW		HV			Urine	HVS	Urine	HVS
				Urine	HVS	Urine	HVS	Urine	HVS	Urine	HVS					
Knowledge of Trichomoniasis	No	294	113	4(1.4)	4(3.5)	12(4.1)	8(7)	3(1)	2(1.8)	0(0)	0(0)	4.620	8.796	0.099	0.012	
	Yes	144	85	2(1.4)	3(3.5)	1(0.7)	0(0)	0(0)	0(0)	0(0)	0(0)					
	I don't know	12	2	0(0)	1(50)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)					
Knowledge of STI	No	46	13	1(2.2)	1(7.7)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0.812	0.029	0.367	0.865	
	Yes	404	187	5(1.2)	7(3.7)	13(3.2)	8(4.3)	3(0.7)	2(1.1)	0(0)	0(0)					
Source of Information	Media	9	6	0(0)	2(33.3)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	5.438	12.701	0.245	0.005	
	HCP	59	27	2(3.4)	1(3.7)	1(1.7)	0(0)	0(0)	0(0)	0(0)	0(0)					
	Books	74	52	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)					
	Fam/friends	1	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)					
	I don't know	307	115	4(1.3)	5(4.3)	12(3.9)	8(7)	3(1)	2(1.7)	0(0)	0(0)					

NE = Number Examined; NP = Number Positive; HVS = High Vaginal Swab; OP = Out Patients; HIV = HIV Positive Persons; PTW = Pregnant Women; HV = Healthy Volunteers. (P<0.05).

Table 2.0: Prevalence of Trichomoniasis based on Personal Hygiene among the study population

Personal Hygiene	Responses	NE		NP (%)								x ²		P-value	
		Urine	HVS	OP		HIV		PTW		HV					
				Urine	HVS	Urine	HVS	Urine	HVS	Urine	HVS	Urine	HVS		
Hand washing Practice	Regularly	373	164	5(1.3)	7(4.3)	9(2.4)	4(2.4)	3(0.8)	2(1.2)	0(0)	0(0)	0.606	1.281	0.738	0.258
	Occasionally	76	36	1(1.3)	1(1.3)	4(5.3)	4(5.3)	0(0)	0(0)	0(0)	0(0)				
	Rarely	1	0	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
Vaginal washing	Regularly	383	176	6(1.6)	8(4.5)	10(2.6)	6(3.4)	2(0.5)	0(0)	0(0)	0(0)	0.198	1.957	0.656	0.162
	Occasionally	67	24	0(0)	0(0)	3(4.5)	2(8.3)	1(1.5)	2(8.3)	0(0)	0(0)				
	Rarely	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
Vaginal wash Method	Water	436	200	6(1.4)	8(4)	13(3)	8(4)	3(0.7)	2(1)	0(0)	0(0)	0.743	NIL	0.389	NIL
	Soaps	14	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
Clean after Urination	Regularly	344	176	4(1.2)	8(4.6)	8(2.3)	6(3.4)	3(0.9)	2(1.1)	0(0)	0(0)	2.028	0.015	0.363	0.903
	Occasionally	95	24	2(2.1)	0(0)	5(5.3)	2(8.3)	0(0)	0(0)	0(0)	0(0)				
	Rarely	11	0	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
Wearing air-tight Underwear	Regularly	97	41	2(2.1)	4(9.6)	5(5.2)	4(9.6)	0(0)	0(0)	0(0)	0(0)	1.712	8.657	0.425	0.013
	Occasionally	187	74	2(1.1)	3(4.1)	5(2.7)	2(2.7)	2(1.1)	2(2.7)	0(0)	0(0)				
	Rarely	166	85	2(1.2)	1(1.2)	3(1.8)	2(2.4)	1(0.6)	0(0)	0(0)	0(0)				
Repeating Underwear	Regularly	23	10	0	1(10)	4(17.4)	0(0)	0(0)	0(0)	0(0)	0(0)	8.369	4.228	0.015	0.121
	Occasionally	120	41	2(1.7)	5(12.2)	3(2.5)	0(0)	1(0.8)	2(4.9)	0(0)	0(0)				
	Rarely	307	149	4(1.3)	2(1.3)	6(2)	8(5.4)	2(0.7)	0(0)	0(0)	0(0)				

NE = Number Examined; NP = Number Positive; HVS = High Vaginal Swab; OP = Out Patients; HIV = HIV Positive Persons; PTW = Pregnant Women; HV = Healthy Volunteers. (P<0.05).

Table 3.0. Prevalence of Trichomoniasis based on Lifestyle among the study population

Lifestyle	Responses	NE		NP (%)								x ²		P-value	
		Urine	HVS	OP		HIV		PTW		HV					
				Urine	HVS	Urine	HVS	Urine	HVS	Urine	HVS	Urine	HVS		
Sexually active	No	57	14	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	3.355	1.489	0.067	0.222
	Yes	393	186	6(1.5)	8(4.3)	13(3.3)	8(4.3)	3(0.8)	2(1.1)	0(0)	0(0)				
Engage in unprotected sex	Regularly	229	93	1(0.4)	5(5.4)	7(3.1)	6(6.5)	3(1.3)	2(2.2)	0(0)	0(0)	5.642	7.22	0.13	0.027
	Occasionally	83	48	2(2.4)	3(6.3)	5(6)	0(0)	0(0)	0(0)	0(0)	0(0)				
	Rarely	73	46	3(4.1)	0(0)	1(1.4)	2(4.4)	0(0)	0(0)	0(0)	0(0)				
	N/A	65	13	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
Protection Used	Condoms	127	80	3(2.4)	3(3.8)	6(4.7)	2(2.5)	0(0)	0(0)	0(0)	0(0)	1.838	1.252	0.399	0.535
	Medication	25	17	1(4)	2(11.8)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
	N/A	298	103	2(0.7)	3(2.9)	7(2.3)	6(5.8)	3(1)	2(1.9)	0(0)	0(0)				
Number of sexual Partners	1	374	176	6(1.6)	8(4.6)	13(3.5)	8(4.6)	3(0.8)	2(1.1)	0(0)	0(0)	4.7	0.095	2.697	0.26
	2	7	5	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
	N/A	69	19	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
STI Screening on Partners	No	307	129	4(1.3)	6(4.7)	11(3.6)	6(4.7)	3(1)	2(1.6)	0(0)	0(0)	3.457	2.679	0.178	0.262
	Yes	53	50	1(1.9)	2(4)	2(3.8)	2(4)	0(0)	0(0)	0(0)	0(0)				
	N/A	90	21	1(1.1)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				
STI Screening on self	Regularly	43	39	1(2.3)	2(5.1)	1(2.3)	0(0)	0(0)	0(0)	0(0)	0(0)	0.933	1.163	0.627	0.559
	Occasionally	104	52	0(0)	0(0)	6(5.8)	4(7.7)	1(1)	2(3.8)	0(0)	0(0)				
	Rarely	277	99	4(1.4)	6(6.1)	6(2.2)	4(4)	2(0.7)	0(0)	0(0)	0(0)				
	N/A	26	10	1(3.4)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)	0(0)				

NE = Number Examined; NP = Number Positive; HVS = High Vaginal Swab; OP = Out Patients; HIV = HIV Positive Persons; PTW = Pregnant Women; HV = Healthy volunteer. (P<0.05)

Discussion

Trichomoniasis is an unpleasant parasitic infection that can go undiagnosed for years and is mainly transmitted by asymptomatic carriers. The prevalence of *T. vaginalis* infection, co-infection with bacterial infection, bacterial vaginosis other sexually transmitted infections (HIV) and attendant high risk of adverse pregnancy outcomes, pelvic inflammatory diseases, and tubal infertility make trichomoniasis an infection of compelling public health concern [2,10].

Comment [A20]: Present reference of this phrase.

The study showed that lack of awareness of *Trichomonas vaginalis* is a risk factor contributing to the spread of the disease as a prevalence rate of 50% (P=0.012) was recorded among subjects who have no knowledge of trichomoniasis. There was also an association between source of information and trichomoniasis among the women. This implies that both knowledge of trichomoniasis and source of information are notable risk factors for trichomonal infection. Women who wear tight underwear and repeat underwear recorded a high prevalence rate of 9.6% (P=0.013) for HVS 17% (P=0.015) for urine specimen. Subjects who clean the vulva after urinating occasionally recorded a prevalence of 8.3% but is not statistically significant (P>0.05). These findings are in agreement with study recorded by Amadi and Nwagbo (2013) where high prevalence was observed among women who did not clean after urinating, wear damp underwear, and tight underwear which irritates the vaginal and upsets bacterial balance predisposing women to trichomoniasis.[11] Hand washing practice and vagina washing is not closely associated with trichomoniasis infection. The prevalence of trichomoniasis among females who are not sexually active was low (0%) which implies that there negligible tendencies for women who are not sexually active to get infected with *Trichomona vaginalis*, therefore, it is not considered as a risk factor for the infection. This study recorded high prevalence among subjects who engage in unprotected sex regularly (6.5%) and a significant (P=0.027) association with the infection. This implies that unprotected sex is a notable risk factor for trichomoniasis among women. This study agrees with Crucitti *et al.*, (2011) that multiple sexual partners and lifestyle are determinants of trichomoniasis infection.[12] Prevalence of subjects who used medications as preventive measure during sexual intercourse was (11.8%), those who had one sexual partner was (4.6%), those who do not carry out STI screening on their partners was (4.7%) and those screening themselves occasionally was (7.7%). There was no significant association between those risk factors and trichomoniasis. It is however a concern that self-reported data on sexual behaviour are usually not reliable which may in-turn affect the outcome and decision of such study.

Conclusion

Comment [A21]: CONCLUSION (...)

This work has demonstrated that amongst selected possible risk factors studied, engaging in unprotected sex, wearing air-tight underwear, repeating underwears, knowledge of trichomoniasis and source of information were identifiable significant risk factors for trichomonal infection among women attending RSUTH.

Comment [A22]: You could conclude by mentioning whether educational studies will be carried out and the importance of further studies on the topic.

References

1. Ghosh, , Muwonge, R., Mittal, S., Banerjee, D., Kundu, P. and Mandal R. (2017). Association between high risk human papilloma virus infection and co-infection with *Candida spp.* and *Trichomonas vaginalis* in women with cervical premalignant and malignant lesions. *Journal of clinical virology*, 87, 43-48.
2. Toth, M. D. (2013). Trichomoniasis and infertility. *Journal of Infectious Diseases*, 19, 68-70.
3. Beri, D., Yadav, P., Devi. H. R. N., Narayana, C., Gadara, D. and Tatu, U. (2020). Demonstration and characterization of cyst-like structures in the life cycle of *Trichomonas vaginalis*. *Frontiers in cellular and infection microbiology*. 9, 430.
4. Okoko, F. J. (2011). Prevalence of trichomoniasis among women at Effurun metropolis, Delta State, Nigeria. *Continental Journal of Biological Sciences*, 4(2), 45-48.
5. Ogomaka, I. A., Nwachinemere, A. V. and Emmanuel, I. O. (2018). Prevalence of Trichomoniasis among adults in Oru-East L.G.A, Imos state, Nigeria. *Archives of Clinical Microbiology*, 9(2), 79.
6. Adekunle, O. N., Mogaji, H. O., Adeleke, M. T., Adesetan, T. O., Agbolade, O. A. and Agbolade, O. M. (2021). Prevalence of trichomoniasis and associated risk factors among female attendees of primary health care centres in Ijebu-North, Southwest Nigeria. *Journal of Innovative Research in Life Sciences*, 3(2), 7-13.
7. Rhoda, N., Onosakponome, E. O., Brenda, A. N. and Tamunonengiye-Ofori, L. (2022) Evaluation of Malaria Parasitaemia among COVID-19 Patients in Rivers State, Nigeria. *Journal of Applied Life Sciences International* 25(4): 12-18, DOI: 10.9734/JALSI/2022/v25i430297
8. Onosakponome, E. O., Ezeanyagu O. C., Ejinaka O. R., Obeta M. U. and Agbalaka P. I. (2022) Impact of Water Sources on Schistosomiasis Transmission and Urine Indicators. *African Journal of Health Sciences* 35(6) : 697-704 .doi.org/10.4314/ajhs.v35i6.4
9. Michael, W. and Onosakponome, E. O. (2021): Evaluating Prevalence and Misdiagnosis of Plasmodium Using Microscopy Compared With Polymerase Chain Reaction Technique in Two Tertiary Care Hospitals in Rivers State, Nigeria *International journal of infection*; 8(1):e109411. doi: 10.5812/iji.109411.
10. Akinbo, F. O., Makobia, C. N. and Ande, A. B. A. (2017). Prevalence of trichomoniasis among pregnant women in Benin city. *Sahel Medical Journal*, 20(2), 67-71.
11. Amadi, A. N. C. and Nwagbo, A. K. (2013). *Trichomonas vaginalis* infection among women in Ikwuano, Abia State, Nigeria. *Journal of Applied Sciences and Environmental Management*, 17, 389-393.
12. Crucitti, T., Jespers, V., Mulenga C., Khondowe S. and Vandepitte J. (2011) Non-Sexual Transmission of *Trichomonas vaginalis* in Adolescent Girls Attending School in Ndola, Zambia. *Public Library of Science*, 6(1), 311-316