

The comparison of antioxidative and cytotoxic activities of fresh and dried *Piper betle* L. leave extracts on MCF-7, HELA and SK-LU-1

ABSTRACT

The use of herbal medicines produced from plants and herbs, as one element of complementary and alternative medicine, is still increasing worldwide. In this study, *Piper betle* L. leaf in two states, fresh and dried, was tested to explore its compounds and total phenolic and flavonoid contents. From there evaluated its antioxidant, and cytotoxic activities *in vitro*. Gas chromatography–mass spectrometry is a method for identifying compounds (GC-MS). The total phenolic content was measured by using Folin-Ciocalteu method and the total flavonoid content was based on a complex formation with $AlCl_3$. The antioxidant activity was evaluated by DPPH and scratch assays, respectively. The cytotoxic activity was performed against MCF7, Hela, and SKLU1 cancer cell lines using sulforhodamine B assay. Toxicity testing for normal cells was also investigated. Results show the dried betel leaf extract (DBLE) contains eugenol (37.56%) as the primary compound, while 2,5-dimethylbenzoic acid (89.58%) was the primary compound in fresh betel leaf extract (FBLE). The extracts are rich polyphenols and flavonoids and have strong antioxidant activity. The polyphenol content in DBLE (240.9 mg GAE/g) is higher than FBLE (165.2 mg GAE/g). Especially, the DBLE has shown significant DPPH scavenging activity ($IC_{50} = 3.21 \mu\text{g/mL}$) when compared with standard ascorbic acid ($IC_{50} = 6.71 \mu\text{g/mL}$) and FBLE ($IC_{50} = 22.97 \mu\text{g/mL}$). The cytotoxic activity of FBLE against MCF-7 and Hela cell lines shows IC_{50} value above $50 \mu\text{g/mL}$, while DBLE shows IC_{50} value of $21.88 \mu\text{g/mL}$ and $26.68 \mu\text{g/mL}$, respectively, with no difference for the SKLU1 line (IC_{50} values between 35.0 – $38.0 \mu\text{g/mL}$). There is a positive correlation between the phenolic content, antioxidant, and cytotoxic activities of the extracts. In conclusion, DBLE exhibits higher antioxidant and cytotoxic activities than FBLE.

Keywords: Betel leaf (*Piper betle*), fresh and dried leaves, antioxidant, cytotoxicity

INTRODUCTION

“Cancer is the leading cause of death worldwide. Among them, lung cancer, breast cancer, and cervical cancer are common cancer types, especially affecting women the most” (Sun et al., 2021).

Currently, failures in cancer treatment are associated with drug resistance and toxicity. There is growing evidence that natural compounds are highly specific to tumor cells and have fewer side effects on normal cells; as a result, they offer great promise for the future treatment of cancer. The use of natural compounds as an adjuvant treatment in targeting the apoptotic pathway and inhibiting the self-renewal pathway of cancer cells (Hh, Notch, WNT, and BMI1) inhibits tumor recurrence and chemotherapy drug resistance (Hashem et al., 2022).

“Approximately 80% of the population in developing countries is dependent on natural or herbal medicines to treat various diseases. *Piper betle* is a traditional herb of the Piperaceae family which is used by many countries due to its wealthy content of macronutrients and micronutrients such as vitamin C, vitamin A, minerals, riboflavin, thiamine, tri-terpenoids, steroids, alkaloids, amino

acids, tannins, essential oils, phenolics, flavonoids... There have been reports of biological benefits of betel leaf including platelet suppression, antidiabetic activity, immunomodulatory properties, and anti-allergic activity. These activities are due to the high antioxidant effect of the compounds contained in this plant through DPPH, superoxide, and nitric oxide free radical scavenging activities" (Rintu et al., 2015). "Studies have shown that the leaf extract of *Piper betle* has phenolic compounds such as hydroxychavicol, chavibetol, eugenol, etc. Hydroxychavicol in betel leaf induces cell cycle arrest at the S phase or G2/M and apoptosis in pancreatic cancer cells" (Majumdar & Subramanian, 2019). Water extract from betel leaf activates ATM, p73, and JNK pathway expressions, and inhibits *in vivo* mouse tumor growth via the MAPK-p73 pathway in human

hepatocellular carcinoma (Wu et al., 2014); hexane extract exhibits significant cytotoxicity to HeLa cells (Sanubol et al., 2017). "Extracting betel leaves with different extraction techniques (soxhlet, sonication, and maceration) and different solvents (water, methanol, ethyl acetate, and hexane) showed various results of total phenolic and flavonoid content, and substance content. Therefore, the antioxidant and cytotoxic activities were also different" (Das et al., 2019; Abraham et al., 2012). From the studies, it was found that different extraction methods play an important role in the presence and activity levels of phytochemicals. This is what accounts for their different pharmacological properties, which is essential for assessing their potential for human health benefits.

To study the differences between betel leaf extract extracted from dried leaves and fresh leaves affecting the biological activity, we analyzed the changes of the main phytochemicals, polyphenol and flavonoid content. Then compare the antioxidant and cytotoxic activities *in vitro* of the extracts. The results indicated that the content of phytochemicals, polyphenols and flavonoids, antioxidant and cytotoxic activity against cancer cells have significant differences between fresh and dried extracts of betel leaves.

MATERIALS

Plant

Fresh betel leaves (*Piper betle* L.) in the mature stage were collected from 8 a.m. to 10 a.m. in Thua Thien Hue province, Viet Nam. Choosing leaves that are intact, fresh, and free from pests.

Cell lines

Three human cancer cell lines including breast cancer (MCF-7), cervical cancer (HeLa) and lung cancer (SK-LU-1) cells are provided by the Institute of Biotechnology, Vietnam Academy of Science and Technology.

MSC was isolated from Wharton's jelly of human umbilical cords provided by Hue Central Hospital (Hue, Vietnam).

METHODS

Extract preparation

Betel leaves were cleaned and drained at room temperature. To collect the maximum possible content of bioactive phytochemicals for developing and applying in specific areas, betel leaf extract was examined in two states, fresh and dry.

For fresh betel leaf extract (FBLE), the leaves were crushed and soaked with ethanol 70% v/v in a ratio of 1:10 (w/v) for five hours at 50°C; stirring occasionally (Thi et al., 2021). For dried betel leaf extract (DBLE), the fresh leaves were dried in an oven at 40°C and ground into fine powder. The powder (150 g) was soaked with absolute ethanol (500 mL) for seven days at room temperature; stirring occasionally (Valle et al., 2016). The samples were then filtered and evaporated at 40-50°C until getting the dark brown concentrated solution (Murugesan et al., 2017). Store samples at 4°C.

Gas chromatography-mass spectrometric (GC-MS) analysis

The analysis of extract compounds was performed on an Agilent GC-MS instrument equipped with a GC7890A gas chromatograph and an MS 5975C mass spectrometer detector. The column used in the analysis was the DB-XLB capillary column (60m x 0.25 mm x 0.25 µm). The extract was diluted with 3% dichloromethane solvent and filtered through a 0.45 µm PTFE filter. Analyses using Isocratic modes were carried out using high-purity helium as carrier gas at a column flow rate of 1 mL/min with a ratio of 100:1. The temperature of the injection port was 250°C. The column temperature program was as follows: an initial temperature of 40°C, raised to 140°C at a rate of 20°C/min and held for five minutes, the final rate was raised from 4°C/min to 270°C. The mass spectrometry conditions included an ion source temperature of 230°C, ionization energy of 70 eV, and a mass scan range of 40–500 amu. Peak areas were used for quantifying the constituent percentage in total betel leaf extracts.

Determination of total polyphenol content

The total phenolic content (TPC) in FBLE and DBLE were estimated using the Folin–Ciocalteu (F-C) reagent described by Fedura et al. (2019).

The F-C 10% reagent was diluted with distilled water just before the experiment. Different solutions of gallic acid (50, 100, 150, and 200 µg/mL) and betel leaf extracts (1 mg/mL) were prepared in methanol. A mixture of 0.5 mL of extract or standard solution at each concentration was mixed with 2.5 mL of F-C 10% reagent for several seconds. After 4 min, 1 mL of 7.5% Na₂CO₃ was added and incubated at room temperature in the dark. Absorbance was measured at 760 nm after two hours.

The TPC was calculated based on the gallic acid calibration curve using the formula:

$$C = c \times V / m$$

where C = TPC (mg GAE/g), c = concentration of gallic acid established from the calibration curve (mg/mL), V = volume of extract (mL) and m = weight of plant extract.

Results were expressed as milligrams of gallic acid equivalents per gram of extract (mg GAE/g extract). All experiments were performed in triplicate.

Determination of total flavonoid content

With slight modifications, the total flavonoid content (TFC) in the extracts was estimated by the aluminum chloride (AlCl₃) colorimetric method described by Marinova *et al.* (2005).

In methanol, different solutions of quercetin standard (20, 40, 60, 80, and 200 µg/mL) and betel leaf extracts (1 mg/mL) were produced. Following the addition of 0.3 mL of 5% NaNO₂, a sample containing 1 mL of extract or standard solution at each concentration was added separately to 4 mL of distilled water. After five minutes, 0.3 mL of 20% AlCl₃ is added and kept for 6 min. Finally, 2 mL of 1 M NaOH solution was added, then up to 10 mL with distilled water and mixed well. The optical absorbance of the reaction solution was measured at 510 nm after 10 minutes. Similar to the determination of TPC, TFC was calculated based on the quercetin calibration curve according to the formula:

$$C = c \times V / m$$

where C = TFC (mg QE/g), c = concentration of quercetin established from the calibration curve in (mg/mL), V = volume of extract (mL) and m = weight of plant extract.

Results were expressed as milligrams of quercetin equivalents per gram of extract (mg QE/g extract). All experiments were performed in triplicate.

DPPH radicals scavenging assay

The antioxidant activity of betel leaf extracts was determined through DPPH free radical scavenging according to Jahan *et al.* (2010) with some modifications.

A 100 µM DPPH solution was prepared in methanol solution. Diluting the extract into different concentrations (5-25 µg/mL). The mixture consisting of 1.5 mL of 100 µM DPPH and 1.5 mL of extract was shaken well and then incubated in the dark at room temperature for 30 min. The absorbance of the mixture was spectrophotometrically measured at $\lambda_{\max} = 517$ nm and compared to the standard antioxidant (ascorbic acid).

Radical scavenging activity is expressed as inhibitory concentration (IC₅₀), i.e., extract concentration necessary to decrease the initial concentration of DPPH by 50%. The percentage of DPPH inhibition was calculated by using the following formula:

$$\text{DPPH inhibition (\%)} = [(A_0 - A) / A_0] \times 100\%$$

Where: A₀ = the absorbance of blank sample solution without the compound to be tested A = the absorbance of the tested sample.

Cell culture

MCF-7 breast cancer, Hela cervical cancer, and SK-LU-1 lung cancer cells were maintained in a DMEM growth medium containing 10% FBS, supplemented with 2 mM L-glutamine, 1% penicillin/streptomycin, and 1 mM sodium pyruvate. Mesenchymal stem cells were cultured in the growth medium of DMEM/F12 medium containing 10% FBS and 1% antibiotic. Cells were grown to confluence in a humidified atmosphere containing 5% CO₂ at 37°C. Morphology was examined under inverted optical microscopy.

Testing the effect of *Piper betle* L. extract on MSC

Mesenchymal stem cells (MSCs) are multipotent stem cells found in bone marrow that are important for making and repairing skeletal tissues. In addition, MSCs have a migration ability when activated, they can migrate into other injured and inflammatory sites to repair and regenerate. In this study, MSC was used for toxicity testing of betel leaf extracts on normal cells.

These secondary MSC were cultured on 6-well plates with a density of 1.000 cells/cm^2 in the growth medium. When the cells were approximately 60% confluent, the betel leaf extracts at various concentrations were added to the dishes. Cell monitoring was extended to 48 hours. Images of the MSC at each test concentration were captured in three different fields to estimate error bars. MSC numbers were assessed by using the ImageJ software.

Cytotoxicity test using Sulforhodamine B (SRB) assay

The cytotoxicity test in cancer cells was performed according to the method of Skekan *et al.* (1990) by sulforhodamine B (SRB) assay.

Trypsinization to collect cell pellets. Adjust the cell concentration with medium to obtain the appropriate cell culture density per well of the 96-well plate at a volume of 190 μl . The extracts were dissolved in DMSO and diluted with culture medium (FBS-free) at concentrations of 2000 $\mu\text{g/mL}$, 400 $\mu\text{g/mL}$, 80 $\mu\text{g/mL}$, and 16 $\mu\text{g/mL}$. Aspirate 10 μl of the diluted extract introduced into wells containing the prepared cancer cell fluid mixture to achieve a final concentration of 100 $\mu\text{g/mL}$, 20 $\mu\text{g/mL}$, 4 $\mu\text{g/mL}$, and 0.8 $\mu\text{g/mL}$. After culturing for 72 hours, 20% TCA solution was added to immobilize cells for 60 min at 4°C . DMSO 1% was used as a control at day 0 (the final concentration in the well was 0.05%). Cells in this well were incubated for only one hour. After fixation, cells were stained with 100 μl of 0.4% SRB for 30 min at 37°C . Remove residual dye by washing it three times with 1% acetic acid and then drying it at room temperature. Then, the SRB was dissolved in 10 mM unbuffered Tris base, gently shaken for 10 min, and read the OD results at 540 nm on an ELISA Plate Reader (Biotek). Ellipticine at final concentrations in each well of 10, 2, 0.4, and 0.08 $\mu\text{g/mL}$ was used as a positive control. The test was repeated 3 times. The IC_{50} value (concentration that inhibits 50% of growth) was determined using TableCurve 2Dv4 computer software.

The percentage inhibition of cell growth was determined by the following formula:

$$\% \text{ cell inhibition} = 100 - \left[\left\{ \frac{A_t - A_b}{A_c - A_b} \right\} \times 100 \right]$$

Where: A_t = Absorbance value of the test compound; A_b = Absorbance value of blank; A_c = Absorbance value of the control.

Statistical analysis

The collected data were expressed as mean \pm standard deviation (mean \pm SD). Microsoft Excel software has been used to analyze data and draw graphs. The $p < 0.05$ indicates statistically significant differences. The significant difference between the experimental samples was performed by one-way ANOVA.

RESULTS

Obtaining *P. betle* extract

The distillation method extracted fresh and dry betel leaves on a laboratory scale. After being extracted, both extracts were in a concentrated state with a dark brown color and the characteristic pungent smell of betel. In terms of solubility, DBLE was more soluble in water than FBLE. All extracts were easily soluble in alcohol. The yield extracted from dried betel leaves (8.89%) was higher than that of fresh ones (4.95%) (Table 1).



Figure 1. Betel leaf

Table 1. The recovery efficiency of two types of extracts

Extract	Weight (g)		Recovery efficiency (%)
	Material	Extract	
FBLE	200	9.90	4.95
DBLE	100	8.89	8.89

Gas chromatography-mass spectroscopic (GC-MS) analysis

GC-MS results have identified FBLE contained the chemical compositions including 2,5-dimethylbenzoic acid (89.58%), 2-Methoxy-1-hydroxy-4-allylbenzene (3.18%), Eugenol (2.89%), Allylguaiacol (2.85%) and Chavicolacetate (1.49%). Chemical compositions of DBLE identified were Eugenol (37.56%), Chavicol acetate (37.34%), Caryophyllene (12.71%), Acetyleneugenol (10.17%) and Humulene (2.22%) (Table 2). The result showed that DBLE's main compounds Chavicolacetate and Eugenol were not presented in FBLE. Meanwhile, the main chemical constituent identified in FBLE was 2,5-dimethylbenzoic acid which accounted for 89.58% and was not presented in DBLE.

Table 2. Chemical compositions of FBLE and DBLE

FBLE			DBLE		
No	Chemical compositions	%	No	Chemical compositions	%
1	2,5-Dimethylbenzoic acid	89.58	1	Eugenol	37.56
2	2-Methoxy-1-hydroxy-4-allylbenzene	3.18	2	Chavicolacetate	37.34
3	Eugenol	2.89	3	Caryophyllene	12.71
4	Allylguaiacol	2.85	4	Acetyleneugenol	10.17
5	Chavicolacetate	1.49	5	Humulene	2.22

Some studies also reported compounds in betel leaves similar to our study. GC-MS analysis of *P. betle* leaf extract collected from Denpasar, Indonesia contained 31 compounds including eugenol (25.03%), and 2,5-

dimethylbenzoic acid (12.08%) which were two main components (Pratiwi & Muderawan, 2016). Methanol extracts from *P. betle* leaves in India revealed that the main active compounds were 2,5-dimethylbenzoic acid, 3,5-dimethylbenzoic acid, methionine, ... (Selvaraj et al., 2022). Betel leaf extract from Indonesia analyzed components such as chavicol, eugenol, caryophyllene, xylene and chalcone (Jenie & Apriyanto, 2008). The presence of caryophyllene, eugenol, and acetyl eugenol in the ethanol extract from the leaves of *P. betle* from Malaysia was also confirmed by HPTLC and GCMS analysis (Annegowda et al., 2013). The chemical composition of betel leaves in different areas of Odisha also contained components such as eugenol, caryophyllene, acetyl eugenol, humulene, cadinene, 2-aminocarbonylbenzoic acid, etc (Das et al., 2016).

Total phenolic and flavonoid contents

TPC and TFC were calculated based on the respective standard linear of gallic acid ($y = 0.009x + 0.1222, R^2 = 0.9931$) and quercetin ($y = 0.0134x + 0.0078, R^2 = 0.9995$). The results are presented in Table 3.

Table 3. TPC (mg GAE/g) and TFC (mg QE/g) of DBLE and FBLE.

Extract	TPC (mg GAE/g)	TFC (mg QE/g)
FBLE	165.2 ± 4.6	76.4 ± 2.1
DBLE	240.9 ± 0.11	82.1 ± 2.9

TPC and TFC of DBLE were higher than that of FBLE, especially the TPC of both extracts was considerably different, ranging from 165 to 241 mg GAE/g. The respective TPC and TFC found in FBLE were 165.2 ± 4.6 mg GAE/g and 76.4 ± 2.1 mg QE/g. Meanwhile, TPC and TFC recorded in DBLE were 240.9 ± 0.11 mg GAE/g and 82.1 ± 2.9 mg QE/g.

Our study shows that dried betel leaf extract has similar TPC and higher TFC than the betel leaf study in Ho Chi Minh City (249.96 mg GAE/g and 27.82 mg RE /g) and Binh Duong province (386,343 mg GAE/g and 55,073 mg QE/g) in Vietnam (Nguyen et al., 2020), (Duron et al., 2022). The dried betel leaf from Odisha extracted by sonication, Soxhlet, and maceration methods using acetone as solvents found TPC from 50 – 57.6 mg GAE/10 mg and TFC from 32.1 – 49.79 mg QE/mg (Das et al., 2019). The extract from fresh betel leaves in Indian regions had TPC from 95–128 mg GAE/g and TFC from 51–62 mg catechin which was lower than in our study (Harini et al., 2018). The difference in total phenolic and flavonoid contents of betel leaf extract in distinct studies depends on geographical location as well as an extraction method.

Antioxidant activity

The antioxidant effect of FBLE and DBLE was determined based on DPPH radical scavenging activity. With the aim of comparison, ascorbic acid was selected to be a positive control. The DPPH assay results are shown in Table 4 and Figure 2. The DPPH scavenging ability is proportional to the concentration of extract. The higher the concentration is, the better the free radical scavenging capacity is. The inhibition of DPPH of DBLE was considerably lower than that of FBLE with IC₅₀ values of 3.21 µg/mL and 22.97 µg/mL.

respectively. It can be seen that the antioxidant activity of DBLE is 7 times higher than that of FBLE. Furthermore, the IC₅₀ value of DBLE was lower than that of the positive control-ascorbic acid (3.21 µg/mL and 6.71 µg/mL respectively), which indicates that the crude extract from the dried betel leaf has a strong antioxidant capacity.

Table 4. DPPH radical scavenging activity of both extracts and ascorbic acid


Concentration (µg/ml)	Inhibition(%)		
	FBLE	DBLE	Ascorbic acid
5	4.83 ± 0.59	47.92 ± 1.74	37.38 ± 0.56
10	10.99 ± 0.47	69.9 ± 1.55	66.15 ± 0.85
15	29.51 ± 0.88	82.37 ± 0.84	78.23 ± 0.99
20	39.31 ± 1.24	92.47 ± 0.75	84.06 ± 0.72
25	57.99 ± 1.10	95.04 ± 1.05	95.71 ± 0.87
IC₅₀	22.97 µg/mL	3.21 µg/mL	6.71 µg/mL

Several studies have also confirmed the high antioxidant activity of betel leaf extract. For instance, the dried leaf *Piper betle* extract with ethanol collected in Phayao Province in Thailand exhibited antioxidant activity with IC₅₀ of 30.0 ± 0.1 µg/mL (Boontha et al., 2019); the ethyl acetate extract from Malaysian dried leaf betel showed the radical scavenging activities against DPPH with IC₅₀ of 40.0 µg/mL (Abrahim et al., 2012); and the fresh betel leaf extract from Indonesian with ethanol is found to exhibit antioxidant activity with the IC₅₀ of 17.4 ppm against DPPH (Sartini et al., 2020); fresh betel leaves in Kuala Lumpur demonstrated the DPPH scavenging activity with an IC₅₀ value of 179.5 ± 93.1 µg/mL (Alam et al., 2023).

Testing the effect of *Piper betle* L. extract on MSC

The toxicity of betel leaf extracts on MSC by adding dilutions of betel leaf extract to the culture medium to determine cytotoxic concentrations. MSC with 60% confluency was exposed to different concentrations (25 µg/mL-125 µg/mL) of FBLE and DBLE, monitoring for cytotoxicity for 48 hours at concentrations of FBLE and DBLE (Figure 2).

A







Figure 2. Effect of *Piper betle*L. extracts at different concentrations on MSC was presented in photos(A) and graph(B). Percentage of cell viability at significant concentrations compared to control (* $p < 0.005$; ** $p < 0.001$)

As shown in Figure 2, both FBLE and DBLE did not induce any significant cytotoxicity at concentrations up to 50 µg/mL after 48 hours. The results also indicated that the cell viability at concentrations greater than 75 µg/mL significantly decreased with increasing the exposure concentration, indicating their less efficiency for the growth of the cells. At a 125 µg/mL concentration, betel leaf extract was dramatically toxic to MSC cells. DBLE exhibited higher cytotoxicity to MSC than FBLE.

Cytotoxicity of extract toward cancer cell lines

According to the standards of the US National Cancer Institute, the extract is considered to have cytotoxic activity on cancer cells with $IC_{50} \leq 20$ µg/mL, while the purified substance is $IC_{50} \leq 5$ µM. After 72 hours of extract treatment, compared with the controls samples, cells shrank, rounded and lost cellular adhesion to the substrate. Most cells were even detached from the surface of the tissue cultured dishes plate and appeared floating in the culture medium which reduces the density of cells and cell clusters (Figure 3). These are characteristics of apoptosis. At a concentration of 100 µg/mL, FBLE was able to inhibit and kill 75.27% of MCF-7 cells, 85.5% of HeLa cells and 87.71% of SK-LU-1 cells. Meanwhile, DBLE at 100 µg/mL can inhibit and kill 96.35% of MCF-7 cells,

99.29% of Hela cells (almost completely) and 97.56% of SK-LU-1 cells (Table 5).





Figure 3. Cell proliferation inhibitory effect of FBLE and DBLE on MCF-7, Hela and SK-LU-1 cells at different concentrations after 72 hours in photos (A) and a graph (B)

Table 5. Inhibitory growth effects against different cancer cells of FBLE and DBLE compared with Ellipticine

Samples	Concentration (µg/ml)	% inhibition		
		MCF-7	Hela	SK-LU-1
FBLE	100	75.27±1.92	85.50±1.19	87.71±1.74
DBLE	100	96.35±2.29	99.29±1.71	97.56±1.42
Ellipticine	10	92.13±2.06	98.63±2.24	97.94±3.79

Table 6. IC₅₀ value of extracts in different cancer cell lines compared with Ellipticine

Samples	IC ₅₀ (µg/ml)		
	MCF-7	Hela	SK-LU-1
FBLE	56.05±5.46	56.7±4.16	35.07±1.39
DBLE	21.88±1.13	26.68±1.55	38.45±2.94
Ellipticine	0.34±0.03	0.37±0.02	0.39±0.02

FBLE and DBLE show high cytotoxic activity against MCF-7 and Hela cell lines, in which the cytotoxicity of DBLE is twice that of FBLE. Specifically, the cytotoxicity of DBLE and FBLE with IC₅₀ were 21.88 µg/mL and 56.05 µg/mL, respectively on MCF-7 cell line. Toward Hela cell line, the IC₅₀ of DBLE and FBLE were 26.68 µg/mL and 56.7 µg/mL, respectively. However, the IC₅₀ for cytotoxicity against SK-LU-1 of two extracts were almost the same at 38.45 µg/mL and 35.07 µg/mL, respectively (Table 6).

In this experiment, Ellipticine was the positive control to compare the cytotoxic activity of the extract. Ellipticine is a potent cancer cell inhibitor. However, it is not used for medical purposes due to its high cytotoxicity causing side effects such as nausea, high blood pressure, dry mouth, and fungal infections in the tongue and esophagus, etc. In our research, DBLE at 100 µg/mL concentration caused inhibition on MCF-7 (96.35%) and Hela (99.29%) cells similar to Ellipticine at 10 µg/mL (92.13% for MCF-7 cells and 98.63% for Hela cells) (Table 5).

The results of cancer-cell cytotoxicity in our study were higher than some studies in the world. The result of cytotoxicity on breast cancer by SRB revealed that ethanol extract from Thai betel leaves exhibited a cytotoxic effect with IC₅₀ = 114 µg/mL (Boontha et al., 2019). Among the four distinct extracts (water, methanol, ethyl acetate and hexane) of *Piper betle* leaves, the ethyl acetate extract exhibited the highest inhibitory effect on the proliferation of MCF-7 cells (IC₅₀ = 65 µg/mL) by the MTT assay (Abraham et al., 2012). The extract of fresh *Piper betle* leaf only

effectively inhibited the proliferation of the KB cell line without affecting the proliferation of HeLa cells (Fathilah et al., 2010). Pre-clinical evaluation in cancer treatment of extracts from *Piper* species based on their cytotoxicity by MTT assay, *Piper betle* exhibited antitumor activities on HeLa cells with $IC_{50} = 49.66$ mg/mL (Sanubol et al., 2017). Thus, with the ability of cytotoxic activity in our study, dried betel leaves can be a potential therapeutic agent in cancer treatment.

DISCUSSION

Scientists are constantly looking for natural drugs to replace chemical drugs in cancer therapy with the goal of less toxicity and side effects. Cytotoxic and antioxidant activities *in vitro* of *P. betle* have been reported in many studies but mainly focused on the aqueous extracts or polar solvents.

However, variations in composition and bioactivity can still occur depending on cultivar, location, growth conditions, and fresh or dried state, so data about antioxidant and cytotoxic activities remain relevant and crucial. In this study, we used ethanol and distilled water to extract fresh and dried leaves of *P. betle* in Vietnam to provide further insight into the antioxidant properties and cytotoxic effects on breast cancer (MCF-7), cervical cancer (HeLa), lung cancer (SK-LU-1) cells *in vitro*.

In our study, the cytotoxicity of DPLE was 2 times higher than that of the FBLE for MCF-7 and HeLa. This result may be due to the effect of the eugenol compound (EUG) with higher concentrations. Studies of eugenol showed its different effects on cancer cells; firstly, the effect of preventing cancer through its antioxidant effect, and secondly, the effect of killing cancer cells by affecting several signaling pathways. Eugenol regulated multiple molecular targets to mediate cytotoxicity against cancer types through inhibition of Nuclear factor-kappa B (NF- κ B) activation, regulation of prostaglandin synthesis, reduction of Cyclooxygenase-2 (Cox-2), B-cell lymphoma-2 (Bcl-2), interleukin-1 beta, and Proliferation Cell Nuclear Antigen (PCNA) activity; decreases in mitochondrial potential ($\Delta\Psi_m$) and inflammatory cytokines; increasing the generation of reactive oxygen species (ROS) and the expression of caspase-3, caspase-9, Bax, cytochrome (Cyt-c); S-phase cell cycle arrest; resulting in apoptotic cell death (Fathy et al., 2019; De Sa Junior et al., 2016). These series of findings suggest that EUG may induce apoptosis in cancer cells, e.g., breast cancer (MDA-MB-231, MCF-7), cervical cancer (SIHA, HeLa), glioblastoma (DBTRG-05MG), lung cancer (A459), colon cancer (NCM-460), other melanoma cells (SK-Mel-28, A2058) (Liang et al., 2015; Zari et al., 2021). Eugenol, one of the main components of betel leaf, has also been shown to possess anti-inflammatory effects in various animal models with different inflammatory agents (Aara et al., 2020).

In addition, the components of DBLE possess powerful antioxidant compounds such as acetyl eugenol, caryophyllene and humulene (Gushiken et al., 2022; Vanin et al., 2014). Caryophyllene inhibited the proliferation of glioblastoma, non-small-cell lung cancer, breast cancer, and myeloma cells (Irrera et al., 2020; Lei et al., 2021). Humulene inhibited hepatocellular carcinoma cell proliferation (Chen et al., 2019). Especially, the combination of humulene and caryophyllene was more effective in reducing the proliferation of MCF-7 cells (Legault & Pichette, 2007). Therefore, we

propose that the active constituents of the extract may interact additively or synergistically to protect against cancer.

Furthermore, the polyphenol content in DBLE (240.9 mg GAE/g) was higher than in FBLE (165.2 mg GAE/g). Polyphenols have been reported to have preventive effects against tumor initiation through numerous mechanisms such as the avoidance of genotoxic molecule formation and the blockade of mutagenic transforming enzyme activity; regulation of heme-containing phase I metabolic enzymes such as cytochrome P450s (CYPs); regulation of phase II metabolic enzymes to detoxify carcinogens, such as NADPH-quinone oxidoreductase-1 (NQO1), quinone reductase (QR), glutathione S-transferase (GST), as well as preventing DNA damage (Losada-Echeberría et al., 2017). These implied that the polyphenol content may contribute to the cytotoxicity of breast and cervical cancer in this study at the tested concentrations of betel leaf. The DPPH free radical scavenging ability of DBLE was also higher ($IC_{50}=3.21\ \mu\text{g/mL}$ compared to $22.97\ \mu\text{g/mL}$). It can be seen that the cytotoxicity against cancer cell potential of betel leaf extracts was correlated with their polyphenol content and antioxidant activity.

Although there was a difference in cytotoxic activity on MCF-7 breast cancer and Hela cervical cancer cells of both extracts, their cytotoxic effect was similar on SK-LU-1 lung cancer cells. That may be due to the dimethylbenzoic compound with high content in FBLE affected the apoptosis of SK-LU-1 cells by a mechanism that has not yet been discovered. The resulting difference between fresh and dried betel leaves may be due to the thermal effect of the drying process on the product, which leads to the degradation of some phytochemicals, namely 2,5-dimethylbenzoic acid, and optimal conditions for the formation of eugenol compounds in dried leaves. These results can serve as a premise for further studies on the molecular mechanisms and signaling pathways in anti-cancer of fresh and dried betel leaves.

CONCLUSION

The results revealed that the different states of betel leaf, dry or fresh, affected the chemical composition, the polyphenols and flavonoids content, antioxidant and cytotoxic activities. Dried betel leaf extract showed significantly better antioxidant DPPH radical scavenging activity than fresh betel leaf extract. The cytotoxic activity of fresh leaf extract and dried leaf extract against lung cancer line SK-LU-1 was the same. However, this activity on MCF-7 breast cancer and Hela cervical cancer cell lines of dried betel leaf extract was two times higher than that of fresh betel leaf extract. These results may be due to differences in polyphenol and flavonoid content and main components in each extract.

Our results of cytotoxicity on cancer cell lines provided a scientific basis for using betel leaf as a potential source of chemotherapeutic agents for the treatment of breast cancer, cervical cancer and lung cancer *in vivo*, especially using dried betel leaf. At the same time, it also serves as a premise for further studies about the cancer inhibition mechanism of dimethylbenzoic compounds present in the fresh betel leaf extract.

References

- Aara, A., Chappidi, V., Ramadas, M. N., & Care, P. (2020). Antioxidant activity of eugenol in Piper betle leaf extract. *Journal of Family Medicine and Primary Care*, 9(1), 327.
- Abraham, N. N., Kanthimathi, M., & Abdul-Aziz, A. J. B. c. (2012). Piper betle shows antioxidant activities, inhibits MCF-7 cell proliferation and increases activities of catalase and superoxide dismutase. *BMC complementary alternative medicine*, 12(1), 1-11.
- Alam, M. B., Park, N. H., Song, B.-R., & Lee, S.-H. J. A. (2023). Antioxidant Potential-Rich Betel Leaves (Piper betle L.) Exert Depigmenting Action by Triggering Autophagy and Downregulating MITF/Tyrosinase In Vitro and In Vivo. 12(2), 374.
- Annegowda, H., Tan, P., Mordi, M., Ramanathan, S., Hamdan, M., Sulaiman, M., & Mansor, S. (2013). TLC–bioautography-guided isolation, HPTLC and GC–MS-assisted analysis of bioactives of Piper betle leaf extract obtained from various extraction techniques: In vitro evaluation of phenolic content, antioxidant and antimicrobial activities. *Food Analytical Methods*, 6(3), 715-726.
- Boontha, S., Taowkaen, J., Phakwan, T., Woruaicha, T., Kamonnate, P., Buranrat, B., & Pitaksuteepong, T. (2019). Evaluation of antioxidant and anticancer effects of Piper betle L. (Piperaceae) leaf extract on MCF-7 cells, and preparation of transdermal patches of the extract. *Tropical Journal of Pharmaceutical Research*, 18(6), 1265-1272.
- Chen, H., Yuan, J., Hao, J., Wen, Y., Lv, Y., Chen, L., & Yang, X. (2019). α -Humulene inhibits hepatocellular carcinoma cell proliferation and induces apoptosis through the inhibition of Akt signaling. *Food Chemical Toxicology*, 134, 110830.
- Das, S., Parida, R., Sandeep, I. S., Kar, B., Nayak, S., & Mohanty, S. (2016). Chemical composition and antioxidant activity of some important betel vine landraces. *Biologia*, 71(2), 128-132.
- Das, S., Ray, A., Nasim, N., Nayak, S., & Mohanty, S. (2019). Effect of different extraction techniques on total phenolic and flavonoid contents, and antioxidant activity of betel vine and quantification of its phenolic constituents by validated HPTLC method. *3 Biotech*, 9(1), 1-8.
- De Sa Junior, P. L., Câmara, D. A. D., Costa, A. S., Ruiz, J. L. M., Levy, D., Azevedo, R. A., . . . Pessoa, N. D. S. (2016). Apoptotic effect of eugenol involves G2/M phase abrogation accompanied by mitochondrial damage and clastogenic effect on cancer cell in vitro. *Phytomedicine*, 23(7), 725-735.
- Dương, H. T., Loan, N. H., Ngân, P. T. K., Thư, L. H. A., Dũng, P. T., Thành, N. V. K., . . . Kiều, N. K. T. (2022). Đánh giá hoạt tính sinh học của cao chiết lá trầu không (Piper betle L.) thun hận bằng phương pháp chiết siêu âm. *Tạp chí Khoa học và Công nghệ Việt Nam*, 64(3), 37-42.
- Fathilah, A., Sujata, R., Norhanom, A., & Adenan, M. (2010). Antiproliferative activity of aqueous extract of Piper betle L. and Psidium guajava L. on KB and HeLa cell lines. *Journal of Medicinal Plants Research*, 4(11), 987-990.
- Fathy, M., Fawzy, M. A., Hintzsche, H., Nikaido, T., Dandekar, T., & Othman, E. M. (2019). Eugenol exerts apoptotic effect and modulates the sensitivity of HeLa cells to cisplatin and radiation. *Molecules*, 24(21), 3979.
- Gushiken, L. F. S., Beserra, F. P., Hussni, M. F., Gonzaga, M. T., Ribeiro, V. P., De Souza, P. F., . . . Takahira, R. K. (2022). Beta-caryophyllene as an antioxidant, anti-inflammatory and re-epithelialization activities in a rat skin wound excision model. *Oxidative medicine cellular longevity*, 2022, 9004014.
- Harini, S. S., Sougandhi, P. R., TENKAYALA, D. S. R., & Gopinath, K. R. (2018). Antioxidant activity (Phenol and flavonoid content) of three different cultivars of Piper Betle L. (Piperaceae). *Journal of Drug Delivery Therapeutics*, 8(5-s), 143-148.
- Hashem, S., Ali, T. A., Akhtar, S., Nisar, S., Sageena, G., Ali, S., . . . Elfaki, I. (2022). Targeting cancer signaling pathways by natural products: Exploring promising anti-cancer agents. *Biomedicine Pharmacotherapy*, 150, 113054.
- Irrera, N., D'Ascola, A., Pallio, G., Bitto, A., Mannino, F., Arcoraci, V., . . . Metro, D. (2020). β -caryophyllene inhibits cell proliferation through a direct modulation of CB2 receptors in

- glioblastoma cells. *Cancers*, 12(4), 1038.
- Jenie, B., & Apriantono, A. (2008). Antibacterial activity of green sirih (Piper betle L) extract towards food pathogens. *Jurnal Teknologi dan Industri Pangan*, 19(1), 1-1.
- Legault, J., & Pichette, A. (2007). Potentiating effect of β -caryophyllene on anticancer activity of α -humulene, isocaryophyllene and paclitaxel. *Journal of Pharmacy Pharmacology*, 59(12), 1643-1647.
- Lei, J., Wang, Q., Li, G., Li, Y., Zhang, P., & Xu, G. (2021). β -Caryophyllene from Chilli Pepper Inhibits the Proliferation of Non-Small Cell Lung Cancer Cells by Affecting miR-659-3p-Targeted Sphingosine Kinase 1 (SphK1). *International Journal of General Medicine*, 14, 9599.
- Liang, W.-Z., Chou, C.-T., Hsu, S.-S., Liao, W.-C., Shieh, P., Kuo, D.-H., ... Jan, C.-R. (2015). The involvement of mitochondrial apoptotic pathway in eugenol-induced cell death in human glioblastoma cells. *Toxicology Letters*, 232(1), 122-132.
- Losada-Echeberría, M., Herranz-López, M., Micol, V., & Barrajón-Catalán, E. (2017). Polyphenols as promising drugs against main breast cancer signatures. *Antioxidants*, 6(4), 88.
- Majumdar, A. G., & Subramanian, M. (2019). Hydroxychavicol from Piper betle induces apoptosis, cell cycle arrest, and inhibits epithelial-mesenchymal transition in pancreatic cancer cells. *Biochemical pharmacology*, 166, 274-291.
- Murugesan, K., Ilanchezian, T., & Vijayan, H. (2017). The Effect of Hydroethanolic Solvents Extracts of Piper betel and Plectranthus amboinicus on Food Poison Disease Causing Pathogenic Bacteria. *International Journal of Pharmaceutical Sciences Review and Research*, 45(1), 257-261.
- Nguyen, L. T. T., Nguyen, T. T., Nguyen, H. N., & Bui, Q. T. P. (2020). Simultaneous determination of active compounds in Piper betle Linn. leaf extract and effect of extracting solvents on bioactivity. *Engineering Reports*, 2(10), e12246.
- Pratiwi, N. P. R. K., & Muderawan, I. W. (2016). Analisis kandungan kimia ekstrak daun sirih hijau (Piper betle) dengan GC-MS. Prosiding Seminar Nasional MIPA,
- Rintu, D., Shinjini, M., Kaustab, M., Pramathadhip, P., Umesh, P., & Banerjee, E. (2015). Anti-oxidant and anti-inflammatory activities of different varieties of Piper leaf extracts (Piper betle L.). *Nutrition and Food Sciences*, 5, 415.
- Sanubol, A., Chaveerach, A., Tanee, T., & Sudmoon, R. (2017). Pre-clinical evaluation of extracts and essential oils from betel-like scented Piper species identified potential cancer treatment. *Afr J Tradit Complement Altern Med*, 14(1), 89-102.
- Sartini, S., Khaerawati, N., Kamril, R. A., & Febriani, N. J. P. (2020). The effects of fresh leaf-to-water ratio and heating time on the antifungal and antioxidant activities of betel leaf (Piper betle L.) extract. *10(1)*, 117-124.
- Selvaraj, G. K., Wilson, J. J., Kanagaraj, N., Subashini, E., & Thangavel, S. (2022). Enhanced antifungal activity of Piper betle against candidiasis infection causing Candida Albicans and In silico analysis with its virulent protein. *Biomedical and Biotechnology Research Journal*, 6(1), 73-80.
- Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021). Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*, 71(3), 209-249.
- Thị, C. H. C., Nguyễn, H. Đ., & Lê Hoàng, D. M. J. H. (2021). Influence of Piper betle L. extract on umbilical cord cells in vitro and potential treating cutaneous wound. *Heliyon* 7(3).
- Valle, D. L., Puzon, J. J. M., Cabrera, E. C., & Rivera, W. L. (2016). Thin Layer Chromatography-Bioautography and Gas Chromatography-Mass Spectrometry of Antimicrobial Leaf Extracts from Philippine Piper betle L. against Multidrug-Resistant Bacteria. *Evidence-Based Complementary and Alternative Medicine*, 2016, 4976791.
- Vanin, A. B., Orlando, T., Piazza, S. P., Puton, B., Cansian, R. L., Oliveira, D., & Paroul, N. (2014). Antimicrobial and antioxidant activities of clove essential oil and eugenyl acetate produced by enzymatic esterification. *Applied Biochemistry Biotechnology*, 174(4), 1286-1298.
- Wu, P.-F., Tseng, H.-C., Chyau, C.-C., Chen, J.-H., & Chou, F.-P. (2014). Piper betle leaf extracts

induced human hepatocellular carcinoma Hep3B cell death via MAPKs regulating the p73 pathway in vitro and in vivo. *Food function*, 5(12), 3320-3328.
Zari, A. T., Zari, T. A., & Hakeem, K. R. (2021). Anticancer Properties of Eugenol: A Review. *Molecules*, 26(23), 7407.

UNDER PEER REVIEW