

Original Research Article

Assessing the diversity of metazoan parasites of *Sarotherodonmelanotheron* (Cichlidae) from inland waters in Southern Benin (west Africa)

ABSTRACT

Aims: This paper presents a snapshot of the metazoan parasite fauna of *Sarotherodonmelanotheron* collected from two stations across lake Nokoué.

Study design: Fish samplings were done at the landing stage of Lac Nokoué (station 1) and Tota (station 2) from October 2016 to March 2017. Water physicochemical parameters such as pH, salinity, temperature, and conductivity were measured monthly.

Methodology: A total of 246 specimens of *Sarotherodonmelanotheron* comprising 149 males and 97 females were investigated. Size and weight were measured. Then, fish were dissected and parasites were checked in several organs. Collected parasites were identified using adequate keys of identification. The chi-square test was used to compare the prevalence of different parasite groups whereas Student T-test was applied to assess difference between parasite mean abundance within the two sampling stations. And the Pearson correlation coefficient was used to appraise possible influence of water parameters on parasite prevalence.

Results: The total prevalence was 86.58%. Eight parasite species were collected belonging to Myxosporidia, Monogenea, Digenea, Cestoda, Acanthocephala and Copepoda. Epidemiological parameters were calculated in order to appreciate the diversity of the parasitofauna according to sampling site, sex, and fish size. Females are significantly more infected than males ($\chi^2 = 10.093$; $p < .05$). Among the parasites, only monogenea prevalence (41.46%) is significantly different ($\chi^2 = 149.645$; $p < .05$). Correlation between physicochemical parameters and the prevalence of parasite groups showed that monogenea and cestoda were the most abundant in rainy season in opposite to the Digenea that are mainly recorded in the wet season.

Keywords: *Sarotherodonmelanotheron* – Parasite fauna – Prevalence – Monogenea – Abundance – Lake Nokoué.

1. INTRODUCTION

Freshwater aquatic organisms are posing health challenges. Fish in their several specificity, could serve as host of parasites that are harmful to human and animals. In Benin, some parasitological investigations were made on cichlids [18, 48, 60, 54, 38, 21, 22]; Silurids [48, 56, 55, 22, 21] and Channids [48, 52, 53]. Concerning the Cichlids, the main works have focused on the genus *Oreochromis* [39, 40, 2] within the continent. Indeed, it appears that *Sarotherodonmelanotheron* received little attention despite of its capacity of adaptation to environmental stress [59]. Only the works proposed by [17] and [54] presented Myxosporean parasites occurring in this species from Benin. As a fish living either in freshwater or in brackish water, we do have to focus on its parasite fauna according to the different habits. Hence, the present study was conducted to provide the first

snapshot on the identity and infestation level of parasites and to evaluate the relationship between the infection and the sex or size of *S. melanotheron*. The physicochemical parameters were also collected in order to appreciate the possible influence on the parasite prevalence.

2. MATERIAL AND METHODS

2.1. Study area and sampling

A survey of parasite fauna in a Cichlid fish was carried in the complex river Sô and Lake Nokoué in southern Benin from October 2016 to March 2017. Lake Nokoué is located between 6°26'04"N and 2°27'06"E. It joined Porto-Novo lagoon on the East by the channel of Totché and communicated southerly with Atlantic Ocean (figure 1).

Sampling A total of 246 individuals of *Sarotherodon melanotheron* was collected from two stations : the landing stage of Lake Nokoué (169 specimens) and Tota (77 specimens). Fish were collected twice a month in the morning between 7h and 10h and kept alive till the laboratory. The physicochemical parameters such as dissolved Oxygen, temperature, pH, salinity and conductivity were measured *in situ* using a multiparameter of type pH/Oxi340i/SET.

2.2. Parasitic investigation

Then, size and weight were measured for each fish specimen while the sex was determined after dissection. Parasites were checked with eye naked and by microscopic observation. The collected parasites were conserved in alcohol 70%. Some smears were instantaneously made with May Grunwald Giemsa, and Canada balsam. Parasites were identified with usual adequate keys of identification. For the parasite community structure, the common measures were evaluated belonging to Bush *et al.*, 1997.

2.3. Statistical analysis

The chi square test permitted to compare the prevalence according to the different parasite groups and the sampling stations whereas Student T-test was used to compare the parasite mean abundance within the stations. All tests were performed with the statistical program MINITAB 16. All results were considered to be significant at 95% ($P < .05$).




Figure 1 : Map of the complex river Sô and Lake Nokoué showing fish sampling sites ST1 (Lake Nokoué) and ST2 (Tota)

3. RESULTS AND DISCUSSION

3.1. Results

Out of 246 specimens of *Sarotherodon melanotheron* investigated, 213 are infected by at least one parasite species. The overall prevalence of infestation was 86.58%. Organs like skin, gills, lamellae, opercula, and intestine were infected (Table 1). The parasite subcommunity is composed of 8 species of metazoans distributed into six (06) parasite groups. There are Myxosporidia, Monogenea, Digena, Cestoda, Acanthocephala and Copepoda. Prevalence by parasite species in fish varies from 1.63 to 41.46% (Table 1).

Table 1. Prevalence of parasites species and their location in the host

Parasites groups	Parasites species	Location	Prevalence (%)
Myxosporidia	<i>Myxobolus fotoi</i>	Gills	34.96
	<i>M. heterospora</i>	Kidney	
Monogenea	<i>Cichlidogyrus tilapiae</i>	Gills	41.46
Digena	<i>Clinostomum</i> sp.	Gills	1.63
		Opercula	
Cestoda (Gryporhynchidae)	Undetermined	Intestine	23.98

Acanthocephala	<i>Acanthogyrussp.</i>		22.36
Copepoda	<i>Ergasilus sp.</i>	Gills	4.90
	<i>Lernaeasp.</i>	Gills	6.89

Table 2 shows the different parasite prevalence obtained according to the fish sex, the study stations. For all the fish examined, parasites were found in 122 /149 males, i.e. a prevalence of 81.88%, compared with 93.81% for females (91 /97 individuals) were parasitized. Females were significantly more parasitized than males ($\chi^2 = 7.205$; $p < .05$). Taking stations into account, a prevalence of 78% was recorded for males (78/100) from station1, compared with 95.65% (66/69) for females meaning that females were significantly more parasitized than males ($\chi^2 = 10.093$; $p < .05$). At station 2, males appear to be more parasitized (89.79% prevalence) than females (89.28%); but this difference remained non-significant ($\chi^2 = 0.005$; $p > .05$). The calculated sex ratio indicates that the populations at the two study stations do not appear to be in equilibrium, with males almost doubling the number of available, breeding females.

Table 2. Overall and parasite wise prevalence by sex of *S. melanotheron*

	Pool of samples			Station 1			Station 2		
	Female	Male	Total	Female	Male	Total	Female	Male	Total
Examined	97	149	246	69	100	169	28	49	77
Infected	91	122	213	66	78	144	25	44	69
Prevalence (%)	93.81	81.88	86.58	95.65	78	85.2	89.28	89.79	89.61
Sex ratio(F/M)	0.65			0.69			0.57		
Chi square (p)	7.205 (0.007)			10.093 (0.001)			0.005 (0.944)		

Figure 2 shows the general prevalence of infestation of *S. melanotheron* individuals according to size, depending on the station. In general, large individuals are more parasitized (91.67%) than medium-sized individuals (89.04%), which in turn are more parasitized than small individuals (81.81%). Nevertheless, the differences between these prevalences are not significant ($\chi^2 = 2.747$; $p > .05$). At station1, medium-sized individuals are the most parasitized (88.68%), while at station2, large-sized individuals are the most parasitized (100%). For both stations, small-sized individuals show the lowest prevalence. There was no significant difference between the prevalence of the different size classes of *S. melanotheron* at either station1 ($\chi^2 = 3.201$; $p > .05$) or station 2 ($\chi^2 = 0.083$; $p > .05$). Infestation is not distributed according to the size class of the *S. melanotheron* individuals examined.




Figure 2: Prevalence of *S. melanotheron* parasitic infestation according to size

S. melanotheron specimens collected at station 1 (ST1) are more parasitized by Monogeneans, Cestodes, Acanthocephalans and Copepods than specimens collected at station 2 (ST2). This trend is significant only for Cestodes ($\chi^2 = 4.337$; $p < .05$). *S. melanotheron* specimens collected at station 2 (ST2) were more infected with Myxosporidia and Digena (fig 3). This trend was significant only for Myxosporidia ($\chi^2 = 65.561$; $p < .001$).




Figure 3. Prevalence of major parasitic groups according to the stations. ST1 in black band and ST2 in gray band.

Depending on the station, fish sex and size, the prevalence of the parasite groups indicate that the *S. melanotheron* specimens collected at station 2 are significantly more infected with

Myxosporidiathan those at station 1. On the other hand, Cestodes and Copepods infested *S. melanotheron* specimens from Station 2 significantly more than those from Station 1, and infestation by Cestodes and Copepods was evenly distributed according to sex. Depending on the size, Copepods infested more the large fish while Cestoda were more encountered in small and medium-sized fish. There was no cestoda infection in large fish.

Table 3. Prevalence (%) of major parasite groups of *S. melanotheron* by station, sex and size

	Myxosporidia	Monogenea	Digena	Cestoda	Acanthocephala	Copepods
Stations						
ST1	18.34	42.01	1.18	27.81	23.08	15.98
ST2	71.43	38.96	2.6	15.58	20.78	2.67
Khi -2 (P)	65.561 (.00)	0.289 (.59)	0.661 (.41)	4.337 (.03)	0.161 (0.68)	9.105 (.003)
Fish sex						
Male	43.27	40.26	1.34	23.49	18.12	12.08
Female	42.27	43.3	2.06	24.74	28.86	11.34
Khi -2 (P)	3.762 (.05)	0.222 (.63)	0.19 (.66)	0.051 (.82)	3.908 (.04)	0.031 (.86)
Fish size						
[6-10[39.77	38.64	1.14	23.86	13.64	4.54
[10-14[38.88	42.46	1.37	26.03	26.71	15.75
[14-18[25	50	8.33	0	33.33	20
Khi -2 (P)	1.699 (.42)	0.711 (.70)	3.567 (.09)	4.121 (.12)	6.284 (.04)	6.921 (.03)

Among the main parasites groups, Monogenea were the most abundant (1.6), while Digenes were the relative least abundant (0.05) (Table 4). In terms of stations, the abundance of the various parasite groups was higher at station 1 than at station 2. The abundance of the different parasite groups was also higher in female than in male. The abundance of Cestodes, Acanthocephala and Copepod is higher in medium-sized individuals, while large individuals have a higher rate of Monogene and Digene. It should be noted, however, that the different trends observed with regard to station, sex and size are only significant for Monogenes. Also, the mean intensity of Digenes is higher in the largest size class (14-18 cm), whereas the medium-sized individuals (10-14 cm) showed a high level of mean intensity compared to the two others size classes considering all the parasite groups.

Table 4. Overall and wise parasite Abundance (Ab) of the parasite groups according to sampling station and fish sex and size

Parameters	Monogenea	Digena	Cestoda	Acanthocephala	Copepods
Overall					
NP	394	12	299	102	48
Ab	1.6	0.05	1.22	0.41	0.20
Station					
Station 1	1.63	0.06	1.63	0.45	0.27
Station 2	1.53	0.03	0.31	0.34	0.03
Student T-test of (P)	31.60 (.02)	3.00 (.205)	1.47 (.380)	7.18 (.088)	1.25 (.430)
Fish Sex					
Male	1.54	0.05	1.10	0.36	0.17
Female	1.70	0.05	1.39	0.51	0.23
T-test of Student (P)	20.25 (.031)	-	8.59 (.074)	5.80 (.109)	6.67 (.095)
Fish Size					
[6-10[1.27	0.01	0.91	0.22	0.07
[10-14[1.78	0.05	1.50	0.53	0.27
[14-18[1.83	0.33	0.00	0.50	0.17
T-test of Student (P)	9.09 (.012)	1.29 (.326)	1.84 (.207)	4.22 (.052)	2.94 (.099)

NP: Number of parasites collected for each group; Ab = Mean abundance

Overall, Cestoda presented the highest mean intensity and the lowest MI was recorded with Copepods (Table 5). The average intensity of the parasite groups is higher at station 1 than at station 2. This trend is significant for Monogenea ($p < 0.001$) only. Mean intensity of Monogeneans, Cestoda is higher in females than in males with a significant difference, while the contrary is observed with Digenea and Acanthocephala, even it revealed a significant difference with Acanthocephala.

Table 5. Mean intensity (MI) of the parasite groups according to sampling station and fish sex and size

	Monogenea	Digena	Cestoda	Acanthocephala	Copepoda
Overall					
NP	394	12	299	102	48
MI	3.94	3	4.98	1.85	1.66
Station					
Station 1	3.94	5.00	5.73	1.95	1.70
Station 2	3.93	1.00	2.00	1.63	1.00
T-test of Student (P)	787.00 (.001)	1.50 (.374)	2.07 (.286)	11.19 (.057)	3.86 (.161)
Fish sex					
Male	3.88	3.50	4.56	1.96	1.44
Female	4.02	2.50	5.63	1.75	2.00
T-test of Student (P)	56.43 (.011)	6.00 (.105)	9.52 (.067)	17.67 (.036)	6.14 (.103)
Size					
[6-10[3.29	1.00	3.64	1.58	1.50

[10-14[4.33	3.50	5.76	1.97	1.74
[14-18[3.67	4.00	0.00	1.50	1.00
T-test of Student (P)	12.39 (.006)	3.05 (.093)	1.86 (.204)	11.59 (.007)	6.48 (.023)

NP: Number of parasites collected for each group; MI = Mean Intensity

The correlation between physico-chemical parameters and the prevalence of parasite groups in *S. melanotheron* (Table 6) shows that among the recorded parameters, only salinity has an influence on parasite distribution. Of the six (06) parasite groups collected in *S. melanotheron*. Myxosporidia and Cestoda are strongly negatively and significantly correlated with salinity. On the other hand, Digena are positively and significantly correlated with salinity.

Table 6. Correlation between physico-chemical parameters and the prevalence of the parasite groups of *S. melanotheron*

Parameters		pH	Temperature	Salinity	Conductivity	Dissolved Oxygen
Myxosporidia	r	-0.657	-0.970	-0.999	-0.963	-0.570
	P	0.544	0.157	0.024	0.174	0.351
Monogena	r	0.859	0.846	0.938	0.831	0.711
	P	0.342	0.358	0.226	0.376	0.657
Digena	r	0.663	0.968	0.999	0.961	-0.216
	P	0.539	0.162	0.030	0.179	0.074
Cestoda	r	-0.582	-0.989	-0.998	-0.984	0.929
	P	0.604	0.096	0.036	0.113	0.060
Acanthocephala	r	0.987	0.592	0.746	0.570	0.657
	P	0.104	0.596	0.464	0.614	0.238
Copepoda	r	0.995	0.537	0.699	0.514	-0.186
	P	0.061	0.639	0.507	0.657	0.992

Values in bold are different from 0 at a significant level equal to 0.05; r = coefficient of correlation, p = probability.

Variation in physico-chemical parameters on the prevalence *S. melanotheron*.

Among all of the physico-chemical parameters, only salinity showed significant differences ($P < 0.05$) between the sampling stations. Nevertheless, the parameters did fluctuate depending on the station (figure 4). Thus, they exist great amplitude between minimum (6.34) and maximum (7.95) of pH in station 2 than in station 1 where pH ranged from 7.33 to 7.63. The average temperature at station 2 is of 29.13°C however it is of 28.68°C at station 1. The maximum (31.3°C) and minimum (27.47°C) values of temperature were recorded at station 2 and station 1 respectively.

Salinity values at station 1 (6.88 g/L) are higher than at station 2 (2.12 g/L). The amplitude of salinity variation was trending as the same. The minimum salinity value (1.26 g/L) is observed at station 2 whereas the maximum (13.69 g/L) was recorded at station 1. As for conductivity, the mean value recorded at station 2 (93.07 $\mu\text{S/cm}$) is significantly higher ($p < 0.008$) than that observed at station 1 (13.40 $\mu\text{S/cm}$). This trend is confirmed by the amplitude of variation. where the lowest conductivity values are grouped together at station1 (5.93-26.03 $\mu\text{S/cm}$) and the highest at station 2 (76.2-117 $\mu\text{S/cm}$). Whereas, Dissolved oxygen in the water varies widely at each

station. The average dissolved oxygen value observed at station 1 (3.04) is slightly higher than at station 2 (2.86) without being significantly different ($= 0.068$).




Figure 4: Variation in water physico-chemical parameters in the two sampling station of *S. melanotheron*. ST = station

3.2. Discussion

The present study gives us the opportunity to encounter a species of myxosporidia never before recorded on fish in Benin. Regarding to the myxosporidian parasite checklist, *Myxobolusfotoi* is recorded for the first time in this area. Described in Cameroon by [15] in the fatty tissues of gill arch, kidney and spleen of *Oreochromis niloticus*, it was already reported by [27] and never in *Sarotherodonmelanotheron*, nor in other Cichlids species. Whereas *Myxobolusheterospora* in the viscera of *Coptodon esculenta*, *C. variabilis*, *O. niloticus* and in the spleen of *Haplochromis* sp. [27] and either reported by [26; 27] in the kidney of *Oreochromis niloticus*.

Myxosporean kidney infestation (34.96%) obtained in this study is much lower than that obtained by [27]. Authors justify this high prevalence by a decrease in the defense capacity of the hematopoietic organs of the fish. They agree with [15] and [1] who suggested that spores could be transported and accumulate in the host's macrophage cells [32]. Indeed, large groups of spores accumulated in the melanomacrophage cells of the kidney, resulting in a high prevalence of infestation in fish kidneys at that precise moment. Conversely, taking into account the role of melanomacrophages in host defense reactions, these cells are capable in some cases of attacking small cysts and large mature spores where the parasites are isolated and progressively destroyed [13] melanomacrophage centers of the hematopoietic organs. Due to the defense mechanism of melanomacrophages, high infestation by myxospores accumulation in the kidneys could be reduced justifying the prevalence of 34.96% observed in this study. Several studies have been carried out on monogenea of inland fish in Africa [9, 25, 12, 35]. In Benin, *Cichlidogyrustilapiae* was described by [12] in *Chromidotilapia guentheri* and by [4] in farmed *Coptodon zillii*. In 2014, [28] attested the presence of *C. tilapiae* in the gill filaments of *Oreochromis niloticus* from fish farms in Brazil. In contrast, [9] revealed the presence of *Cichlidogyrus acerbus* and *C. halli* in *S. melanotheron*. Despite multiple studies on monogenea, only few results displayed the presence of *Cichlidogyrustilapiae*. Out of the six (06) helminth species collected, Monogenea are highly prevalent. This high prevalence of Monogenea may be due to their short developmental cycle. Indeed, monogenea are easy contagiousness, representing a major threat for fish [14]. The mechanical action exerted by the opisthaptor could induce extensive damage and exposed the host to secondary infections by bacteria, fungi and viruses [47] whatever Monogenea are considered one of the important parasites sensitive to any change in water quality [62].

The present work reported Gryporhynchidaemetacercariae. Information based on taxonomy, development cycle, ecology and pathology are very scarce or non-existent on this helminth [29]. Identifying the Gryporhynchidae down to genus and species level according to morphological characteristics of the larvae is very difficult as the adult parasites is located in piscivorous birds [49]. Gryporhynchidae are parasitic on several groups of fish [29], but studies on the damage caused by Gryporhynchidae on their hosts are very rare [49]. Fish infested with metacestodes showed ungrowth compared to the uninfested or low infested fish. Several species of the *Acanthogyrus (Acanthocephalus)* genus have been described in African Cichlids particularly in Tanzania [7], Congo [43] and Madagascar [19]. Nkwengulila and Mwita [37] investigated the organic preferences of Acanthocephalans. For these authors, differences in the physical environment of the gut and food supply availability and quantity are the factors that limit the distribution of parasites in different sections of the alimentary tract. It is reported that *A. tilapiae* is better adapted to some cichlid hosts than others [5]. For we have no especial studies focused on Acanthocephalans and copepoda, but several authors have reported serious economic losses caused these parasites [45, 10, 51]. According to some authors, gill infestations by Ergasilidae cause depletion of fish populations and weight loss.

According to [33], life traits, size, habitat and feeding behavior could directly or not influence the parasite richness. Whereas prevalence variation could depend on the genetics, proximity living environment, host age, energy expenditure to potential hosts or the presence of other parasites [12]. According to [8], euryhaline fish cross various endemiotopes and could tolerated great variations in salinity. Indeed. the euryhaline parasitofauna would be affected by their displacement. On the other hand, fish that migrate without changing environmental conditions. would be less affected as physicochemical parameters are of very low amplitude as reported by [52]

Changes in diet and habitat requirements at different developmental stages of a species can strongly affect the composition, abundance and richness of the parasitic community. For [11], tropical fish whose diets are based on macroinvertebrates have helminth-rich digestive tracts. [30] attributes this richness of gastrointestinal helminths to physio-reproductive and feeding behavior factors.

Considering the pollution criteria, [38] admitted that the distribution of Monogenea and Digenea was limited to unpolluted and moderately polluted lakes. [24] revealed that fish parasite community is known to respond directly to the abiotic environment of the host. The correlation between physico-chemical parameters and the prevalence of the main groups of parasites of *S. melanotheron* in the present work showed that only salinity does influence parasite distribution.

According to the present study, fish with great size were more parasitized than smaller ones. This finding is agreed with [58] and [34] who noted that parasite richness is positively correlated with host size. Indeed. [61] and [34] assured that long sized fish carry more parasite species or a great abundance of parasites. However, we can concluded with [42] that the significant difference between prevalence and intensity does not necessarily indicate that larger hosts have a higher parasite load than the smaller one. Like [44], the prevalence in *S. melanotheron* according to sex reveal that females are more parasitized than males. Sex-related differences in parasite infestation rates may be attributed to the host immune response. due to differences in the endocrine gland activities of host fish. [50] [23] and [17] suggested that failure in resistance to parasitic infection in female fish may be due to investment in gonadal development.

4. CONCLUSION

The Cichlid *Sarotherodon melanotheron* hosted a great diversity of parasites from many organs. Parasite prevalence was elevated and dominated by monogeneans. Salinity remained a chemical determinant that either control the live of this fish or influence its prevalence in parasites.

ETHICAL APPROVAL

Non applicable.

REFERENCES

1. Abakar-Ousma AF, Ngassam P, Bouix G. Myxosporidian (Myxozoa) parasites of fresh water fish teleostean from Chad: New or few known species. *Annales Université de N'Djamena*. 2006;1:111-121.

2. Akoll P, Konecny R, Mwanja WW, Nattabi JK, Agoe C, Schiemer F. Parasite fauna of farmed Nile tilapia (*Oreochromis niloticus*) and African catfish (*Clarias gariepinus*) in Uganda. *Parasitol. Res.* 2012; 110:315-323.
3. Albaret JJ. Fish populations in Casamance (Senegal) during periods of drought. *Journal of Tropical Hydrobiology.* 1987; 20:291-310.
4. Al-Samman A, Molnar K, Székely C. Infection of cultured and freshwater fishes with monogeneans in Syria. *Bull. Eur. Ass. Fish Pathol.* 2006; 26(4):170.
5. Amin OM, Oosterhout CV, Blais J, Robinson RL. On the Ecology and Host relationships of *Acanthogyrus* (*Acanthosentis*) *tilapiae* (*Acanthocephala*: *Quadrigyridae*) from Cichlids in Lake Malawi. *Comp. Parasitol.* 2008; 75(2):278-282.
6. Amoussou TO, Toguyeni A, Toko II, Chikou A and Youssao IAK. Biological and zootechnical characteristics of the African tilapia *Oreochromis niloticus* (Linnaeus, 1758) and *Sarotherodon melanocheilus* Rüppell, 1852. A review. *Int. J. Biol. Chem. Sci.* 2016; 10(4):1869-1887.
7. Baylis HA. A new acanthocephalan from an East African Freshwater Fish. *DNA. Mag. nat. Hist.* 1947; 14:861-868.
8. Benhassine OK. Parasitic copepods of mugilid fish in the western Mediterranean (French and Tunisian coast). Morphology, bio-ecology, evolutionary cycle. State Doctorate Thesis, U.S.T.L. Montpellier. 1983; 452.
9. Blahoua KG, N'Douba V, N'Guessan J. Seasonal variations in the epidemiological indices of three Monogenean parasites of *Sarotherodon melanocheilus* (Pisces: Cichlidae) in Lake Ayamé I (Côte-d'Ivoire). *Science and Nature.* 2009;(6): 39-47.
10. Boualleg C, Kaouachi N, Bensouilah M, Ternengo S. The Copepod parasite of the gills of four teleost fishes caught from the gulf of Annaba (Algeria). *Advanc. J. Microb. Res.* 2010: 189. https://www.researchgate.net/publication/281954329_The_Copepod_parasite_of_the_gills_of_four_teleost_fishes_caught_from_the_gulf_of_Annaba_Algeria
11. Choudhury A, Dick T. Richness and diversity of helminth communities in tropical freshwater fishes: Empirical evidence. *J. Biogeogr.* 2000; 27:935-956.
12. Combes C. Sustainable interactions Ecology and evolution of parasitism. Masson. 1995; 52:4.
13. Dossou C. Monogenean parasites of freshwater fish in Benin (West Africa). Montpellier - France: University of Sciences and Technology of Languedoc. 1985: 116.
14. Dyková I. The role of melanomacrophage centers in the tissue reaction to myxosporean infections of fish cyprinid species, myxosporeans, *Sphaerosporarenicola*, *Mitrasporacyprini*,

Thelohanellus, *Hoferellus*cyprini, *Myxobolus*. Bulletin of the European Association of Fish Pathologists. 1984; 4:65.

15. El Madhi Y, Belghi D. Distribution of twomonogenes in host individuals of *Trachinotusovatus* (L 1758) from the coast of Mehdiya Biology and Health. 2006; 6:2.

16. Fomena A, Marques A, Boiux G. Myxosporodies (Myxozoa) of *Oreochromisniloticuslinnaeus*, 1757 (TeleostCichlidae) in fish-rearing ponds in Melen, Yaoundé; CameroonParasitol Res.1993; 78:201-208.

17. Gbankoto A, Anago E, Houndjo PA, Adjahouinou DC, Gbaguidi F. Effect of Aqueous and EthanolicExtracts of *Caesalpinia bonduc* Root on SexualBehavior of Male Wistar Rats. Int J of MultidisciplinaryCurrentresearch..<http://ijmcr.com/wp-content/uploads/2015/11/Paper61137-11412.pdf>. 2015; 3:1137-41.

18. Gbankoto A, Pampoulie C, Marques A, Sakiti GN. Occurrence of myxosporean parasites in the gills of two tilapia speciesfrom Lake Nokoué (Benin, West Africa): effect of host size and sex, and seasonal patterns of infection. Say AquatOrg. 2001; 44:217-222.

19. Gbankoto A, Pampoulie C, Marques A, Sakiti GN and Dramane KL. Infection patterns of *Myxobolusheterospora* in two tilapia species (Teleostei: Cichlidae) and itspotential effects. Say AquatOrg. 2003; 55:125-131.

20. Golvan YJ. Acanthocephala of Fishes. Explor. Hydrobiol. Lakes Kivu, Edouard and Albert 1952-1954. Brussels. 1957; 3:5-64.

21. HouénouSèdogbo DM, Zannou TB, Siko EJE, Tossavi ND, Togla AI, Fiogbé ED, Ibikounlé M. Fauna of metazoan parasites of *Clarias gariepinus* (Clariidae) and *Oreochromisniloticus* (Cichlidae), twowedosfishes of the upper delta of the Ouémé River in southern Benin. International Journal of Biological and Chemical Sciences. 2019a; 13(2):983-997.

22. Houénousèdogbo M, Zannou BT, Siko JE, Tossavi ND, Togla I, Fiogbé ED and Ibikounle M. Parasite faunaassociatedwith the Nile Tilapia *Oreochromisniloticus* L. (Cichlidae) fromfishfarms in Southern Benin (West Africa). International Journal of Multidisciplinary and CurrentResearch. 2019b; 7:624-630.

23. Ibrahim MM, Soliman MF. Prevalence and site preferences of heterophyidmetacercariae in *Tilapia zillii*fromIsmaliafresh water canal, Egypt. Parasite. 2011;17:233-239.

24. Karvonen A, Kristjánsson BK, Skúlason S, Lanki M, Rellstab C, Jokela J. Water temperature, not fishmorph, determines parasite infections of sympatricIcelandicthreespinesticklebacks (*Gasterosteusaculeatus*) <https://doi.org/10.1002/ece3.568>. 2013.

25. Lambert A. The Monogeneans*Monopisthocotylea* parasites of freshwaterfishfromMediterranean France. Bulletin of the National Museum of Natural History. 1997; 3(429):39.

26. Landsberg JH. Myxosporean infections in cultured Tilapias in Israël. J Protozool. 1985; 32:194-201.
27. Lekeufackfoleack GB, Malakengne C, Feudjiodongmo B, Fomena A. Prevalence and Mean Intensity of Myxobolus spp. parasitizing *Oreochromis niloticus* in Cameroon. Int. J. Biol., 2019; 11(2):916-9671.
28. Martins ML, Santana AR, Tomas JG, Tancredo KR, Tavares GEL, Bampi D, Mattos SA. Microhabitat preference and seasonality of gill Monogeneans in Nile Tilapia reared in Southern Brazil. Neotropical Helminthology. 2014: 47-58.
29. Mashego SN, Saayman JE, Mokgalong NM. Parasites of the fish population with notes on the helminth parasites of the water birds of Middle Letaba Dam. South Africa University of the North. Microbiol. Res 4.1991; (9): 801-807.
30. Mohamed MI. Variation in parasite infracommunities of *Tilapia zillii* in relation to some biotic and abiotic factors. Int J Zool Res. 2012; 8:59-70.
31. Moller H, Anders K. Diseases and Parasites of Marine Fishes. Kiel: Moller. 1986:365.
32. Molnár K, Kovács Gayer E. The pathogenicity and development within the host fish of *Myxobolus cyprinidoflein*. 1898. Parasitology. 1985; 90(3):549 -555.
33. Morand S, Poulin K, Hayward C. Aggregation and species coexistence of ectoparasites of marine fishes. Int J Parasitol. 1999; 29:663-672.
34. Munoz G, Zamora L. Ontogenetic variation in parasite infracommunities of the Clingfish *Scyasessanguineus* (Pisces: Gobiesocidae). J. Parasitol. 2011; 97:14-19.
35. N'Douba V, Lambert A. Un nouveau *Macrogyrodactylus* (Monogenea, Gyrodactylifae) parasite de *Heterobranchus longifilis* Valenciennes, 1840 (Téléostéen, siluriformes) en Côte d'Ivoire. Zoosystema. 1999;21:7-11.
36. N'Douba V, Lambert A. Deux monogènes nouveaux parasites branchiaux de *Clarias ebriensis pellegrin*, (1920) (Siluriformes, Clariidae) en Côte d'Ivoire Zoosystema. 2001; 23(3):411-416.
37. Nkwengulila G, Mwita C. Spatial distribution of parasites along the gut of the catfish *Clarias gariepinus* (Burchell, 1822) (Clariidae) from the Mwanza gulf, lake Victoria. Tanzan J Sci. 2004;30(1):63-70.
38. Houénou Sédogbo M, Aboki-Onzo A, Imorou-Toko I, Siko EJE, Zannou B, Tossavi ND, Fiogbé E, Ibikounlé M. Parasite fauna associated with the tilapia *Oreochromis niloticus* L. (Cichlidae) from fish farms in Southern Benin (West Africa). Int. J. Multidiscipl. Curr. Res. 2019;624-630
39. Opara KN, Okon AO. Studies on the parasites of cultured *Oreochromis niloticus* (Cichlidae) in a rainforest fish pond in southeastern Nigeria. J Aquat Sci. 2002; 17(1): 17-20.

40. Pariselle A, BilongBilong CF, Euzet L. Four new species of Cichlidogyrus Paperna, 1960 (Monogenea, Ancyrocephalidae), all gill parasites from African mouthbreeder tilapias of the genera Sarotherodon and Oreochromis (Pisces, Cichlidae), with a redescription of *C. thurstonae* Ergens, 1981. *Systematic parasitology*. 2003; 56:201-210.
41. Pèlèbè ROE, ImorouToko I, Verheyen E, Van Steenberge M. Molecular Identification of an Invasive Sarotherodon species from the Atchakpa Freshwater Reservoir (Ouémé River Basin, Benin) and Comparison within *S. melanotheron* Using COI. Markers. *Diversity*. 2021; 13(7), 297; <https://doi.org/10.3390/d13070297>.
42. Poulin R. Sexual inequalities in helminth infections: A cost of being male. *Am. Nat.* 1996; 147:287-295.
43. Prudhoe S. Trematoda, Cestoda et Acanthocephala. *Result. Scient. Explor. Hydrobiol. Lake Tanganyika*. 1951;3:1-10.
44. Ramadan HH. Effect of host species, sex, length, diet and different seasons on the parasitic infection of Tilapia fish in Lake Manzalah. *J. King Abdulaziz Univ. Marine Sci.* 1991;2:81-91.
45. Ramdane Z and Trilles JP. Parasitic Copepods (Crustacea: Copepoda) from Algerian marine fishes. *Zootaxa*. 2007;1574:49-68.
46. Ravinchandran S, Rameshkumar G, Balasubramanian T. Infestation of isopod parasites in commercial marine fishes. *Journal of Parasitic Diseases*. 2010;34(2):97-98.
47. Reed PA, Francis-Flyord R and Klinger RE. Monogenean parasites of fish. *FAO28/FAO33*. 2012.
48. Sakiti GN. Doctorat d'Etat des Sciences de Biologies Animale, Option Parasitologie (Université Nationale du Bénin. 1997.
49. Scholz T, Bray RA, Kuchta R and Repova R. Larvae of Gryporhynchid cestodes (Cyclophyllidea) from fish: a review. *Folia Parasitol*. 2004;51:131-152.
50. Simokova A, Jarkovsky J, Koubkova B, Barus V and Prokes M. Association between fish reproduction cycle and the dynamics of metazoan parasite infection. *Parasitol*. 2005;95:985-989.
51. Ternengo S. Caractérisation des communautés des parasites de poissons de la réserve naturelle des bouches de Bonifacio, Thèse de Doctorat, Fac. Sci. Techn., Univ. Corse. 2004: 209p.
52. Togla AI, Zannou TB, Tossavi ND, Bouko B, HouénouSèdogbo DM, Ibikounle M, Diversité des métazoaires parasites de *Parachanna obscura* (Gunther 1861) de la rivière Sô au Sud du Bénin. 2020;16(6):14-24.
53. Togla AI, Bouko B, Zannou TB, Tossavi ND, HouenouSedogbo DM, Ibikounle M. Incidence des Nématodes parasites chez *Parachanna obscura* (Gunther, 1861) dans la rivière Sô au sud du Bénin. 2021; (9):2321-3

[PMC free article] [PubMed] [Cross Ref]

54. Tossavi ND, Gbankoto A, Siko EJE, Christophe C, Florence V and Tareck R. Pattern and potential pathophysiological effects of myxosporean infections in the gills of Tilapia species (Teleostei:Cichlidae) from Benin. *Int J Multidisciplinary Curt Res. Rev.* 2016;4:1229–1238.

[PMC free article] [PubMed]

55. Tossavi ND, Bankoto A, Yessoufou A, Escandes ML, Dimitri G, Ribou AC, Moutairou K and Sakiti GN. Histopathological and ultrastructural studies of a Henneguya species (Myxozoa:Myxosporea) infesting the intestine of Clarias gariepinus from Benin (West Africa). *Parasitology Research. Rev.* 2015;114:861–8

[PubMed]

56. Tossavi ND, Gbankoto A, Adite A, Ibikounle M, Grunau C, Sakiti GN. Metazoan parasite communities of catfishes (Teleostei:Siluridae) in Benin (West Africa) DOI 10.1007/s00436-014-4063-x. 2014:1-1

57. Villena I. Parasites and food, surveillance and means of control in France. 2023; (550): 3-6

[PMC free article] [PubMed] [Cross Ref]

58. Violante-Gonzalez J, Mendoza-Franco EF, Rojas-Herrera A And Gil-Guerrero S. Factors determining parasite community richness and species composition in black snook *Centropomus nigrescens* (Centropomidae). 2010; 107:59-6

[PubMed]

59. Yoboue AN, Adepo-Gourene AB and Agnese JF. Diversity and Genetic Structure of *Sarotherodon melanotheron* (Pisces:Cichlidae) Revealed by Microsatellites. *European Scientific Journal.* 2014;10(33):1857-7881.

60. Zannou TB. A comparative study of the diversity of Metazoans parasitic to *O. niloticus* in controlled and natural environments in southern Benin. *Memo Master Applied Hydrobiology/ FAST/ UAC.* 2014:4

[PMC free article] [PubMed]

61. Zapata AB, Ten T, Cejalvo C, Gutierrez de Frias C and Corts A. Ontogeny of the immune system of fish. *Fish Shellfish Immunol.* 2006;20:126–136.

[PMC free article] [PubMed]

62. Zargar UR, Chishti MZ, Yousuf AR, Fayaz A. Infection level of monogenean gill parasite, *Diplozoon kashmirensis* (Monogenea, Polyopisthocotylea) in the Crucian Carp, *Carassius carassius* from lake ecosystems of an altered water quality an impact on the Diplozoon infection. *Vet. Parasitol.* 2012; 189(2-4):218-2