

## Original Research Article

# ANTIOXIDANT ACTIVITY, PHENOLIC AND FLAVONOID CONTENT OF *TECOMA STANS*(L.) KUNTH AND *ZINGIBER OFFICINALES* ROSCOE

**Comment [J1]:** Contents, let your topic align with your aim please

**Comment [J2]:** Italicize and space

### ABSTRACT

**Aim:** The purpose of this study was to determine the total phenolic and flavonoid content and antioxidant activity in the three medicinal plants *Tecoma stans* and *Zingiber officinales*.

**Methods:** Aqueous, methanolic, and ethanolic solvent extracts were prepared; the DPPH assay was used to measure the extracts' antioxidant activity. Standard techniques were used to determine the total phenolic and flavonoid contents.

**Results:** Methanolic solvent contained the highest total phenolic (16.67±0.04, 11.56±0.09 mg GAE/g dw) and flavonoid contents (14.21±0.10, 7.96±0.06 mg QUE/g,) as well as antioxidant activity (68.91%, 53.09% at 300 µg/ml) in *Z. officinales* and *T. stans* extracts respectively.

**Conclusion:** Our study's screened extracts of can be exploited as possible antioxidant agents as well as resources for the production of novel drugs.

**Comment [J3]:** Let your aim follow the same trend as ur title

**Comment [J4]:** Space

**Comment [J5]:** Space

**Comment [J6]:** Space

**Comment [J7]:** Space

**Comment [J8]:** Recast

**Keywords:** Antioxidant activity, DPPH assay, Total phenolic content, Total flavonoid content, *Tecoma stans*, *Zingiber officinales*.

### 1. INTRODUCTION

Plants and natural products constitute the foundation of both contemporary and traditional medicines [1]. Medicinal plants are particularly significant in facilitating the development of alternative remedies that do not have the negative side effects of synthetic pharmaceuticals [2,3]. Herbs are used in approximately 25% of prescription medications globally [4,5].

Ginger (*Zingiber officinale* Roscoe) is a member of the Zingiberaceae family and contains a variety of active compounds including terpenes (sesquiterpene hydrocarbons), alkaloids, and polyphenols [6].

Phenolic compounds are linked to a wide range of biological activities, one of which is their antioxidant ability [7], which may aid in protecting cells from oxidative damage induced by free radicals [8, 9]. Studies have reported that ginger has antioxidant properties [10,11]. From ancient times, ginger has been used as a medicine to cure a variety of conditions, including fever, indigestion, hypertension, sprains, sore throats, cramps, and rheumatism [12]. Ginger has a variety of biological properties, including antibacterial activity, antifungal activity, anti-inflammatory activity, and anticoagulant impact [13,14].

A decorative shrub from the *Bignoniaceae* family, is *Tecoma stans* (L.) Kunth also known as yellow elder, has been used in traditional medicine. Aerial parts of the plant are used for the treatment of diabetes, to control fungal infections, and as a tonic, diuretic and dewormer [15].

Teas made from flowers and leaves are commonly used to treat diabetes and digestive problems, and the decoction of flowers has been used to relieve stomach pain [16,17]. The presence of several bioactive compounds in species of *Tecoma* genus have been reported, which include saponins, flavonoids, alkaloids, phenols, steroids, anthraquinones, tannins, terpenes, hydrocarbons, volatile oils, glycosylated flavonoids and phenolic acids [17,18]. These compounds have been shown antioxidant, antibacterial and antifungal activities [19].

These plants have key bio-natural components that are quite effective in giving critical medical advantages. The present study aims to determine the total phenolic and flavonoid content and antioxidant activity in the medicinal plants *Tecoma stans* and *Zingiber officinale*.

## 2. MATERIAL AND METHODS

The selected medicinal plants / parts were collected as follows: leaves of *Tacoma stans* were obtained from the plants growing at Banasthali Vidyapith campus, and rhizome of *Zingiber officinale* were procured from the market and kept in a refrigerator at 4°C.

### 2.2 Extraction of Leaf Samples

20g of leaf sample was dissolved in 100 ml of solvent (ethanol, methanol, and water) for 24–42 hours at room temperature. Then, the extracts were filtered through sterile muslin cloth and Whatman filter paper. The filtrates from each solvent were evaporated in a desiccator. The dried extracts were packaged in airtight containers, labelled, and kept in a refrigerator (2–4°C) for further use.

### 2.3 Total phenolic and total flavonoid content

The total phenolic [20,21] and flavonoid [22,23] content of various fractions of extracts (methanol, ethanol, and aqueous) were assessed using standard procedures.

### 2.4 Determination of Antioxidant Activity:

To determine DPPH (1, 1-dihydroxy-2-picrylhydrazyl, Sigma Aldrich) free radical scavenging activity, 1.0ml of DPPH solution (0.135mM DPPH in methanol) was added to (100–300µg/ml) plant extract [19]. The reaction mixture was kept in the dark for 30 minutes and its absorbance at 517nm was measured with a spectrophotometer. Ascorbic acid was utilized as a reference.

### 2.6 Statistical Analysis:

The experiments were repeated three times, and the findings were provided as Mean ± S.E. All tests were designed using a random block design and ANOVA at p<0.05. Appropriate post-hoc tests were performed, including the "Duncan Multi-Distance Test" (DMRT) and the "Least Significant Difference" (LSD).

## 3. RESULTS AND DISCUSSION

### 3.1 Total Phenolic and Flavonoid Contents

Table 1 shows the amounts of phenolic and flavonoid content in several medicinal plants. Among these three solvents, MetOH extraction yielded the most phenolic acid in all three extracts, but Aq. solvent extraction yielded the least phenolic content. TPC (16.67, 17.78 and 10.59 mg GAE/g extract respectively) and TFC (14.21, 12.75 and 8.10 mg QE/g dw extract respectively) are abundant in *Z. officinale* extracts in MetOH, EtOH, and Aq. Solvent.

Comment [J9]: Space

Comment [J10]: Follow the same trend as your title

Comment [J11]: Space

Comment [J12]: Space

Comment [J13]: Space

Comment [J14]: Space

Comment [J15]: Space

Comment [J16]: Which one comes first in your title between DPPH and total phenolic and flavonoid

Comment [J17]: thrice

Comment [J18]: Space

Comment [J19]: Rndomised Block Design

Comment [J20]: The correct meaning is Duncan Multiple Range Test. Please check

**Table 1. Total phenolic and flavonoid content of plant extracts. (Results are mean  $\pm$  standard deviation (SD); significance value according to at  $P < 0.05$  following ANOVA and LSD)**

Plants/ Extracts	TPC (mg GAE/g dw)			TFC (mg QUE/g dw)		
	MetOH	EtOH	Aq.	MetOH	EtOH	Aq.
TS	11.56 $\pm$ 0.09	9.73 $\pm$ 0.07	4.30 $\pm$ 0.14	7.96 $\pm$ 0.06	6.20 $\pm$ 0.11	3.66 $\pm$ 0.06
ZO	16.67 $\pm$ 0.04	12.78 $\pm$ 0.06	10.59 $\pm$ 0.07	14.21 $\pm$ 0.10	12.75 $\pm$ 0.13	8.10 $\pm$ 0.10


Results are mean  $\pm$  standard deviation (SD), EtOH= Ethanolic extract, MetOH= Methanolic extract, Aq.= Aqueous extract; *T. stans*=TS; *Z. officinales* = ZO.


**Comment [J21]:** Did you use DMRT and LSD or LSD alone?

### 3.2 Total antioxidant activity:

The efficacy of three distinct solvent extracts to scavenge DPPH free radicals was tested and compared to the standard, ascorbic acid. Figure 1 shows the antioxidant properties of medicinal plant extracts as assessed by DPPH scavenging experiments. The present study's results demonstrated that *Z. officinales* sample DPPH $\cdot$  scavenging activity considerably increased in methanol solvent. However, the extracts' DPPH radical scavenging capacities were lower than those of ascorbic acid (82.29%) at 300 $\mu$ g/ml. This result clearly shows that the extracts have proton donating potential and might be used as free radical inhibitors or scavengers, possibly functioning as primary antioxidants

**Comment [J22]:** Space





**Figure 1. Total antioxidant activity of *Z. officinales* and *T. stans* in different solvents (A) methanolic (B) ethanolic (C) aqueous.**

In the current investigation, high TPC and TFC values, as well as antioxidant activity, were found in methanolic solvent extract. Plant phenolic compounds are potent antioxidants and antimicrobials [24,25]. Several investigations have shown that flavonoids in plant extracts have antioxidant and antifungal properties[26,27].

The presence of phenolic and flavonoid compounds in these plants may enhance their biological capabilities when compared to other therapeutic plants evaluated. The antioxidant capabilities of phenolic and flavonoids compounds are determined by the structural interaction between various elements of their chemical structure. Natural polyphenols have the ability to eliminate free radicals, bind metal prooxidants, decrease radicals, inhibit oxidases, and terminate the oxidation chain reaction [28]. Based on in vitro findings, these medicinal plants are intriguing and promising as a possible source for new antioxidant medications.

**Comment [J23]:** Space

**Comment [J24]:** You may lengthen your discussion a little bit by stating their relevance in drug industry

## REFERENCES

1. Khare T, Anand U, Dey A, Assaraf YG, Chen ZS, Liu Z, Kumar V. Exploring phytochemicals for combating antibiotic resistance in microbial pathogens. *Frontiers in Pharmacology*. 2021 Jul 21;12:720726.
2. Vaou N, Stavropoulou E, Voidarou C, Tsigalou C, Bezirtzoglou E. Towards advances in medicinal plant antimicrobial activity: A review study on challenges and future perspectives. *Microorganisms*. 2021 Sep 27;9(10):2041.
3. Bisso BN, Kayoka-Kabongo PN, Tchuengem RT, Dzoyem JP. Phytochemical analysis and antifungal potentiating activity of extracts from loquat (*Eriobotrya japonica*) against *Cryptococcus neoformans* clinical isolates. *Advances in Pharmacological and Pharmaceutical Sciences*. 2022 Apr 14;2022.
4. Rasool A, Bhat KM, Sheikh AA, Jan A, Hassan S. Medicinal plants: Role, distribution and future. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(2):2111-4.
5. Savadi S, Vazifedoost M, Didar Z, Nematshahi MM, Jahed E. Phytochemical analysis and antimicrobial/antioxidant activity of *Cynodon dactylon* (L.) Pers. rhizome methanolic extract. *Journal of Food Quality*. 2020 Apr 1;2020:1-0.

6. Bonilla J, Poloni T, Lourenço RV, Sobral PJ. Antioxidant potential of eugenol and ginger essential oils with gelatin/chitosan films. *Food Bioscience*. 2018 Jun 1;23:107-14.
7. Gouveia S, Castilho PC. Antioxidant potential of *Artemisia argentea* L'Hér alcoholic extract and its relation with the phenolic composition. *Food Research International*. 2011 Jul 1;44(6):1620-31.
8. van Breemen RB, Tao Y, Li W. Cyclooxygenase-2 inhibitors in ginger (*Zingiber officinale*). *Fitoterapia*. 2011 Jan 1;82(1):38-43.
9. Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC, Seal T, Mukherjee B. Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases. *Journal of Ethnopharmacology*. 2003 Feb 1;84(2-3):131-8.
10. Stoilova I, Krastanov A, Stoyanova A, Denev P, Gargova S. Antioxidant activity of a ginger extract (*Zingiber officinale*). *Food Chemistry*. 2007 Jan 1;102(3):764-70.
11. Maizura M, Aminah A, Wan Aida WM. Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract. *International Food Research Journal*. 2011 May 1;18(2).
12. Ali BH, Blunden G, Tanira MO, Nemmar A. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food and Chemical Toxicology*. 2008 Feb 1;46(2):409-20.
13. Abdalla WE, Abdallah EM. Antibacterial activity of ginger (*Zingiber officinale* Rosc.) rhizome: A mini review. *International Journal of Pharmacognosy & Chinese Medicine*. 2018;2(4):000142.
14. Jabborova D, Annapurna K, Fayzullaeva M, Sulaymonov K, Kadirova D, Jabbarov Z, Sayyed RZ. Isolation and characterization of endophytic bacteria from ginger (*Zingiber officinale* Rosc.). *Annals of Phytomedicine*. 2020;9:116-21.
15. Gonçalves TP, Parreira AG, dos Santos Zanuncio VS, de Souza Farias K, da Silva DB, dos Santos Lima LA. Antibacterial and antioxidant properties of flowers from *Tecoma stans* (L.) Juss. ex Kunth (Bignoniaceae). *South African Journal of Botany*. 2022 Jan 1;144:156-65.
16. Lozoya-Meckes M, Mellado-Campos V. Is the *Tecoma stans* infusion an antidiabetic remedy? *Journal of Ethnopharmacology*. 1985 Sep 1;14(1):1-9.
17. Singh A, Nagori BP, Mathur K. *Tecoma stans*: an important medicinal plant. *Journal of Pharmaceutical Erudition*. 2013 Aug;3:13-21.
18. Anburaj G, Marimuthu M, Rajasudha V, Manikandan R. In vitro anti-cancer activity *Tecoma stans* against human breast cancer yellow elder (*Tecoma stans*). *Journal of Pharmacognosy and Phytochemistry*. 2016;5(5):331-4.
19. Taher MA, Dawood DH, Sanad MI, Hassan RA. Searching for anti-hyperglycemic phytomolecules of *Tecoma stans*. *European Journal of Chemistry*. 2016 Dec 31;7(4):397-404.
20. Al-Azzawi A, Al Dibsawi A, Talath S, Wali AF, Sarheed O. Method Development: The antioxidant and antifungal activity of the isolated component from the ethanolic extract of *Tecoma stans* leaves using flash chromatography. *Separations*. 2022 Oct 18;9(10):317.
21. Chakraborty D, Mandal SM. Fractional changes in phenolic acids composition in root nodules of *Arachishypogaea* L. *Plant Growth Regulation*. 2008 Jul; 55:159-63.
22. CI KC, Indira G. Quantitative estimation of total phenolic, flavonoids, tannin and chlorophyll content of leaves of *Strobilanthes kunthiana* (Neelakurinji). *Journal of Medicinal Plants*. 2016; 4:282-6.
23. Aliyu AB, Ibrahim MA, Musa AM, Musa AO, Kiplimo JJ, Oyewale AO. Free radical scavenging and total antioxidant capacity of root extracts of *Anchomanes difformis* Engl. (Araceae). *Acta Poloniae Pharmaceutica* 2013 Jan 1;70(1):115-21.
24. Alam MN, Bristi NJ, Rafiquzzaman M. Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*. 2013; 21(2): 143-152. doi: <https://doi.org/10.1016/j.jsps.2012.05.002>

25. Adedapo AA, Jimoh FO, Koduru S, Masika PJ, Afolayan AJ. Assessment of the medicinal potentials of the methanol extracts of the leaves and stems of *Buddlejasaligna*. BMC Complementary and Alternative Medicine. 2009 Dec;9(1):1-8.
26. Adesegun SA, Fajana A, Orabueze CI, Coker HA. Evaluation of antioxidant properties of *Phaulopsisfascisepala* CB Cl. (Acanthaceae). Evidence-Based Complementary and Alternative Medicine. 2009 Jun 1; 6:227-31.
27. Yoshida T, Konishi M, Horinaka M, Yasuda T, Goda AE, Taniguchi H, Yano K, Wakada M, Sakai T. Kaempferol sensitizes colon cancer cells to TRAIL-induced apoptosis. Biochemical and Biophysical Research Communications. 2008 Oct 10;375(1):129-33.
28. Carocho M, Ferreira IC. A review on antioxidants, prooxidants and related controversy: Natural and synthetic compounds, screening and analysis methodologies and future perspectives. Food and chemical toxicology. 2013 Jan 1;51:15-25.

UNDER PEER REVIEW