

## Case report

### ANEURYSMAL BONE CYST OF MAXILLA MIMICKING DENTIGEROUS CYST: A RARE CASE REPORT

#### Abstract:

Aneurysmal bone cysts are not real cysts or aneurysms; they are rare non-neoplastic bone lesions occurring mainly in the long bone metaphyses, they are rarely seen in the facial bones. Aneurysmal bone cyst (ABC) is a rare benign intraosseous lesion characterized by blood-filled spaces of various sizes. The incidence of aneurysmal bone cyst in the craniofacial skeleton is rare which may remain undiagnosed for a long period prior to becoming symptomatic which may cause associated issues secondary to compression by extending to the surrounding vital anatomical areas. In the jawbones, the mandible is involved more commonly than the maxilla. We present a rare case of aneurysmal bone cyst in maxilla in 16-year-old male patient reporting to our department with complaint of painless swelling in upper right anterior region of jaw. Based upon clinical examination and radiographic examination this case was diagnosed as dentigerous cyst. Enucleation was done but histopathological features were in accordance with aneurysmal bone cyst. An aggressive course can lead to bony destruction with intracranial extension and are more likely to pose a diagnostic challenge as they are more likely to be confused clinically with other expansile radiolucent bone lesions such as dentigerous cyst, ameloblastoma, osteoblastoma, and giant cell tumor among many others. Hence we conclude that practitioners must have knowledge of aneurysmal bone cyst for correct diagnosis, investigation and effective management.

## **Keywords:**

Aneurysmal bone cyst, Case report, Cyst, Enucleation, Impacted tooth, Maxilla,

## **Introduction :**

“Aneurysmal bone cyst (ABC) has been described for the first time in 1942 by Jaffe and Lichtenstein”.<sup>[1]</sup> In 2002, an ABC was defined by the World Health Organisation (WHO) as ‘a benign cystic lesion of bone composed of blood filled spaces separated by connective tissue septa containing fibroblasts, osteoclast-type giant cells and reactive woven bone’<sup>[2]</sup> “Aneurysmal bone cysts (ABC) are typically found in long bones and the spine, but rarely can be seen in the craniofacial region. The incidence is 2% and 1.3% in the craniofacial region and maxillary region, respectively”.<sup>[3]</sup> “Available literature suggests that the mandible is affected more frequently than the maxilla, the proportions varying from 2:1 to 11:9”.<sup>[4]</sup> “The etiology and pathogenesis of ABCs are still uncertain but are thought to be reactively related to previous trauma or pathology. The leading hypothesis etiology in ABC pathology is vascular and affected by local bone changes”.<sup>[5]</sup> “It may be clinically asymptomatic or may arise as a quickly advancing lesion enlarged to cause a pathological fracture”.<sup>[6]</sup> “Diagnosis should base on clinical symptoms, radiological, and histopathological examinations. A diagnosis of dentigerous cyst is usually entertained when impacted tooth is associated within a cyst. However, we report a rare case of aneurysmal bone cyst of maxilla with impacted maxillary right central incisor tooth within the cyst causing diagnostic dilemma. The purpose of this case report is to emphasize the importance of ABC diagnosis and effective treatment.

## **Case Report:**

**Patient information:** A 16 years old male patient reported to the department of oral medicine and radiology with complaint of painless swelling in upper right front region of jaw since 2 -3 months. Swelling was gradually increasing in the size to attain the present size. There was no history of trauma and no significant past medical or family history or known allergy.

**Clinical findings:** on extra oral examination, there was a diffuse swelling of right mid face of normal skin colour extending mediolaterally from ala of the nose towards tragus of ear and superoinferiorly from 1.5 cm below inferior orbital fissure till 1.5 cm below alaragus line. With obliterated nasolabial fold. Absence of extraoral fistula. On palpation swelling was nontender, soft in consistency, with smooth surface, afebrile, nonfluctuant, noncompressible, nonreducible, nonpulsatile. No signs of paraesthesia reported. Intra oral hard tissue examination showed missing teeth with 11, palatally displaced with 12, grade ii mobility with 11, 21, crowding with 21,22,23 and over retained with 63. Soft tissue examination revealed solitary, dome shaped, bluish red coloured swelling of size approximately 3 x 3 cm present on attached gingiva of 11,12,13. Obliteration of right upper labial vestibule was evident. Absence of any discharge with smooth surface texture and surrounding tissue appears to be normal. On palpation swelling was smooth, sessile, nontender, firm in consistency, nonfluctuant, noncompressible, nonreducible, nonpulsatile. [fig.1]



**Figure 1: Preoperative image**

Based on the history and clinical examination a provisional diagnosis of dentigerous cyst with 11 was made. The differential diagnoses included AOT, ameloblastoma, aneurysmal bone cyst, CGCG, AV shunt.

**Diagnostic assessment:** FNAC was done in which 5ml brownish fluid smear revealed numerous foamy macrophages against the background of peripheral blood smear. No epithelial cells seen. [fig. 2]




**Figure 2: FNAC with 5ml brownish fluid**

Routine panoramic radiograph revealed well-defined unilocular radiolucency in maxillary right anterior region, roughly oval in shape of size approximately 4cm mesiodistally and 3 cm superoinferiorly. Internal structure of lesion shows impacted 11. Displacement of 12, 13 is evident. [fig 3]



**Figure 3: OPG**

CBCT scans overall radiographic impression was osteolytic lesion w r t right maxillary anterior and posterior region. [fig 4]




**Figure 4: CBCT**

D/d-adenomatoid odontogenic tumor, dentigerous cyst, aneurysmal bone cyst, unicystic ameloblastoma

**Therapeutic intervention:** haematological investigations were within normal range. The treatment planned included extraction of 12 followed by enucleation of cyst under general anaesthesia. Resected tissue was then sent for histopathological examination which revealed no epithelial lining and multiple blood-filled spaces with stroma consisting of spindle shaped cells and multinucleate giant cells which was suggestive of aneurysmal bone cyst. [fig 5] Hence a final diagnosis of ABC was made.

**Follow up and outcomes:** Patient condition was improved in follow up after 20 days.



**Figure 5: (a) Excised tissue, (b) Histopathology**

**Discussion:**

“An aneurysmal bone cyst (ABC) is a pseudocyst due to the absence of a cyst epithelial. Bernier and Bhaskar reported the first ABC case containing the maxillofacial skeleton in 1958”.<sup>[5]</sup> “Jaffe and Lichtenstein defined it in 1942 as blood-filled lesions in which giant cells and irregular bone trabeculae”.<sup>[1]</sup> “More than 90% of ABC found in the jaws occur in the posterior areas, especially in the mandible”.<sup>[7]</sup> in our case it was found in maxillary anterior region. “The pathogenesis of ABC is still controversial. The role of trauma has been

suggested by few authors”.<sup>[1]</sup>“Some authors favoured intra medullary hematoma as the cause of ABC”.<sup>[8]</sup>“Local circulatory abnormalities leading to increased venous pressure and resulting in dilation of the local vascular network is one of the most widely accepted theories available in literature”.<sup>[9]</sup>“Recently, chromosomal translocation  $t(16;17)(q22;p13)$  as a recurrent cytogenetic abnormality has also been suggested as a responsible factor for primary ABCs”.<sup>[10]</sup>“Aneurysmal bone cysts are located most commonly in the shaft of a long bone or in the vertebral column. It is generally found in patients younger than the age of 20. No significant gender predilection is noted”.<sup>[11]</sup>present case was 16 year old male patient. These lesions can have multiple clinical features, from an asymptomatic lesion discovered on routine X-ray to a sometimes-expansive lesion. “However, the main symptom is a painless swelling of the jaw. Additional symptoms, such as limited mouth opening, loosening of teeth, nasal obstruction and lip paraesthesia are described in literature”.<sup>[12]</sup>“ABC usually presents as a rapidly growing swelling. Paraesthesia is a very rare feature. Crepitus and perforation is also uncommon. On occasion, malocclusion, mobility, migration of involved teeth may be present. Maxillary lesions often bulge into the adjacent sinus; nasal obstruction, nasal bleeding, proptosis, and diplopia are noted uncommonly. Radiology is suggestive but not diagnostic for aneurysmal bone cyst. These lesions are usually unilocular, but long standing lesions may show a “soap-bubble” appearance and may become progressively calcified. Destruction of the bone cortex and periosteal reaction is rare”.<sup>[11]</sup>“Radiographically it may appear as expansive, osteolytic, unilocular or multilocular radiolucent lesion, with expansion and thinning of the surrounding cortical bone. This variegated appearance in imaging could be due to the various types of aneurysmal bone cyst (solid, vascular and mixed)”.<sup>[13]</sup>“The radiographic appearance of maxillary aneurysmal bone cysts is even less characteristic than that of mandibular lesion. CT scan ABC may appear unicystic, multilocular, or moth eaten, causing expansion, perforation, or extensive destruction of the bony cortices”.<sup>[14]</sup>

“Histologically ABC is characterized by blood-filled spaces of various sizes and channels divided by connective tissue septa, which can contain osteoid tissue, osteoclast-like giant cells and woven bone”.<sup>[7]</sup> “Three types of ABC have been described based on histopathological features.<sup>[15]</sup> Solid type (5% of the cases) is characterized by a dense stroma, scanty sinusoids, few blood vessels and caverns, bone expansion (instead of perforation), and without severe bleeding during surgery. Vascular variant (95% of cases) is characterized by a loose scanty stroma, numerous engorged blood-filled sinusoids and caverns. Brisk bleeding during surgery and extensive bony destruction with spread in the soft tissues are also obvious. The mixed type lies between the 2 previous variants”.<sup>[15]</sup> Our case was of solid variant type as it was not associated with severe bleeding during surgery. “Surgical therapy is the most frequently applied treatment of ABCs,<sup>[6]</sup> although embolization, cryotherapy, and wait and see strategy have also been used as other treatment modalities”.<sup>[4]</sup> “Recurrence rates range from 20% to 30% according to different series and seems to occur most frequently within the first year after surgery”.<sup>[4,11,16]</sup> “Concerning treatment options for ABCs, it has been shown that, the extent of the lesion is usually used in deciding whether the patient should undergo either surgical resection or curettage”. [18] The first treatment approach was described in the work of Aiba *et al.* on ABCs, which included “curettage and reconstructing the defect with bone graft, which remains the mainstay of modern treatment”. “However, this approach was later found to be associated with high rates of recurrence. Self-healing after long-term follow-up of patients with ABCs has also been reported. Surgical resection is also another option for treating patients with ABCs”.<sup>[16]</sup> “Several authors recommend immediate reconstruction of the defect with autogenous grafts in cases of aesthetic deformity and in cases with high risk of fractures”.<sup>[11]</sup> “Considering the aggressive nature of the lesion and age of the patient, a conservative surgical resection of the lesion was done. Novel, less invasive treatment alternatives are under research. Denosumab—monoclonal antibody suppressing osteoclast

activity—administration significantly improved outcomes in 9 patients with ABCs but with a higher recurrence rate compared to surgical resection” [9]. “Another study used autologous bone-marrow-derived mononuclear cells with  $\beta$ -tricalcium phosphate and absorbable atelocollagen to form new bone within the cysts”.<sup>[17]</sup>

### **Conclusion:**

ABCs of maxilla are rarer. Impacted tooth within aneurysmal bone cyst leads to diagnostic dilemma. biopsy and aspiration may help in a definitive diagnosis over its confusing clinical and radiological features which are similar to various other pathologic entities of the jaws. Conservative surgical resection is preferable to curettage as the latter is associated with much higher rate of recurrence.

### **Ethical Approval:**

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

### **Consent**

As per international standards, parental written consent has been collected and preserved by the author(s).

### **References :**

1. JAFFE HL, Lichtenstein L. Solitary unicameral bone cyst: with emphasis on the roentgen picture, the pathologic appearance and the pathogenesis. *Archives of Surgery*. 1942 Jun 1;44(6):1004-25.
2. Barnes L, editor. *Pathology and genetics of head and neck tumours*. IARC; 2005.
3. Sharma BB, Ramchandran P, Sharma S, Sharma S. Aneurysmal bone cyst of the maxilla rare presentation with radiological and pathological correlation. *Plastic and Aesthetic Research*. 2015 Mar 13;2:73-5.
4. Debnath SC, Adhyapok AK, Hazarika K, Malik K, Vatsyayan A. Aneurysmal bone cyst of maxillary alveolus: A rare case report. *Contemporary Clinical Dentistry*. 2016 Jan;7(1):111.
5. Alparslan ES, BAŞTÜRK F, GÜRSES G, DOLANMAZ D. Aneurysmal bone cyst of the mandible: A case report of 5 years follow-up. *Current Research in Dental Sciences*. 2022 Apr 4;32(2):185-8.
6. Motamedi MH. Aneurysmal bone cysts of the jaws: clinicopathological features, radiographic evaluation and treatment analysis of 17 cases. *Journal of Cranio-Maxillofacial Surgery*. 1998 Feb 1;26(1):56-62.
7. Urík M, Šlapák I, Máchalová M, Jančíková J, Šíkolová S, Pavlovská D, Jabandžiev P, Ježová M. An expansive aneurysmal bone cyst of the maxillary sinus in an 8 year old child: Case report and review of literature. *Acta Oto-Laryngologica Case Reports*. 2020 Jan 1;5(1):86-90.
8. Hillerup S, Hjørting-Hansen E. Aneurysmal bone cyst—simple bone cyst, two aspects of the same pathologic entity?. *International journal of oral surgery*. 1978 Feb 1;7(1):16-22.
9. Biesecker JL, Marcove RC, Huvos AG, Mike V. Aneurysmal bone cysts. A clinicopathologic study of 66 cases. *Cancer*. 1970 Sep;26(3):615-25.

10. Panoutsakopoulos G, Pandis N, Kyriazoglou I, Gustafson P, Mertens F, Mandahl N. Recurrent t (16; 17)(q22; p13) in aneurysmal bone cysts. *Genes, Chromosomes and Cancer*. 1999 Nov;26(3):265-6.
11. Iyengar AR, Patil S, Uppal MK, Jason A, Karibasappa GG. A Case Report of Aneurysmal Bone Cyst of the Mandible.
12. Sun ZJ, Zhao YF, Yang RL, Zwahlen RA. Aneurysmal bone cysts of the jaws: analysis of 17 cases. *Journal of Oral and Maxillofacial Surgery*. 2010 Sep 1;68(9):2122-8.
13. Bataineh AB. Aneurysmal bone cysts of the maxilla:: A clinicopathologic review. *Journal of oral and maxillofacial surgery*. 1997 Nov 1;55(11):1212-6.
14. Verma RK, Kumar R, Bal A, Panda NK. Aneurysmal bone cyst of maxilla with ectopic molar tooth–A case report. *Otolaryngologia Polska*. 2013 Nov 1;67(6):302-7.
15. Rosai J. Stromal tumors (GISTs and related lesions) *Ackerman's Surgical Pathology*.
16. Yahaya JJ, Morgan ED, Abraham ZS, Othieno E. Aneurysmal bone cyst of the mandible: a rare case report and literature review. *Annals of Medicine and Surgery*. 2023 Oct 1;85(10):5133-7.
17. Khan JA, Saleh T, Shafqat A, Albalkhi I, Saleh W. Unusual presentation of an aneurysmal bone cyst: A case report and literature review. *Radiology Case Reports*. 2023 Mar 1;18(3):1320-3.
18. Yahaya JJ, Morgan ED, Abraham ZS, Othieno E. Aneurysmal bone cyst of the mandible: a rare case report and literature review. *Annals of Medicine and Surgery*. 2023 Oct 1;85(10):5133-7.