

Original Research Article

Preliminary Inventory Of Stomoxes, Potential Vectors Of Pathogens In the Minkébé Forest Massif In the North-East Of Gabon

ABSTRACT

Aims: This study aims to characterize Stomoxes populations, potential vectors of pathogens in the north-east of Gabon.

Place and Duration of Study: This research was conducted in the Minkébé forest massif, Woleu-Ntem Province, Gabon from April to May 2023 (rainy season).

Methodology: The daily captures were conducted at Mikouka, Massoko and Minkébé using Vavoua and Nzi traps. All captured specimens were morphologically identified to species using binocular microscope (Leica Microsystems ©). The identification was aided by “specific” taxonomic keys. The collected data were organized from Microsoft Office Excel 2013. This database was used to calculate numbers and proportions of specimens observed by species. Several parameters including Apparent Density per Trap per day (ADT), Shannon, Simpson, Pielou and Bray-Curtis indices were evaluated.

Results: A total of 1552 Stomoxes were caught, including 610 at Massoko, 533 at Mikouka and 409 at Minkébé. These stomoxes belonged to two species: *Stomoxys nigerriger* and *Stomoxys omega*. *Stomoxys nigerriger* was the most abundant with 1176 specimens caught and an ADT of 4 stomoxes/trap/day (s/p/d). On the other hand, *Stomoxys omega* was poorly captured (n= 376; ADT= 1.28). As for the biotopes surveyed, Massoko (n= 619; ADT= 6.42) and Mikouka (n= 533; ADT= 5.44) were the most infested with Stomoxes. However, the lowest catches were recorded at Minkébé (n= 409; ADT= 4.17). The Shannon, Pielou and Simpson index values indicate relatively low levels of diversity between sites. The Shannon index reached higher values at Minkébé than at the other study sites. The Simpson and Pielou indices also had higher values at Minkébé. As for the Bray-Curtis index, its values were low between Minkébé and Mikouka (0.592), between Mikouka and Massoko (0.302) and between Minkébé and Massoko (0.384).

Conclusion: Our findings show the coexistence of *Stomoxys nigerriger* and *Stomoxys omega* in Minkébé. Their presence constitutes a health risk, because these vectors could play a role in the circulation of certain pathogens in the Minkébé forest massif. So, a longitudinal study of these vectors and a screening of their infectious agents are urgently needed.

Keywords: *Stomoxys nigerriger*, *Stomoxys omega*, insect vectors, Minkébé, Gabon

1. INTRODUCTION

The vector-borne diseases are among the major causes of morbidity and mortality in many developing countries of tropical Africa [1, 2]. According to the World Health Organization (WHO), they account for 17% of infectious diseases worldwide and are responsible for more than one million deaths by year [2]. Among these diseases, trypanosomiasis and filariasis represent one of the main obstacles to the development of livestock farming, food security and human health [3-5]. In the livestock sector, the economic losses due to these diseases have been estimated at around 400 million dollars [6, 7]. These two vector-borne diseases are transmitted by numerous groups of insects, including the stomoxes [5]. The stomoxes are hematophagous insects [8] that play a vectorial role in the transmission of several pathogens [4, 5, 9, 10]. These flies occupy an important place in the epidemiological system and therefore constitute a major health problem for domestic and/or wild fauna, as well as for humans [11-16]. According to Raymond [17], stomoxes are involved in the transmission of *Trypanosoma evansi* (Balbiani, 1888) in Africa, America and Asia.

In Gabon, stomoxes populations are relatively well known in certain regions [16, 18-22].

However, in many other localities, they are still poorly known. This is the case of the Minkébé forest massif in North-East of Gabon [23] and where the meteorological and environmental conditions are

favourable to the development and maintaining of stomoxes, potential vectors of trypanosomes and filariae. To fill this gap, an entomological study was carried out at three sites in the Minkébé region in order to characterize the stomoxes populations present in the Minkébé forest massif.

2. MATERIAL AND METHODS

2.1 Study area

This study was conducted in the Minkébé forest massif located in the Woleu-Ntem province in North-East of Gabon (Figure 1). This region has an equatorial climate with two rainy seasons and two dry seasons [23]. The rainy seasons extend from October to December (short rainy season) and from March to May (long rainy season). On the other hand, the dry seasons extend from January to February (short dry season) and from June to August (long dry season). Average annual rainfall varies between 1,600 and 1,800 mm [23]. In general, average temperatures are between 23°C and 24°C, while humidity levels usually exceed 80%.




Fig. 1. Map of study area showing the three sites surveyed in Woleu-Ntem province

The vegetation of this study area is composed of Guinean-Congolian dense forests [23]. These forests cover a large part of the region and form an almost continuous vegetation carpet marked by a mosaic of evergreen and semi-deciduous plant formations [24]. Waterbodies are often bordered by vast areas of floodplain and swamp forests [23]. Regarding savannahs, they are found only at Mvadi and in the south-west of the Minkébé National Park [24].

Moreover, the Minkébé forest massif has undergone several human disturbances, especially in the Minkébé area, due to the increase of the gold-washing between the 1950s and the 2000s [24]. This activity has had important repercussions on this forest massif through the establishment of gold-washing camps and the significant decline of wildlife in areas of human activities [24].

2.2 Capture of stomoxes at the study surveyed

The stomoxes were collected using 7 Vavoua and 7 Nzi traps [25, 26]. These traps have been used many times in the studies of hematophagous dipterans carried out in Gabon [19, 20, 22, 27]. They were

installed at three different forest sites, namely Mikouka, Massoko and Minkébé, according to the following sequence: Mikouka-Massoko-Minkébé.

These traps were placed both in the forest environments and along the waterbodies, in order to maximize the quality and quantity of the catches. In this fact, each site surveyed was equipped with seven Vavoua traps and seven Nzi traps.

These traps were labelled with a corresponding number and were installed at given collection points. These traps were equidistant at least 500 m in order to better distribute the traps across the study sites. They were set up at 07:00 a.m. on the first day of trapping and removed at 06:00 p.m. on the last day. During the present study, these traps were continually in use and were checked at 6:00 p.m. daily. When the traps were visited, the captured insects were brought back to the camp and packaged for later identification in the laboratory in Libreville. All stomoxes captures were carried out between 28th April and 18th May 2023 (rainy season), with 7 days of capture for each study site.

2.3 Morphological identification of stomoxes

All the captured insects were identified morphologically using binocular microscope at the Laboratoire d'Ecologie Vectorielle of Institut de Recherche en Ecologie Tropicale (LEV-IRET). To this end, an initial separation was made between the stomoxes and the other insects. The stomoxes were then carefully identified at the species level using the keys developed by Zumpt [12] and Garros et al. [28] in order to distinguish more easily *S. calitrans* (Linnaeus, 1758) and *S. nigerniger* (Macquart, 1851).

2.4 Data analysis

The collected data were meticulously entered and organized from Microsoft Office Excel 2013. This spread sheet was used to calculate numbers and proportions of specimens observed by species. The data were then presented as the graphs and the tables to make it easier to visualize the different results. Moreover, several parameters including Apparent Density per Trap per day, Shannon's index, Simpson's index, Pielou's index and Bray-Curtis's index, were also estimated in order to better understand the biodiversity of the stomoxes populations colonizing the study sites.

The Apparent Density per Trap per day (ADT) corresponds to the relative abundance of stomoxes species in each surveyed site. It was calculated using the following formula:

$$ADT = \frac{\text{Number of captured stomoxes}}{\text{Number of traps} \times \text{number of trapping days}}$$

The Shannon's Diversity Index (H') was also calculated in order to assess the specific diversity of stomoxes at the surveyed sites. This index is expressed according to the following formula: $H' = -\sum (Ni/N) \times \log (Ni/N)$; where Ni : represents the number of individuals of a given species and N : the total number of individuals.

Regarding the Simpson's Dominance index (D), it was estimated in order to determine the dominance of particular *Stomoxys* species in the community according to the following formula: $D = \sum Ni (Ni-1)/N(N-1)$; where D : represents Simpson's index, Ni : the number of individuals of the species and N : the total number of individuals.

In addition, Pielou's index (J) and Bray-Curtis's index were evaluated to provide further information on the study area. Indeed, the Pielou index enables to measure the equitability of the distribution of individuals between the different species of stomoxes present at each study site. Nevertheless, the Bray-Curtis's index is used to determine the similarity or dissimilarity between the species of stomoxes captured in the different study sites.

Spatial analyses were performed using Quantum GIS version 3.10.7 (<https://www.qgis.org/>, accessed on 10 September 2023).

3. RESULTS AND DISCUSSION

3.1 Species composition of stomoxes

A total of 1552 stomoxes were sampled in the study area with an apparent density per trap per day (ADT) of 5.28 stomoxes/trap/day (s/t/d). Of these stomoxes, 619 (39.30%) specimens were captured at Massoko with an ADT of 6.42 s/t/d and 533 (34.30%) at Mikouka with an ADT of 5.44 s/t/d. The lowest number of individuals was caught at Minkébé with 409 (26.40%) stomoxes with an ADT of 4.17 s/t/d (Figure 2).




Fig. 2. Apparent density of stomoxes per trap according to the sites surveyed

In total, 2 species of stomoxes, including *Stomoxysnigerniger* and *Stomoxys omega*, were identified during this study, with variable ADT. Indeed, *Stomoxysnigerniger* was the most abundant species with 1176 individuals caught and an ADT of 4 s/t/d. Nevertheless, *Stomoxys omega* was weakly collected (n= 376; ADT= 1.28).

Moreover, the ADTs of these two species vary according to the sites surveyed (Figure 3). Indeed, *Stomoxysnigerniger* was caught in abundance (461 individuals) at Massoko with an ADT of 4.70 s/t/d. In the other two environments, this species presented almost similar abundances with ADTs of 3.65 s/t/d and 3.64 s/t/d respectively.

Regarding *Stomoxys omega*, it was abundant at Mikouka and Massoko, with ADT of 1.80 s/t/d and 1.52 s/t/d respectively. On the other hand, at Minkébé, *Stomoxys omega* was collected very little, with 51 individuals and an ADT of 0.52 s/t/d (Table 1).




Fig. 3. Distribution of stomoxes species according to the sites surveyed

Table 1. Diversity and distribution of stomoxes according to the sites surveyed

The Shannon's index, Simpson's index and Pielou's index showed higher values at Minkébé ($H=0.829$; $D=0.759$; $J=0.625$) than at the other study sites (Table 2). However, the Bray-Curtis's index recorded the highest values between Minkébé and Mikouka (0.592), while between Mikouka and Massoko (0.302), then Minkébé and Massoko (0.384), the values of this index were low.

Table 2. Indices of diversity of stomoxes in the sites surveyed

4. DISCUSSION

This study revealed the concomitant presence of *Stomoxysnigerniger* and *Stomoxys omega* with variable abundances. The coexistence of these two stomoxes species could represent a danger for animals and humans, because stomoxes are known to be mechanical and/or biological vectors of bacteria, viruses, helminths or trypanosomes [5, 29, 30]. Globally, the collections of stomoxes were not abundant in the study area. This low abundance of catches could be explained by the reduced number of trapping days in this study compared to the trapping duration in previous studies [27]. However, the Nzi and Vavoua traps have confirmed their effectiveness in the capture of stomoxes in Gabon as reported by other authors [8, 19, 20, 27, 31]. However, the combination of these attractants would certainly have made it possible to increase the size of captures and improve species diversity [32]. In addition, the low number of collections of stomoxes could be attributed to the frequency of heavy rains and thunderstorms during the study period. These unfavourable weather conditions probably reduced the attractiveness of the traps used, causing thus fewer captures of these insects. From Raymond [33], the heavy rains have a negative impact on the development and the survival of hematophagous flies, as well as on the effectiveness of the traps. Besides, previous studies conducted

by Mavoungou et al. [34] and DoumbaNdaleboulyet al. [31] also reported a decrease in the abundance of hematophagous insects in response to rainfall in Makokou and Ndendé in Gabon. Among the species of stomoxes identified, *Stomoxysnigerniger* was more abundant than *Stomoxys omega*. This observation could be linked, on the one hand, to the ubiquity of *Stomoxys omega* and on the other hand, to the dependence of *Stomoxysnigerniger* on forest environments [12, 27]. Various authors have also observed this predominance of *Stomoxysnigerniger* compared with *Stomoxys omega* in Gabon and Cameroon [16, 35, 36]. Moreover, these two *Stomoxys* species showed a heterogeneous distribution, as attested by the values of the ecological indices. This observation could be attributed to ecological and environmental factors, as well as to the specific characteristics of the sites surveyed. Indeed, in Mikouka and Massoko sites, characterized by a relatively intact forest ecosystem, we noted a higher abundance of *Stomoxys omega*. On the other hand, in the Minkébé site, which is heavily impacted by gold-washing and therefore submitted to environmental disturbances, this species was rather low in abundance. These results are in line with those reported by Mavoungou [27] and Desquesnes et al. [37, 38], who showed that there was a correlation between the abundance of certain species of hematophagous insects, the presence of the forest and human disturbances on their abundance. Furthermore, according to Mavoungou [27], at Makokou (north-east of Gabon), variation of the abundance of *Stomoxys omega* was strongly correlated with the presence of primary forest, whereas *Stomoxysnigerniger* was closely linked to the presence of secondary forest and disturbed environments [39]. Consequently, it would be important that future studies incorporate an in-depth analysis of environmental factors to obtain a more complete understanding of the abundance and distribution of stomoxes in the Minkébé forest massif. In the global context of vector-borne infectious diseases, the identification of environmental factors determining the distribution of stomoxes has a great importance. The analysis of the ecological indices revealed low levels of diversity between the sites surveyed. Although the minimum number of individuals was obtained at Minkébé, this environment seems to present the greatest diversity, which is reflected in the higher values of these indices at this site. The Bray-Curtis's index revealed similarities between the communities of stomoxes in the sites surveyed. This observation is coherent, given that the three study sites are forest environments with variable levels of anthropization.

4. CONCLUSION

This study provides the first data on the populations of stomoxes present in the Minkébé forest massif. We identified two species of stomoxes including *Stomoxysnigerniger* and *Stomoxys omega*, having a heterogeneous distribution. These stomoxes are already known to be biological and/or mechanical vectors of trypanosomes in humans and animals.

Therefore, a particular attention must be paid to monitoring these insects in order to better understand the biology and the ecology of these insects. This is essential to develop the strategies of vector control which are specific to this region. The results of this study constitute a useful information base for implementing of vector control measures with in order to mitigate the associated risks and to prevent the spread of the diseases they transmit to their hosts. They also contribute to a better understanding of the abundance and distribution of stomoxes in the Minkébé forest massif in Gabon, where hematophagous flies have never been recorded.

In order to better control these vectors, bioecological and microbiological studies are needed. This will make it possible to assess the epidemiological importance of these insects in the Minkébé forest massif and to identify the environmental and anthropogenic factors determining the dynamics of their population. It would also be interesting to examine the genetic diversity of stomoxes and their pathogens, in order to have a better understanding of the mechanisms involved in the vector transmission.

REFERENCES

1. World Health Organization. Human African Trypanosomiasis (sleeping sickness): epidemiological update. *Weekly Epidemiological Record*. 2006;8:69–80.
2. World Health Organization. Vector-borne diseases. World Health Organization. WHO. 2020. Accessed 25 July 2023. Available: <https://www.who.int/fr/news-room/fact-sheets/detail/vector-borne-diseases>. French.
3. Rodhain F, Perez C. Summary of medical and veterinary entomology, Maloine. Paris. 1985;13:210-223.
4. Dumas J M. CIRDES international conference: Demographic developments and climate change: impacts on vector-borne diseases in West Africa. *Parasite*. 2009;16:75-78.
5. Duvallet G, Fontenille D, Robert V. Medical and Veterinary Entomology. IRD Éditions, Edition QUAE. Marseilles, Versailles. 2017;688p. French.
6. Kunz S E, Murell K D, Lambert G, James L F, Terill C E. Estimated losses of livestock to pests, In: D. Pimentel (Editor), *CRC Handbook of pest management in agriculture*, CRC Press, Boca Raton, FL. 1991;1:69-98.
7. Foil L D, Hogsette J A. Biology and control of tabanids, stable flies and horn flies. *Scientific and Technical Review of the Office International des Epizooties*. 1994;13(4):1125-1158.
8. Zinga Koumba C R, Mbang Nguema O A, Acapovi Yao G L, Shango M, M'batchi B, Mavoungou J F. Preliminary study of the infestation of tsetse flies by trypanosomes in the Mombabaï (North-eastern Gabon). *African Agronomy*. 2015;27(2):95-104. French.
9. Mavoungou J F. Ecology and vector role of stomoxes (Diptera: Muscidae) in Gabon. Unique Doctorate Thesis in Medical Entomology. University of Montpellier III Paul Valery, Montpellier, France. 2007;137p. French.
10. Dia M L, Desquesnes M, Hamadou S, Bouyer J, Yoni W, Gouro A S. Tetra Trap: Evaluation of a small model for the capture of animal trypanosomiasis vectors. *Journal of Veterinary Medicine*. 2008;159(1):17-21. French.
11. Leclercq M. Contribution to the study of Palaearctic Trypetidae (Diptera) and their relationships with plants. *Gembloux Agronomic Research Bulletin*. 1967;2(1):64-105. French.
12. Zumpt F. The Stomoxyinae biting flies of the world: Taxonomy, biology, economic importance and control measures. Gustav Fischer Verlag, Stuttgart. 1973;175p.
13. Desquesnes M, Dia M L. Mechanical transmission of *Trypanosoma vivax*: in cattle by the African Tabanid *Atylotus fuscipes*. *Veterinary Parasitology*. 2003;119:9-19.
14. Desquesnes M, Dia M L. Mechanical Transmission of *Trypanosoma congolense* in Cattle by the African Tabanid *Atylotus agrestis*. *Experimental Parasitology*. 2003;105:221-226.
15. Mramba F, Broce A B, Zurek L. Vector competence of stable flies, *Stomoxys calcitrans* L. (Diptera: Muscidae), for *Enterobacter sakazakii*. *Journal of Vector Ecology*. 2007;32(1):134-139.

16. Mavoungou J F, Jay-Robert P, Gilles J, Atsame Edda A, Duvallet G. Ecology of stomoxes (Diptera: Muscidae) in Gabon. I. First inventory in different ecological zones. *Parasite*. 2008;15:27-34. French.
17. Raymond H L. Insects harmful to livestock and climate. In Influence of climate on grazing animals. INRA SAD Guyana, Theix, March 31-April 1. 1982;169-183. French.
18. Mavoungou J F, Makanga B, Acapovi-Yao G L, Desquesnes M, M'batchi B. Chorology of Tabanidae (Diptera) in the Ipassa-Makokou Biosphere Reserve (Gabon) in the rainy season. *Parasite*. 2012;19:165-171. French.
19. Zinga-Koumba C R, Acapovi-Yao G L, Mavoungou J F, Kohagne Tongue L, Mbang-Nguema O A, Obame Ondo P K, Shango M. Influence of the season on the ecodistribution of tsetse flies, tabanids, stomoxes from the Baï of MombaMakokou , Gabon. *African Agronomy*. 2013;25:149-158. French.
20. Bitome Essono P Y, François-Xavier Dechaume-Moncharmont F-X, Mavoungou F, Obiang Mba R, Duvallet G, Bretagnolle F. Distribution and abundance of hematophagous flies (Glossinidae, Stomoxys, and Tabanidae) in two national parks of Gabon. *Parasite*. 2015;22:23-10. DOI: 10.1051/parasite/2015023.
21. Mavoungou J F, Mintsa Nguema R, Acapovi G L, Zinga Koumba R, Mounioko F, Sevidzem S L, Bakakas I K, Gilles J, Duvallet G, M'Batchi B, Picard N. Breeding Sites of Stomoxys spp (Diptera: Muscidae), a Preliminary Study in the Makokou Region (North-East Gabon). *Vector Biology Journal*. 2017;2:1. DOI: 10.4172/2473-4810.1000115.
22. Zinga Koumba C R, Koumba A A, Mouniko F, Sevidzem S L, Sima Owono L R, Ovono A P M, Djogbenou L S, Acapovi-Yao G L, Mavoungou J F. Daily activity of Stomoxys spp. (Diptera: Muscidae) in the cattle farm of the Nyanga ranch (Southern Gabon). *Journal of Applied Biosciences*. 2023;188:19847-19859. French.
23. Vande Weghe J P. The national parks of Gabon: Minkébé, Wildlife Conservation Society, Libreville, Gabon. 2013;344p.
24. Sally A L. Gold panning in northeastern Gabon: History and socio-ecological analysis. Multi-disciplinary study for the National Center for Scientific and Technical Research and the Central African Regional Program for the Environment, Libreville, Gabon. 2002;136p.
25. Laveissière C, Grébaud P. Research on tsetse traps (Diptera: Glossinidae). Development of an economic model: the "Vavoua" trap. *Tropical Medicine and Parasitology*. 1990;41(2):185-192.
26. Mihok S. The development of a multipurpose traps (the Nzi) for tsetse and other biting flies. *Bulletin of Entomological Research*. 2002;92:385-403.
27. Mavoungou J F. Ecology and vector role of stomoxes (Diptera: Muscidae) in Gabon. Doctoral thesis. Montpellier III Paul Valéry University, Montpellier, France. 2007;137p.
28. Garros C, Gilles J, Duvallet G. A new morphological character to distinguish *Stomoxys calcitrans* and *S. niger* (Diptera: Muscidae). Comparison of populations on Reunion Island. *Parasite*. 2004;11:329-332. French.
29. Baldacchino F, Muenworn V, Desquesnes M, Desoli F, Charoenviriyaphap T, Duvallet G. Transmission of pathogens by *Stomoxys* flies (Diptera, Muscidae): a review. *Parasite*. 2013;20:26. DOI: 10.1051/parasite/2013026.
30. Mounioko F, Maganga G D, Mavoungou J F, Zinga Koumba C R, Koumba A A, Sevidzem S L et al. Molecular Screening of *Trypanosoma* spp. in *Glossina*, *Stomoxys* and *Tabanids* in the Moukalaba Doudou National Park (South-West, Gabon). *World Journal of Veterinary Science*. 2018;6:52-61.

31. DoumbaNdalebouly A G, Zinga Koumba C R, Mounioko F, Maroundou A P, Mbang Nguema O A, Acapovi-Yao G L et al. Contribution to the study of Stomoxes and Tabanids, mechanical vectors of trypanosomoses in the Ndené region in southern Gabon. *Faunistic Entomology-Faunistic Entomology*. 2016;69:111-123.
32. Mihok S, Lange K. Synergism between ammonia and phenols for *Hybomitra* tabanids in northern and temperate Canada. *Medical and Veterinary Entomology*. 2012;26:282–290.
33. Raymond H L. Relative abundance and seasonal dynamics of Tabanidae (Diptera) in a savannah of French Guiana. *Canadian naturalist*. 1988;115:251-259. French.
34. Mavoungou J F, Makanga B, Acapovi-Yao G L, Desquesnes M, M'batchi B. Chorology of Tabanidae (Diptera) in the Ipassa-Makokou Biosphere Reserve (GABON) in the rainy season. *Parasite*. 2012;19:165-171. French.
35. Sevidzem S L, Affiri O J M, Zinga-Koumba C R, Mounioko F., Koumba, A. A., Nguema, R. M et al. Abundance and Diurnal Activity Rhythm of *Stomoxys* spp. in a Wildlife-human Interface in Makokou (North East-Gabon). *Asian J. Res. Zoo*. 2019;2:1-10.
36. Sevidzem S L, Mavoungou J F, Zinga-Koumba C R, Koumba A A, Duvallet G. Factors Influencing Seasonal and Daily Dynamics of the Genus *Stomoxys* Geoffroy, 1762 (Diptera: Muscidae), in the Adamawa Plateau, Cameroon. *International Journal of Zoology*. 2019;3636943:9.
37. Desquesnes M, Dia L M, Acapovi Yao G L, Yoni W, Foil L, Pin R. Mechanical vectors of animal trypanosomoses: Generalities, morphology, biology, impacts and control. Identification of the most abundant species in West Africa. Bobo-Dioulasso, Burkina Faso. CIRDES Editions. 2005;68p. French.
38. Desquesnes M, Dia M L, Dargantes A. Mechanical transmission of *Trypanosoma evansi* and *T. equiperdum* by *Stomoxys niger* and *S. taeniatus* in the Philippines. *Veterinary Parasitology*. 2005;127(1):23-29.
39. Gilles J, David J F, Duvallet G, De La Rocque S, Tillard E. Efficiency of traps for *Stomoxys calcitrans* and *Stomoxys niger* on Reunion Island. *Medical and Veterinary Entomology*. 2007;21:65-69.