

"Influence of land uses on soil carbon mineralization in selected agro-ecological units of south Kerala"

ABSTRACT

Soil is a major reservoir of terrestrial carbon and it plays an important role in the global carbon cycle. The soil organic carbon (SOC) is the fundamental factor for sustainable agriculture because of the ability in maintaining the soil fertility which is critical to soil productivity. Soil carbon pool has gained paramount importance in the recent decades owing to the alarming issues of climate change and global warming. Land use change was found to have a larger net effect on SOC storage than projected climate change. Depending on the land use management practices, soils can be a net sink or source for CO₂. In this regard the present study was carried out to assess the impact of different land uses on carbon storage and mineralization in soils of selected agro-ecological units of south Kerala. The agro-ecological units (AEUs) of south Kerala namely, southern coastal plain (AEU 1), Onattukara sandy plain (AEU 3), southern laterites (AEU 8), south central laterites (AEU 9) and southern and central foothills (AEU 12) were selected. In each AEU, different agricultural land use categories as described by IPCC for the carbon inventory such as, garden land (coconut), wet land (rice), fallow (uncultivated) and plantation (rubber) were also selected for the study. The potential for carbon mineralization in soil was assessed by evaluating factors such as Total organic carbon, mineralizable carbon, particulate organic carbon, global warming potential, carbon distribution, and turnover. The total organic carbon and particulate organic carbon were higher in rubber land use and AEU 12. The mineralizable C content in soil varied from 1.40 to 3.45, 1.18 to 3.41, 1.04 to 3.04, 1.23 to 3.35 and 1.01 to 3.02 mg g⁻¹ in AEU 1, 3, 8, 9 and 12 respectively. The highest value was observed from AEU 1 and rice land use. Similar trend was obtained for global warming potential of soils based on CO₂ evolution which varied from 31.82 to 78.41, 26.89 to 79.02, 22.89 to 69.02, 28.03 to 76.06 and 22.96 to 68.64 in AEU 1, 3, 8, 9 and 12 respectively. The C proportion and turnover rates were in the range of 0.25 to 0.77 and 0.04 to 0.17 respectively. The C proportion was the highest in AEU 12 and rubber land use whereas the C turnover was the highest in AEU 1 and rice land use.

Keywords- Land use, agro-ecological units, potential carbon mineralization, global warming potential, carbon proportion, carbon turnover

1. Introduction

Soil serves as a significant repository of carbon within terrestrial ecosystems and holds a pivotal position in the overall global carbon cycle. Approximately 2500 Pg of carbon is stored in the soil, composed of 1550 Pg as organic carbon (SOC) and 950 Pg as inorganic carbon (SIC) within the top 1 meter of the soil profile [1]. According to a recent global assessment, soils within the 0-30 cm

depth hold an estimated 680 Pg of SOC stock, playing a crucial role in soil biodiversity, nutrient regulation, management of greenhouse gases, and maintaining water balance [2]. The SOC plays a major role in improving soil fertility, increasing cation exchange capacity of soil, enhancing water holding capacity, promoting soil aggregation and maintaining soil flora and fauna. It helps to increase the diversity of various soil microbes, which play an important role in maintaining soil quality and functionality [3].

Increasing atmospheric concentration of greenhouse gases especially CO₂ is an important issue of the twenty-first century. Atmospheric CO₂ concentration has reached 400 ppm in 2013 [4]. Soil carbon sequestration is a promising strategy for the mitigation of global climate change by preventing building up of atmospheric CO₂. Any attempt to enrich the soil reservoir through sequestration of C will help to offset atmospheric CO₂ concentration and helps to abate global warming to a greater extent. Changes in land use and land cover significantly impact the SOC pool. This is primarily due to variations in the rates of input (such as plant litter) and output (like SOC mineralization) of soil organic matter (SOM) resulting from alterations in plant communities and land management practices [5]. Improper land use practices can result in the depletion of SOC, contributing to a decline in soil quality. Additionally, this depletion can lead to the release of carbon emissions into the atmosphere. A global assessment of SOC revealed a total loss of 116 Pg of carbon within the 2 meter depth of soil due to agricultural land use changes spanning 1200 years [6]. Land use management practices affects the rate of soil C mineralization [7]. Changes in land use have a significant impact on global CO₂ emission and land use changes along with the management practice are regarded as the major source of GHG emission after fossil fuels [8].

Agro-ecological units (AEUs) are broad spatial divisions based on climate variability, land forms and soils. The FAO introduced agro-ecological delineations, emphasizing similar agro-climatic parameters, to identify agriculturally promising areas ideal for specific crops or combinations thereof. This approach aims to maximize production potential for optimal agricultural outcomes. AEUs with different land uses have typical signature on soil organic carbon pools and its storage in different soil layers with unique recalcitrant, lability and carbon management index. There are Twenty- three AEUs in Kerala. In this context the present study was proposed to assess the impact of different land uses on carbon mineralization potential in soils of selected AEUs of south Kerala. The main objectives of the study were to determine the mineralizable C content, global warming potential, total organic carbon, particulate organic carbon, carbon turnover and carbon proportion in soils of selected AEUs under different land use categories in order to determine the source/ sink capacity of the selected land uses and their vulnerability status.

2. MATERIAL AND METHODS

2.1.1. Study area and soil sampling

The agro-ecological units (AEUs) of south Kerala namely, southern coastal plain (AEU 1), Onattukara sandy plain (AEU 3), southern laterites (AEU 8), south central laterites (AEU 9) and southern and central foothills (AEU 12) were selected. In each AEU, different agricultural land use categories as described by IPCC for the carbon inventory such as, garden land (coconut), wet land (rice), fallow (uncultivated) and plantation (rubber) were also selected for the study. Survey was

conducted in the selected AEU and land uses, and identified locations for soil sampling. Three sites were selected randomly from each land use representing each AEU and considered as replications for the study. Soil samples were collected from various locations including Thuravoor, Nedumpaikulam, Kundra and Magalapuram in AEU 1, Bharanikkavu, Mynagappalli and Karunagappali in AEU 3, Ookkod, Kalliyoor, Venganoor, Balaramapuram and Perumkadavila in AEU 8, Karavaram, Pooyappalli and Madappalli in AEU 9, and Karyom, Thalavoor, Altharamoodu, Kuttikkadu, and Vellarada in AEU 12. Geo referenced soil samples were collected from 0 to 25 cm depth and the samples were stored in plastic covers and labelled. Soil sampling was done during February to March, 2020 and November, 2020 to February, 2021. The soil samples were shade dried, powdered, sieved through a 2 mm sieve and stored in a moisture free environment for laboratory analysis. The location map of sampling sites in the AEU selected for the study is depicted in figure 1.




Fig. 1 Location map of study area and sampling points

2.1.2 Soil Analysis

2.1.2.1 Total organic carbon

Total organic carbon (TOC) content in the samples was determined by weight loss on ignition method [9].

2.1.2.2 Particulate organic carbon

The assessment of particulate organic carbon (POC) utilized the sodium hexametaphosphate

dissolution method [10]. This involved dispersing ten grams of air-dried soil in 30 ml of a 0.5% sodium hexametaphosphate solution, agitating the mixture for 15 hours on a reciprocal shaker. Subsequently, the dispersed soil sample was filtered through a 53 µm sieve. The retained material on the sieve underwent multiple rinses with water, after which it was collected, dried at 50°C in a hot air oven, and analyzed for carbon content using the wet oxidation method [11].

2.1.2.3 Mineralizable carbon

Mineralizable carbon was determined by CO₂ evolution method following laboratory incubation study as outlined by Ladd et al., [12]. Hundred gram soil sample was taken in a conical flask. The moisture content of the sample was maintained at field capacity throughout the incubation period. The CO₂ evolved were trapped in vials containing 0.1 N NaOH, which were hung inside the conical flask using a thread and the flask was sealed tightly using a rubber stopper to prevent any CO₂ loss. The CO₂ absorbed by NaOH was precipitated using BaCl₂ and estimated by titrating against 0.1 N HCl using phenolphthalein as indicator. The CO₂ measurement was made on daily basis during the initial period and the interval was fixed based on the evolution and the subsequent measurements were continued up to 50 days depending on the amount of CO₂ evolved.

2.1.2.4 Global warming potential

The global warming potential (GWP) of soils under various land uses was assessed by considering the amount of CO₂ released during the incubation experiment [12]. According to IPCC [13], GWP of 1 µmol CO₂ was assumed to be 1 and GWP of selected land use was calculated as described by Cai [14]. The GWP value is unit-less.

$$\text{GWP}(\text{CO}_2) = [m(\text{CO}_2)/44] \times 1, \text{ where 'm' is the mass of the gas}$$

2.1.2.5 Carbon proportion and turnover

Carbon proportion and turnover was worked out by the method given by Chacko et al., [15]. The carbon proportion was computed by the ratio of particulate organic carbon (POC) to total organic carbon (TOC). Carbon turnover was computed by the ratio of C mineralization to TOC.

2.1.3 Statistical Analysis

The data generated were subjected to statistical analysis using GRAPES software [16]. The statistical design followed was two way ANOVA.

3. Results and discussion

3.1 Total organic carbon

The primary elements influencing the dynamics of TOC in soil encompass land uses, climatic conditions, soil types, vegetation and management practices. In the present investigation, the TOC content varied from 1.27 to 3.56 per cent in different AEUs under various land use systems (Table 1). Among the different AEUs, TOC content followed the order AEU 9 (2.53 %)> AEU 12 (2.46 %)> AEU 8 (2.45 %)> AEU 3 (2.24 %)> AEU 1 (2.09 %) and in land uses it varied in the order rubber (3.18 %)> coconut (2.56 %)> rice (2.16 %)> uncultivated land (1.52 %). The TOC build up in soil under any land uses or soil type is the balance of the carbon input (litter, crop residue, root exudation & manure) and output (respiration, oxidation, leaching & erosion losses). In all the AEUs higher TOC was registered from rubber land use. The elevated organic carbon levels found in rubber plantations in this study might be attributed to high levels of organic residue

input in the form of leaf litter, root biomass and exudation, and its recalcitrant nature that slows down the rate of microbial decomposition. The results are in conformity with the findings of Mandal et al., [17]. Lesser TOC content in other land use types implies the considerable depletion of SOC through the nature of crops and its management practices. This happens particularly in areas subjected to intensive cultivation, leading to disruptions in soil structure and increased exposure of soil carbon fractions to oxidation. Similar results were obtained by Hussain et al., [18] who reported that the absence of organic matter integration into the soil, the removal of crop residues, and the use of traditional tillage techniques in agricultural production systems can result in a reduction of carbon content in the soil and tillage accelerates the oxidation of organic carbon by disrupting the soil structure, thereby exposing the organic matter to soil microorganisms.

Table 1. Effect of agricultural land uses on total organic carbon (%) content in soil

| AEUs (A) | Land use (L) | | | | Mean |
|--------------|---------------------|--------------------|--------------------|--------------------|-------------------|
| | Coconut | Rice | Rubber | Uncultivated land | |
| AEU 1 | 2.32 ^{fg} | 1.98 ^h | 2.80 ^{cd} | 1.27 ⁱ | 2.09 ^C |
| AEU 3 | 2.54 ^{def} | 1.97 ^h | 2.89 ^c | 1.56 ⁱ | 2.24 ^B |
| AEU 8 | 2.64 ^{cde} | 2.23 ^{gh} | 3.46 ^a | 1.47 ^{ij} | 2.45 ^A |
| AEU 9 | 2.79 ^{cd} | 2.51 ^{ef} | 3.18 ^b | 1.63 ⁱ | 2.53 ^A |
| AEU 12 | 2.51 ^{ef} | 2.08 ^{gh} | 3.56 ^a | 1.68 ⁱ | 2.46 ^A |
| Mean | 2.56 ^B | 2.16 ^C | 3.18 ^A | 1.52 ^D | |
| | A | L | A x L | | |
| S.E(m) | 0.046 | 0.041 | 0.093 | | |
| CD (P= 0.05) | 0.132 | 0.118 | 0.265 | | |

*Interaction values represented by same lower case superscript letters are not significantly different

* Mean values represented by same upper case superscript letters are not significantly different

3.2 Particulate organic carbon

Particulate organic carbon (POC) is a stable C fraction as it is physically protected within the aggregates and is the most contributing factor to TOC content in soil. The POC fraction varied significantly among different systems and is depicted in figure 2. Among the land uses rubber land use recorded the highest POC (2.07 %) which was found to be significantly superior to coconut (1.57 %) and rice (1.13 %) whereas uncultivated land recorded the lowest POC (0.61 %). With respect to different AEUs, the highest value was observed for AEU 12 (1.74 %) followed by AEU 9 (1.66 %), AEU 8 (1.33 %) and AEU 3 (1.24 %) which were differed significantly and the lowest value was recorded for AEU 1 (0.75 %). It is also revealed that in all AEUs rubber land use recorded higher POC compared to other land uses. The high POC value under rubber land use might be due to high biomass input and minimum soil disturbances and soil aggregation in the land use. Land management practices involving minimal soil disturbance can result in POC accumulation whereas intense cultivation practices reduce the concentration particularly in the surface soil layers. Hence POC is a sensitive indicator of land use management practices in soil. Also lignins and tannins from

decomposition of litter from rubber plantations enhances soil aggregation and POC is physically protected within the aggregates. The results are in line with the findings of Kalambukattu et al., [19].




Fig. 2. Effect of agricultural land uses on particulate organic carbon (%) in soil

3.3 Mineralizable carbon

The results of cumulative amount of C mineralized for 50 days of incubation in soils of different agro-ecological units under different land use systems is given in table 2. The mineralizable C content varied between 1.40 to 3.45 mg g⁻¹, 1.18 to 3.41 mg g⁻¹, 1.04 to 3.04 mg g⁻¹, 1.23 to 3.35 mg g⁻¹ and 1.01 to 3.02 mg g⁻¹ in AEU 1, 3, 8, 9 and 12. The higher amount of mineralizable carbon in AEU 1 might be attributed to the sandy nature of the soil which possess low sequestration capacity for carbon that augmented mineralization of carbon. AEU 12 recorded the lowest mineralizable carbon content which enhanced the SOC storage in the system. While comparing the mean value of cumulative mineralizable carbon in different land uses, the highest value was observed in rice land use (3.25 mg g⁻¹), which was found to be significantly different from coconut (2.36 mg g⁻¹) and rubber (1.74 mg g⁻¹) and the lowest value registered in uncultivated land (1.17 mg g⁻¹). The increased carbon mineralization observed in rice land use may be linked to the extensive tillage practices that disrupt soil aggregation and promote faster mineralization as reported by (15 Chacko et al. (2014). Among the land uses uncultivated land showed lowest mineralizable carbon which could be due to reduced soil disturbance that lowers carbon mineralization. The results are similar to the findings of Gladis et al., [20].

Table 2. Cumulative mineralizable carbon (mg g⁻¹) in soils of different AEU_s under various agricultural land use systems for 50 days

| AEUs (A) | Land use (L) | | | | Mean |
|-----------|--------------------|-------------------|-------------------|-------------------|-------------------|
| | Coconut | Rice | Rubber | Uncultivated land | |
| AEU 1 | 2.61 ^c | 3.45 ^a | 1.98 ^f | 1.40 ⁱ | 2.36 ^A |
| AEU 3 | 2.67 ^c | 3.41 ^a | 1.98 ^f | 1.18 ^j | 2.31 ^A |
| AEU 8 | 2.32 ^d | 3.04 ^b | 1.41 ⁱ | 1.04 ^k | 1.95 ^C |
| AEU 9 | 2.16 ^e | 3.35 ^a | 1.73 ^g | 1.23 ^j | 2.12 ^B |
| AEU 12 | 2.06 ^{ef} | 3.02 ^b | 1.58 ^h | 1.01 ^k | 1.92 ^C |
| Mean | 2.36 ^B | 3.25 ^A | 1.74 ^C | 1.17 ^D | |
| | A | L | A x L | | |
| S.E(m) | 0.023 | 0.020 | 0.045 | | |
| CD (0.05) | 0.065 | 0.058 | 0.130 | | |

*Interaction values represented by same lower case superscript letters are not significantly different

* Mean values represented by same upper case superscript letters are not significantly different

3.4 Global warming potential

Soil carbon mineralization directly impacts the potential release of CO₂, which, in turn, contributes to and influences global warming. Different land uses have varying ability to store and release carbon. The CO₂ production and GWP of soils of various AEU_s and land uses followed the same trend of mineralizable carbon. The GWP of various AEU_s ranged in the descending order AEU 1 > AEU 3 > AEU 9 > AEU 8 > AEU 12 and land uses in the order rice > coconut > rubber > uncultivated land (Figure 3). In all the AEU_s, the highest value of GWP was noticed in rice which is attributed to the higher carbon dioxide evolution from soil due to the rapid mineralization of organic matter facilitated by intense tillage practices that contributes to the higher GWP (74.03). The results are in agreement with the findings of Chacko et al. [15] and Sreekanth et al., [21]




Fig.3 Global warming potential of soils in different AEU under various agricultural land use systems

3.5 Carbon proportion

The ratio of particulate organic carbon to total organic carbon (POC/TOC) represents carbon proportion in soil. The sink capacity of soil for carbon storage can be represented by soil carbon proportion (Fig. 4). Highest POC/TOC was observed in AEU 12 (0.70) and rubber land use (0.64) which indicate the potential of these systems to store carbon and act as C sink. The physiography, climatic conditions, vegetation, soil type and management practices might have contributed to the higher carbon proportion in the systems [15,20].




Fig.4 Carbon proportion in soils of different AEU under various agricultural land use systems

3.6 Carbon turnover

The mineralizable carbon content in TOC (mineralizable C/TOC) represents the carbon turnover in soil. The source capacity can be represented by potential carbon mineralization and carbon turnover rate (Fig.5). Highest C turnover was observed in AEU 1 (0.12) and rice land use (0.15) which might be due to more carbon mineralization prevailing in the systems. The highest C turnover indicates the vulnerability to release C from the soil indicating it as a potential C source as reported by Chacko et al., [15]. Intensive puddling and tillage methods employed in rice cultivation result in the disruption of soil structure, ultimately causing decomposition and mineralization of organic matter in the soil. This makes the land use system more vulnerable as a potential carbon source. The results were comparable with the findings of Paramesh et al., [22].




Fig.5 Carbon turnover in soils of different AEU under various agricultural land use systems

4. Conclusion

Higher levels of TOC, POC, and carbon proportion were notably found in rubber cultivation areas, particularly within AEU 12. This suggests a conducive environment for accumulating organic carbon in the soil. Consequently, it hints at the potential for soil in these systems to act as a significant carbon sink. Among the different land uses, the highest values for mineralizable carbon, GWP and carbon turnover were recorded in the rice land use specifically in AEU 1. The highest carbon turnover signifies the susceptibility of the soil to release carbon, indicating its potential as a carbon source. In rice cultivation, the use of intensive puddling and tillage methods disrupts soil structure, leading to the decomposition and mineralization of organic matter. This renders the land use system more vulnerable as a potential carbon source of atmospheric carbon dioxide. The study underscores the significance of land uses and management practices in storing soil carbon and reducing the accumulation of CO₂ in the atmosphere.

References

- 1). Lal R. Soil carbon sequestration impacts on global climate change and food security. *Science*. 2004; 304(5677):1623-1627.
- 2). Ahirwal J, Gogoi A, Sahoo, UK. Stability of soil organic carbon pools affected by land use and land cover changes in forests of eastern Himalayan region, India. *Catena*. 2022; 215:106308.
- 3). McBratney AB, Stockmann U, Angers DA, Minasny B, Field DJ. Challenges for soil organic carbon research. *Soil carbon*. 2014; 3-16.
- 4). Lal R, Negassa W, Lorenz K. Carbon sequestration in soil. *Current Opinion in Environmental Sustainability*. 2015; 15:79-86.
- 5). Poeplau C, Don A. Sensitivity of soil organic carbon stocks and fractions to different land-use changes across Europe. *Geoderma*; 2013; 192:189-201.
- 6). Sanderman J, Hengl T, Fiske, GJ. Soil carbon debt of 12,000 years of human land use. *Proceedings of the National Academy of Sciences*. 2017; 114(36):9575-9580.
- 7). Ren C, Wang T, Xu Y, Deng J, Zhao F, Yang G, Han X, Feng Y, Ren G. Differential soil microbial community responses to the linkage of soil organic carbon fractions with respiration across land-use changes. *Forest Ecology and Management*. 2018; 409:170-178.
- 8). Zhong Q, Wang K, Nie M, Zhang G, Zhang W, Zhu Y, Fu Y, Zhang Q, Gao Y. Responses of wetland soil carbon and nutrient pools and microbial activities after 7 years of experimental warming in the Yangtze Estuary. *Ecological Engineering*. 2019; 136:68-78.
- 9). FAI (Fertilizer Association of India), The Fertilizer (Control) Order, 1985. The Fertilizer Association of India, New Delhi; 2017. Available: <http://www.astaspice.org/food-safety/astaspice-analytical-methods-manual>
- 10). Cambardella CA, Elliott ET. Particulate soil organic-matter changes across a grassland cultivation sequence. *Soil science society of America journal*. 1992; 56(3):777-783.
- 11). Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science*. 1934; 37(1): 29-38.
- 12). Ladd JN, Amato M, Grace PR, Van Veen JA. Simulation of ¹⁴C turnover through the microbial biomass in soils incubated with ¹⁴C-labelled plant residues. *Soil Biology and Biochemistry*. 1995; 27(6):777-783.
- 13). IPCC (Intergovernmental Panel on Climate Change). *Climate Change 2001: The Scientific Basis*. Cambridge University Press, New York, USA; 2001.
- 14). Cai Z. Effect of water regime on CO₂, CH₄, and NO₂ emissions and overall potential for greenhouse effect caused by emitted gases. *Acta Pedol. Sinica*. 1999; 36 (4):484-491.

- 15). Chacko FM, Sreekanth NP, Shanthi PV, Padmakumar B, Thomas AP. Soil carbon dynamics and global warming potential of selected soil series and landuse categories. *Octa Journal of Environmental Research*. 2014; 2(1).
- 16). Gopinath PP, Parsad R, Joseph B, Adarsh VS, GRAPES: General R shiny Based Analysis Platform Empowered by Statistics. 2020.
- 17). Mandal A, Toor AS, Dhaliwa, SS. Assessment of sequestered organic carbon and its pools under different agricultural land-uses in the semi-arid soils of south-western Punjab, India. *Journal of Soil Science and Plant Nutrition*. 2020;20:259-273.
- 18). Hussain S, Sharma V, Arya VM, Sharma KR, Rao CS. Total organic and inorganic carbon in soils under different land use/land cover systems in the foothill Himalayas. *Catena*. 2019; 182:104104.
- 19). Kalambukattu JG, Singh R, Patra AK, Arunkumar K. Soil carbon pools and carbon management index under different land use systems in the Central Himalayan region. *Acta Agriculturae Scandinavica, Section B–Soil & Plant Science*. 2013; 63(3):200-205.
- 20). Gladis R, Dhanya KR, Joseph B, Aparna B, Rehana MR. Soil Carbon Stock and Pools in Acid Sulphate Soils of Kerala. *Current Journal of Applied Science and Technology*. 2020; 39(21):135-147.
- 21). Sreekanth NP, Shanthi P, Babu P, Thomas AP. Effect of land use conversion on soil carbon storage in a tropical grassland. *Annals of Environmental Science*, 2013; 7:101-112.
- 22). Paramesh V, Singh SK, Mohekar DS, Arunachalam V, Misra, SD, Jat SL, Kumar P, Nath AJ, Kumar N, Mahajan GR, Bhagat T. Impact of sustainable land-use management practices on soil carbon storage and soil quality in Goa State, India. *Land Degradation & Development*, 2022; 33(1):28-40.