

Review Article

Role of salicylic acid in mitigating stress and improving productivity of crops

Abstract

Salicylic acid (SA) is a pivotal plant hormone known for its significant contributions to crop stress mitigation and productivity enhancement. This review provides an in-depth analysis of SA's role in abiotic and biotic stress tolerance, shedding light on its diverse mechanisms of action. SA's positive impact on crop yield is explored, offering promising prospects for sustainable agriculture. The molecular pathways and biochemical underpinnings of SA-mediated stress alleviation are examined, offering insights into the intricate networks governing these processes. Additionally, the review discusses practical applications and potential challenges associated with integrating SA into agricultural practices. In a rapidly changing agricultural landscape, understanding the potential of SA to optimize crop productivity and combat stressors is paramount. The review underscores SA's promise as a key player in sustainable agriculture and outlines future research directions. This abstract provides a concise overview of the key points covered in your review article on the role of salicylic acid in crop stress mitigation and productivity enhancement.

Keywords: salicylic acid, stress, crops, molecular pathways, biochemical parameters

1. Introduction

Agriculture faces the formidable challenge of ensuring food security and sustainability in a world marked by a growing population and changing climatic conditions. The ability to mitigate the impact of abiotic and biotic stressors on crop productivity has become a critical pursuit in contemporary agriculture. Among the multifaceted strategies employed, the role of salicylic acid (SA) has emerged as a subject of increasing importance. SA, a naturally occurring plant hormone, has gained recognition for its pivotal contributions to enhancing stress tolerance and productivity in crops.

The significance of SA in plant physiology lies in its capacity to orchestrate a myriad of responses to environmental stress. Abiotic stress factors, including drought, salinity,

temperature extremes, and heavy metal toxicity, have been known to exact a toll on crop yields. Similarly, biotic stressors such as pathogens and pests pose substantial threats to agriculture. It is in the context of these adversities that SA has demonstrated its remarkable potential.

This comprehensive review seeks to explore the multifaceted dimensions of SA's involvement in mitigating stress and improving crop productivity. We delve into the molecular and biochemical mechanisms that underlie SA's ability to enhance stress tolerance, shedding light on its influence on gene expression, antioxidant defense, and osmotic regulation. Moreover, the interaction between SA and the complex network of biotic stress responses is a subject of in-depth investigation, revealing its capacity to enhance disease resistance.

As we embark on this exploration, we aim to provide a detailed account of SA's role in crop management, considering its practical applications and potential challenges. The agricultural sector's continual evolution demands a thorough understanding of SA's potential to optimize productivity and combat the detrimental effects of stress. This review not only elucidates SA's current role but also identifies future research directions and potential impediments in the translation of SA-based strategies into real-world agricultural practices. In doing so, we hope to contribute to the growing body of knowledge that can inform sustainable and resilient agricultural systems for a changing world.

2. Salicylic Acid: A Plant Hormone

Salicylic acid (SA) stands as a fundamental player in the intricate signaling network of plants. As a naturally occurring phenolic compound, SA functions as a vital plant hormone, orchestrating various physiological responses critical for plant growth, development, and adaptation to environmental challenges. It is a low-molecular-weight compound derived from the phenylpropanoid pathway and is found in varying concentrations within different plant species.

2.1 Biosynthesis and Distribution SA biosynthesis involves the conversion of chorismic acid, a precursor in the shikimate pathway, into SA via a series of enzymatic reactions. These reactions take place primarily in the chloroplasts and mitochondria. Notably, SA production

is often stimulated by stress conditions, both abiotic and biotic, as part of the plant's defense mechanisms.

SA is not uniformly distributed throughout the plant but accumulates in specific tissues and cellular compartments. Its distribution is responsive to developmental stages and environmental cues. Understanding the spatial and temporal distribution of SA is crucial in elucidating its multifaceted roles in stress response and defense.

2.2 Salicylic Acid Signaling Pathways The biological functions of SA are mediated through intricate signaling pathways. Key components of SA signaling include Nonexpressor of Pathogenesis-Related genes 1 (NPR1), a transcriptional coactivator that plays a central role in regulating the expression of genes associated with defense responses. NPR1 is maintained in an oligomeric state in the cytoplasm under non-stress conditions but translocates to the nucleus upon SA induction to activate defense-related gene expression.

In addition to NPR1, SA interacts with other components of the signaling cascade, including mitogen-activated protein kinases (MAPKs) and WRKY transcription factors. The crosstalk between SA and other phytohormones, such as jasmonic acid (JA) and ethylene (ET), further modulates plant responses to stress and pathogens. Understanding the interactions and interplay between these signaling molecules is essential in comprehending the complexity of SA's role in plant defense.

2.3 Role in Systemic Acquired Resistance (SAR) One of the most remarkable features of SA is its ability to induce systemic acquired resistance (SAR), a heightened state of defense that protects not only the locally affected tissue but the entire plant. SA acts as a priming agent, enabling the plant to respond more effectively to subsequent pathogen attacks. The establishment of SAR is crucial in the context of biotic stress management in crops.

In summary, SA is a pivotal plant hormone with a remarkable capacity to influence plant responses to stress and pathogens. Its biosynthesis, distribution, and signaling pathways, including its role in SAR, make it a subject of great interest in the context of crop stress mitigation and productivity enhancement. Understanding the intricacies of SA's functions lays the foundation for exploring its applications in agriculture and improving the resilience of crops to environmental challenges.

2. Salicylic Acid: A Plant Hormone

Salicylic acid (SA) stands as a fundamental player in the intricate signaling network of plants. As a naturally occurring phenolic compound, SA functions as a vital plant hormone, orchestrating various physiological responses critical for plant growth, development, and adaptation to environmental challenges. It is a low-molecular-weight compound derived from the phenylpropanoid pathway and is found in varying concentrations within different plant species.

2.1 Biosynthesis and Distribution SA biosynthesis involves the conversion of chorismic acid, a precursor in the shikimate pathway, into SA via a series of enzymatic reactions. These reactions take place primarily in the chloroplasts and mitochondria. Notably, SA production is often stimulated by stress conditions, both abiotic and biotic, as part of the plant's defense mechanisms.

SA is not uniformly distributed throughout the plant but accumulates in specific tissues and cellular compartments. Its distribution is responsive to developmental stages and environmental cues. Understanding the spatial and temporal distribution of SA is crucial in elucidating its multifaceted roles in stress response and defense.

2.2 Salicylic Acid Signaling Pathways The biological functions of SA are mediated through intricate signaling pathways. Key components of SA signaling include Nonexpressor of Pathogenesis-Related genes 1 (NPR1), a transcriptional coactivator that plays a central role in regulating the expression of genes associated with defense responses. NPR1 is maintained in an oligomeric state in the cytoplasm under non-stress conditions but translocates to the nucleus upon SA induction to activate defense-related gene expression.

In addition to NPR1, SA interacts with other components of the signaling cascade, including mitogen-activated protein kinases (MAPKs) and WRKY transcription factors. The crosstalk between SA and other phytohormones, such as jasmonic acid (JA) and ethylene (ET), further modulates plant responses to stress and pathogens. Understanding the interactions and interplay between these signaling molecules is essential in comprehending the complexity of SA's role in plant defense.

2.3 Role in Systemic Acquired Resistance (SAR) One of the most remarkable features of SA is its ability to induce systemic acquired resistance (SAR), a heightened state of defense that protects not only the locally affected tissue but the entire plant. SA acts as a priming agent, enabling the plant to respond more effectively to subsequent pathogen attacks. The establishment of SAR is crucial in the context of biotic stress management in crops.

In summary, SA is a pivotal plant hormone with a remarkable capacity to influence plant responses to stress and pathogens. Its biosynthesis, distribution, and signaling pathways, including its role in SAR, make it a subject of great interest in the context of crop stress mitigation and productivity enhancement. Understanding the intricacies of SA's functions lays the foundation for exploring its applications in agriculture and improving the resilience of crops to environmental challenges.

4. Role of Salicylic Acid in Abiotic Stress Tolerance

Salicylic acid (SA), a versatile plant hormone, has emerged as a key player in bolstering abiotic stress tolerance in crops. The multifaceted roles of SA encompass its ability to mitigate the detrimental effects of various stressors, including drought, salinity, extreme temperatures, and heavy metal toxicity. Understanding how SA influences stress tolerance mechanisms is pivotal in harnessing its potential for improving crop resilience.

4.1 Modulation of Gene Expression One of the primary ways SA enhances abiotic stress tolerance is by modulating gene expression. SA-induced changes in gene expression patterns influence the synthesis of stress-related proteins, enzymes, and osmoprotectants. This, in turn, helps plants cope with stress by maintaining cellular homeostasis and promoting stress adaptation.

4.2 Antioxidant Defense Abiotic stress often leads to the overproduction of reactive oxygen species (ROS) in plant cells, causing oxidative damage. SA acts as an inducer of antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD). These enzymes neutralize ROS, thus reducing oxidative stress and preserving cell integrity.

4.3 Osmotic Regulation In the face of abiotic stress, plants frequently experience water deficits and osmotic imbalances. SA plays a pivotal role in regulating osmotic balance by

controlling the synthesis of osmoprotectants like proline, sugars, and polyamines. These osmoprotectants contribute to maintaining turgor pressure and cellular hydration.

4.4 Enhancement of Photosynthesis Under abiotic stress conditions, photosynthesis is often hampered, leading to reduced carbon assimilation and decreased crop yields. SA has been shown to improve photosynthetic efficiency by enhancing chlorophyll content, maintaining stomatal conductance, and protecting photosynthetic machinery from stress-induced damage.

4.5 Hormonal Crosstalk The interplay between SA and other plant hormones, such as jasmonic acid (JA) and abscisic acid (ABA), is crucial for orchestrating the plant's response to stress. SA-JA crosstalk, in particular, plays a significant role in balancing defense responses with growth and development. SA's ability to modulate hormonal interactions contributes to stress tolerance in crops.

In summary, the role of SA in abiotic stress tolerance is underpinned by its influence on gene expression, antioxidant defense, osmotic regulation, photosynthesis, and hormonal crosstalk. The ability of SA to mediate these responses is a testament to its potential as a tool for enhancing crop resilience in the face of environmental challenges.

In the subsequent sections, we will delve into the specific mechanisms through which SA operates in response to various abiotic stressors and explore the implications of SA application in agriculture as a means of mitigating stress and improving crop productivity.

5. Salicylic Acid and Biotic Stress Responses

While the role of salicylic acid (SA) in abiotic stress tolerance is significant, its involvement in biotic stress responses is equally noteworthy. SA is a central player in the plant's defense mechanisms against various pathogens and pests. Understanding how SA interacts with the plant's immune system and influences disease resistance is essential for comprehending its potential in agriculture.

5.1 Plant Defense Against Pathogens Pathogens, such as fungi, bacteria, viruses, and oomycetes, pose a constant threat to crop health and productivity. Plants have evolved a sophisticated defense system to ward off these invaders. SA is a critical component of the plant's immune response, particularly against biotrophic and hemibiotrophic pathogens.

5.2 Induction of Systemic Acquired Resistance (SAR) One of the hallmark features of SA's role in plant defense is the induction of systemic acquired resistance (SAR). When a plant is exposed to a pathogen, SA is synthesized locally at the infection site and is transported systemically throughout the plant. This "priming" of the plant's immune system enables it to respond more robustly to future pathogen attacks, even in distant tissues.

5.3 Activation of Defense Genes SA plays a key role in the activation of defense-related genes. Upon pathogen recognition, SA induces the expression of pathogenesis-related (PR) proteins, antimicrobial compounds, and cell wall reinforcement proteins. These gene products collectively contribute to the plant's defense mechanisms, inhibiting pathogen proliferation.

5.4 Crosstalk with Other Defense Pathways In addition to its direct influence on pathogen defense, SA engages in complex crosstalk with other defense pathways. The interaction between SA and jasmonic acid (JA), a hormone associated with defense against herbivores and necrotrophic pathogens, is a prominent example. The delicate balance between SA and JA signaling determines the plant's ability to defend against different types of attackers while minimizing negative trade-offs.

5.5 Role in Disease Resistance The pivotal role of SA in enhancing disease resistance in crops has been demonstrated in numerous studies. SA-mediated resistance is particularly effective against diseases caused by biotrophic and hemibiotrophic pathogens, such as rusts, mildews, and certain bacterial pathogens. The application of SA or SA analogs has shown promise in protecting crops against these diseases.

In summary, SA is a central component of the plant's defense system against pathogens. Its induction of SAR, activation of defense genes, and crosstalk with other defense pathways collectively contribute to disease resistance in plants. The implications of SA's role in biotic stress responses extend to crop protection and the development of sustainable disease management strategies in agriculture.

The subsequent sections will delve into the specific mechanisms through which SA operates in response to different biotic stressors and explore the potential applications of SA in crop protection and pathogen management.

6. Mechanisms of Salicylic Acid-Mediated Stress Mitigation

Salicylic acid (SA) exerts its role in mitigating both abiotic and biotic stress through a complex array of molecular and biochemical mechanisms. Understanding these mechanisms is essential for appreciating the diverse ways in which SA enhances stress tolerance in crops and improves their overall productivity.

6.1 Signaling Pathways At the heart of SA-mediated stress mitigation lies its interaction with signaling pathways. SA acts as a pivotal signal molecule, initiating a cascade of responses that enable plants to withstand environmental stress. These pathways include SA-specific signaling, the mitogen-activated protein kinase (MAPK) cascade, and hormonal crosstalk.

6.2 Gene Expression Modulation One of the fundamental mechanisms through which SA enhances stress tolerance is the modulation of gene expression. SA triggers the expression of stress-related genes, which results in the production of stress-responsive proteins and enzymes. These gene products play a crucial role in stress adaptation and the defense against pathogens.

6.3 Antioxidant Defense Under stressful conditions, plants often accumulate reactive oxygen species (ROS), leading to oxidative stress. SA helps counteract this by stimulating the synthesis and activation of antioxidant enzymes. Superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD) are among the enzymes that neutralize ROS and reduce oxidative damage.

6.4 Osmotic Regulation Maintaining water balance and cellular turgor pressure is essential for coping with abiotic stress. SA contributes to osmotic regulation by promoting the synthesis of osmoprotectants like proline, sugars, and polyamines. These compounds help to retain water within plant cells and prevent dehydration.

6.5 Defense Pathways In the context of biotic stress, SA activates defense pathways that culminate in the production of antimicrobial compounds, pathogenesis-related (PR) proteins, and other defense-related molecules. This armament is essential for preventing pathogen invasion and propagation.

6.6 Hormonal Crosstalk The intricate interplay between SA and other plant hormones, such as jasmonic acid (JA) and abscisic acid (ABA), is critical in balancing the trade-offs between

growth, development, and stress responses. Hormonal crosstalk ensures that the plant can adapt to different stressors without compromising its overall fitness.

In summary, SA-mediated stress mitigation involves signaling pathways, gene expression modulation, antioxidant defense, osmotic regulation, activation of defense pathways, and hormonal crosstalk. The synergy between these mechanisms enables SA to enhance the resilience of crops in the face of various stressors, ultimately contributing to improved productivity.

In the subsequent sections, we will delve into the specific applications of SA in agriculture and its potential as a tool for sustainable crop management, considering its multifaceted effects on stress tolerance.

7. Enhancing Crop Productivity with Salicylic Acid

The potential of salicylic acid (SA) to improve crop productivity has garnered increasing attention in the realm of sustainable agriculture. SA's multifaceted roles in mitigating both abiotic and biotic stress, as well as its influence on growth and development, make it a promising candidate for enhancing crop yields.

7.1 SA's Impact on Crop Yield Studies across a range of crop species have demonstrated the positive impact of SA on crop yield. SA-treated plants often exhibit increased biomass, larger fruit sizes, and higher grain or seed yields. These effects are particularly pronounced under stress conditions, where SA's stress-mitigating mechanisms come into play.

7.2 Stress Mitigation and Productivity The stress-mitigating effects of SA are closely linked to its role in enhancing crop productivity. By improving stress tolerance and minimizing the negative impact of abiotic stressors like drought and salinity, SA allows crops to maintain optimal growth and development. This, in turn, leads to increased yields even in adverse environments.

7.3 Disease Resistance and Quality Enhancement SA's influence on disease resistance contributes to improved crop quality and yield. Plants with enhanced disease resistance are less prone to pathogen-induced damage, ensuring that a higher proportion of the crop reaches

harvestable quality. Moreover, SA-treated crops may require fewer chemical inputs for disease management, reducing production costs.

7.4 Optimizing Photosynthesis SA's ability to enhance photosynthetic efficiency has a direct impact on crop productivity. Improved chlorophyll content, stomatal conductance, and protection of photosynthetic machinery under stress conditions result in increased carbon assimilation and higher crop yields.

7.5 Hormonal Balance The hormonal crosstalk regulated by SA also influences crop productivity. By maintaining a balance between SA, jasmonic acid (JA), and other hormones, SA promotes growth while simultaneously enhancing stress tolerance and defense mechanisms.

7.6 Practical Applications in Agriculture The practical application of SA in agriculture is a subject of increasing interest. SA can be applied through various methods, including foliar sprays, seed treatments, and soil drenches. The timing and concentration of SA application may vary depending on the crop, growth stage, and environmental conditions.

7.7 Challenges and Considerations While the potential benefits of SA in enhancing crop productivity are substantial, there are challenges and considerations to be addressed. These include potential phytotoxicity at high concentrations, variability in crop responses, and the need for standardized application protocols. Further research is necessary to optimize SA application methods and dosage for different crops and growing conditions.

In conclusion, salicylic acid holds significant promise as a tool for enhancing crop productivity. Its multifaceted effects on stress mitigation, disease resistance, and growth promotion make it a valuable asset in the pursuit of sustainable and resilient agriculture. The subsequent sections will explore the practical applications of SA in the field and the ongoing research directions aimed at maximizing its potential in crop management.

8. Application of Salicylic Acid in Agriculture

The practical application of salicylic acid (SA) in agriculture has gained attention as a means to enhance crop productivity and resilience in the face of environmental challenges.

Understanding the methods and considerations of SA application is essential for translating its potential into real-world agricultural practices.

8.1 Application Methods SA can be applied to crops through various methods, each with its advantages and considerations:

- **Foliar Sprays:** Foliar application is a common method of delivering SA to plants. SA solutions are sprayed directly onto the leaves, allowing for rapid absorption. This method is especially effective for providing immediate protection against diseases and mitigating stress.
- **Seed Treatments:** Treating seeds with SA solutions or coatings can help establish SA-mediated stress tolerance from the early stages of plant growth. This method is advantageous for promoting uniformity in plant responses and enhancing seedling vigor.
- **Soil Drenches:** Applying SA to the root zone through soil drenches is another option. This method enables the plant to absorb SA through the roots and distribute it systemically. It can be particularly useful for enhancing stress tolerance, especially in cases of soil-borne diseases or salinity stress.

8.2 Timing and Dosage The timing of SA application is critical and varies depending on the crop, growth stage, and the specific stressor. Early applications during critical growth stages, such as flowering and fruit set, can maximize the benefits of SA. Dosage is another crucial consideration, as excessive SA can lead to phytotoxicity, while inadequate amounts may not confer the desired effects.

8.3 Challenges and Considerations The practical application of SA in agriculture is not without challenges:

- **Phytotoxicity:** Excessive SA concentrations can have detrimental effects on plant growth, leading to phytotoxicity. Careful dosage and monitoring are essential to avoid such issues.
- **Crop Variability:** The response to SA varies among different crop species and even cultivars. Research is ongoing to determine optimal application protocols for specific crops.

- **Environmental Factors:** Environmental conditions, including temperature, humidity, and light, can influence the efficacy of SA application. Adapting application strategies to local conditions is crucial.
- **Regulatory and Safety Considerations:** Regulations and safety guidelines related to the use of SA in agriculture may vary by region. Compliance with local regulations and safety standards is essential.

8.4 Potential for Sustainable Agriculture The application of SA in agriculture holds the potential to contribute to sustainable and resilient farming practices. By enhancing crop stress tolerance, improving disease resistance, and optimizing growth and development, SA can aid in reducing the reliance on chemical inputs and fostering environmentally friendly and economically viable agriculture.

8.5 Future Directions Ongoing research is focused on fine-tuning SA application protocols for different crops and stressors. Additionally, efforts are being made to develop SA analogs with improved efficacy and reduced phytotoxicity. These advancements are expected to further expand the practical utility of SA in agriculture.

In conclusion, the practical application of salicylic acid in agriculture is a promising avenue for optimizing crop management. As research continues to refine application methods and dosage considerations, SA's role in promoting sustainable and resilient agriculture is likely to become increasingly prominent.

You can expand upon this section with specific examples, case studies, and data related to the practical application of SA in agriculture, taking into account the focus and depth of your review article.

Conclusion

The role of salicylic acid (SA) in mitigating stress and enhancing crop productivity presents a compelling narrative in the ever-evolving landscape of agriculture. This review has provided

an in-depth exploration of the multifaceted dimensions of SA's influence on plant responses to abiotic and biotic stress, as well as its capacity to improve crop yields.

Salicylic acid, as a central plant hormone, orchestrates a complex network of responses to environmental challenges. It plays a pivotal role in abiotic stress tolerance by modulating gene expression, bolstering antioxidant defense, regulating osmotic balance, and optimizing photosynthesis. These mechanisms collectively enable plants to maintain growth and productivity even under adverse conditions.

In the realm of biotic stress, SA's contribution to plant defense against pathogens is undeniable. It induces systemic acquired resistance (SAR), primes the plant's immune system, and activates defense genes, resulting in enhanced disease resistance and improved crop quality.

The mechanisms through which SA operates are intricately intertwined, involving signaling pathways, gene expression modulation, antioxidant defense, osmotic regulation, and hormonal crosstalk. This interplay ensures that SA not only aids in stress mitigation but also promotes growth and development, making it a valuable asset for crop management.

The practical application of SA in agriculture offers promising prospects. Through foliar sprays, seed treatments, or soil drenches, SA can be integrated into farming practices to enhance stress tolerance, disease resistance, and crop productivity. However, it is essential to carefully consider dosage, timing, and environmental factors to maximize the benefits while avoiding potential phytotoxicity.

Despite the considerable potential of SA in agriculture, challenges remain, including crop variability, regulatory considerations, and the need for optimized application strategies. Ongoing research endeavors aim to address these challenges and further refine the use of SA in farming practices.

As we navigate the complexities of modern agriculture, SA emerges as a valuable tool for sustainable and resilient crop management. Its ability to enhance stress tolerance, bolster disease resistance, and optimize growth while minimizing the reliance on chemical inputs holds promise for the future of farming.

In conclusion, the comprehensive understanding of SA's roles and practical applications presented in this review contributes to the growing body of knowledge that can inform and shape the future of agriculture. The journey towards sustainable and productive farming continues, and salicylic acid stands as a beacon of hope in this endeavor.

Salicylic acid (SA), a key signaling molecule in plants, has garnered substantial attention for its remarkable role in enhancing stress tolerance and crop productivity. In this comprehensive review, we delve into the multifaceted effects of SA on crops exposed to various abiotic and biotic stressors. Abiotic stress factors, such as drought, salinity, and extreme temperatures, significantly impact crop yields. The ability of SA to ameliorate these stressors through the modulation of gene expression, antioxidant defense, and osmotic regulation is examined in detail. Additionally, the interplay between SA and biotic stress responses, particularly in plant defense against pathogens, is explored, shedding light on its potential for enhancing disease resistance.

Understanding the molecular and biochemical mechanisms behind SA-mediated stress mitigation is crucial, and we provide insights into the intricate pathways that underline these processes. The review also highlights the influence of SA on yield enhancement across a range of crop species, demonstrating its potential as a valuable tool for sustainable agriculture. Practical applications of SA in agriculture are discussed, along with associated challenges and considerations.

As the agricultural landscape continues to evolve, exploring the potential of SA in crop management becomes increasingly relevant. We conclude by emphasizing the pivotal role of SA as a promising candidate for optimizing crop productivity and mitigating the detrimental effects of stress, while also identifying avenues for future research and potential challenges in translating SA-based strategies to real-world agricultural practices.

References

1. Khan, M. I. R., Fatma, M., Per, T. S., Anjum, N. A., & Khan, N. A. (2015). Salicylic acid- induced abiotic stress tolerance and underlying mechanisms in plants. *Frontiers in Plant Science*, 6, 462.
2. Miura, K., & Tada, Y. (2014). Regulation of water, salinity, and cold stress responses by salicylic acid. *Frontiers in Plant Science*, 5, 4.
3. Anjum, S. A., Tanveer, M., Hussain, S., Bao, M., Wang, L., Khan, I., & Tung, S. A. (2016). Salinity tolerance in plants: Physiological, biochemical and molecular regulation. In Anjum, S.A., Chan, M.T., & Umar, S. (Eds.), *Salinity Stress in Plants: Signalling, Omics and Adaptations* (pp. 33-69). Springer.
4. Mana, P. W., Wang-Bara, B., Mvondo, V. Y. E., Bourou, S., & Palai, O. (2023). Evaluation of the agronomic and technological performance of three new cotton varieties in the cotton zone of Cameroon. *Acta Botanica Plantae*, 2, 28-39.
5. Essra Ali Safdar, Nida Ali Safdar, Syeda Aifa Fathima, Pathan Amanullah Khan (2023). A Survey to Assess Knowledge Attitude Practice of People towards Vitamin D. *Acta Traditional Medicine*. V2i01, 27-34. DOI: <https://doi.org/10.5281/zenodo.8282626>
6. Vlot, A. C., Dempsey, D. A., & Klessig, D. F. (2009). Salicylic acid, a multifaceted hormone to combat disease. *Annual Review of Phytopathology*, 47, 177-206.
7. Zhang, L., Zhang, F., & Melotto, M. (2017). Salicylic acid represses the biosynthesis of ethylene and improves the resistance of pepper to *Phytophthora capsici*. *Journal of Phytopathology*, 165(6), 357-366.
8. Fatima, S., Nausheed, R., Hussain, S. M., Fatima, I., Begum, N., & Siddi-qua, R. (2023). Assessment of Soil Fertility Status of Mango Orchard at Vikarabad Farm house in Manneguda Village of Telangana State) *Acta Botanica Plantae*.
9. Yang, Y., Zhang, Y., Wang, M., Li, Y., Chen, T., & Sun, Y. (2017). The role of salicylic acid in the systemic acquired resistance induced in tomato by BABA. In Y. Li, R. M. Kelman, & M. Du (Eds.), *Advanced Research on Plant Lipids* (pp. 1-13). Springer.
10. Singh, A. K., Yadav, N., Singh, A., & Singh, A. (2023). Stay-green rice has greater drought resistance: one unique, functional SG Rice increases grain production in dry conditions. *Acta Botanica Plantae*. V02i02, 31, 38.
11. Nanda, R., Ahmed, F., & Sharma, R. NishaBhagat and Kewal Kumar (2022). Ethnobotanical Studies on Some Angiosperms of Tehsil Hiranagar of District Kathua (Jammu and Kashmir), India. *Acta Botanica Plantae*, 01-11.
12. Sabitha, N., Mohan Reddy, D., Lokanadha Reddy, D., Hemanth Kumar, M., Sudhakar, P., Ravindra Reddy, B., & Mallikarjuna, S. J. (2022). Genetic divergence analysis over seasons in single cross hybrids of maize (*Zea mays* L.). *Acta Botanica Plantae*, 1(2), 12-18.
13. Vallad, G. E., & Goodman, R. M. (2004). Systemic acquired resistance and induced systemic resistance in conventional agriculture. *Crop Science*, 44(6), 1920-1934.
14. Horváth, E., Szalai, G., & Janda, T. (2007). Induction of abiotic stress tolerance by salicylic acid signaling. *Journal of Plant Growth Regulation*, 26(3), 290-300.
15. Miura, K., & Tada, Y. (2014). Regulation of water, salinity, and cold stress responses by salicylic acid. *Frontiers in Plant Science*, 5, 4.
16. Khan, M. I. R., Fatma, M., Per, T. S., Anjum, N. A., & Khan, N. A. (2015). Salicylic acid- induced abiotic stress tolerance and underlying mechanisms in plants. *Frontiers in Plant Science*, 6, 462.
17. Ishikawa, S., & Shigeoka, S. (2008). Recent advances in ascorbate biosynthesis and the physiological significance of ascorbate peroxidase in photosynthesizing organisms. *Bioscience, Biotechnology, and Biochemistry*, 72(5), 1143-1154.
18. Verbruggen, N., & Hermans, C. (2008). Proline accumulation in plants: A review. *Amino Acids*, 35(4), 753-759.

19. Ogori, A. F., Eke, M. O., Girgih, T. A., & Abu, J. O. (2022). Influence of Aduwa (*Balanites aegyptiaca*. del) Meal Protein Enrichment on the Proximate, Phytochemical, Functional and Sensory Properties of Ogi. *Acta Botanica Plantae*. V01i03, 22-35.
20. Khan, W., Prithiviraj, B., & Smith, D. L. (2003). Photosynthetic responses of corn and soybean to foliar application of salicylates. *Journal of Plant Physiology*, 160(5), 485-492.
21. Borsani, O., Valpuesta, V., & Botella, M. A. (2001). Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in *Arabidopsis* seedlings. *Plant Physiology*, 126(3), 1024-1030.
22. Nweze, C. C., & Muhammad, B. Y. (2023). Wandoo Tseaa, Rahima Yunusa, Happy Abimiku Manasseh, Lateefat Bisola Adedipe, Eneh William Nebechukwu, Yakubu Atanyi Emmanuel (2023). Comparative Biochemical Effects of Natural and Synthetic Pesticides on Preserved *Phaseolus vulgaris* in Male Albino Rats. *Acta Botanica Plantae*. V02i01, 01-10.
23. Makela, P., Karkkainen, J., & Somersalo, S. (2000). Subcellular localization of peroxidase and the effect of salt stress in two cell lines of *Pisum sativum*. *Physiologia Plantarum*, 109(4), 386-394.
24. Rivas- San Vicente, M., & Plasencia, J. (2011). Salicylic acid beyond defence: its role in plant growth and development. *Journal of Experimental Botany*, 62(10), 3321-3338.
25. Pasternak, T. P., Potters, G., Caubergs, R. J., Jansen, M. A., & Coman, D. (2005). The effect of UV- A and UV- B radiation on the ascorbate and glutathione content of bean (*Phaseolus vulgaris* L.) leaves. *The Plant Cell & Environment*, 28(6), 853-863.