

Abstract

Fish consumption has become an essential phenomenon due to the increasing appreciation of its health and the economic benefits. The current study aims to identify the pattern of fish consumption and knowledge regarding nutritive values of fish in Benghazi city. As well as, it aims to analyse the heavy metals content of lead, cadmium and mercury of the top five most consumed fish's species in Benghazi. A cross-sectional study carried out 29th December 2022 and extended to 30th June 2023. A total of 300 Subjects randomly approached to participate in the study giving a response rate of 85.5%. Fresh fish's samples were collected from Bankina and analysed in term of heavy metals using Atomic Absorption Spectrophotometer. Sparus aurata, Pegellus bogaraveo, Sardina pilchardus, Mullus barbatus and Epinephelus marginatus are the most preferred consumed fishes in Benghazi. Age, female gender and number of family members are the variables that associated with fish consumption pattern in Benghazi. Lead, cadmium and mercury are the analysed heavy metals in the top five most consumed fishes in Benghazi. All heavy metals results are within the recommended level by (WHO). The results of the current study recommend that increasing fish consumption habit should be considered as a case of essential fact for having healthier generations. Furthermore, fish consumption should be increased via giving information by educational institutions, government agencies and non-governmental organizations.

Keywords: Benghazi, Consumption, Pattern, Fish, Heavy Metals

Introduction

Fishes are approximately 34,000 species of vertebrate animals found in the fresh and salt water globally. Fishes species range from the primitive jawless hagfishes and lampreys through the cartilaginous skates, sharks, rays to the ample and varied bony fishes. Most fish species are cold-blooded; however, the opah (*Lampris guttatus*), is warm-blooded. Pollution of water streams may affect aquatic animals, like fishes directly or indirectly. (1, 2) Metals are a group of elements with atomic weights ranging from 63.549 to 200,590 Dalton. Metals are characterized by a similar electronic distribution in the outer shell, such as copper, cadmium, and zinc. Metals can be divided into two groups: transition metals and metal-like elements. The transition metals include manganese, iron, copper, and cobalt. This group is essential for metabolism at low concentration levels. However, they can become toxic at higher concentrations. The metal-like elements include arsenic, cadmium, lead, mercury, selenium, and tin. This group is

not important for metabolic function. This group also becomes toxic at low concentration levels. (3) The term heavy metal refers to any metallic chemical element that has a relatively high density and is toxic or poisonous at low concentrations. Heavy metals include mercury (Hg), cadmium (Cd), arsenic (As), chromium (Cr), thallium (Tl), and lead (Pb). Heavy metals cannot be destroyed or degraded. They enter the humans' bodies via food, drinking water, and air. As trace elements, some heavy metals such as copper, selenium, and zinc are essential to maintaining the human body's metabolism. However, at higher concentrations, they can lead to poisoning. The presence of heavy metals in foods is a timely issue, involving contamination of the food chain and harm to public health. (3). There is a wide range of foods contaminated by heavy metals. In particular, fish is one of the products most affected by heavy metal contamination, because heavy metals are bioaccumulative in the food chain. When heavy metals

- Replace with: Consumption : [1] Comment Pattern and Heavy Metal Content of Commonly Consumed Fish in Benghazi
- Remove : [2] Comment
- Replace with : regarding the nutr: [3] Comment itive
- Capitalize : [4] Comment
- Remove : [5] Comment
- Replace with: metal content of : [6] Comment lead, cadmium, and mercury in the
- Replace with: study was carried : [7] Comment out on December 29, 2022, and extended to June 30, 2023
- Replace with: subjects were ran : [8] Comment donly
- Add a comma. : [9] Comment
- Remove the extra space. : [10] Comment
- Replace with: terms of heavy : [11] Comment metals using an atomic absorption spectrophotometer
- Replace with: Sparus aurata, : [12] Comment Pegellus bogaraveo, Sardina pilchardus, Mullus barbatus, and Epinephelus marginatus
- Replace with: Sparus aurata, : [13] Comment Pegellus bogaraveo, Sardina pilchardus, Mullus barbatus, and Epinephelus marginatus. are the most commonly consumed fish in Benghazi. Age, female gender, and number of family members are the variables associated with the fish consumption pattern in Benghazi. Lead, cadmium and mercury are the three heavy metals analyzed in the top five most consumed fish in Benghazi. All heavy metal results are within the recommended level by the WHO. The results of the current study recommend that increasing fish consumption habits should be considered an essential fact for having healthier generations. Furthermore, fish consumption should be increased through information given by educational institutions, government agencies, and
- Fish approximately : [14] Comment 34,000 species of vertebrate animals found in fresh and salt water globally. Fish
- Follow the referencing style. : [15] Comment Do this where it is necessary

accumulate in waters, fish ingest them through their diet and accumulate them over the course of their lives.⁽⁹⁾ The influence of heavy metals ~~from~~ in fishes varies between different species and different metals. Fishes accumulate heavy metals ~~to~~ at a concentrations many times higher than those present in water streams. Environmental circumstances such as water hardness, salinity, and pH can play a significant role in heavy metals accumulation in the fish's bodies.^(10, 11) Aquaculture is a recent activity in the Libya, beginning with the culture of a variety of freshwater fish species during in the 1970s. Success, however, was not forthcoming as a result of a lack of acceptance of farmed fish products by domestic consumers. The culture of marine fish species followed a decade later, encouraged through several Government initiatives. In terms of human resources, no historical data is available. The main marine species cultured on a commercial basis in the Libya are gilthead seabream (*Sparus aurata*), European seabass (*Dicentrarchus labrax*), and Atlantic bluefin tuna (*Thunnus thynnus thynnus*), common carp (*Cyprinus carpio*), bighead carp (*Hypophthalmichthys nobilis*), silver carp (*Hypophthalmichthys molitrix*) and grass carp (*Ctenopharyngodon idellus*) and catfish. Nile tilapia (*Oreochromis niloticus*) and red tilapia (*Tilapia* species) are the most important freshwater-cultured fish. Libya does not have any endemic freshwater species, and all freshwater species were introduced in the 1970s and 1990s.^(12, 13) Many studies have been conducted in different regions ~~in~~ of the world to evaluate fish consumption patterns and the parameters affecting ~~it~~ them. It was reported that the parameters determining the fish consumption include ~~ing~~ income level, education level, social status, parents influencing ~~in~~ the pattern of fish consumption within the family, religion, and awareness of the nutritional value of fish. Benghazi is the second-biggest city in Libya. However, to the best

of our knowledge, there are almost no previous studies on the consumption trend of fish in Benghazi city. According to the Food and Agriculture Organization (FAO), Libya's Gross domestic product (GDP) from fisheries is 0.13%, while Agricultural GDP from fisheries is 2.69%. Furthermore, the fish available for consumption is 14.9 kg/capita. During 2019, imports of fish and fish products makes made up USD 236 983 550 of the Libyan Economy.⁽¹²⁻¹⁵⁾ Fish is an important food material because of its high protein, vitamins, and minerals, as well as being an important source of omega-3 fatty acids. Fishes have an activity against cancer, and heart diseases. Heavy metal toxicity as a result of fish consumption can result in damage or reduced mental and central nervous system function, lower energy levels, and damage to blood composition, lungs, kidneys, bones, liver, and other vital organs. Long-term exposure may result in Alzheimer's disease, Parkinson's disease, muscular dystrophy, multiple sclerosis, allergies, and cancer.⁽¹⁶⁾ The presence of fish in the diet is very important because it is a source of proteins, vitamins, minerals and unsaturated essential fatty acids, especially omega-3 fatty acids.^(18, 19) Several factors determine the dietary habits of the people in the Arab world. Food consumption patterns has have dramatically changed in some Arab countries as a result of the sudden increase in income from oil revenue. It is believed that the food subsidy policy has adversely affected the food habits in the Arab states by encouraging the intake of fat, sugar, rice, wheat flour, and meat. Socio-cultural factors such as changing lifestyles, food preferences, education, and women's employment all have a noticeable influence on food consumption patterns in this region. Comprehensive studies on social, cultural, and economic factors associated with food consumption patterns in the Arab region are highly recommended. Libya is important part of

Remove extra spaces. :[23]Comment

Remove the extra spaces. :[16]Comment

Remove the extra spaces. :[17]Comment

Remove the extra spaces. :[18]Comment

Remove the extra spaces. :[19]Comment

Remove all extra spaces. :[24]Comment

Remove the extra spaces. :[20]Comment

Remove the extra spaces. :[21]Comment

Remove :[22]Comment

the African agrie-food system. Fish provides 19% of animal protein intake to Africans. The fish consumption pattern in North Africa is 13.5 kg/person / year. Africa lags behind other regions in developing its share in global fish production, consumption, and trade, and Libya is no exception. Currently, however, Africans are in the paradoxical position of being highly dependent on fish for animal protein but ranking low in per capita fish consumption. Globally, Africans are second only to Asians in terms of relying on fish as a major share of the total animal protein intake in their diet. Fish represent over 20% of animal protein intake in twenty African countries. No data are available about the fish consumption pattern in Libya. To the best of our knowledge, this is the first study regarding fish consumption patterns in Libya. ^(12, 20-22)

Several techniques have been used for the determination of metal concentrations in fish species, such as flame atomic absorption spectrometry, graphite furnace atomic absorption spectrometer, electrothermal atomic absorption spectrometry, inductive coupled plasma, and mass spectrometry. Different digestion methods were used as sample preparation methods for the determination of heavy metals in fish samples. ⁽²³⁻²⁷⁾ In 2017, a study was carried out in Sabratha coastal sediments, Mediterranean Sea, Libya, to detect the distribution and enrichment of heavy metals. 30 sediment samples were collected for Fe, Cu, Pb, Mn, Cd, Co, Ni, and Zn analysis using Atomic Absorption Spectrometry. The analysis indicated that the Sabratha's coastal sediments were enriched with Cd, Pb, Cu, Ni, Co, and Zn (EF = 81.48, 17.26, 12.80, 11.42, 9.85, and 8.56 respectively). The highest levels of Mn, Cu, Ni, Pb, and Co were recorded nearby the Mellitah complex oil and gas station in the western Libyan region, while the highest levels of Zn and Cd were recorded at in the central part of the study area nearby the fishing port and Sabratha hospital. ⁽²⁵⁾ The

Average values of Cd, Pb, and Co were mostly higher than the ones recorded from the Arabian and Oman gulfs, the Red Sea, the Gulf of Aqaba, the Caspian Sea, the coast of Tanzania, and the background shale and the earth's crust. The high levels of most of the studied heavy metals suggested significant anthropogenic sources along the Sabratha coast. ⁽²⁸⁾ In 2010, a Libyan study determined the concentrations of Co, Cd, Pb, Fe, and Cu in different tissues (bone, skin, flesh, and tail) of fish species of the Mediterranean Sea (Libyan coastline) using flame atomic absorption spectrometry. The fish species used in this study were *Sardinella aurita*, *Pagellus erythrinus*, *Balistes capricus*, *Trachurus trachurus*, *Synodus saurus*, and *Dactylopterus volitans*. The concentration of Co was ranged from 0.570 mg/kg to 44.693 mg/kg. These concentrations of Co in all examined tissues were more higher than the World Health Organization's (WHO) recommended level. The results of the analysis indicated that the concentrations of the Cd in all examined tissues varied from 0.328 mg/kg to 2.929 mg/kg. These values were greater than the WHO-recommended level. The concentration level of Pb in all examined tissues ranged from 0.246 mg/kg to 2.386 mg/kg. These values were higher than the accepted values by WHO. ⁽²⁹⁾ During In 2008, another Libyan study was published. This study aimed to investigate the relationship between the concentration of some heavy metals (Lead, Cadmium, Mercury, Copper and Zinc) in water and their bioaccumulation in the liver and muscle tissues of some marine fish species at in the El Khoms area, of Libya. The present work revealed a marked increase of in heavy metals concentration in the muscles and liver tissues of investigated fish, and such increases were positively correlated with a corresponding increase in heavy metals concentration in water. ⁽³⁰⁾ The concentration of heavy metals in two tropical fish species from the Ogba River, Nigeria, showed varying concentrations of

Remove the extra spaces. :[25]Comment

Italize :[29]Comment

Italize :[30]Comment

Remove the extra space. :[26]Comment

Replace with: values accepted by the WHO :[31]Comment

Write out the full words before the acronym in a bracket :[27]Comment

Remove :[28]Comment

Remove extra space. :[32]Comment

Cu, Mn, Zn, Pb, Cr, Ni, and Cd in the fish tissues. Concentrations of Cu, Mn, Cr, and Ni in both fish were higher than the permissible reported by the WHO while those of Zn and Pb were lower than permissible concentrations.⁽³¹⁾ Limited literatures are found regarding fish consumption patterns in Benghazi as well as the heavy metals contents of Benghazi fishes. The current study aims to evaluate the heavy metals contents in of the most commonly consumed fishes in Benghazi. The current study seeks to study methods of fish's meals preparation, knowledge regarding nutritive value of fishes, the association between socio-economic variables, and the current fish consumption pattern in Benghazi.

Methodology

This is a cross-sectional study carried out from 29th December 2022 and extended to 30th June 2023. A total of 300 subjects (Male and female) were randomly approached to participate in the study. The study respondents were people aged 18 years and above older, who were fishers, selling, and purchasing fishes from the fish local market (Bankina) in Benghazi city, giving a response rate of 85.5%. The sample size was determined by using the Richard Jaeger equation. Informed consent was obtained from the subjects who were also assured of the confidentiality of the information collected. The research was approved by the administration of the Faculty of Public Health, University of Benghazi. The questionnaire consists of four sections: socioeconomic information, fish's consumption pattern, information regarding Nutritional value, and fish cooking methods. A Questionnaire was developed by deep reading in similar pervious previous studies. The questionnaire's validity was assessed via a pilot study on thirty participants. After

determination of all consumed types of fishes in Benghazi, the top five consumed fishes were nominated to be analyse for heavy metals content. These fishes include *Sparus aurata*, *Pegellus bogaraveo*, *Sardina pilchardus*, *Mullus barbatus*, and *Epinephelus marginatus*. Three fresh samples were collected from each fish type from in the Bankina. A Pphysical inspection was done for on all samples. Fishes samples have been collected from the Benghazi local fish market (Bankina). Collected samples are illustrated in Figure 2.1. Samples were analysed in the Food Chemistry Laboratory, Department of Nutrition, Faculty of Public Health, University of Benghazi. **Reagents:** Water: redistilled or deionised, Nitric acid A.R. 0. 1 M, - diluted 7 mL concentrated concentrated acid to 1 litre, Nitric acid concentrated concentrsted (Sp. Grade 1.40), Standard solutions for cadmium, mercury, and lead. Working standard solutions; for flame analysis, dilute the standard solution with 0.1 M HNO₃ to a range of standards that covers the concentration of the elements to be determined.⁽³²⁾ The researchers weighted accurately 1 gm of each fish type assigned to be analysed. 10 ml of concentrated nitric acid has been added. The dish content is heated with a soft flame using a Bunsen burner until all the volatile or readily combustible matter has been removed. Then, the sample is leaved left to cool. 5 mL of hydrogen peroxide is added and heated for five minutes. The solution is transferred to a volumetric flask to volume be filled with water. All the previous steps of sample preparation are repeated with blanks. Identical procedures are followed even in heat and time. Samples preparation is illustrated in Figure (2) 2.

Remove the extra space. :[33]Comment

Replace with:regarding the nut:[34]Comment
ritional

Replace with December29th, 2 :[35]Comment
022, and

Re-check?? :[40]Comment

Replace with: June 30th, 2023 :[36]Comment

Remove the extra space. :[37]Comment

Remove the extra spaces. :[38]Comment

Remove the extra space. :[39]Comment




Figure (4)2: Samples of analysed fishes species

Determination

Set the instrument as per the previously established optimum conditions as per the guide lines given in the Instruction Manual

(provided along with the instrument). Determine the absorbance of the sample solution(s) and blank. Calculate the heavy metal content from the standard curve.

of what????????? :[41]Comment

Preparation of Standard Curve:

Read the absorbance of a series of standard metal solutions in the Atomic Absorption Spectrophotometer after setting the instrument as per to optimum conditions. Plot absorbance is done against μg of metal/mL solution.⁽³²⁾ All data will be coded prior to being entered into a computer. The description and analysis of the data were done by *Statistical Package for the Social Sciences* (SPSS)

version 22. The Level of significance was set at a p value < 0.05 . The researchers of the current study realize that the observational and cross-sectional design, with a single measurement in one season is a limitation of the current study. A shortage in laboratory materials limits the nominated analyzed samples to only five types and the analyzed heavy metals to only three metals.

UNDER PEER REVIEW



Figure (2): Samples Preparation

Label the pictures and describe the pictures under with the title.

Results

The percentage of subjects from in the age group 18-40 years is the highest percentage, (44%). The percentage of subjects from in the age group 41-60 years made up (37.67) 37.67%. The males made a percentage of (59.33%) 59.33 and the females made a percentage of (40.67%) 40.67 out of the total number 300 of subjects (300). The general average ages of males and females mean \pm SD is about 34 ± 4.6 as shown in Table (1). Socio-economic characteristics and details of the

respondents were presented in Table (2). The marital status shows that, (50.33%) 50.33% are married and (49.67%) 49.67% are not married. Regarding the education level of the subjects, the percentage of subjects with a basic education was (6%) 6%, and the percentage of people with a secondary education was (18%) 18%, while the highest percentage of subjects with a university level was (72.33%) 72.33%. Regarding occupation, (5.33%) 5.33% of the subjects were

students, ~~the housewife~~housewives, employees, freelancers ~~made~~were ~~(40.33%)~~40.33% and ~~(36%)~~36%, and ~~(3.3%)~~3.3% respectively of the total sample. The percentage of subjects whose income was less than 500 Libyan Diner (LD) was about ~~(3.67%)~~3.67%, while the percentage of subjects whose average income was between 500-1000 LD was ~~(42.33%)~~42.33%, which is the highest

percentage of earned income. The subjects whose average income was 1000-1500 LD were ~~(23.33%)~~23.33%. The percentage of subjects whose income was more than 1500 LD was about ~~(30.67%)~~30.67%. Families with less than five members ~~made were~~ (40%)40%, while families with more than five members ~~made were~~ (60%)60% of the total sample.

Table (1): Subject characteristics

Age (Years)		Total		Total
		Male	Female	
20-40	No. (%)	82(27.33)	50(16.67)	132(44)
41-60	No. (%)	78(26)	35(11.67)	113(37.67)
61-80	No. (%)	18(6)	37(12.33)	55(18.33)
Total	No. (%)	178(59.33)	122(40.67)	300(100)
Age (Years)Mean ± SD		35±3	33±6.2	34±4.6

Give a description of the Table :[43]Comment under theTable.

Remove bold :[44]Comment

Table (2): Socioeconomic Characteristics of the Subjects

Socioeconomic Characteristics	No.	%
Marital status		
Married	151	50.33
Not Married	149	49.67
Education Level		
Basic	18	6
Secondary	54	18
University	217	72.33
Higher Education	11	3.67
Occupation		
Student	16	5.33
House wife	121	40.33
Employee	108	36
Freelance	10	3.3
Retired	45	15
Income level		
<500	11	3.67
500-1000	127	42.33
1000-1500	70	23.33
>1500	92	30.67
Family members		
<5	120	40
≥5	180	60

Give a description of the Tabl :[45]Comment e under theTable.

Remove bold :[46]Comment

Remove bold :[47]Comment

Remove bold :[48]Comment

Remov bold :[49]Comment

Fish Consumption Pattern

The percentage of subjects who consume fish is ~~(88%)~~88%. The reasons for not eating fishes varied between tastes and smell (33.33%) and bone (30.56%). Subjects who consume fish from once to

twice a weekly make about ~~(42.33%)~~42.33%. It is has the highest percentage. ~~(8.67%)~~8.67% of subjects consumes fish three to four times a week. ~~(33%)~~33% of subjects who do not eat fish

on a weekly basis. ~~(42.67%)~~42.67% of the subjects consume once to twice a month which is the highest percentage. ~~(25.33%)~~25.33% of the subjects consume fish three to four times a month. Furthermore, subjects who consumed fresh fish made ~~(72%)~~72% of the total sample; which is the highest percentage. ~~(7.3%)~~7.3% of subjects preferred canned fish, frozen fish, salted fish, and smoked fish were preferred by ~~(10%)~~10%, ~~(6.7%)~~ 6.7%, and ~~(4%)~~4% of

the subjects, respectively. Figure ~~(3)~~3 shows the frequency of the most consumed fishes in Benghazi. The top five consumed fishes were *Sparus aurata* (45.33%), *Pegellus bogaraveo* (36.67%), *Sardine pilchardus* (34.67%), *Mullus barbatus* (33.33%), and *Epinephelus marginatus* (29.67%).

Table (3)3: Fish Consumption Pattern

Socioeconomic Characteristics	No.	%
Fish Consumption		
Yes	264	88
No	36	12
Reasons for Not eating fishes		
Taste and Smell	12	33.33
bones	11	30.56
Taste, Smell and Bones	13	36.11
Weekly fish consumption		
No	99	33
1-2	127	42.33
3-4	26	8.67
5-6	4	1.33
≥ 7	8	2.67
Monthly fish consumption		
No	20	6.67
1-2	128	42.67
3-4	76	25.33
5-6	26	8.67
≥ 7	14	4.67
Types of preferred Fishes		
Fresh	216	72
Canned	22	22
Frozen	30	30
Salted	20	20
Smoked	12	12




Figure (3)3: Frequency of Different fishes Consumption in Benghazi

Figure (4)4 presents the reasons for selected the preferred fishes. The fish quality was (9%)9%; it is the lowest percentage. The price of the fish was (+16%)16%, the taste was (-39%)39%; it is the highest percentage, and the nutritive value was (-30%)30%. Figure (5)5 shows the sources of consumed fishes. The highest percentage comes from the local market "Bankina" with (+46%)46%. Supermarkets, fish restaurants, fishermen, and fishing make (-24%)24%, (+11%)11%, (+12%)12%, and (-7%)7%, respectively. Figure (6)6 shows the reasons of for selecting fish's sources. The highest percentage (51%) was for available fishes types, (-30%)30% for place close to home and (-19%)19% for price. Figure (7)7 presents factors affecting fish selection. The highest percentage was for there is no special reason (44.3%), (-14.7%)14.7% for gills color, and (-12%)12% for eyes status. The lowest

percentages were (-10.3%)10.3%, (+10%)10%, and (-8.7%)8.7% for tenderness, smell, and color, respectively. Figure (8)8 shows the most preferred cooking methods of for fishes in Benghazi. Grilling on charcoal is the preferred preparation method of for fish in Benghazi with (35.2%). (-20.9%)20.9% of the subjects prefer eating fried fish. Libyan traditional Tagin and oven roasting are coming after that with a percentage of (-12.36%)12.36 and (-11.11%)11.11, respectively.




Figure (4): Reasons for selected the preferred fishes

Figure 4: Reasons for selectin :[51]Comment
g the preferred fish




Figure (5): Sources of Consumed Fishes




Figure (6): Reasons of Selecting Fishes Sources

Figure 5: Sources of consume :[52]Comment
d fish

Figure 6: Reasons for selectin :[53]Comment
g fish sources




Figure (7): Factors affecting fish Selection




Figure (8): Fish Pprefering Ccooking Mmethods

Figure (9) shows the subjects knowledge regarding the nutritive value of fish. The highest percentage of the subjects (45%) know that fish are good sources of omega 3. Only (19%) 19% of the subjects know that fish are good sources of protein. Furthermore, only (9%) 9% of the subjects know that fish are good sources of healthy types of fats. (10%) 10% and (5%) 5% of the subjects know that fish are good sources of vitamin D and carbohydrate, respectively. (12%) 12% of the subjects state that they do not have any information regarding the nutritive value of the fish. Figure (10) shows the subjects knowledge regarding the adult health benefits of fish.




Figure (9): Subjects knowledge regarding the nutritive value of fish

(29.33%) 29.33% of the subjects know that fish is are significant for bone health. Prevention of chronic diseases and brain health as benefits of fish has response by the subjects (25.33%) and (25.67%), respectively. Only (19.67%) 19.67% of the subjects think that fish are important for general health. Figure (11) shows the sources of information regarding fish. The internet was the biggest source of information with a percentage of (42%) 42%. The radio was the second source of information, with a percentage of (26.7%) 26.7%. Television, and health education programs have similar effects, with a percentage of (15.7%) 15.7% and (15%) 15%, respectively.




Figure (10): Subjects knowledge regarding the nutritive value of fish




Figure (11): Sources of information regarding fish

In *Sparus aurata* and *Epinephelus marginatus*, lead is higher in muscles than guts and grills respectively. In *Pegellus bogaraveo*, *Sardina pilchardus*, and *Mullus barbatus* lead (Pb) is are higher in guts than muscles and grills. Regarding,

cadmium (Cd), it is higher in gills than muscles and guts in all samples of the five fishes. Looking at mercury, all samples provide less than 0.01 ppm except in *Epinephelus marginatus*. *Epinephelus marginatus* contains 0.03 ppm of mercury

Space :[55]Comment

Remove the extra space. :[56]Comment

Remove the extra space. :[54]Comment

in the guts, which is the highest compared to gills and muscles. Muscles and gills of *Epinephelus marginatus* contain 0.027 and 0.016 ppm of mercury, respectively as illustrated in Table 4.

Table (4)4: Heavy metals contents in the most consumed fishes in Benghazi

Fishes Samples	Gills			Guts			Muscles		
	Pb	Cd	Hg	Pb	Cd	Hg	Pb	Cd	Hg
<i>Sparus aurata</i>	0.010	0.012	<0.01	0.015	<0.005	<0.01	0.018	<0.005	<0.01
<i>Pegellus bogaraveo</i>	0.014	0.010	<0.01	0.024	<0.005	0.013	0.020	<0.005	0.011
<i>Sardina pilchardus</i>	<0.005	0.007	<0.01	0.008	<0.005	<0.01	<0.005	<0.005	<0.01
<i>Mullus barbatus</i>	0.008	0.017	<0.01	0.011	0.006	<0.01	0.010	<0.005	<0.01
<i>Epinephelus marginatus</i>	0.019	0.025	0.016	0.027	0.009	0.030	0.032	0.006	0.027

Indicate the full name of the metal acronym below the table. **:[57]Comment**

Italicize the fish name. Do this for the remaining fish name. **:[58]Comment**

The results of the Logit model are shown in Table 5. The age variable has been found significant at the 5% level statistically. As the age increases a unit, fish consumption amount probability increases 1.643. The probability of consuming fish increases by 1.643 for

every unit increase in age. In this study, females consume more fish than males. Also, as family members increase, the probability of consumers' consumption increases at 1.40.

Table (5)5: Association of socioeconomic variables with fish consumption

Variables	B (Coefficient of explanatory variables)	Standard Error	Z-statistics	Severity Rating	Betting odds
Age	0.497	0.193	2.568	0.010	1.643
Gender	-0.715	0.357	-2.003	0.045	0.489
family members	0.337	0.139	2.425	0.015	1.401

Format as the rest. **:[59]Comment**

Discussion

Worldwide, fish consumption and marketing has become an essential phenomenon due to the increasing appreciation of its health and economic benefits. Given this, the current study sought to identify the pattern of fish consumption and knowledge regarding the nutritive values of fish in Benghazi City. Consequently, the top five most consumed fish species in Benghazi have been analyzed in terms of the heavy metals content of lead, cadmium, and mercury. Data obtained have been analyzed using the Statistical Package for Social Sciences (SPSS). *Sparus aurata*, *Pegellus bogaraveo*, *Sardina pilchardus*, *Mullus barbatus*, and *Epinephelus marginatus* are

the most preferred commonly consumed fishes in Benghazi. (1, 17) Fish consumption preferences are affected by individuals' socioeconomic characteristics. The percentage of subjects from the age group of 20-40 years is the highest percentage, and they constituted about 44%. It is followed by subjects from the age group 41-60 years (37.67%). The mean ages of male's \pm SD were about 35 ± 3 , and the mean \pm SD for females was about 33 ± 6.2 . The general average ages of males and females mean \pm SD about 34 ± 4.6 . This mean is almost close to the mean age of a similar study 31 years done by Rhoda Foil's 2020. (33) In Tripoli similar study

Italicize **:[60]Comment**

Italicize **:[61]Comment**

in Tripoli, the age category (36-45) was had the highest age (34%) among the study sample. The age variable has been found significant at the 5% level statistically.⁽³⁴⁾ The coefficient of the age variable took a positive value. ~~When the age increases a unit, fish consumption amount probability increases 1.643.~~ The probability of consuming fish increases by 1.643 for every unit increase in age. This result is similar to previous studies by Ahmad ~~N~~ (2016) and Burger ~~J~~ (2002).^(35, 36) The males made a percentage of ~~(59.33%)~~ 59.33% and the females made a percentage of ~~(40.67%)~~ 40.67% of the total number 300 of the subjects (300). This result is opposite to the Tripoli study of fish consumption patterns. In Tripoli, ~~it was that~~ nearly 60% of respondents were females, while 40% were males. The coefficient of the gender variable takes a negative value. In this case, it has been determined that females consume fish more than males. Females consume more fish ~~in~~ at 0.489 rates ~~more~~ than males. The findings of this study are similar to those of a Turkish study that was carried out during in 2013.^(34, 37) The marital status shows that ~~(50.33%)~~ 50.33% are married. In the current study as well as in previous studies, statistical tests did not show significant differences relating to the fish consumption and marital status of respondents.⁽³⁴⁾ Regarding the education level of the subjects; the percentage of subjects with a basic education was ~~(6%)~~ 6%, and the percentage of people with a secondary education was ~~(18%)~~ 18%, while with the highest percentage of subjects with a university level (72.33%). Comparing to the literature, most of the respondents (36%) indicated that the terminal educational level was Junior High School, while a minority (8%) indicated that primary as their last level of education. Only 8% of the respondents reported not having a formal education. Statistical tests did not show a significant difference relating to the fish consumption

and the level of education.^(34, 37) Regarding occupation; ~~(5.33%)~~ 5.33% of the subjects were students, the ~~house wife~~ housewives (40.33%), employee (36%), and freelance (3.3%), respectively, of the total sample. In the previous study, 17.4% of them ~~are were~~ civil servants, 20.3% ~~are were~~ workers, 23.2% ~~are were~~ homemakers, 19.2% ~~are were~~ other, 10.7% ~~are were~~ self-employed, and 9.2% ~~are were~~ unemployed. Statistical tests did not show a significant difference relating to the fish consumption ~~and the~~ occupation.⁽³³⁻³⁷⁾ The percentage of subjects whose income was less than 500 Libyan ~~Diner~~ dinars (LD) is about ~~(3.67%)~~ 3.67%, and the percentage of subjects whose average income was between 500- and 1000 is ~~(42.33%)~~ 42.33%, which is the highest percentage of earned income. When considering the exchange rate, the results of this study ~~is~~ are similar to those of the Turkish study.⁽³⁷⁾ Families with less than five members make (40%) 40% up to the total sample, while families with more than five members make ~~(60%)~~ 60% of the total sample. In this study, the variable of the individual number in the family takes on a positive value. A unit increases the number of family members; increases the probability of consumers' consumption at by 1.401. This result is similar to the results ~~by~~ of a previous study done by E. Onurlubas during in 2013.⁽³⁷⁾ The current study reveals that the percentage of subjects who consume fish was ~~(88%)~~ 88%. Accordingly, ~~(12%)~~ 12% of the subjects do not consume fish. The consumption rate in Benghazi is higher than the previous African rate in sub-Saharan African, where the majority of the respondents (72%) consumed fishes, while ~~(28%)~~ 28% indicated that they do not consume fishes. The consumption rate in Benghazi is lower than in Tripoli. Tripoli results showed that 98% of the respondents consume fish.^(33, 34) The reasons for not eating fishes in the current study vary between tastes and smell

Remove extra spaces :[62]Comment

Remove the extra spaces :[63]Comment

Remove the extra spaces. :[64]Comment

(33.33%) and bone content (30.56%). In a previous study, subjects ~~do~~ did not like fish, and being very expensive ~~was~~ was one of the main reasons for non-consumption of fish, with a percentage of 16.7% and 16.7%, respectively. It is noteworthy that 44.4% of the respondents did not give a reason for not consuming fish. ⁽³⁴⁾ Subjects who consume fish ~~from~~ once to twice a weekly make up about ~~(42.33%)~~ 42.33% of the total sample. It is the highest percentage (8.67%) of subjects who consumes fish three to four times a week. ~~(33%)~~ 33% of subjects who do not eat fish on a weekly ~~bases~~ basis. Regarding monthly fish consumption, ~~(6.67%)~~ 6.67% of the subjects don't consume, and ~~(42.67%)~~ 42.67% of the subjects consume once to twice a month, which is the highest percentage. ~~(25.33%)~~ 25.33% of the subjects consume fish three to four times a month. The results of the Tripoli study ~~results~~ show that the largest percentage of respondents (66%) consume fresh fish one to two times per month, 27% consume fish once to twice a week, and 5% ~~of respondents~~ consume fresh fish daily. Furthermore, subjects who consumed fresh fish made up ~~(72%)~~ 72% of the total sample, which is the highest percentage. ~~(7.3%)~~ 7.3% of subjects preferred canned fish, and frozen fish (10%), salted fish (6.7%), and ~~for~~ smoked fish (4%) were preferred, respectively. In the previous study, the categories of fish consumed ~~are~~ were, in descending order, include fresh, canned, frozen, and salted, and smoked with percentages of 96.7, 94.7, 21.8 and 0.98%, respectively. ⁽³³⁻³⁷⁾ Nineteen fish types are consumed in Benghazi. The top five consumed fishes were *Sparus aurata* (45.33%), *Pegellus bogaraveo* (36.67%), *Sardine pilchardus* (34.67%), *Mullus barbatus* (33.33%), and *Epinephelus marginatus* (29.67%). In previous Libyan date, more than one-third of the respondents frequently buy and consume sardine, sea bream, and mackerel with a percentage of 57.6, 47.0 and 37.2%,

respectively. The reasons ~~off~~ for the selection of these types include fish quality (9%); price (16%), taste (39%), and nutritive value (30%), respectively. In previous study, the reasons behind respondents' choices towards a certain fish included that 61% of the respondents buy certain types of fish because of their taste, while 38% of them make their choice ~~is~~ based on their belief that this type of fish is good for their health. Meanwhile, 22, 21, and 8% of the respondents buy certain kinds of fish because their prices are affordable, ~~and~~ fit ~~to~~ within budget, high quality, and easy to prepare, respectively. ⁽³⁴⁾ The highest percentage of consumed fish (46%) comes from the local market "Bankina" ~~with~~ (46%). Supermarket, fishers, and fish restaurants, and fishing make ~~(24%)~~, ~~(11%)~~, ~~(12%)~~, and ~~(7%)~~, respectively. Literature showed the that fish market at the seaport is the most popular market for respondents (75%) to buy fish from, followed by the fish store (13.3%), directly from ~~of~~ fishermen (11%), the fish and meat market (6%), the supermarket (5%), and vehicles for selling fish (3%). ⁽³³⁻³⁷⁾ The ~~Reasons~~ behind the selection of these places ~~are~~ varies vary in the current study. The highest percentage (51%) was for available fishes types, ~~(30%)~~ for places close to home, and ~~(19%)~~ for price. In a previous study, the reasons behind the ~~chosen~~ choice of a certain place for buying fish by respondents ~~are~~ were varied. The majority of the respondents (82.2%) choose a place ~~ofto~~ buying fish because of the availability of fresh fish in the chosen place, followed by the reasons of being near ~~my~~ their house, and cheap, with ~~the~~ a percentage of 23.6% and 14.3%, respectively. ⁽³⁴⁾ Grilling on charcoal is the preferred preparation method ~~offor~~ for fish in Benghazi ~~with~~ (35.2%). ~~(20.9%)~~ of the subjects prefer eating fried fish. Libyan traditional ~~T~~ tagine (mixed fish and different types of vegetables such as potato, zucchini, carrot, eggplant, and garlic) and oven roasting are coming after that with a percentage of ~~(12.36%)~~ and

(11.11%), respectively. The most popular methods of fish cooking include frying, grilling, with soup, Tagine, and couscous. The results from the Tripoli Study revealed that frying is the most frequent method of cooking fish (71.1%) by among the respondents in this study, followed by grilling (61.7%).⁽³⁴⁾ The highest percentage of the subjects (45%) knows that fish are good sources of omega 3. Only (19%) of the subjects know that fish are good sources of protein. Furthermore, only (9%) of the subjects know that fish are good sources of healthy types of fats. (10%) and (5%) of the subjects know that fish are good sources of vitamin D and carbohydratesprotein, respectively. (12%) of the subjects state that they do not have any information regarding the nutritive value of the fish. In the previous study, the subjects havehad better knowledge than Benghazi subjects regarding the fish content of omega 3. In a previous study, a lower percentage of the subjects (32%) of the participants in the study were able to identify omega 3 fatty acids as essential fats.^(36, 37) In the current study, (29.33%) of the subjects know that fish is are significant for bone health. Prevention of chronic diseases and brain health as a benefits of fish have response received byfrom the subjects (25.33% and 25.67%), respectively. Only (19.67%) of the subjects think that fish are important for general health. In the previous study, the majority of the respondents (85.6%) agreed that eating fish is good for health. Again, literature indicates that knowledge is much lower in Benghazi regarding the health benefits of fish compared to other regions.⁽³⁴⁾ The internet was the biggest source of information, with a percentage of (42%). The radio was the second source of information, with a percentage of (26.7%). Television, and health education programs have similar effects, with a percentage of

(15.7%) and (15%), respectively. In a previous study, the majority of the respondents (83%) chose television as the most effective medium for getting information about the benefits and use of fish. WhilstWhile 12%, 4%, and 1% of the respondents chose radio, the internet, and newspapers as an effective mediums for obtaining information about the benefits and use of fish. Different sources of information justify the differences in knowledge levels compareding to other regions.^(34, 35) Lead (Pb), cadmium (Cd), and mercury (Hg) are the analysed heavy metals analysed in the top five most consumed fishes in Benghazi in Ppart Pper Mmillion (PPMppm) in the current study using Atomic Absorption Spectrophotometer inside the Food Chemistry and Analysis Laboratory of the Department of Nutrition, Faculty of Public Health, University of Benghazi. In *Sparus aurata* and *Epinephelus marginatus* lead is higher in muscles than guts and grills, respectively. In *Pegellus bogaraveo*, *Sardina pilchardus*, and *Mullus barbatus* lead isare higher in guts than muscles and grills. All these values are higher than previous Libyan studies published by researchers from Tripoli, Alkhomse, and Sebha. However, all these levels are still within the recommended levels by the WHO. Regarding, cadmium it is higher in gills than muscles and guts in all samples of the five fishes. Moreover, these all results for cadmium are still within the recommended levels by the WHO. Looking at mercury, all samples provide < 0.01 ppm except in *Epinephelus marginatus*. *Epinephelus marginatus* contains 0.03 ppm of mercury in guts, which is the highest compareding to gills and muscles. Muscles and gills of *Epinephelus marginatus* contain 0.027 and 0.016 ppm of mercury, respectively. All these results are within the recommended level by the WHO.^(34, 37, 38)

Remove the extra spaces. :[67]Comment

Remove the extra space. :[66]Comment

Remove the extra space. :[68]Comment

Conclusion

Sparus aurata, *Pegellus bogaraveo*, *Sardina pilchardus*, *Mullus barbatus*, and *Epinephelus marginatus* are the most commonly preferred consumed fishes in Benghazi. Age, female gender, and number of family members are the variables that associated with fish the consumption pattern in Benghazi. The current study reveals that the percentage of subjects who consume fish in Benghazi was ~~(88%)~~88%. Accordingly, ~~(12%)~~12% of the subjects do not consume fish. Lead (Pb), cadmium (Cd), and mercury (Hg) are the analyzed heavy metals analyzed in the top five most consumed fishes in Benghazi. All these results are within the recommended level by the WHO.

References

1. Barbarossa V, Schmitt RJ, Huijbregts MA, Zarfl C, King H, Schipper AM. Impacts of current and future large dams on the geographic range connectivity of freshwater fish worldwide. *Proceedings of the National Academy of Sciences*. 2020 Feb 18;117(7):3648-3655.
2. Kilic Z. Water pollution: causes, negative effects and prevention methods. *İstanbul Sabahattin Zaim Üniversitesi Fen Bilimleri Enstitüsü Dergisi*. 2021;3(2):129-132.
3. Kosimovna KS, Omonjonovich NB, Shukhratovich AN, Azamovna KM. ~~CLASSIFICATION OF HEAVY METALS IN MEAT AND DAIRY PRODUCTS BASED ON THE DEFINITION~~ Classification of heavy metals in meat and dairy products based on the definition. *Open Access Repository*. 2022 Nov 25;8(11):260-265.
4. Ali H, Khan E, Ilahi I. Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation. *Journal of chemistry*. 2019 Mar 5;2019.
5. Renu K, Chakraborty R, Myakala H, Koti R, Famurewa AC, Madhyastha H, Vellingiri B, George A, Gopalakrishnan AV. Molecular mechanism of heavy metals (Lead, Chromium, Arsenic, Mercury, Nickel and Cadmium)-induced hepatotoxicity—A review. *Chemosphere*. 2021 May 1;271:129735.
6. Zhu T, Li L, Duan Q, Liu X, Chen M. Progress in our understanding of plant responses to the stress of heavy metal cadmium. *Plant Signaling & Behavior*. 2021 Jan 2;16(1):1836884.
7. Azeez NA, Dash SS, Gummadi SN, Deepa VS. Nano-remediation of toxic heavy metal contamination: Hexavalent chromium [Cr (VI)]. *Chemosphere*. 2021 Mar 1;266:129204.
8. Ahmad K, Shah HU, Ashfaq M, Nawaz H. Removal of decidedly lethal metal arsenic from water using metal organic frameworks: A critical review. *Reviews in Inorganic Chemistry*. 2022 Jun 27;42(2):197-227.
9. Wang J, Shan Q, Liang X, Guan F, Zhang Z, Huang H, Fang H. Levels and human health risk assessments of heavy metals in fish tissue obtained from the agricultural heritage rice-fish-farming system in China. *Journal of hazardous materials*. 2020 Mar 15;386:121627.
10. Bozorgzadeh E, Pasdaran A, Ebrahimi-Najafabadi H. Determination of toxic heavy metals in fish samples using dispersive micro solid phase extraction combined with inductively coupled plasma optical emission spectroscopy. *Food Chemistry*. 2021 Jun 1;346:128916.
11. Töre Y, Ustaoglu F, Tepe Y, Kalipci E. Levels of toxic metals in edible fish species of the Tigris River (Turkey); threat to public health. *Ecological Indicators*. 2021 Apr 1;123:107361.
12. Filogh A. Libya Fishing Industry. *Menba Kastamonu Üniversitesi Su Ürünleri Fakültesi Dergisi*. 2019 Nov 7;5(1):16-26.
13. Anderson E, Blake G. The Libyan fishing industry. In *Libya Since Independence*

- (RLE Economy of Middle East) 2014; **Oct-24** (pp. 85-104). Routledge.
14. Elbaraasi H, Elabar B, Elaabidi S, Bashir A, Elsilini O, Shakman E, Azzurro E. Updated checklist of bony fishes along the Libyan coasts (southern Mediterranean Sea). *Mediterranean marine science*. 2019 **May-10**;20(1):90-105.
 15. Omar M, Nouh F, Younis M, Younis M, Nabil N, Saad M, Ali M. Vitamin D status and contributing factors in patients attending three polyclinics in Benghazi Libya. *J Adv Med Med Res*. 2017 **Nov 6**;24:1-3.
 16. Azaman F, Juahir H, Yunus K, Azid A, Kamarudin MK, Toriman ME, Mustafa AD, Amran MA, Hasnam CN, Saudi AS. Heavy metal in fish: Analysis and human health-a review. *Jurnal Teknologi*. 2015 **Oct-20**;77(1).
 17. Kawser Ahmed M, Baki MA, Kundu GK, Islam S, Islam M, Hossain M. Human health risks from heavy metals in fish of Buriganga river, Bangladesh. SpringerPlus. 2016 **Dec**;5(1):1-2.
 18. Jobling M. Fish nutrition research: past, present and future. *Aquaculture international*. 2016 **Jun**;24(3):767-786.
 19. Authman MM, Zaki MS, Khallaf EA, Abbas HH. Use of fish as bio-indicator of the effects of heavy metals pollution. *Journal of Aquaculture Research & Development*. 2015 **Jan-1**;6(4):1-3.
 20. Rahim HF, Sibai A, Khader Y, Hwalla N, Fadhil I, Alsiyabi H, Mataria A, Mendis S, Mokdad AH, Hussein A. Non-communicable diseases in the Arab world. *The Lancet*. 2014 **Jan 25**;383(9914):356-367.
 21. Hwalla N, Al Dhaheri AS, Radwan H, Alfawaz HA, Fouda MA, Al-Daghri NM, Zaghoul S, Blumberg JB. The prevalence of micronutrient deficiencies and inadequacies in the Middle East and approaches to interventions. *Nutrients*. 2017 **Mar-3**;9(3):229.
 22. Argyriou T, Cook TD, Muftah AM, Pavlakis P, Boaz NT, Murray AM. A fish assemblage from an early Miocene horizon from Jabal Zaltan, Libya. *Journal of African Earth Sciences*. 2015 **Feb 4**;102:86-101.
 23. Pirsabe M, Fattahi N. Trace determination of heavy metals in farmed trout fish using dispersive liquid-liquid microextraction based on solidification of floating organic drop and graphite furnace atomic absorption spectrometry. *Analytical methods*. 2015;7(15):6266-6273.
 24. Barich BE. Northwest Libya from the early to late Holocene: New data on environment and subsistence from the Jebel Gharbi. *Quaternary International*. 2014 **Jan-23**;320:15-27.
 25. Nour HE, El-Sorogy AS. Distribution and enrichment of heavy metals in Sabratha coastal sediments, Mediterranean Sea, Libya. *Journal of African Earth Sciences*. 2017 **Oct 4**;134:222-229.
 26. Soliman NF, Nasr SM, Okbah MA, El Haddad HS. Assessment of metals contamination in sediments from the Mediterranean Sea (Libya) using pollution indices and multivariate statistical techniques. *Global Journal of Advanced Research*. 2015;1:120-136.
 27. Nasr SM, Okbah MA, El Haddad HS, Soliman NF. Fractionation profile and mobility pattern of metals in sediments from the Mediterranean Coast, Libya. *Environmental monitoring and assessment*. 2015 **Jun**;187:1-8.
 28. Al-Kahtany K, El-Sorogy A, Al-Kahtany F, Youssef M. Heavy metals in mangrove sediments of the central Arabian Gulf shoreline, Saudi Arabia. *Arabian Journal of Geosciences*. 2018 **Apr**;11:1-2.
 29. Khalifa KM, Hamil AM, Al-Houni AQ, Ackacha MA. Determination of heavy metals in fish species of the Mediterranean Sea (Libyan coastline) using atomic absorption spectrometry. *International Journal of PharmTech Research*. 2010;2(2):1350-1354.
 30. Metwally MA, Fouad IM. Biochemical changes induced by heavy metal pollution in marine fishes at Khomse

- Coast, Libya. *Global veterinaria*. 2008;2(6):308-311.
31. Obasohan EE, Oronsaye JAO, Eguavoen OI. A comparative assessment of the heavy metal loads in the tissues of a common catfish (*Clarias gariepinus*) from Ikpoba and Ogba Rivers in Benin City, Nigeria. *African Scientist*: 2008; 9, (1), 20-31.
 32. Thongsaw A, Sananmuang R, Udnan Y, Ampiah-Bonney RJ, Chaiyasith WC. Immobilized activated carbon as sorbent in solid phase extraction with cold vapor atomic absorption spectrometry for the preconcentration and determination of mercury species in water and freshwater fish samples. *Analytical Sciences*. 2019 **Nov 10**;35(11):1195-1202.
 33. Foli R, Awuni I, Amponsah SK. Pattern of fish consumption and fish distribution: A Study on Sunyani municipality, Ghana. *Asian J. Fish. Aquat. Res*. 2020;10:28-39.
 34. Abuhlega TA. Fish Consumption and Knowledge of Chemical Pollutants Among a Sample of New Mothers in Tripoli, Libya. *Journal of Agricultural and Marine Sciences [JAMS]*. 2020 **Apr 10**;25:39-47.
 35. Ahmad NI, Wan Mahiyuddin WR, Tengku Mohamad TR, Ling CY, Daud SF, Hussein NC, Abdullah NA, Shaharudin R, Sulaiman LH. Fish consumption pattern among adults of different ethnics in Peninsular Malaysia. *Food & nutrition research*. 2016 **Jan 1**;60(1):32697.
 36. Burger J. Consumption patterns and why people fish. *Environmental research*. 2002 **Oct 1**;90(2):125-135.
 37. Onurlubas E. The factors affecting fish consumption of the consumers in Kesan Township in Edirne. *Bulgarian Journal of Agricultural Science*. 2013;19(6):1346-1350.
 38. Metwally MA, Fouad IM. Biochemical changes induced by heavy metal pollution in marine fishes at Khomse Coast, Libya. *Global Veterinaria*. 2008;2(6):308-311.
 39. Khalifa KM, Hamil AM, Al-Houni AQ, Ackacha MA. Determination of heavy metals in fish species of the Mediterranean Sea (Libyan coastline) using atomic absorption spectrometry. *International Journal of PharmTech Research*. 2010;2(2):1350-1354.