

### Original Research Article

#### Effect of level of nitrogen and phosphorus on growth, yield and economics of lentil (*Lens culinaris* L.).

#### ABSTRACT

The present investigation was carried out in *Rabi* (winter) season 2022-23 at Agronomy Research Farm of Chandra Bhanu Gupta Krishi Snatakot Mahavidyalaya, Bakshika Talab, Lucknow (U.P.). The 16 treatments comprised of four nitrogen levels viz., 0, 15, 30 and 45 kg N ha<sup>-1</sup> with four phosphorus levels viz., 0, 20, 40 and 60 kg P<sub>2</sub>O<sub>5</sub> were tried in Factorial Randomized Block Design (FRBD). The soil was silty-loam texture with slightly alkaline pH 8.1, organic carbon 0.70%, available nitrogen 270.00 kg ha<sup>-1</sup>, available phosphorus 27.0 kg ha<sup>-1</sup>, and available potash 262.0 kg ha<sup>-1</sup> during 2022-2023. The rainfall received during the crop period was 0.00 and 0.02 mm, respectively. The relative humidity was recorded maximum during month of December study while, the sunshine ranged from 2.3 to 9.1 during crop period. Increasing levels of nitrogen and phosphorus significantly increased most of the growth and yield attributing characters viz., plant height, number of branches, dry matter accumulation and yield attributes viz., number of pods plant<sup>-1</sup>, number of grains pod<sup>-1</sup>, grain weight and test weight were significantly higher with 45 kg N ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and superior to other doses of nitrogen and phosphorus. Yield studies viz., Grain yield, straw yield, biological yield and harvest index were significantly higher with 45 kg N ha<sup>-1</sup> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and superior to other doses of nitrogen and phosphorus. The N, P, and protein content (%) and uptake of nutrient increased with increasing level of nitrogen and phosphorus. The highest nutrient content and uptake were recorded with 45 kg N ha<sup>-1</sup> with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Economics analysis viz., gross return, net return and benefit:cost ratio was recorded highest with 45 kg N ha<sup>-1</sup> + 60 kg P<sub>2</sub>O<sub>5</sub> while lowest with control. Thus for obtaining better yield and profit apply 45 kg N ha<sup>-1</sup> in combination with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> for lentil.

**KEYWORDS:** Nitrogen, phosphorus, Growth, Yield, Economics lentil.

#### INTRODUCTION

Lentil (*Lens culinaris* L.) is a herbaceous annual plant which originated from Middle East and a rich source of protein. The genus *Lens* is part of the subfamily Faboideae which is contained in the flowering plant family Fabaceae or commonly known as legume or bean family, of the order Fabales in the kingdom Plantae. As food crop, the majority of world production comes from Canada and India, producing 58% together. India ranked first in area and second in the production with 39.79% and 22.79% of world area and production respectively. Canada ranked first in production (41.16%) due to very high level of productivity (1633 kg ha<sup>-1</sup>) as compared to India (611 kg/ha) (FAO state, 2014). The major lentil producing states are Uttar Pradesh, Madhya Pradesh and Bihar. Uttar Pradesh alone occupies 0.58 million ha area with 0.44 million tons production and productivity of 701 kg ha<sup>-1</sup>. Nitrogen is critical element for increasing the quality of food crops. Nitrogen deficiency occurred frequently in every where, therefore, these elements should be comprised as a fertilizer. Application of optimum amount of nitrogen in legumes increased pod number, seed number and seed weight. Verma and Kalra *et al.* (1983) reported that lentil have good response to nitrogen fertilizer application. Studies have shown that nitrogen fertilizer application rate of 20 to 25 kilograms of fertilizer per hectare as a starter is necessary to achieve maximum plant growth. The abundant N supply through N<sub>2</sub> fixation is thought to exacerbate the indeterminate growth habit of lentil. In addition, when a legume crop like lentil has access to N fertilizer, the crop favours fertilizer N uptake instead of nitrogen fixation, due to nitrate or ammonium uptake being a less energy consuming process for the plant. Inhibiting effects of nitrate on legume N<sub>2</sub> fixation have also been described in the literature (Bremer *et al.*, 1989). Therefore, the N supply to lentil particularly due to N<sub>2</sub> fixation could play a role in delayed maturity. (Togay *et al.*, 2005), (Deol, 2007).

Comment [AM1]: Where are these treatments?

Comment [AM2]: How is factorial? Look at the tables.

Comment [AM3]: ??????????

Comment [AM4]: and phosphorus

Comment [AM5]: ?????????

Comment [AM6]: level of

Comment [AM7]: recorded highest

Comment [AM8]: and profit

Comment [AM9]: kg ha<sup>-1</sup>

Comment [AM10]: delete

Comment [AM11]: hectares area

Comment [AM12]: Source

Comment [AM13]: delete

Comment [AM14]: Very old

Comment [AM15]: fertilizer N

Phosphorus is an important macro element for growth of legumes. It has important role information of root nodules and this opinion has an important role in nitrogen fixation. Phosphorus (P) is a non-renewable and second most important macronutrient which is required for young tissues and performs several functions related to growth, development, and metabolism of the plant and also regulates many metabolic activities of the plant life. Application of 20-80 kg P/ha in combination with 40 kg N ha<sup>-1</sup> increased the number of filled pod lentil (Sharma *et al.*, 2004). Determining the appropriate amount and mix of fertilizer nitrogen and phosphorus is important because that have a large role in increasing the performance of crops. Other studies showed that percentage of lentil seed protein was increased by applying 40-60 kg P ha<sup>-1</sup>.

#### METHODS AND MATERIALS

A field experiment was conducted at Shradhay Bhagwati Singh Agriculture Research Farm (Hajipur), Chandra Bhanu Gupta Krishi Snatkotah Mahavidyalaya, Bakshi-Ka-Talab, Lucknow University, Lucknow (U.P.) during Rabi season 2022-2023. The experimental site is situated at 26.50° North latitude and 80.50° East longitude with an altitude of 123 meters above mean sea level. The soil of experimental field was silty-loam texture, slightly alkaline in reaction (8.0 p<sup>H</sup>), medium in organic carbon (0.70%) and available nitrogen (270 kg ha<sup>-1</sup>) phosphorus (27 kg ha<sup>-1</sup>) and potassium (262 kg ha<sup>-1</sup>). Sixteen treatments comprised of four levels of Nitrogen (0, 15, 30 and 45 N/ha) and four levels of phosphorus (0, 20, 40 and 60 P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) were tested in Factorial Randomized Block Design with three replications. Nitrogen and phosphorus were applied through urea and SSP, respectively. A full dose of Nitrogen and Phosphorus as per treatment was applied at the time of sowing. A common dose of potassium (20 kg K<sub>2</sub>O ha<sup>-1</sup>) was applied at sowing time to all plots through Singlesuper phosphate and muriate of potash, respectively. The Lentil variety (DPL-62) was sown third week of November 2022, using 45 kg ha<sup>-1</sup> seed at 25 cm apart rows and harvested in second week of March 2023. All improved packages of practices were followed to raise the crop. The data on plant height and branches were recorded from the area already marked by tagged. Sample for dry matter accumulation was recorded by cutting of plants from ground level. The fresh samples were first sun dried and then kept in electric oven at 65-70°C till the constant dry weight attained. Yield attributes were recorded from 10 pods selected randomly from each plot. Grain and straw yields of lentil were recorded at harvest the harvest index was calculated as grain yield divided by total biological yield and multiplied by hundred. The uptake of nutrients was calculated as nutrient content in grain and straw multiplied by respective yield. Economics of different treatments was worked out on the basis of prevailing market prices. The data so obtained on various parameters were analysed as per standard statistical procedures. The content of N, P, and K in grain and straw was determined using standard laboratory procedures.

#### RESULTS AND DISCUSSION

##### Growth attributes

Plant height, number of branches, and dry matter accumulation of lentil were affected significantly due to application of different levels of Nitrogen and phosphorus (Table 1). Increasing levels of Nitrogen up to 45 kg N/ha increased the plant height, number of branches, and dry matter accumulation significantly over rest of the Nitrogen levels. The maximum plant height (34.47 cm), number of branches (12.20 plant<sup>-1</sup>) and dry matter accumulation (13.16 g ha<sup>-1</sup>) was recorded significantly with 45 kg N ha<sup>-1</sup> over rest of nitrogen levels but being at par with 30 kg N ha<sup>-1</sup>. Significant improvement in growth parameters with increasing the levels of nitrogen might be due to increased availability of nitrogen and its uptake by plant increased the metabolic activity and formation of meristematic tissues which improved the cell elongation and cell division which in turn improved the plant height, number of branches, and dry matter accumulation of crop. These results are supported by the findings of (Khare *et al.*, 2017), (Methew *et al.*, 2002). Plant height increased significantly with increase in phosphorus dose up to 40

- Comment [AM16]: Source
- Comment [AM17]: P ha<sup>-1</sup>
- Comment [AM18]: and Singh
- Comment [AM19]: important because
- Comment [AM20]: Where is the aim of study?

- Comment [AM21]: N ha<sup>-1</sup>
- Comment [AM22]: phosphorus
- Comment [AM23]: Randomized Complete Block Design with
- Comment [AM24]: Make sure. Look the tables.
- Comment [AM25]: n
- Comment [AM26]: p
- Comment [AM27]: singlesuper

- Comment [AM28]: Where is the statistical analysis?
- Comment [AM29]: significantly due
- Comment [AM30]: n
- Comment [AM31]: n
- Comment [AM32]: N ha<sup>-1</sup>
- Comment [AM33]: Delete because there is no significant among treatments.
- Comment [AM34]: of branch numbers
- Comment [AM35]: delete
- Comment [AM36]: delete
- Comment [AM37]: due to
- Comment [AM38]: delete
- Comment [AM39]: source
- Comment [AM40]: (Khare *et al.*, 2017; Methew *et al.*, 2002)

kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. This may be because of the fact that phosphorus encourages the cell division and root development thereby resulting in improved growth and development of the plant. The results are in close conformity with the findings of (Niriet al., 2010), (Rabbi et al., 2011), (Verma et al., 1983)

Application of phosphorus @ 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> being at par with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> but recorded significantly taller plant, higher number of branches, and dry matter accumulation as compared to without phosphorus treatment. The highest plant height (33.37 cm), number of branches (10.68 plant<sup>-1</sup>) and dry matter accumulation (9.76 q/ha) was recorded with the application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. The favourable influence of phosphorus application on growth may be due to its role in various enzymatic reactions, growth processes, hormone production and protein synthesis and also translocation of photosynthates in various plant parts leading to higher growth of crop. Similar results were obtained by (Gan et al., 2005), (Joudi et al., 2008).

#### Yield attributes

Yield attributing characters like No. of pod plant<sup>-1</sup>, number of grains pod<sup>-1</sup>, grain weight (g) and test weight (g) were affected significantly due to application of Nitrogen and phosphorus (Table 2). Crop fertilized with 45 kg N/ha produced significantly the highest number of pod plant<sup>-1</sup> (57.13), number of grains pod<sup>-1</sup> (2.80), grain weight (4.74 g) and test weight (21.80 g) over rest of the nitrogen levels, however the difference between 60 kg N ha<sup>-1</sup> and 40 kg N ha<sup>-1</sup> was recorded non-significant. Dry matter accumulation showed significant and consistent improvement with nitrogen application up to 45 kg N ha<sup>-1</sup> at all stages of crop growth. Similar findings were reported by (Hoque et al., 1994), (Kumar et al., 1993). Increase in number of pod plant<sup>-1</sup> with increasing level of nitrogen might be due to increased availability of nitrogen which enhanced the cell division. The higher availability of nitrogen with supply of N-levels enhance the meristematic tissue activity located at initiation of branching point which improved with supply of nitrogen and recorded higher no. of pod plant<sup>-1</sup> with increasing levels of nitrogen resulted more branches with increasing levels of N. Similar results were also reported by (Verma et al., 1983)

The increase supply of nitrogen leads to improve pod. The higher values of yield attributes with application of N was mainly due to increase availability of nitrogen to plant which enhanced the photosynthetic activity and its translocation to since resulted in improved yield attributes. These results are in agreement with the findings of (Kumar et al., 1993), (Methew et al., 2002) However, (Joudi et al., 2008) reported non-significant effect of N fertilizers on grain number plant<sup>-1</sup>, grain number/pod and filled pod number.

Application of phosphorus had remarkable effects on yield attributes of lentil (Table 2). The maximum values of yield attributes like no. of pod plant<sup>-1</sup> (58.10), number of grains/pod (2.65), grain weight/pod (4.11 g) and test weight (21.83 g) was obtained with application of phosphorus @ 60 kg P<sub>2</sub>O<sub>5</sub>/ha, however being on par with 40 kg ha<sup>-1</sup> phosphorus. Significant response of phosphorus application on yield attributes of lentil was due to its favourable influence on various enzymatic reactions, growth processes, hormone production, protein synthesis and also the translocation of photosynthates to reproductive parts thereby leading to higher growth and yield attributes.

Application of Nitrogen up to 45 kg N ha<sup>-1</sup> improved significantly the grain and straw yield and test levels of nitrogen (Table 2) However, the difference between 45 kg ha<sup>-1</sup> and 30 kg/ha was recorded significant. Crop fertilized with 45 kg N ha<sup>-1</sup> produced significantly highest grain yield (8.74 q ha<sup>-1</sup>) and straw yield (12.37 q ha<sup>-1</sup>) over rest of the Nitrogen levels except 30 kg N ha<sup>-1</sup>. Application of Nitrogen to crop enhanced the growth and yield attributes of wheat thus resulted higher yield.

Application of phosphorus from 0 to 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> increased significantly the grain yield and straw yield. The highest grain yield (8.74 q ha<sup>-1</sup>) and straw yield (11.60 q ha<sup>-1</sup>) was recorded with application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, however the difference between 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> was found significant regarding to grain yield and straw yield. The seed yield was also increased significantly up to 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> as the seed yield is directly dependent on the yield attributing characters. The straw yield also showed the same trend like the seed yield. These results are in conformity with the findings of (Sharma et al., 2004), (Mandal et al., 2001), (Gan et al.,

Comment [AM41]: and Kalra

Comment [AM42]: Move down.

Comment [AM43]: (Verma et al., 1983; Niriet al., 2010; Rabbi et al., 2011)

Comment [AM44]: Without phosphorus

Comment [AM45]: delete

Comment [AM46]: qha<sup>-1</sup>

Comment [AM47]: source

Comment [AM48]: (Gan et al., 2005; Joudi et al., 2008).

Comment [AM49]: n

Comment [AM50]: N ha<sup>-1</sup>

Comment [AM51]: except in number of bods

Comment [AM52]: delete

Comment [AM53]: and Haq

Comment [AM54]: (Kumar et al., 1993; Hoque et al., 1994).

Comment [AM55]: source

Comment [AM56]: and Kalra

Comment [AM57]: and Agarwal

Comment [AM58]: rearrange. Rewrite.

Comment [AM59]: The maximum

Comment [AM60]: ??????

Comment [AM61]: Source.

Comment [AM62]: Move up

Comment [AM63]: ???????

Comment [AM64]: Where is this characteristic in tables?????

Comment [AM65]: ???????

Comment [AM66]: and Sing

Comment [AM67]: and Majumdar

2005], (Sharma *et al.*, 1999), (Rabbi *et al.*, 2011).

**Comment [AM68]:** and Singh

**Comment [AM69]:** Same note above.

**Comment [AM70]:** Where is the interaction treatments between N and P levels???

UNDER PEER REVIEW

**Table1. Growth attributes of lentil affected by levels of nitrogen and phosphorus**

Treatments	Plant height(cm)	Number of Branches plant <sup>-1</sup>	Dry matter accumulation(q ha <sup>-1</sup> )
<b>Level of Nitrogen (Kg ha<sup>-1</sup>)</b>			
N0-0	28.98	8.34	10.39
N1-15	30.29	9.10	12.05
N2-30	33.54	10.84	12.87
N3-45	34.47	12.20	13.16
SEm±	1.20	0.48	0.10
CDat5%	NS	1.34	0.30
<b>Level of phosphorus (Kg ha<sup>-1</sup>)</b>			
P0-0	30.34	8.10	9.22
P1-20	32.99	9.30	9.32
P2-40	33.02	1.51	9.60
P3-60	33.37	10.68	9.76
SEm±	0.94	0.48	0.10
CDat5%	2.83	1.34	0.30

Comment [AM71]: 10.51

**Table2. Yield attributes characters and yield of lentil affected by levels of nitrogen and Phosphorus**

Treatments	No. of pod/plant(g/m)	No. of grain/pod	Grain weight/plant(g)	Test weight(ht(g))	Grain yield(d(q/ha))	Straw yield(q/ha)
<b>Level of Nitrogen (Kg ha<sup>-1</sup>)</b>						
N0-0	30.90	1.61	2.50	16.76	5.30	7.55
N1-15	53.71	2.04	3.11	18.79	7.42	10.21
N2-30	56.87	2.76	4.31	21.71	9.28	11.35
N3-45	57.13	2.80	4.74	21.80	9.30	12.37
SEm±	0.08	0.018	0.16	0.037	0.86	1.14
CDat5%	0.25	0.053	0.48	0.107	2.52	3.31
<b>Level of phosphorus (Kg ha<sup>-1</sup>)</b>						
P0-0	34.95	1.41	2.57	16.90	5.70	7.36
P1-20	26.03	2.34	3.56	18.16	7.04	9.03
P2-40	57.93	2.51	3.85	21.79	8.72	11.58
P3-60	58.10	2.65	4.11	21.83	8.74	11.60
SEm±	0.089	0.01	0.09	0.037	0.86	1.14
CDat5%	0.30	0.30	0.27	0.107	2.52	5.31

### Nutrient Content and uptake-

Data revealed that crop fertilized with 45 kg N ha<sup>-1</sup> being at par with 30 kg N ha<sup>-1</sup> but recorded significantly the maximum uptake (kg ha<sup>-1</sup>) of N, P, in grain (19.62 kg ha<sup>-1</sup>, 4.60 kg ha<sup>-1</sup>), and in straw (21.23 kg ha<sup>-1</sup>, 5.65 kg ha<sup>-1</sup>), respectively over rest of phosphorus levels (Table 3) The uptake of nitrogen and phosphorus by seed and straw increased significantly at higher levels of nitrogen. This could be attributed to better fixation of nitrogen by the rhizobium at higher nitrogen levels. Earlier (Singh *et al.*, 2005), also reported similar findings. They further reported that there was an increase in root growth and development distributed widely up to different depths and thus creating a proper gradient for N absorption at the root soil interface.

Comment [AM72]: source

Comment [AM73]: source

### Economics

The economics of different treatment combinations was worked out on the basis of input-output analysis. The data pertaining to economics have been presented in Table 4 various components of economics are being described here as under.

The data presented in table 4 regarding the cost of cultivation make it clear that the highest cost of cultivation (Rs. 20254.58), Gross return (Rs. 56256), Net return (Rs. 36001.42) and Benefit cost ratio (1.77) was recorded in treatment combination where 45 kg N ha<sup>-1</sup> was applied with 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>.

The lowest cost of cultivation (Rs. 19435.58), Gross return (26422), Net return (6986.42) and Benefit cost ratio (0.35) was recorded in treatment combination where 0 kg N ha<sup>-1</sup> was applied with 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>.

UNDER PEER REVIEW

**Table 3** N and P uptake by lentil crop was affected by different levels of nitrogen and phosphorus

Treatment	NUptake(kg ha <sup>-1</sup> ) by grain	NUptake(kg ha <sup>-1</sup> ) by straw	PUptake(kg ha <sup>-1</sup> ) by grain	PUptake(kg ha <sup>-1</sup> ) by Straw
<b>Level of Nitrogen (Kg ha<sup>-1</sup>)</b>				
N0	18.03	16.42	2.48	4.34
N15	18.29	19.03	3.91	4.03
N30	18.56	20.20	4.29	4.25
N45	19.62	21.23	4.60	5.65
Sem±	0.53	0.51	0.17	0.20
CD at 5 %	1.10	1.05	0.35	0.42
<b>Level of Phosphorus (Kg ha<sup>-1</sup>)</b>				
P0	18.01	17.31	3.38	3.61
P20	18.38	19.08	4.09	4.80
P40	18.52	20.12	5.36	5.45
P60	19.05	20.31	6.67	5.64
Sem±	0.53	0.51	0.17	0.20
CD at 5 %	1.10	1.05	0.35	0.42

**Table 4** Economics was affected by different treatment combination of lentil crop

Treatments	Total cost of cultivation (Rs. ha <sup>-1</sup> )	Gross income (Rs. ha <sup>-1</sup> )	Net income (Rs. ha <sup>-1</sup> )	B.C. Ratio
N0P0(T1)	19435.58	26422	6986.42	0.35
N0P1(T2)	19615.58	28920	9304.42	0.47
N0P2(T3)	19795.58	31880	12084.42	0.61
N0P3(T4)	19975.58	35418	15442.42	0.77
N1P0(T5)	19528.58	41150	21621.42	1.10
N1P1(T6)	19708.58	42372	22663.42	1.14
N1P2(T7)	19888.58	44100	24211.42	1.21
N1P3(T8)	20068.58	45260	25191.42	1.25
N2P0(T9)	19621.58	46408	26786.42	1.36
N2P1(T10)	19801.58	47560	27758.42	1.40
N2P2(T11)	19981.58	48700	28718.42	1.43
N2P3(T12)	20161.58	50376	30214.42	1.49
N3P0(T13)	19714.58	52718	33003.42	1.67
N3P1(T14)	19894.58	53890	33995.42	1.70
N3P2(T15)	20074.58	55098	35023.42	1.74
N3P3(T16)	20254.58	56256	36001.42	1.77

## CONCLUSION

It can be concluded from the above result that lentil crop should be fertilized with 45 kg N/ha along with 60 kg P<sub>2</sub>O<sub>5</sub>/ha<sup>-1</sup> to obtain high yield and more net income under central plain zone of U.P.

## REFERENCE

1. Bremer, E., Van Kessel, C. and Karamanos, R. (1989). Inoculant, phosphorus and nitrogen responses of lentil. *Can. J. Plant Sci.*, 69 :691-701.
2. Deol, M.S. (2007). Effect of Phosphate Solubilizing Bacteria (PSB) farmyard manure and phosphorus on growth and yield of lentil (*Lens culinaris* M.). Ph.D. Thesis, Department of Agronomy, G.B. Pant University of Agriculture & Technology, Pantnagar, India.
3. Hoque, M.M. and Haq, M.F. (1994). Rhizobial inoculation and fertilization in lentil in Bangladesh. *Lens Newsletter*, 21(2) :29-30
4. Kumar, P. and Agarwal, J.P. (1993). Response of lentil (*Lens esculentus*) to rhizobium inoculation, nitrogen and phosphorus fertilization. *Indian J. Agronomy*, 38:318-20.
5. Gan, Y., Hanson, K. G., Zentner, R. P., Selles, F. and McDonald, C. L. (2005). Response of lentil to microbial inoculation and low rates of fertilization in the semiarid Canadian prairies. *Canadian Journal of Plant Science*, 847-855.
6. Joudi, F., Tobeh, A., Ebadi, A., Jamaati Soomarin, S. and Zabihi Mahmoodabad, R. (2008). Study of yield and nitrogen use and recovery efficiencies in lentil genotypes as affected by nitrogen. *Plant Ecophysiology*, 2 :87-90.
7. Khare, M. M. S., Singh, V. P. and Kumar, A. (2017). Studies on effect of phosphorus levels on growth and yield of rabi pulse. *International Journal Pure App. Biosci.* 5 (4): 800-808
8. Methew, H., Shah, S. H. and Nazir, S. M. (2002). Differential genotypic response to phosphorus application in lentil (*Lens culinaris* Medic). *International Journal of Agriculture and Biology*, 4(1):61-63.
9. Mandal, K.G. and D.K. Majumdar. (2001). Agro-physiological characteristics of lentil in relation to irrigation, nitrogen and plant density. *J. Interacademia*, 5(2) :156-161.
10. Muhammad, H., Shah, S.H. and Nazir, S.M. (2002). Differential genotypic response to phosphorus application in lentil (*Lens culinaris* Medic). *International Journal of Agriculture and Biology*, 4(1):61-63
11. Nakhzari Moghadam, A., and Ramroudi, M. (2003). Effect of planting date and nitrogen rate on yield and yield components of lentil. *Journal of Agriculture and Resource Science*, 9(4):33-41.
12. Niri, H.H., Tobeh, A., Gholipouri, A., Mostafaei, H. and Jamaati-e-Somarin, S. (2010). Effect of nitrogen and phosphorus rates on fertilizer use efficiency in lentil. *World Appl Sci. Journal*, 9:1043-1046.
13. Rabbi, A. K. M. Z., Paul, A. K. and Sarker, J. R. (2011). Effect of nitrogen and molybdenum on the growth and yield of garden pea (*Pisum sativum* L.). *IJBSM*, 2(2):230-235
14. Singh, D., 2017. Effect of sources of phosphorus on growth, yield and nutrient uptake in pea (*Pisum sativum* L.) *Annals of Plant and Soil Research*. 19(2):240-242.
15. Singh, A.K.; Singh, B. and Singh, H.C. (2005). Response of chickpea (*Cicer arietinum* L.) to fertilizer phosphorus and phosphorus application under rainfed condition of eastern U.P. *Indian J. Dryland Agric. Res. & Dev.*, 20 (2):114-117.

Comment [AM74]: *Lens culinaris*

Comment [AM75]: *Lens esculentus*

Comment [AM76]: Plant Ecophysiology

Comment [AM77]: *Lens culinaris*

Comment [AM78]: I didn't find it in manuscript body.

Comment [AM79]: I didn't find it in manuscript body.

Comment [AM80]: *Pisum sativum*

Comment [AM81]: I didn't find it in manuscript body.

Comment [AM82]: *Cicer arietinum*

16. Sharma, B.C. and Singh, S.C. (2004). Integrated nutrient management in lentil. *Advances in Plant Sciences*, 17(1):19-5-19
17. Sharma, B. B. and Singh, R.R. (1999). Rooting and yield pattern in lentil under different rates of seedling, seed inoculation, nitrogen and phosphorus fertilization. *Legume Research*, 9(2):69-72
18. Togay, Y., Togay, N., Dogan, Y. and Ciftci, V. (2005). Effects of nitrogen levels forms on the yield and yield components of lentil (*Lens culinaris Medic.*). *Asian Journal of Plant Sciences*, 4(1):6466.
19. Verma, V.S. and G.S. Kalra, 1983. Effect of different levels of irrigation, N and P on growth and yield of lentil. *Indian J. Agric. Sci.*, 17(3): 124-128
20. Verma, V. and Kalera, G.S. (1983). Effect of different levels of irrigation, N and P on growth and yield of lentil. *Indian Journal of Agricultural Science*, 17(3):124-128
21. Veeresh, N.K. (2003). Response of French bean (*Phaseolus vulgaris L.*) to fertilizer levels in Northern Transitional Zone of Karnataka. M.Sc. (Agri.) Thesis, Univ. Agric. Sci., Dharwad. p.37-79.
22. Zeidan, M.S. (2007). Effect of organic manure and phosphorus fertilizers on growth, yield and quality of lentil plants in sandy soil. *Research Journal of Agriculture and Biological Sciences*, 3(6):748-752.

**Comment [AM83]:** Repeater.

**Comment [AM84]:** I didn't find it in manuscript body.

**Comment [AM85]:** I didn't find it in manuscript body.