

Impact of Crop Load on Growth, Yield and Physical Parameters in apple cv. Gala Redlum

ABSTRACT

The study was conducted at the experimental field of the Division of Fruit Science, SKUAST-Kashmir, during the 2021-2022 period. The objective was to assess the influence of varying crop loads on growth, yield, quality, and return bloom, with a focus on trunk cross-sectional area (TCSA). Additionally, the study aimed to determine the effects of different crop loads on leaf and fruit nutritional status. The experiment, organized in a randomized complete block design (RCBD) with three replications, utilized 4-year-old "Gala Redlum" plants. The trees were hand-thinned to 4, 6, 8, 10 fruits per cm² of TCSA, with a control group undergoing no thinning. Results indicated that the highest values for annual shoot length (36.40 cm), return bloom (111.50 flower clusters/tree), fruit length (68.49 mm), fruit diameter (75.07 mm), and fruit weight (191.62 g) were observed in the C2 treatment (4 fruits per cm² of TCSA). Following closely, the C3 treatment (6 fruits per cm² of TCSA) exhibited substantial values for the aforementioned parameters. Furthermore, the S2 treatment (10-12 cm² TCSA) recorded maximum annual shoot length (30.63 cm), return bloom (64.13 flower clusters/tree), fruit length (61.74 mm), fruit diameter (69.18 mm), and fruit weight (168.16 g). Notably, the combined treatment C2S2 yielded the highest return bloom (114.00 flower clusters/tree) and fruit characteristics (fruit length, fruit diameter, and fruit weight).

Keywords: Gala Redlum, hand thinning, TCSA, return bloom, yield.

1. INTRODUCTION

The cultivated apple (*Malus x domestica* Borkh.) holds the distinction of being the premier table fruit globally, often referred to as the "King of temperate fruits." Originating in the temperate region of western Asia, between the Black Sea and Caspian Sea, the cultivated apple belongs to the Rosaceae family and is an interspecific hybrid complex of allopolyploid origin, with *Malus sieversii* identified as a progenitor species. A vital temperate fruit in the northwestern Himalayan region, apple cultivation is significant in over 90 countries worldwide, covering 4,717,384 ha and producing 79,139,368 MT annually [1]. In India, it spans 309,000 ha, with Jammu and Kashmir leading in production [2]

Despite being the primary fruit of the Kashmir valley, the productivity of apples in Jammu and Kashmir, at 11 MT/ha, falls short of international benchmarks. Transitioning from low to high-density orcharding, particularly with the introduction of exotic varieties, poses challenges in achieving desired size and quality if not thinned properly. High-density orcharding, coupled with excessive crop load, contributes to biennial bearing, leading to economic losses through reduced fruit size and sporadic yields especially in early years. The reliance on chemical thinning has been prominent, but the need for environmentally acceptable alternatives prompts a focus on precision crop load management, integrating precision pruning, chemical, and hand thinning.

The prevailing industry norm involves chemical thinning during bloom and post-bloom, followed by hand thinning. However, the weather-dependent response to chemical thinning, coupled with varying cultivar sensitivity, underscores the need for environmentally friendly approaches. Crop manipulation methods, such as mechanical thinning, hand flower thinning, and bud extinction during winter, have been explored [3]. The study aims to assess the impact of different crop loads on growth, yield, and physical parameters concerning trunk cross-sectional area in high-density apple cultivar Gala Redlum, considering the significance of effective thinning practices for optimizing fruit quality, marketable size, and addressing alternate bearing. Keeping in view the importance and economic value of thinning the investigation therefore was carried out to study the effect of different crop loads on growth, yield and physical parameters vis-à-vis trunk cross-sectional area in high density apple cv. Gala Redlum.

MATERIALS AND METHODS

The experiment was conducted on four-year-old Gala Redlum apple trees grafted onto M-9 rootstock at the Experimental Farm of the Division of Fruit Science, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, Shalimar campus, during the agricultural year 2021-22. Adhering to recommended package practices, including fertilizer application, fertigation, and irrigation, the trial aimed to assess the impact of two key factors: hand thinning and trunk cross-sectional area (TCSA). A Randomized Complete Block Design (RCBD) with three replications

was employed for the factorial experiment. The factorial design incorporated various combinations of hand thinning and TCSA as distinct treatment factors. This approach enhances the experimental robustness by systematically varying these factors and considering their individual and interactive effects on the growth, yield, and quality parameters of Gala Redlum apple trees. The meticulous application of recommended horticultural practices ensured the experiment's precision and reliability. The experiment involved crop loads of 4, 6, 8, 10 fruits per cm² of TCSA, labeled as C2, C3, C4 and C5 respectively, with a non-thinned control group as C1. The orchard trees were categorized into two groups labeled as S1 (8-10 cm² TCSA) and S2 (10-12 cm² TCSA) based on trunk cross-sectional area measurements. The Trunk Cross-sectional Area was determined as per Westwood [4]:

$$\text{Trunk Cross sectional Area (T.T.S.C.A) in cm}^2 = [\text{Trunk girth of scion (cm)}]^2 / 3.14.$$

Return Bloom was assessed during the month of April in the subsequent year, with a comprehensive evaluation of return bloom was conducted by quantifying the total number of flower clusters per tree at full bloom. This meticulous count provides valuable insights into the regenerative capacity of the apple trees, indicating the potential for subsequent flowering and fruiting. The return bloom metric serves as a pivotal indicator of the orchard's ability to consistently yield floral clusters, influencing the overall productivity and sustainability of the high-density apple cultivar Gala Redlum. Plant Height (m) was measured from base to terminal growth point. Annual shoot length (cm) was determined by recording the growth of the current season of the four selected branches on four sides and the mean increment growth was calculated. Fruit length (mm) of fifteen fruits per tree was measured with the help of a digital vernier calliper and average fruit length was expressed in millimetres (mm). Fruit diameter (mm) of fifteen fruits per tree was measured with the help of a digital vernier calliper and average diameter was expressed in millimetres (mm). Length: Diameter ratio was calculated by dividing length by respective diameter of fruit sample. Fruit weight from different treatments was weighed individually on a sensitive monopan balance and the average fruit weight was recorded in grams (g) per fruit. Precision in these measurements was ensured through the use of digital vernier calipers and a sensitive monopan balance. The assessment of all the key parameters in the study is grounded in scientific rigor.

RESULTS AND DISCUSSION

Plant Height (m):

The impact of crop load on plant height in apple cultivar Gala Redlum was assessed and presented in Table 1. Notably, the results indicate that crop load does not exert a significant influence on plant height (Table 1). However, plant height does exhibit a significant correlation with trunk cross-sectional area (TCSA). The highest plant height (3.27 m) was observed in S2 (10-12 cm² TCSA), while the lowest (2.86 m) was recorded in S1 (8-10 cm² TCSA).

Table 1: Effect of crop load on plant height (m) in apple cv. Gala Redlum

	TCSA		Mean
	S1 (8-10 cm ² TCSA)	S2 (10-12 cm ² TCSA)	
C1 (No thinning)	2.82	3.27	3.04
C2 (4 fruits per cm² of TCSA)	2.91	3.34	3.12
C3 (6 fruits per cm² of TCSA)	2.95	3.28	3.11
C4 (8 fruits per cm² of TCSA)	2.81	3.26	3.03
C5 (10 fruits per cm² of TCSA)	2.85	3.21	3.01
Mean	2.86	3.27	
C.D (p≤0.05)			
Trunk cross sectional area (S) = 0.1239			
Crop load (C) = N.S (S×C) = N.S			

The Table further, illustrates that the interaction effect of crop load and trunk cross-sectional area (TCSA) on plant height was non-significant. These findings align with the research conducted by Serra [5], who similarly reported that plant height remained unaffected by variations in crop load.

The examination of plant height (in meters) further revealed a significant association with trunk cross-sectional area (TCSA). Specifically, the maximum plant height was observed under S2 (10-12 cm² TCSA), whereas the minimum was recorded under S1 (8-10 cm² TCSA). This correlation suggests that an increase in plant height corresponds to an increase in trunk cross-sectional area (TCSA). The rationale behind this phenomenon may be attributed to the direct relationship between TCSA and the transport of nutrients from the root to various parts of the plant. Additionally, TCSA influences the distribution of food materials from the production site to the utilization site, thereby impacting vegetative growth [6]. These outcomes align with the findings of Srivastava *et al.* [7] who reported a positive correlation between plant height and TCSA.

While crop load does not appear to significantly affect plant height in the studied apple cultivar Gala Redlum, trunk cross-sectional area emerges as a crucial factor influencing plant height. The positive correlation between plant height and TCSA underscores the importance of nutrient transport and food material distribution in shaping vegetative growth.

Return Bloom (Number of Flower Clusters per Tree):

Table 2 presents the significant impact of both crop load and trunk cross-sectional area on return bloom. The data reveals that C2 (4 fruits per cm² of TCSA) exhibits the highest return bloom, recording 111.50 flower clusters per tree, followed closely by C3 (6 fruits per cm² of TCSA) with 88.16 flower clusters per tree. In contrast, the control group (no thinning) shows the lowest return bloom at 12.16 flower clusters per tree. Furthermore, return bloom is significantly influenced by varying trunk cross-sectional areas, with S2 (10-12 cm² TCSA) yielding the maximum at 64.13 flower clusters per tree, and S1 (8-10 cm² TCSA) resulting in the minimum at 60.66 flower clusters per tree.

Table 2: Effect of crop load on return bloom (number of flower clusters per tree) in apple cv. Gala Redlum

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C ₁ (No thinning)	10.33	14.00	12.16
C ₂ (4 fruits per cm ² of TCSA)	109.00	114.00	111.50
C ₃ (6 fruits per cm ² of TCSA)	87.33	89.00	88.16
C ₄ (8 fruits per cm ² of TCSA)	72.00	75.66	73.83
C ₅ (10 fruits per cm ² of TCSA)	24.66	28.00	26.33
Mean	60.66	64.13	
C.D (p≤0.05)			
Trunk cross sectional area (S) = 0.98			
Crop load (C) = 1.56 (S×C) = N.S			

The analysis of the interaction effect between crop load and trunk cross-sectional area (TCSA) on return bloom indicates a non-significant impact. However, specific treatment combinations do exhibit distinct effects. The C2S2 treatment combination records the highest return bloom at 114.00 flower clusters per tree, followed by C2S1 with 109.00 flower clusters per tree. Conversely, the minimum return bloom of 10.33 flower clusters per tree is observed in the C1S1 treatment combination.

The observed increase in return bloom with an increase in trunk cross-sectional area (TCSA) can be attributed to the direct relationship between TCSA and the transport of nutrients from the root to various parts of the plant. Additionally, TCSA influences the distribution of food materials from the production site to the utilization site [6]. This, in turn, ultimately impacts return bloom. These findings underscore the importance of understanding the physiological mechanisms associated with trunk cross-sectional area in influencing the flowering patterns of apple trees.

The results highlight the significant roles of both crop load and trunk cross-sectional area in shaping return bloom in the studied apple cultivar. The intricate relationship between these factors emphasizes the need for a holistic understanding of orchard management practices to optimize flowering and, consequently, fruit production.

Annual Shoot Length (cm):

Table-3 presents the findings on annual shoot length, indicating that both crop load and trunk cross-sectional area significantly influence this parameter. The highest annual shoot length, measuring 36.40 cm, was observed in C2 (4 fruits per cm² of TCSA), followed by C3 (6 fruits per cm² of TCSA) with an annual shoot length of 32.98 cm. Conversely, the control group (no thinning) exhibited the minimum annual shoot length at 23.49 cm. Additionally, annual shoot length was significantly affected by different trunk cross-sectional areas, with S2 (10-12 cm² TCSA) producing the maximum annual shoot length at 30.63 cm, and S1 (8-10 cm²TCSA) resulting in the minimum at 30.10 cm.

Table 3: Effect of crop load on annual shoot length (cm) in apple cv. Gala Redlum

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C₁ (No thinning)	23.15	23.84	23.49
C₂ (4 fruits per cm² of TCSA)	36.15	36.66	36.40
C₃ (6 fruits per cm² of TCSA)	32.86	33.11	32.98
C₄ (8 fruits per cm² of TCSA)	31.32	31.53	31.42
C₅ (10 fruits per cm² of TCSA)	27.04	28.03	27.54
Mean	30.10	30.63	

C.D (p≤0.05)
Trunk cross sectional area (S) = 0.017
Crop load (C) = 0.027 (S×C) = 0.038

It is evident that the interaction effect of crop load and trunk cross-sectional area (TCSA) on annual shoot length was significant. The C2S2 treatment combination yielded the maximum annual shoot length at 36.66 cm, followed by C2S1 with an annual shoot length of 36.15 cm. In contrast, the minimum annual shoot length of 23.15 cm was recorded in the C1S1 treatment combination.

The observed increase in annual shoot length with an increase in trunk cross-sectional area (TCSA) is consistent with the idea that TCSA is directly linked to nutrient transport from the root to different parts of the plant, as well as the distribution of food materials from the site of production to the site of utilization [6]. This relationship ultimately influences annual shoot length. These results align with the findings of Srivastava et al. [8], supporting the notion that annual shoot length tends to increase with an augmentation of TCSA.

The study underscores the significant roles played by both crop load and trunk cross-sectional area in shaping annual shoot length in the studied apple cultivar. The interaction effect further emphasizes the need for a nuanced understanding of these factors for effective orchard management practices, aligning with the broader goal of optimizing tree growth and productivity.

Fruit Length (cm):

Table 4 presents the significant impact of both crop load and trunk cross-sectional area on fruit length in the studied apple cultivar. The data reveals that C2 (4 fruits per cm² of TCSA) exhibits the maximum fruit length, recording 68.49 mm, followed closely by C3 (6 fruits per cm² of TCSA) with a fruit length of 65.37 mm. In contrast, the control group (no thinning) shows the minimum fruit length at 52.31 mm. Additionally, there is a significant difference in fruit length across different trunk cross-sectional areas, with S2 (10-12 cm² TCSA) producing the maximum fruit length at 61.74 mm, and S1 (8-10 cm² TCSA) resulting in the minimum at 59.88 mm.

1 **Table 4: Effect of crop load on fruit length (cm) in apple cv. Gala Redlum**

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C ₁ (No thinning)	50.31	54.31	52.31
C ₂ (4 fruits per cm ² of TCSA)	67.87	69.12	68.49
C ₃ (6 fruits per cm ² of TCSA)	64.36	66.39	65.37
C ₄ (8 fruits per cm ² of TCSA)	62.54	63.48	63.01
C ₅ (10 fruits per cm ² of TCSA)	54.31	55.42	54.86
Mean	59.88	61.74	

C.D (p≤0.05)
 Trunk cross sectional area (S) = 0.327
 Crop load (C) = 0.516 (S×C) = 0.730

2

3 The influence of both crop load and trunk cross-sectional area on fruit length, highlights the importance of these
 4 factors in determining the size and morphology of the fruit. The observed variation in fruit length can be attributed to
 5 the direct impact of crop load on the resources available for fruit development, as well as the role of trunk cross-
 6 sectional area in nutrient transport and distribution within the tree.

7 Results of the interaction studies regarding different crop loads and trunk cross-sectional areas (TCSA) with respect
 8 to fruit length were found to be significant. The data indicates that the C2S2 treatment combination resulted in the
 9 maximum fruit length, recording 69.12 mm. This was followed closely by the C2S1 treatment combination with a fruit
 10 length of 67.87 mm. In contrast, the minimum fruit length of 50.31 mm was recorded in the C1S1 treatment
 11 combination.

12 Understanding the interaction effects between crop load and trunk cross-sectional area is essential for devising
 13 effective orchard management strategies aimed at achieving desirable fruit characteristics. The identified treatment
 14 combinations with optimal fruit length can guide future cultivation practices for maximizing fruit quality and overall
 15 orchard yield.

16 **Fruit Diameter (cm):**

17 Results from Table 5 demonstrate the significant impact of both crop load and trunk cross-sectional area (TCSA) on
 18 fruit diameter in the studied apple cultivar. The maximum fruit diameter of 75.07 mm was observed in C2 (4 fruits per
 19 cm² of TCSA), followed by 72.13 mm in C3 (6 fruits per cm² of TCSA), while the minimum diameter of 60.98 mm was
 20 recorded in the control group (no thinning). Similarly, different trunk cross-sectional areas significantly influenced fruit
 21 diameter, with S2 (10-12 cm² TCSA) yielding the maximum diameter of 69.18 mm and S1 (8-10 cm² TCSA) resulting
 22 in the minimum at 67.62 mm.

23 **Table 5: Effect of crop load on fruit diameter (cm) in apple cv. Gala Redlum**

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C ₁ (No thinning)	59.31	62.64	60.98
C ₂ (4 fruits per cm ² of TCSA)	74.45	75.70	75.07
C ₃ (6 fruits per cm ² of TCSA)	71.12	73.15	72.13
C ₄ (8 fruits per cm ² of TCSA)	69.57	70.27	69.92
C ₅ (10 fruits per cm ² of TCSA)	63.66	64.14	63.90
Mean	67.62	69.18	

C.D (p≤0.05)
 Trunk cross sectional area (S) = 0.134

$$\text{Crop load (C)} = 0.212 \quad (\text{S} \times \text{C}) = 0.299$$

Importantly, the interaction effect of crop load and trunk cross-sectional area on fruit diameter was found to be significant. The C2S2 treatment combination resulted in the maximum fruit diameter of 75.70 mm, followed by 74.45 mm in the C2S1 treatment combination. Conversely, the minimum fruit diameter of 59.31 mm was recorded in the C1S1 treatment combination. These results emphasize the need to consider both crop load and trunk cross-sectional area in orchard management to optimize fruit diameter. The identified treatment combinations with optimal fruit diameter can guide cultivation practices for enhancing fruit quality and overall orchard yield.

Length: Diameter Ratio

The data presented in Table 6 on the length-to-diameter ratio of fruits, indicates the significant influence of both crop load and trunk cross-sectional area (TCSA). The maximum length-to-diameter ratio (0.912) was observed in C2 (4 fruits per cm² of TCSA), while the minimum ratio (0.858) was recorded in the control group (no thinning). Additionally, different trunk cross-sectional areas significantly affected the length-to-diameter ratio, with S2 (10-12 cm² TCSA) showing the maximum ratio of 0.891, and S1 (8-10 cm² TCSA) resulting in the minimum at 0.883.

Table 6: Effect of crop load on Length: Diameter ratio in apple cv. Gala Redlum

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C ₁ (No thinning)	0.848	0.867	0.858
C ₂ (4 fruits per cm ² of TCSA)	0.912	0.913	0.912
C ₃ (6 fruits per cm ² of TCSA)	0.905	0.908	0.906
C ₄ (8 fruits per cm ² of TCSA)	0.899	0.903	0.901
C ₅ (10 fruits per cm ² of TCSA)	0.853	0.864	0.859
Mean	0.883	0.891	
C.D (p≤0.05)			
Trunk cross sectional area (S) = 0.005			
Crop load (C) = 0.008 (S×C) = 0.011			

The interaction studies reveal significant effects of different crop loads and TCSA on the length-to-diameter ratio. The C2S2 treatment combination displayed the maximum ratio (0.913), followed by 0.912 in the C2S1 treatment combination. In contrast, the minimum ratio (0.848) was recorded in the C1S1 treatment combination.

These findings highlight the importance of considering both crop load and trunk cross-sectional area in orchard management to optimize the length-to-diameter ratio of fruits. The identified treatment combinations with optimal ratios can guide cultivation practices for enhancing fruit quality and overall orchard yield.

Fruit weight (g):

Table-7 illustrates the significant impact of both crop load and trunk cross-sectional area on fruit weight in the studied apple cultivar. Maximum fruit weight (191.62 g) was observed in C2 (4 fruits per cm² of TCSA), followed by 180.78 g in C3 (6 fruits per cm² of TCSA), while the minimum weight (137.14 g) was recorded in the control group (no thinning). Different trunk cross-sectional areas also significantly influenced fruit weight, with S2 (10-12 cm² TCSA) yielding the maximum weight of 168.16 g, and S1 (8-10 cm² TCSA) resulting in the minimum at 165.78 g.

Table 7: Effect of crop load on fruit weight ratio in apple cv. Gala Redlum

Crop load	TCSA		Mean
	S ₁ (8-10 cm ² TCSA)	S ₂ (10-12 cm ² TCSA)	
C ₁ (No thinning)	136.09	138.19	137.14
C ₂ (4 fruits per cm ² of TCSA)	190.36	192.89	191.62
C ₃ (6 fruits per cm ² of TCSA)	179.07	182.49	180.78
C ₄ (8 fruits per cm ² of TCSA)	171.33	172.19	171.76
C ₅ (10 fruits per cm ² of TCSA)	152.08	155.05	153.56
Mean	165.78	168.16	
C.D (p≤0.05)			

Trunk cross sectional area (S) = 0.48
Crop load (C) = 0.76 (S×C) = 1.08

52

53 The interaction effect of crop load and trunk cross-sectional area on fruit weight was found to be significant. The
54 C2S2 treatment combination resulted in the maximum fruit weight (192.89 g), followed by 190.36 g in the C2S1
55 treatment combination. In contrast, the minimum fruit weight (136.09 g) was recorded in the C1S1 treatment
56 combination.

57 Thinning, as evidenced by the increase in fruit size and weight, is attributed to factors such as a reduced number of
58 fruits per tree, an increased leaf to fruit ratio, enhanced availability of photosynthates, and decreased nutritional
59 competition among developing fruits. This leads to greater translocation of assimilates to the remaining fruits, resulting
60 in increased size and weight. These findings align with prior research, including Henriod et al. [9], Link [10], Knight
61 [11], Koike and Ono [12], and Serra [5], who similarly reported the positive impact of thinning on fruit size and weight.

62 CONCLUSION

63 In conclusion, the study underscores the significance of thinning, particularly in high-density orchards, for achieving
64 both quality yield in the current year and satisfactory return bloom in the following year for the "Gala Redlum" apple
65 cultivar. Moreover, the data suggests that maintaining an optimal fruit quality and consistent bloom is achievable at a
66 crop load threshold of around 6 fruits per cm² of TCSA.

67 Conflict of Interest:

68 The authors declare no conflict of interest.

69 REFERENCES

70

- 71 [1]. FAOSTAT, 2019. Agricultural Statistical database, available online at <http://faostat.fao.org>.
72 [2]. Anonymous, 2020. District wise area and production of major horticultural crops in Jammu and Kashmir state for
73 the year 2018-19. Department of Horticulture, Jammu and Kashmir.
74 [3]. Costa, G. D., Blanke, M. M. and Widmer, A. 2013. Principles of thinning in fruit tree crops Needs and novelties.
75 *Acta Horticulturae* **998**: 17–26.
76 [4]. Westwood, M. N. 1993. Temperate Zone Pomology. Timber Press, Portland, Oregon. Revised Edition pp. 223.
77 [5]. Serra, S. 2016. Crop load influences fruit quality, nutritional balance, and return bloom in 'Honeycrisp' apple. *Hort*
78 *Science* **51**(3): 236-244.
79 [6]. Hartmann, H. T. and Kester, D. E. 2002. Plant propagation: principles and practices (4th Ed.). Prentice Hall of
80 India Private Limited, New Delhi, pp. 377-79.
81 [7]. Srivastava, K. K., Kumar, D., Singh, S. R. and Sharma, O. C. 2019b. Effect of cultivars on tree growth, yield and
82 quality attributes of apple on espalier architecture under high density planting system. *Journal of Horticultural*
83 *Sciences* **14**(1): 20-25.
84 [8]. Srivastava, K. K., Kumar, D., Singh, S. R., Chand, O. and Pandey, S. 2019a. Effect of tree architecture and variety
85 on growth and yield attributes of apple under UHDP. *Journal of Crop and Weed* **15**(1): 50-55.
86 [9]. Henriod, R. E., Tustin, D. S., Breen, K. C., Oliver, M., Dayatilake, G. A., Palmer, J. W., Seymour, S., Diack, R. and
87 Johnston, J. 2008. Thinning effects on "Scifresh" apple fruit quality at harvest and after storage. In IX International
88 *Symposium on Integrating Canopy, Rootstock and Environmental Physiology in Orchard Systems* **903**: 783-788.
89 [10]. Link, H. 2000. Significance of flower and fruit thinning on fruit quality. *Plant Growth Regulators* **31**(1): 17-26.
90 [11]. Knight, J. N. 1980. Fruit thinning of the apple cultivar Cox's Orange Pippin. *Journal of Horticultural Science*
91 **55**(39): 267-273.
92 [12]. Koike, H. and Ono, T. 1998. Optimum crop load for Fuji apples in Japan. *Compact Fruit Tree* **31**(1): 1-9.
93