

Original Research Article

FACTORS ASSOCIATED WITH FEBRILE TREATMENT-SEEKING BEHAVIOURS AMONG EXPECTANT MOTHERS IN SSEKANYONYI SUB-COUNTY, MITYANA DISTRICT, UGANDA

Abstract

Background: Malaria manifested in febrile illnesses is a serious health problem that contributes greatly to morbidity and mortality in Uganda. This is no different from the situation in Mityana District where 45% of pregnant mothers still do not seek treatment in health facilities when they experience febrile illnesses while 4 in 10 pregnancy deaths are malaria-related. Therefore, against this background, a study was undertaken to identify factors associated with treatment-seeking behavior among pregnant women suffering from febrile illnesses assumed to be malaria in Ssekanyonyi Sub County Mityana District, Uganda.

Methods: A cross-sectional study design in which questionnaires and key informant guides were administered to study subjects as part of the data collection process. SPSS software version 20.0 was used for data analysis and a Logistic Regression model was fitted to identify factors that independently influenced the health-seeking behavior of pregnant mothers.

Results: 198 expectant mothers were enrolled in the study. Out of these, 42.9% were aged 15-25 years, 73.7% had achieved Secondary education, and 46.5% were married. The treatment-seeking behavior was found to be standing at only 56.6%. Among the different factors studied, health education on malaria (AOR = 3.68, P = 0.000), the attitude of midwives (AOR = 1.45, P = 0.003), patient care (AOR = 0.33, P = 0.030), and attitude of the pregnant mother (AOR = 5.38, P = 0.000) were found to be statistically significantly associated with treatment seeking behavior among pregnant mothers with febrile illnesses.

Conclusions: From the above results, it was concluded that the following factors: health education on malaria, the attitude of midwives, patient care and attitude of the pregnant mothers need to be given attention in a bid to reduce febrile illnesses. The Ministry of Health and other responsible stakeholders need to reinforce awareness programs on health education among women of reproductive age about the danger of febrile illnesses during pregnancy and also help midwives address issues related to their attitude towards pregnant mothers.

Keywords: Health Based and Personal Factors on Treatment Seeking Behaviour, pregnant women, febrile illnesses.

Background

Treatment-seeking behavior for febrile illnesses assumed to be malaria among pregnant women is a great concern than other population groups (other women, men, and children) (UNICEF, 2009 and WHO, 2016). Febrile illnesses can lead to abortion, intrauterine fetal death, premature delivery, and even maternal death in case pregnant mothers do not seek treatment in time (Mvondo, James & Cambell, 2012; WHO, 2016).

Previous studies (Chepkemoi & Mutulei, 2014) indicate that factors such as insufficient health education, the attitude of midwives, and myths about sickness during pregnancy are the major hindrances to seeking treatment among pregnant mothers even though they are exposed to febrile illnesses. The majority of the mothers depend financially on their husbands for health support,

yet the majority of the vulnerable mothers are found in poverty-stricken countries and families where many use herbal medication before seeking conventional treatment (Chepkemoi & Mutulei, 2014; UNICEF, 2009 and WHO, 2016).

According to UNICEF (2009) and Chepkemoi & Mutulei (2014), health education in rural areas is said to be unconvincing yet mothers are aided to shun health units due to the bad attitude of the midwives during the antenatal visits. Most mothers in rural areas cover long distances to the health centers. There are several other factors that seem to influence treatment-seeking behavior among pregnant mothers when they experience febrile illnesses (Chepkemoi & Mutulei 2014).

WHO, (2007) estimates that globally, 70% of women access antenatal care at least once in pregnancy. This presents an opportunity for pregnant women to access several health care services including treatment of febrile illnesses assumed to be malaria. According to UNICEF, (2009) in East Asia and the Pacific, only 90% of expectant mothers seek necessary health treatment from a skilled provider while others seek help from herbalists and drug shops.

Available epidemiologic studies of malaria and febrile illnesses in pregnancy (MIP) in India by Sholapurkar, Gupta, Mahajan, (2015) found that many mothers were not seeking treatment due to low income and distance, yet malaria was associated with adverse maternal and neonatal outcomes including maternal anemia, preterm labor, stillbirths, and low birth weight. In Ethiopia, more than three-quarters of the landmass (altitude <2000 m) of the country is malarious, and pregnant women and under-five years old children are the most vulnerable groups (Musa, Salaudeen & Jimoh, 2009).

According to UNICEF, (2009), the overall burden of febrile illnesses assumed to be malaria among pregnant mothers is high and its adverse outcomes to the pregnant mother and the unborn child are widespread. Despite the growing awareness about the pregnancy-associated febrile illness and the need for treatment, research has revealed that community members still attribute malaria to bed bugs, mangoes, maize, and poor nutrition yet many mothers seem to think febrile illnesses among pregnant women is normal and may not require medical treatment (MoH, 2013). Malaria manifested in febrile illnesses is a serious health problem that contributes greatly to morbidity and mortality in Uganda (MoH, 2014). This is no different from the situation in Mityana District where 45% of pregnant mothers still do not seek treatment in health facilities when they experience febrile illnesses while 4 in 10 pregnancy deaths are malaria-related (Mityana District 2018).

Methods

Study design

The study adopted descriptive survey design with a cross sectional design to collect behavioral reactions from different respondents when they experience febrile illnesses. Both qualitative and quantitative research approaches were used. The study was descriptive to allow the researcher to discover patterns in the respondents' thinking and also to describe issues from their own point of view. Quantitative approach was used to analyze primary data from the field using descriptive statistics while qualitative approach was used to describe the state of reactions to febrile illnesses assumed to be malaria in Ssekanyonyi Sub County and discussion in relationship to interview results. The correlations design was used to establish relation between variables.

Setting

The study was carried out in Ssekanyonyi Sub County in Mityana District. Ssekanyonyi is located 20 kilometers by road after Mityana Municipality and Busunju Town council.

Sample

Sample size of 321 pregnant women was used, as derived from the target population (9532) who were estimated to be visiting the health units in the study area. Proportionate allocation sampling formula by Kothari (2004) was used to obtain the sample size from each health unit, based on the formula derived by Krejcie and Morgan (1970).

Table 1: Target Population and Sample Size

Health Center	Target Pp	Sample Size
Ssekanyonyi HC IV	4396	162
St Padre Pio HV III	2066	70
Bussunju HC III	1086	32
Bussunju Police HC II	998	29
Kassikombe HC II	986	28
Total	9532	321

However, only 198 respondents who turned up were interviewed.

Sampling

A simple random sampling procedure was employed to select 198 pregnant women suffering from febrile illnesses, while purposive sampling was used to select 20 health workers at Health center II, three (3) VHTs, and the District Health officer who acted as key informants in Ssekanyonyi Sub County Mityana District, Uganda.

Data collection

Instruments

The study used both a questionnaire and key informant interview guide as a supplementary tool for this study such that additional information was obtained from the opinion leaders and relevant people within the locality in relation to the subject of study. The key informant included; two (2) health workers at Health Center II, three (3) VHTs, and the District Health officer.

Data analysis

Data was collected, checked, and edited for completeness and accuracy. The data was then analyzed using the statistical package for social scientists (SPSS) version 17.0. Data was analyzed and interpreted in line with the objectives of the study. Descriptive and inferential analyses were made using SPSS v.17.0. The descriptive analyses were made, and results were presented using information of frequencies and percentages. While for the case of inferential statistics, the Chi-squared test and Logistic Regression analyses were appropriate for testing for association between the outcome variable and categorical factors, establishing the factors associated with treatment-seeking behavior among pregnant women suffering from febrile illnesses assumed to be malaria in Ssekanyonyi Sub-county, with the level of statistical significance set at $p \leq 0.05$; as this would help to compare the observed values in the data to the expected values that we would see if the null hypothesis is true (Ugoni & Walker, 1995). All factors with a p-value of ≤ 0.05 were considered as significantly associated with treatment-seeking behaviour, and the Null hypothesis was rejected. Factors with a P-value of >0.05 were considered not to be associated with alcohol over consumption, and the Null hypothesis was accepted.

Ethical considerations

Informed consent was obtained from the respondents prior to data collection then the researcher establish rapport and proceeded with the interview in a private quiet room. The anonymity and privacy of the participants was observed. The participants remained anonymous during the whole process of the study. The participants' information was kept confidential and only used for the purpose of the research, the local Research Ethics Committee (REC) and Uganda National Council for Science and Technology (UNCST) as entities which may have access to private information that identifies the research participants by name.

Results

A total of 189 pregnant women suffering from febrile illnesses were sampled.

Table 2: Social Demographic Characteristics of Respondents

Item	Frequency (n = 189)	Percent (%)
Age (years)		
15-25	85	42.9
26-35	71	35.9
36-45	42	21.2
Level of education		
Non formal	18	9.1
Primary	34	17.2
Secondary	146	73.7
Marital Status		
Single	12	6.1
Married	92	46.5
Divorced/Separated	10	5.1
Widowed	8	4.0
Cohabiting	76	38.4

Source: Primary Data (2020)

From results in Table 2, a higher proportion (42.9%) were in the age bracket of 15 – 25 years, majority (73.7%) were secondary level graduates, and a higher proportion (46.5%) were married.

Table 3: Factors Associated with Treatment Seeking Behaviours

Items	Frequency (n = 189)	Percent (%)
Health Education on Malaria		
Have you ever attended village seminars on health and malaria?		
Yes	44	22.2
No	154	77.8
Have health workers ever visited your home and spoke about malaria issues		
Yes	12	6.1
No	186	93.9
Attitude of midwives		

Do midwives have positive attitude towards expectant mothers?		
Yes	172	86.9
No	26	13.1
Patient Care		
Do you get warm welcome from the health workers?		
Yes	172	86.9
No	26	13.1
Do you feel you should always come back to the health center whenever you get malaria?		
Yes	168	84.8
No	30	15.2
Medical Supplies and Equipments		
Do medical workers test malaria with a kit?		
Yes	126	63.6
No	72	36.4
Have you been told that there were no malaria medicines in the health unit?		
Yes	112	56.6
No	86	43.4
Distance (km)		
less than 1 km	75	37.9
1-2 km	73	36.9
2-5 km	50	25.3
Income Level		
How much do you earn in a month?		
Less than 49,000/=	124	62.6
50,000-99,000/=	74	37.4
Attitude of the Pregnant Mother		
Adverse effects on pregnancy		
Yes	170	85.9
No	28	14.1

Source: Primary Data (2020)

From Table 3 results, majority (77.8%) had never attended any village seminar on health and malaria, while most of the respondents (93.9%) had never had health workers visiting their homes to talk about malaria issues. Majority (86.9%) reported positive attitude of midwives towards expectant mothers, majority (86.9%) reported receiving warm welcome from health workers, while majority (84.8%) reported feeling like they should be coming back to the health centre whenever they get malaria. Also, most of the respondents (63.6%) reported that the medical workers usually test malaria with a kit. Slightly above average (56.6%) reported that they had ever been told that the health unit has no malaria medicine. A higher proportion (37.9%) reported traveling less than one kilometer to the health unit to receive malaria treatment, followed by those who reported travelling 1 – 2 kilometers (36.9%), while those traveling more than 2 kilometers to the health unit were the minority (25.3%). Majority (62.6%) reported having a monthly income level of less than Uganda shillings 49,000; while most of the mothers (85.9%) reported that malaria has adverse effects on pregnancy.

Table 1: Treatment Seeking Behaviour among Pregnant Women

Item	Frequency (n = 189)	Percent (%)
Seeking Treatment		
Have you had malaria symptoms when pregnant?		
Yes	120	60.6
No	78	39.4
If yes, did you go for medical treatment immediately you felt feverish?		
Yes	112	56.6
No	8	4.0
Health care service provider		
Have you ever found no health workers at the health center?		
Yes	51	25.8
No	147	74.2
Adherence		
I observe the dosage as prescribed by the medical worker for malaria treatment.		
All the time	88	44.4
Most of the time	66	33.3
Some time	40	20.2
Never	4	2.0

Source: Primary Data (2020)

From table 4, results show that majority (60.6%) of the respondents reported having ever had malaria symptoms when pregnant, while, among these, slightly above average (56.6%) reported that they went for medical treatment immediately they felt feverish. Majority (74.2%) reported that they had never missed to find health workers at the health center when they go for medication; whereas a higher proportion (44.4%) reported that all the time they observe the dosage as prescribed by the medical worker for malaria treatment.

Table 2: Factors Associated with Treatment Seeking Behaviour

Variables	Treatment Seeking		UOR (95% CI)	p-value	AOR (95% CI)	p-value
	Sought n (%)	Not Sought n (%)				
Health education on Malaria						
Attended training	104 (86.6)	16 (13.4)	3.46 (2.56 – 3.98)	0.000**	3.68 (2.91 – 4.39)	0.000**
Did not attend training	50 (64.1)	28 (35.9)	1		1	
Attitude of midwives						
Positive	110 (91.7)	10 (8.3)	2.57 (1.88 – 3.01)	0.000**	1.45 (0.93 – 2.05)	0.003**
Negative	74 (94.9)	4 (5.1)	1		1	
Patient Care						

	Good care	115 (95.8)	5 (4.2)	2.55 (1.75 – 3.23)	0.000**	8.22 (6.45 – 10.96)	0.030**
	Not caring	70 (89.7)	8 (10.3)	1		1	
Medical Supplies and Equipment							
	Have equipment	94 (78.4)	26 (21.6)	2.52 (1.67 – 3.42)	0.000**	0.79 (0.26 – 1.31)	0.650
	No equipment	32 (41)	46 (59)	1		1	
Distance							
	Short	113 (76.3)	35 (23.6)	0.314 (0.112 – 0.754)	0.052		
	Long	36 (72)	14 (28)	1			
Income level							
	High	14 (21.8)	50 (78.1)	0.047 (0.013 – 0.101)	0.828		
	Low	113 (84.3)	21 (15.6)	1			
Attitude of the pregnant mother							
	Positive	100 (83.3)	20 (16.7)	5.53 (4.87 – 5.97)	0.000**	5.38 (4.87 – 6.17)	0.000**
	Negative	50 (64.1)	28 (35.9)	1		1	

*p < 0.05

**p < 0.01

RC=1

**Significant at 5%

From table 6, findings revealed that health-based factors had a significant association to treatment seeking behaviours among pregnant women suffering from febrile illnesses assumed to be malaria. Health education on malaria, attitude of midwives and patient care had a significant association with treatment seeking behaviours among pregnant women suffering from febrile illnesses assumed to be malaria. The adjusted odds ratio of health education on malaria [AOR = 3.68, (CI 95% = 2.91 – 4.39), p=0.000] implies that the odds of treatment seeking behaviours among pregnant women who had attended training on malaria were about 4 times higher compared to those who did not attend trainings on malaria. While the adjusted odds ratio of attitude of midwives [AOR = 1.45, (CI 95% = 0.93 – 2.05), p=0.003] implies that the odds of the odds of treatment seeking behaviours among pregnant expectant mothers who reported positive attitude of midwives were 1.45 times higher as compared to those who reported negative attitude of midwives. On the other hand, the adjusted odds ratio of patient care [AOR = 8.22, (CI 95% = 6.45 – 10.96), p=0.030] implies that the odds of treatment seeking behaviours among pregnant expectant mothers who they were given good care by the medical workers at the health units were 8 times higher compared to those who were not given good care by the medical workers at the health units. Other health-based factors such as medical supplies and equipment and distance to health unit had no significant association to treatment seeking behaviours among pregnant expectant mothers.

Personal factors also had a significant association with treatment seeking behaviours among pregnant expectant mothers. Attitude of pregnant mother was the only personal factor that had a significant association with treatment seeking behaviours among pregnant expectant mothers. The adjusted odds ratio malaria [AOR = 5.38, (CI 95% = 4.87 – 6.17), p=0.000] imply that the odds of treatment seeking behaviours among pregnant expectant mothers who had had positive attitude towards the midwives were about 5 times higher compared to those who had negative attitudes. Other personal factors such as income level had no significant association to treatment seeking behaviours among pregnant expectant mothers. The quantitative results were consistent with the qualitative results.

Discussion

The study found that health education on malaria is significantly associated with treatment-seeking behavior for febrile illnesses. This implies that when pregnant mothers are given varied sessions of training to acquire knowledge about malaria, they are more likely to seek treatment immediately after they experience changes in their body temperature. This can contribute to fighting deaths among mothers caused by other febrile illnesses. The findings above are consistent with Tyagi, Roy, and Malhotra (2005) who reported that the failure to establish the level of knowledge of community members regarding malaria appeared to be responsible for the inability of intervention programs to achieve sustainable control.

Findings show that the attitude of midwives is significantly associated with treatment-seeking behavior for febrile illnesses assumed to be malaria among pregnant mothers in the area of study. This implies that when pregnant mothers are handled well, they will always seek treatment for febrile sicknesses assumed to be malaria. These findings are supported by Adeyemo (2013), and Relatedly, et al., (2012) who stressed that nurses should be good to their patients and in case they are rude, shout at their clients, refuse to offer assistance, and sometimes threaten pregnant women during ANC and labour. It is thought that such bad customer handling may be a big factor to scare away mothers even though they fall sick with malaria.

As observed from the findings, patient care was significantly associated with treatment-seeking behavior among respondents. The findings suggested that the health workers in the research area were still aware of the obligations to their clients. Conrad et al. (2012) noted that it is important that expectant mothers are given enough patient care that can propel them to always seek hospital treatment whenever they are seeking it. According to Mgawadere, (2009), if mothers have changed considerably their normal moods, the health workers should be in a position to give comfort and contain their emotions.

Also, the attitude of pregnant mothers was found to be significantly associated with treatment-seeking behavior for pregnant mothers having febrile illnesses assumed to be malaria. This was a very contributing factor in health promotion aspects of the community that health workers should try to be diligent when serving special populations. In support of the findings Mubyazi, et al., (2014) Sholapurkar et al., (2015) inform that expectant mothers are simply negligent about seeking health treatment when they are sick of malaria but they are actually aware of the challenges caused by the delayed treatment of malaria. Sholapurkar et al., (2015) inform that it is not certain whether mood swings alone can deter mothers from seeking malaria treatment but what is known is that mother's attitude towards the treatment of malaria is undesirable for health outcomes.

Limitations

The study was largely cross sectional in nature and data was collected at a single point in time which may not allow changes in behavior to surface, the qualitative longitudinal studies with in-depth interviews could have provided more insights after passage of few months about how the expectant mothers change in treatment-seeking with changes in various situations.

Conclusions

From the study, health education on malaria, the attitude of midwives, patient care, and the attitude of the pregnant mothers were found to be significantly associated with treatment-seeking

behaviors among pregnant women suffering from febrile illnesses assumed to be malaria. This calls for more attention by the Ministry of Health and other responsible stakeholders to reinforce awareness programs on health education among women of reproductive age about the danger of febrile illnesses during pregnancy and help midwives address issues affecting their attitude towards pregnant mothers.

Abbreviations

CI: Confidence Interval; HBM: Health Belief Model; MoH: Ministry of Health; SYAHD: Society for Youth Awareness and Health Development; TPB: Theory of Planned Behavior; WHO: World Health Organisation.

Ethics approval and consent to participate

Before the study was conducted, the protocol was reviewed and approved by the local Research Ethics Committee (REC) and Uganda National Council for Science and Technology (UNCST) as entities that may have access to private information that identifies the research participants by name. Verbal informed consent was obtained from the pregnant others who participated in the study.

Consent for publication

Not applicable.

References

- Adefioye, O.A. Adeyeba, O. A. Hassan W.O. Oyeniran and O.A. (2007) Prevalence of Malaria Parasite Infection among Pregnant Women in Osogbo, Southwest Nigeria. *American-Eurasian Journal of Scientific Research* 2 (1): 43-45.
- Adeyemo (2013) Comparative Analysis of Health Institutions on the Attitude and practice of Midwives towards pregnant women during child delivery in Ogbomoso, Oyo State, Nigeria. Retrieved from <http://www.iosrjournals.org/iosr-jnhs/papers/vol1-issue3/D0131419.pdf> on 18/5/2014
- Amin, M. (2005). *Social Research Conception "Methodology and Analysis*. Kampala: Makerere Printery.
- Aziem A., Osman M., Abbaker O., & Ishag A. (2010). Use of antenatal care services in Kassala, eastern Sudan. Retrieved from <http://www.biomedcentral.com/1471-2393/10/67> on 13th March 2016.

- Banda, C. (2013). Barriers to Utilization of Focused Antenatal Care among Pregnant Women in Ntchisi District in Malawi: Master's Thesis University of Tampere, Tampere School of Health Sciences. Tampere School of Health Sciences/Public Health. University of Tampere.
- Chepkemoi A & Mutulei N (2014) Factors Influencing the Uptake of Intermittent Preventive Treatment for Malaria in Pregnancy: Evidence from Bungoma East District, Kenya. Science and Education Publishing. Retrieved from <http://pubs.sciepub.com/ajphr/1/5/2/index.html> on 3rd March 2016.
- Chimankar, A. and Sahoo, H. (2011). Factors influencing the Utilization of Maternal Health Care Services in Uttarakhand. Department of Sociology, Marathwada University, Aurangabad, Maharashtra, India.
- Chimere A, Wellington A and Anorlu R (2010) Prevalence of Malaria in Pregnant Women in Lagos, South-West Nigeria. *Parasitology* 47(2): 179–183.
- Conrad, P., De Allegri, M., Moses, A., Larsson, E.C., Neuhann, F. Muller, O. & Sarker, M. (2012) Antenatal care services in rural Uganda: missed opportunities for good-quality care. *Qualitative Health Research* 22, 619-629.
- Cronbach, L. (1946). Response Sets and Tests Validity. *Journal for Educational & Psychological Measurement*, 475 - 494.
- Dey, D. (2009). The Reproductive and Child Health (RCH) status in West Bengal: Observations from NFHS surveys in 1992-93 and 1998-99. In: DK Adak, A Chattopadhyay, P Bharati (Eds.): People of Contemporary West Bengal. New Delhi: Mohit Publications.
- Emelumadu, O., Ukegbu, A., Ezeama, N., Kanu, O., Ifeadike, C., Onyeonoro, U. (2014). Socio-demographic determinants of maternal health-care service utilization among rural women in anambra state, South East Nigeria. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/24971212> on 13th January 2016.
- Federal Ministry of Health (2008). Malaria situation analysis document. Nigeria: Federal Ministry of Health. p. 14.
- Hay S.I., Okiro E. & Gething P.W. (2010) Estimating the global clinical burden of Plasmodium falciparum malaria. Retrieved from <http://www.malariajournal.com/content/11/1/274#B1> on 20th December 2015.
- Krejcie, R., & D.W, M. (1970). Determining Sample Size for Research Activities. *Economic Psychological Measurement*, 607 - 610.
- Meghna D & Feiko O (2009) Epidemiology and burden of malaria in pregnancy. The Latent Infectious Diseases. Retrieved from <http://www.thelancet.com/journals/laninf/article/PIIS1473-3099%2807%2970021-X/fulltext?rss=yes> on 27th December 2015.

- Mgawadere F. (2009) Assessing the Quality of Antenatal Care at Lungwena Health Centre in Rural Malawi. Retrieved from www.medcol.mw/mph/dissertations/.../Florence%20Mgawadere.pdf on 22/11/2014
- Ministry of Health (2013), “Implementing Guidelines for the National Home Based Management of Fever Strategy,” Uganda.
- Mityana District (2018). Local Government Performance Assessment Mityan District. Downloaded from: https://budget.finance.go.ug/sites/default/files/mityana_district_assessment_lgpa_full_report-compressed.pdf
- Mubyazi G, Bloch P, Kamugisha M, Kitua A, Ijumba J (2014). Intermittent preventive treatment of malaria during pregnancy: a qualitative study of knowledge, attitudes and practices of district health managers, antenatal care staff and pregnant women in Korogwe District, north-eastern Tanzania. *Malar J.* 2015;4:31.
- Musa OI, Salaudeen GA, Jimoh RO (2009). Awareness and use of insecticide treated nets among women attending ante-natal clinic in a northern state of Nigeria. *J*
- Mvondo JL, James MA, Cambell CC (2012). Malaria and pregnancy in Cameroonian women. Effect of pregnancy on Plasmodium falciparum parasitaemia and the response to chloroquine. *Tropical medicine parasitology*, 43(5)1-5.
- Namusoke, Fasti, N Kironde F (2010) Malaria Burden in Pregnancy at Mulago National Referral Hospital in Kampala, Uganda. *Malaria Research and Treatment Volume 2010 (2010)*, Article ID 913857, 10.
- Okutu (2011). Access to and utilization of antenatal care services in Uganda. Retrieved from <http://uaps2011.princeton.edu/papers/110707> on 27/6/2015
- Onasoga (2012) Perceived effects of midwives attitude towards women in labour in Bayelsa State, Nigeria. Retrieved from <http://scholarsresearchlibrary.com/aasr-vol4-iss2/AASR-2012-4-2-960-964.pdf> on 18/5/2014
- Onen, W., & Oso. (2008). *A General Guideline to Writing Research Proposal and Report*. Kampala: Makerere University Printers.
- Sekaran, U. (2000). *Research Methods for Business*. John Wiley & Sons Inc.
- Sholapurkar S, Gupta A, Mahajan R (2015) Clinical course of malaria in pregnancy a prospective controlled study from India. *Trans R Soc Trop Med Hyg.* 82:376– 379.
- Terefe G. F, Samuel Y. Ayanto & Fiseha L (2015) Gurmamo Assessment of knowledge, attitude and practice about malaria and ITNs utilization among pregnant women in Shashogo District, Southern Ethiopia. *Malaria Journal* 2015 14:235

- Titaley Christiana R, Hunter Cynthia L, Heywood Peter and Michael J Dibley (2010) Why don't some women attend antenatal and postnatal care services? West Java Province, Indonesia. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2964562/> on 15th March 2014.
- Tyagi, P., Roy, A., & Malhotra, M.S. (2005). Knowledge, awareness and practices towards malaria in communities of Rural, semi-rural and bordering areas of east Delhi, India. *Journal of Vector Borne Disease*, 42, 30-35.
- United States Embassy in Nigeria. (2011). *Nigeria malaria fact sheet*. Retrieved June 5, 2014, from <http://photos.state.gov/libraries/nigeria/23177/Public/December.Malaria.FactSheet2.pdf>
- WHO (2008). World Malaria Report 2008. Switzerland: World Health Organization; 2008. pp. 99–101.
- WHO (2012) World malaria report. Geneva, Switzerland: World Health Organization; 2012.
- WHO (2015), Malaria: Fact Sheet, Media Centre. Retrieved from <http://www.who.int/mediacentre/factsheets/fs094/en/> on 5th January 2016.
- WHO (2016) Malaria: Global Health Observatory (GHO) data. Retrieved from <http://www.who.int/gho/malaria/en/> on 4th January 2016.
- WHO and Ministry of Health Uganda (2016), WHO Country Cooperation Strategy, Uganda, 2016–2020. Down loaded from <https://www.afro.who.int/sites/default/files/2017-08/3rd%20WHO%20Country%20Cooperation%20Strategy%20in%20Uganda.pdf>
- Yeka ABanek., K., & Bakyaita N (2009) “Artemisinin versus nonartemisinin combination therapy for uncomplicated malaria: randomized clinical trials from four sites in Uganda,” *PLoS Medicine*, vol. 2, (7),45.