

A promising coriander (*Coriandrum sativum* L.) genotype WFPS 48-1 for high yield and quality for South Eastern Rajasthan

Abstract:

A promising coriander (*Coriandrum sativum* L.) genotype WFPS 48-1 has been identified for high yield and essential oil, on the basis of multilocation testing across the coriander growing states and specifically for Zone V of Rajasthan state. It bears white flowers and recorded average yield of 1933 kg/ha with around 11, 19 and 25 percent superiority over the zonal, state and national checks viz., RKD 18, RCr 436 and Hisar Anand, respectively. Its seeds contain higher essential oil content of >0.55% showing an advantage of 6 and 18 percent, respectively over the national checks RCr 728 and Hisar Anand.

Introduction:

Coriander (*Coriandrum sativum* L.) is an important seed spice crop cultivated for both seed and leaf, used for flavouring and seasoning of food and food products. India is the largest producer, consumer and exporter of coriander in the world. The total production of coriander in the country was 811 thousand tons from an area of 640 thousand hectare (2021-22). The major coriander producing states are Rajasthan, Gujarat and Madhya Pradesh. It is cultivated in an area of 124286 ha in Rajasthan and produces 182705 m tonnes (2020-21) of coriander seeds. Rajasthan contributes nearly 19 and 20 percent to the country's total area and production, respectively. The south eastern part of Rajasthan i.e., humid south eastern plain zone or zone V covering districts of Kota, Bundi, Baran and Jhalawaris the largest producer of coriander in the state where it is cultivated in an area of 118030 ha contributing the production of 173764 m. tonnes (2020-21), thereby contributing around 95 percent to the total area and production of the state.

Development of high yielding varieties is the first and foremost objective of any crop improvement programme to meet the crop's domestic as well international demand. Since coriander is valued for its aroma owing to the presence of essential oil in its seeds and leaves, therefore, besides high yield, high quality also assumes priority. Yield is a complex character dependent on various yield contributing factors. Similarly, essential oil content also varies in varying environmental conditions. In general, seed size has negative correlation while maturity duration has positive correlation with quality suggesting that small sized and late maturing types tend to possess higher content of essential oil and therefore, fetch higher price in the market but late maturing genotypes are generally not preferred for Zone V where there's a sudden rise in temperature and heat stress during the terminal stage of the crop. The essential oil being volatile in nature diminishes fast due to high temperature. Keeping the above facts in view, an attempt was made in the present study to identify a coriander genotype for high yield and quality suitable for the agroclimatic conditions of Zone V of the state.

Material and methods:

The experiment was conducted at multilocations of zone V viz., ARS, Kota, ATC, Bundi, Nanta, Digod, ARSS, Aklera, Khanpur (Jhalawar) during *rabi* 2018-19 to evaluate

the coriander genotype against the existing checks for yield, yield contributing and quality traits to identify its suitability for the agroclimatic conditions of Zone V of the state. The trials were conducted following all the recommended package of practices for zone V. The field data were recorded at maturity and statistically analysed while the essential oil content of the genotypes from all the locations was estimated from the seed by Clevenger method in Central Laboratory, ARS, Kota.

Results and discussion:

The data recorded on yield in multilocation trials under zone V is provided in table 1. On the basis of yield data recorded at six locations, the coriander genotype WFPS 48-1 recorded a mean yield of 1933 kg/ha with around 11, 19 and 25 percent superiority over the zonal, state and national checks *viz.*, RKD 18, RCr 436 and Hisar Anand, respectively (table 1). This genotype was earlier evaluated under coordinated trials for three consecutive years during 2015-16 to 2017-18. The data on other yield parameters *viz.*, plant height, days to maturity and test weight is provided in table 2. The genotype WFPS 48-1 is early in maturity as compared to checks RCr 436 and Hisar Anand while it is at par to the popular zonal check RKD 18. The plants are medium tall as compared to taller varieties with height of nearly 100 cm, making them prone to lodging. The seeds are medium bold with an average test weight of 14.9g which is also desirable as more bold seeds are said to be poor in essential oil. The seeds contain essential content of more than 0.44-0.55% showing an advantage of 5, 6, 18 and 50 percent, over the zonal check RKD 18, national checks RCr 728 and Hisar Anand and state check RCr 436, respectively (table 3).

The popular variety of coriander presently under cultivation in zone V is RKD 18 (Pratap Raj Dhania 1) notified in the year 2015. The superiority and suitability of this variety for the zone has been documented by various researchers in the past (Meena *et al.*, 2013, Doshi *et al.*, 2014, Giridhar *et al.*, 2014 and Verma *et al.* 2015). But the varieties more than ten years old need to be replaced by new varieties in the seed chain. The seed replacement rate in coriander is low, therefore, efforts are needed for development and popularization of new genotypes/varieties in the zone.

Environmental and genetic parameters can affect plants performance (Golam *et al.*, 2014, Medipally *et al.*, 2014, Shams *et al.*, 2014). The production of secondary metabolites in plants depends on different factors such as temperature, light, plant genotype, altitude, slope, and hormones (Veeresham and Kokate, 2004, Mahzoon *et al.*, 20014, Omidbeygi, 2000). Shams *et al.* (2016) reported that the quantity and quality of active ingredients is influenced by genetic and environmental factors and there is a direct relation between production of plant metabolites and climate factors. Among the climate parameters, altitude and precipitation have the most impact on the essential oil content and dry matter yield of coriander. Low altitudes could increase the essential oil content of coriander. Keeping in view the effect of environmental factors on quality of coriander, the essential oil content in the seed of the genotype WFPS 48-1 along with checks grown at multilocations was estimated. Among all the genotypes, the highest essential oil content was observed in the

genotype WFPS 48-1, irrespective of the locations, however, the quantity varied as per the locations indicating the effect of environmental factors on essential oil.

The distinct morphological traits help in the easy identification of varieties in seed production and certification programmes. The genotype WFPS 48-1 is quite distinct in view of its white flower colour as compared to pink flowers of the existing varieties. The superiority of the white flower coriander genotypes was reported earlier also by Verma et al. (2014). It is concluded that the genotype WFPS 48-1 showing superiority in terms of seed yield and quality could be a suitable replacement of the existing varieties in the zone to increase coriander production and productivity.

References

1. Doshi Vibha, Verma Preeti and Khan Ishan Ullah (2014). Comparative analysis of coriander (*Coriandrum sativum* L.) varieties for quality traits. *The Journal of Rural and Agricultural Research* 14 (2): 57 – 58.
2. Giridhar K., Surya Kumari S., Rajani A., Sarada C. and Naram Naidu L. (2014). Identification of potential genotypes of coriander (*Coriandrum sativum* L.) suitable for rainfed vertisols. *Applied Biological Research* 16 (2): 00-00.
3. Golam F, Zaidi K, Nezhadahmadi A, Osman M. (2014). Genetic analysis of F1 hybrids derived from aromatic (Exotic) × aromatic (Malaysian) rice crosses and their callus induction performance for haploid production. *Indian Journal of Science and Technology*. 7(11):1852–60.
4. Mahzooni-Kachapi SS, Mahdavi M, Jouri MH, Akbarzadeh M, Roozbeh-Nasira'ei L. The effect of altitude on chemical composition and function of essential oils in *Stachys lavandulifolia* Vahl. (Iran). *Int J Med Arom Plants*. 2014; 4(2):107–16.
5. Medipally SR, Naresh B, Manoj Kumar S, Yusoff FM, Devi P. Somatic Embryogenesis from Leaf and Shoot tip Explants of *Jatropha curcas* L. *Indian Journal of Science and Technology*. 2014; 7(11):1842–6.
6. Meena, B.S., Dhaka, B.L. and Poonia M.K. (2013). Assessment of yield, quality and economics of coriander (*Coriandrum sativum* L.) genotypes in south-eastern plains of Rajasthan under irrigated condition. *International J. Seed Spices* 3(1): 58-60
7. Omidbeygi R. Approaches of production and output of medicinal plants. Fekre now publications. 2000.
8. Preeti Verma, Vibha Doshi, S.S. Punia, Minakshi Dheer and Mashiat Ali (2014). White flower coriander (*Coriandrum sativum* L.): Yet to be explored. *Annals of Biology* 30(1): 162-165.
9. Shams, Mostafakamal, Ramezani, Mahdi, Esfahan, Sasan Zandi, Esfahan, Ehsan Zandi, Dursun Atilla and Yildirim Ertan. (2016). Effects of Climatic Factors on the Quantity of Essential Oil and Dry Matter Yield of Coriander (*Coriandrum sativum* L.). *Indian Journal of Science and Technology*, Vol 9(6).
10. Shams M, Zandi Esfahan E, Ramezani M, Ghandkanlou M. Allelopathic effects of weeds extract on seed germination of *Trachyspermum copticum* L., *Foeniculum vulgare* and *Cuminum cyminum*. *J. Bio. and Env. Sci*. 2014; 5 (1): 82-90.
11. Veeresham C, kokate CK. Medicinal plant biotechnology: CBS Publishers and Distributors 2004.
12. Verma Preeti, Ali Mashiat, Doshi Vibha and Solanki R.K. (2015). Stability analysis in coriander (*Coriandrum sativum* L.). *Indian J. Hort.* 71(1): 126-129.

UNDER PEER REVIEW

Table 1: Yield performance (kg/ha) of WFPS 48-1 at multilocations of zone V during 2018-19

Entries/Checks	ARS, Kota	ATC, Bundi	Nanta	Digod	ARSS, Khanpur	ARSS, Aklera	Mean	%increase over check
WFPS 48-1	1815	3810	1546	1499	1265	1660	1933	
RCr 436 (C)	1539	3274	1290	1254	1117	1262	1623	19.09
RKD-18 (C)	1591	3333	1472	1405	1110	1516	1738	11.20
HisarAnand (C)	1244	2902	1412	1348	1083	1280	1545	25.09
CD (5%)	172.95	595.37	217.69	263.12	270.18	200.08		

Table 2: Ancillary data of WFPS 48-1 under multilocation testing in zone V during 2018-19

Entries/Checks	Plant height (cm)			Mean	Days to maturity				Mean	Test weight (g)				Mean
	ARS, Kota	ATC, Bundi	ARSS, Aklera		ARS, Kota	ATC, Bundi	ARSS, Aklera	ARSS, Khanpur		ARS, Kota	ATC, Bundi	ARSS, Aklera	ARSS, Khanpur	
WFPS 48-1	99.82	93.26	58.00	83.69	108	116	112	118	113	14.03	16.3	16.1	13.20	14.90
RCr 436 (C)	85.00	86.08	61.75	77.61	108	126	116	118	117	12.6	15.2	15.5	14.00	14.32
RKD-18 (C)	82.00	88.04	62.75	77.60	105	116	113	119	113	11.62	15.7	14.6	13.20	13.78
HisarAnand (C)	106.35	127.02	69.25	100.87	118	130	118	117	121	11.50	17.7	14.5	13.60	14.32

Table 3: Essential oil (%) of WFPS 48-1 in multilocations of zone V during 2018-19

Entries/Checks	Testing in coordinated trials					Testing in Zone V					
	2015-16	2016-17	2017-18	Mean	%increase over checks	ATC, Bundi	ARSS, Khanpur	ARSS, Aklera	ARS, Kota	Mean	%increase over checks
WFPS 48-1	0.47	0.8	0.4	0.556		0.51	0.52	0.40	0.36	0.4475	
RCr 436 (C)	-	-	-	-	-	0.31	0.32	0.28	0.28	0.2975	50.42
RKD-18 (C)	-	-	-	-	-	0.47	0.45	0.40	0.38	0.425	5.29
HisarAnand (C)	0.38	0.6	0.43	0.47	18.43	0.30	0.30	0.28	0.28	0.29	54.31
RCr 728 (C)	0.37	0.6	0.6	0.52	6.36	-	-	-	-	-	-

UNDER PEER REVIEW