

Review article

RISK FACTORS OF ACUTE MALNUTRITION AMONG CHILDREN UNDER FIVE YEARS OF AGE

Abstract

Childhood malnutrition remains a significant global public health concern, and despite numerous advances and improvements in child health care; acute malnutrition among children still one of the main public health challenges in the 21st century, particularly in developing countries and is associated with a high risk of morbidity and mortality. Children are the most vulnerable group to the effects of acute malnutrition during the period of their most rapid physical growth and development, because of the extra nutritional requirements for growth and development at this time. Acute malnutrition results from the complex interplay of a range of distal and proximal factors. This review article gives a comprehensive background on the causes of acute malnutrition among children under five years of age.

KEYWORDS: Malnutrition, acute malnutrition, causes of acute malnutrition, children

Introduction

Childhood malnutrition remains a significant global public health concern (1), and “despite numerous advances and improvements in child health care; acute malnutrition among children remain one of the main public health challenges in developing countries” (2), “and is associated with a high risk of morbidity and mortality (3). “Children are the most vulnerable group to the effects of acute malnutrition during the period of their most rapid physical growth and development, because of the extra nutritional requirements for growth and development at this time” (4). “Malnutrition defined as cellular imbalance between the supply of nutrients and energy and body’s demand to ensure growth, maintenance, and specific functions” (5), “causes measurable adverse effects on body form, function and clinical outcome” (6).

“The term of acute malnutrition has been defined in various ways and names with partially overlapping definitions; including protein-energy malnutrition, wasting, marasmus and kwashiorkor, with different clinical and pathological characteristics” (7, 8). “Acute malnutrition occurs when the quantity of one or more nutrients available to body tissues is inadequate to maintain optimal bodily functions” (9). “It develops as a result of recent rapid weight loss or a failure to gain weight” (10). “It can manifest over a short period of time when the body does not receive adequate amounts of micronutrients or energy, either as a result of insufficient dietary intake or through malabsorption of nutrients or loss of appetite due to illness” (11). “It increases dramatically in emergencies and in developing countries generally, where these settings are plagued by chronic poverty, poor hygiene, lack of education, poor diets and limited access to food” (12).

Components of nutrition:

Good nutrition is essential for the growth, health, development, and economic well-being of individuals and populations (3). It is crucial for optimal child development

throughout the first years of life and beyond (13). It is divided to two main groups which include the following:

Macronutrients: Macronutrients refer to carbohydrates, proteins and fats, which make up the bulk of a diet and supply the body with energy. Carbohydrates such as starches and sugars are often a large part of the diet (80%) and the main source of energy. Fats, also an essential component in the diet, make up about 10% of the diet and supply energy and are important in cell formation. Proteins are required to build new tissue, it is derived from animal origin such as milk, meat and eggs, which products contain essential amino acids (14).

Micronutrients: Micronutrients include about forty types from different minerals and vitamins that are essential for good health. Most micronutrients divide to two types; type I, which includes iodine, iron and Vitamins (Vit A and Vit C), its deficiency do not affect growth, but leading to compromised body functions such as impaired vision, reduced intelligence, anemia, scurvy and impaired immunity. It is not determined by anthropometric measurement (14, 15). Type II micronutrients are essential for growth and tissue repair, it includes phosphorus, zinc, potassium, magnesium, nitrogen, essential amino acids, sulphur, sodium and chloride. "It is required in small quantities, but the correct balance is essential for good health. A deficiency in any of the type II micronutrients will lead to growth failure in children" (14).

Types of malnutrition:

Malnutrition is a condition that results from deficiencies, excesses, or imbalances in a person's intake of nutrients or energy. According to classification of malnutrition by classified as three broad groups of conditions such as the following (16):

Undernutrition: Undernutrition is a consequence of absorption, inadequate nutrient intake and illness or disease (3, 17). It is including:

- Acute malnutrition refers to low weight for height (wasting)
- Chronic malnutrition refers to low height for age (stunting)
- Underweight refers to low weight for age

Micronutrient deficiencies: Micronutrient deficiencies refer to lack of important vitamins and minerals such as vitamin A, iodine, iron, and zinc. It is assessed using biochemical and clinical methods, not by anthropometric measurements (3).

Overweight (obesity): "Overweight is described as an imbalance between the calories ingested and the calories expended. Childhood obesity is associated with health problems in adulthood such as insulin resistance, type 2 diabetes, hyperlipidemia, hypertension, and psychological problems like depression and eating disorders" (18).

Acute malnutrition:

Acute malnutrition results from acute food shortages or an insufficient intake of protein, energy or other nutrients (19, 20). "According to UNICEF definition, acute malnutrition is characterized by a rapid deterioration in nutritional status over a short period of time. Among children, it can be identified using the weight for height nutritional index or mid upper arm circumference (MUAC), or presence of bilateral pitting oedema" (17).

Classification and definitions of acute malnutrition:

In children less than five years of age, acute malnutrition can be classified as either moderate or severe based on level of wasting for individual, and according to specific cut offs and reference standards. At the population level, acute malnutrition is categorized in three levels as the following (10, 17, 21):

Moderate acute malnutrition (MAM): MAM in children 6–59 months is defined as a weight for height Z-score (WHZ) < -2 and ≥ -3 standard deviations of the WHO standards, and/or MUAC from 115 mm to less than 125 mm (10, 17).

Severe acute malnutrition (SAM): SAM is the most extreme and visible form of acute malnutrition that requires urgent treatment to survive (UNICEF, 2015a). “It results from sudden reductions in diet quality or food intake and is often combined with pathological causes, and is associated with loss of body fat and wasting of skeletal muscle, develops as a result of recent rapid weight loss or a failure to gain weight” (4, 7, 21). “SAM in children 6–59 months of age defined as a weight for height/length < -3 Z-score of the WHO growth standard, and/or MUAC of less than 115 mm, or the presence of bilateral pitting oedema” (10, 17, 22).

Global acute malnutrition (GAM): “GAM refers to MAM and SAM together; it used as a measurement of nutritional status on the level of population and as an indicator of the severity of an emergency situation” (7, 10, 23).

Risk factors of acute malnutrition:

“Acute malnutrition results from the complex interplay of a range of distal and proximal factors, as described by UNICEF conceptual framework for malnutrition. The framework defines immediate, basic and underlying causes of malnutrition and demonstrates how these causes are interconnected” (1, 24). “Based on scientific literature investigating the relationships among specific individual, household, and environmental factors and the development of acute malnutrition in children, the following are significant risk factors for SAM” (1, 13, 24-27):

Immediate causes of acute malnutrition:

The common immediate causes of acute malnutrition include the following:

1- Inadequate dietary intake: Inadequate dietary intake is one of the most immediate causes of child malnutrition (28). Inadequate feeding lacking in quality and quantity can restrict growth and jeopardize child survival and development (29). Case control study conducted in Chad by Dodos, Altare (30) showed that the type of complementary meal introduced for children was significantly associated with acute malnutrition.

2- Disease (health status): “Acute malnutrition results of complications of repeated infections, which may further worsen the child’s nutritional status at a time of greater nutritional needs. This interaction between acute malnutrition and infection creates a potentially lethal cycle of worsening illness and deteriorating nutritional status” (24). Acute malnutrition is associated with increased severity of infectious diseases (31). Childhood illness is a contributing causes with acute malnutrition, and it is a major immediate cause of child malnutrition (7, 28). Children in the first years of life are more susceptible to recurrent infectious such as diarrhea and respiratory infection, which adversely affect metabolism, appetite and nutritional status. Being exposed recurrent infections; children can easily enter a vicious circle of weight loss and ever worsening nutritional status. Diarrhea, in turn, not only leads to further loss of micronutrients such as potassium, zinc and other nutrients, but also

deteriorates the absorption capability of the intestine, an addition to that the fever increases energy expenditure; thus putting the child at an increased risk of acute malnutrition. (2, 4, 9, 22, 32). “The relationship between acute malnutrition and diarrhea is bidirectional, acute malnutrition predisposes children to a greater incidence and duration of diarrhea, while diarrhea can, in turn, precipitate or worsen acute malnutrition. Also significant water losses from diarrhea can lead to dehydration and electrolyte imbalance” (22).

Underlying causes of acute malnutrition:

There are many major underlying causes of acute malnutrition which include the following (1, 24):

1- Inappropriate feeding practices: Inappropriate infant and young child feeding (IYCF) practices increase vulnerability to undernutrition (25). Inappropriate IYCF practices such as inappropriate breastfeeding and complementary feeding is one of the essential reasons associated with acute malnutrition (33). Inappropriate IYCF is characterized with poorly timed introduction of complementary foods as too early or late introduced, infrequent feeding and poor feeding methods, poor hygiene and care practices (34). “IYCF practices, such as early onset of weaning, inadequate nutritional content of weaning foods, inadequate feed rations, insufficient breastfeeding and poor hygiene behaviors, have been identified as the leading causes of acute malnutrition, growth faltering, diarrhea, increased rate of infections, vitamin-mineral deficiency, poor cognitive development and increased mortality among children” (35). “Combination of nutritionally inferior diets and improper feeding practices and suboptimal breastfeeding are major contributing factors for the development of childhood malnutrition and put children at high risk for acute malnutrition” (29, 34).

2- Household food insecurity: Household food insecurity or limited access or availability of food is one of the major causes of acute malnutrition (9, 25). It is an important factor contributing to poor nutritional status during conflict and associated with the risk of acute malnutrition in children (36). Young and Marshak mentioned that the acute malnutrition is nevertheless still strongly associated with situations of food insecurity (37). “Food shortages may be acute as sudden/sharp or chronic as long lasting, arise as a result of poverty, natural disaster or conflicts, which may lead to the displacement of people from their homes and disruption of food supplies which results in accelerate and increase of acute malnutrition among children” (4, 6).

3- Unhealthy household environment and limited access to health services: “Limited access to adequate health services or inadequate environmental health conditions such as safe water and sanitation facilities are associated with develop of acute malnutrition” (37). “Poor water, sanitation and hygiene have serious consequences on health and nutritional status among the most vulnerable population groups” (27). A case control study done in Chad by Dodos, Altare showed that absence of toilet in the household and caretakers’ hand washing habits were significantly associated with both acute malnutrition and diarrhea (30).

4- Other causes: Incomplete vaccinations one of the underlying cause of malnutrition, a study conducted in Burkina Faso founds incomplete vaccinations was a risk factor for acute malnutrition relapse (7). Birth order of children and number of family members are significantly associated with SAM (38). Lenters, Wazny (7), refer to the large size of the family is associated with an increased risk of acute malnutrition. Purohit, Sahu demonstrated significant association between birth weight and acute malnutrition (38). Also, study

conducted in Pakistan by Lenters, Wazny (7), found that low birth weight was risk factor for acute malnutrition in CU5.

5- Poverty and Famine: Famine is fundamentally linked to alleviating hunger and malnutrition (37). Poverty is the main underlying cause of acute malnutrition and its determinants (9, 39). “Economically disadvantaged families are less likely to have access to improved sources of food, drinking water and less likely to have access to latrines” (7).

Basic causes of acute malnutrition:

Young and Marshak indicated that the several basic causes of acute malnutrition include the root causes that impact on all underlying causes such as describe in UNICEF conceptual framework for malnutrition (37), it is include the following:

1- Sociocultural, economic, political, and environmental instability: “The basic causes of acute malnutrition in a community originate at the regional and national level, where strategies and policies that affect the allocation of resources such as human, economic, political and cultural. In addition, geographical isolation and lack of access to markets or services due to poor infrastructure, displacement or conflict can have a huge negative impact on health and food security that lead to child malnutrition” (1, 28).

2- Lack of education: Lack of mother's knowledge has an important role lead to child malnutrition (40). Children may not have receive to the variety of foods that will provide all the necessary vitamins and minerals in their diet, this often happens as a result of a lack of mother's knowledge about appropriate feeding practice (4). Lenters, Wazny indicated to a several studies conducted in Bangladesh, India, and Pakistan demonstrated a correlation between low parental education and increased risk of acute malnutrition in children (7). Also, study conducted in India by Purohit, Sahu concluded that maternal education was significantly associated with acute malnutrition among children (38).

Conclusion:

Acute malnutrition among children remains one of the main public health challenges, particularly in developing countries and is associated with a high risk of morbidity and mortality. There are many factors associated with the occurrence of malnutrition among children such as basic, underlying and immediate causes.




Fig 1 :Conceptual framework of the risk factors of child malnutrition (1, 24)

References

1. Groot Rd, Handa S, Ragno LP, Spadafora T. Child malnutrition, consumption growth, maternal care and price shocks: new evidence from Northern Ghana. *Development Studies Research*. 2020;7(1):18-30.
2. Bhutta ZA, Salam RA. Global nutrition epidemiology and trends. *Annals of Nutrition and Metabolism*. 2012;61(Suppl. 1):19-27.
3. Cashin K, Oot L. Guide to anthropometry: a practical tool for program planners, managers, and implementers. *Food Nutr Tech Assist III Proj (FANTA)/FHI*. 2018;360:93-115.
4. Picot J, Hartwell D, Harris P, Mendes D, Clegg A, Takeda A. The effectiveness of interventions to treat severe acute malnutrition in young children: a systematic review. *Health Technology Assessment* 2012;16(19):1–316.
5. Kasio Iboyi A, Zha L. Severe acute malnutrition (SAM) evaluation of associated risk factors, screening tools, and therapeutic management among South Sudanese children age 6< 59 months in emergency settings: case reports and review of literatures. *J Trop Dis*. 2019;7(298):2.
6. Younis K, Ahmad S, Badpa A. Malnutrition: causes and strategies. *J Food Process Technol*. 2015;6(434):2.
7. Lenters L, Wazny K, Bhutta ZA. Management of severe and moderate acute malnutrition in children.

<https://books.google.com.eg/books?hl=ar&lr=&id=Nh0LDAAAQBAJ&oi=fnd&pg=PA2&dq=Management+of+severe+and+moderate+acute+malnutrition+in+children&ots> 2016.

8. Trehan I, Manary MJ. Management of severe acute malnutrition in low-income and middle-income countries. *Archives of disease in childhood*. 2015;100(3):283-7.
9. Schoonees A ,Lombard M, Musekiwa A, Nel E, Volmink J. Ready-to-use therapeutic food for home-based treatment of severe acute malnutrition in children from six months to five years of age. *Cochrane Database of Systematic Reviews*. 2013(6.(
10. WFP. Nutrition at the World Food Programme. <https://documents.wfp.org/stellent/groups/public/documents/communications/wfp258650.pdf> 2012.
11. James PT, Van den Briel N, Rozet A, Israël AD, Fenn B, Navarro-Colorado C. Low-dose RUTF protocol and improved service delivery lead to good programme outcomes in the treatment of uncomplicated SAM: a programme report from Myanmar. *Maternal & child nutrition*. 2015;11(4):859-69.
12. UNICEF. Severe acute malnutrition. https://www.unicef.org/nutrition/index_sam.html .2015
13. Black, Victora CG, Walker SP, Bhutta ZA, Christian P, De Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The lancet*. 2013;382(9890):427-51.
14. WHO, UNICEF. National guideline for integrated management of acute malnutrition 2009.
15. Okello FA. Assessment of the performance of the community based outpatient therapeutic program in abyei administrative area, south sudan. https://d1wqtxts1xzle7.cloudfront.net/50946137/Frank_Thesis_Final_2016-with-cover-page-v2.pdf: Uganda Christian University; 2016.
16. WHO. Malnutrition. [Fact sheets]. <https://www.who.int/news-room/fact-sheets/detail/malnutrition2020>.
17. UNICEF. Management of severe acute malnutrition in children: working towards results at scale. UNICEF programme guidance document. New York (NY): UNICEF. https://reliefweb.int/attachments/45f163e5-526f-3a09-ac8a-dc9f91c10bd3/unicef_2015_management_of_severe_acute_malnutrition.pdf 2015.
18. Ayala-Marín AM, Iguacel I, De Miguel-Etayo P, Moreno LA. Consideration of social disadvantages for understanding and preventing obesity in children. *Frontiers in Public Health*. 2020;8.
19. Fanzo J, Hawkes C, Udomkesmalee E, Afshin A, Allemandi L, Assery O, et al. 2018 global nutrition report. <https://globalnutritionreport.org/reports/global-nutrition-report-2018/> 2019.
20. Shanka N, Lemma S, Abyu D. Recovery rate and determinants in treatment of children with severe acute malnutrition using outpatient therapeutic feeding program in Kamba District, South West Ethiopia. *J Nutr Disord Ther*. 2015;5(2):155.

21. WFP, UNICEF, SCI. Guidelines for community management of acute malnutrition. <https://reliefweb.int/report/south-sudan/community-management-acute-malnutrition-cmam-guidelines2017>.
22. WHO. Guideline: updates on the management of severe acute malnutrition in infants and children. <https://www.who.int/publications/i/item/9789241506328>: World Health Organization; 2013.
23. Force MT. Moderate acute malnutrition: a decision tool for emergencies. Geneva: IASC Global Nutrition Cluster. 2012.
24. UNICEF. Improving child nutrition—the achievable imperative for global progress. <https://data.unicef.org/resources/improving-child-nutrition-the-achievable-imperative-for-global-progress/> 2013.
25. Sphere. Sphere Handbook: Humanitarian charter and minimum standards in humanitarian response. fourth edition, Geneva, Switzerland, 2018. <https://spherestandards.org/wp-content/uploads/Sphere-Handbook-2018-EN.pdf>: PRACTICAL ACTION; 2018.
26. ENN, GNC, UNICEF. Scaling up nutrition services and maintaining service during conflict in Yemen: Lessons from the Hodeidah sub-national Nutrition Cluster. <https://www.ennonline.net/fex/52/nutritionservicesduringconflict> 2016. 57.[
27. WFP, FAO, UNICEF, FSC, GOVT Y. Yemen - food security and nutrition assessment (EFSNA). <https://reliefweb.int/report/yemen/yemen-food-security-and-nutrition-assessment-efsna-june-20172017>: [124 p.].
28. JMoH. Jordan guidelines for management of acute malnutrition. The Jordanian Ministry of Health in 2013. 2013:74.
29. UNICEF. Infant and young child feeding: programming guide. <https://www.ennonline.net/attachments/1470/unicef-iyfc-programming-guide-may-26-2011.pdf> 2011.
30. Dodos J, Altare C, Bechir M, Myatt M, Pedro B, Bellet F, et al. Individual and household risk factors of severe acute malnutrition among under-five children in Mao, Chad: a matched case-control study. *Archives of Public Health*. 2018;76(1):35.
31. Jones KD, Berkley JA. Severe acute malnutrition and infection. *Paediatrics and international child health*. 2014;34(sup1):S1-S29.
32. Rodríguez L, Cervantes E, Ortiz R. Malnutrition and gastrointestinal and respiratory infections in children: a public health problem. *Int J Environ Res Public Health*. 2011;8(4):1174-205.
33. Mahmood H, Suleman Y, Hazir T, Akram DS, Uddin S, Dibley MJ, et al. Overview of the infant and young child feeding policy environment in Pakistan: Federal, Sindh and Punjab context. *BMC Public Health*. 2017;17(2):474.
34. Abeshu MA, Adish A, Haki GD, Lelisa A, Geleta B. Assessment of caregiver's knowledge, complementary feeding practices, and adequacy of nutrient intake from homemade foods for children of 6–23 months in food insecure Woredas of Wolayita zone, Ethiopia. *Frontiers in nutrition*. 2016;3:32.

35. Arikpo D, Edet ES, Chibuzor MT, Odey F, Caldwell DM. Educational interventions for improving primary caregiver complementary feeding practices for children aged 24 months and under: ohn Wiley & Sons, Ltd.; 2018.
36. Carroll GJ, Lama SD, Martinez-Brockman JL, Pérez-Escamilla R. Evaluation of nutrition interventions in children in conflict zones: a narrative review. *Advances in Nutrition*. 2017;8(5):770-9.
37. Young H, Marshak A. Persistent global acute malnutrition. https://www.ecbproject.org/system/files/content/resource/files/main/FIC-Publication-Persistent-Global-Acute-Malnutrition_web_2.26s.pdf: Boston: Feinstein International Center, Tufts University,; 2018.
38. Purohit L, Sahu P, Godale LB. Nutritional status of under-five children in a city of Maharashtra: a community based study. *Int J Community Med Public Health*. 2017;4(4):1171-8.
39. Müller O, Krawinkel M. Malnutrition and health in developing countries. *Cmaj*. 2005;173(3):279-86.
40. Sangra S, Nowreen N. Knowledge, attitude, and practice of mothers regarding nutrition of under-five children: a cross-sectional study in rural settings. *Int J Med Sci Public Health*. 2019;8(5):392-4.